A Redundant Function for the N-Terminal Tail of Ndc80 in Kinetochores–Microtubule Interaction in *Saccharomyces cerevisiae*

Pinar B. Demirel,* Brice E. Keyes,† Mandovi Chatterjee,* Courtney E. Remington,* and Daniel J. Burke*,¹

*Department of Biochemistry and Molecular Genetics, University of Virginia School of Medicine, University of Virginia, Charlottesville, Virginia 22906, †Department of Cell Biology, University of Virginia School of Medicine, University of Virginia, Charlottesville, Virginia 22906, and ¹Laboratory of Mammalian Cell Biology and Development, The Rockefeller University, New York, New York 10021-6399

ABSTRACT The N-terminal tail of Ndc80 is essential for kinetochore–microtubule binding in human cells but is not required for viability in yeast. We show that the yeast Ndc80 tail is required for timely mitotic progression and accurate chromosome segregation. The tail is essential when cells are limited for Dam1, demonstrating a redundant function for the Ndc80 and Dam1 complexes *in vivo*.

THE evolutionarily conserved Ndc80 complex is composed of Ndc80, Nuf2, Spc24, and Spc25, integral kinetochore (KT) proteins responsible for mediating microtubule (MT) attachments from yeast to humans (Wigge and Kilmartin 2001; McClelland *et al.* 2003; Ciferri *et al.* 2008). Recombinant Ndc80 complex binds MTs *in vitro*, and the activity is restricted to the head domains of Nuf2 and Ndc80 (Wei *et al.* 2005; Ciferri *et al.* 2008). Yeast has an additional protein complex, Dam1, composed of 10 proteins, that is also required for KT–MT attachment (Westermann *et al.* 2006; Lampert *et al.* 2010; Tien *et al.* 2010). The Dam1 complex binds MTs *in vivo* and *in vitro* and localizes to the KT in a MT-dependent fashion (Li *et al.* 2002). The Dam1 complex, at high concentrations *in vitro*, oligomerizes into a closed-ring structure that encircles MTs (Miranda *et al.* 2005; Lampert *et al.* 2010; Tien *et al.* 2010). There is an interaction between Dam1 and Ndc80 complexes on MTs *in vitro* that enhances the ability of Ndc80 to maintain MT attachments (Lampert *et al.* 2010; Tien *et al.* 2010). The molecular details of the interaction are unclear.

The Ndc80 protein has an extended N-terminal tail that varies among different species in both length and sequence

but is highly positively charged in all species. The N-terminal tail is unstructured and not present in the crystal structure of Ndc80 complexes or in cryo-electron microscopy images of the Ndc80 complex bound to MTs. The N-terminal tail of Ndc80 is enriched for Ipl1/AuroraB phosphorylation sites, and MT binding can be regulated by the kinase (Guimaraes *et al.* 2008; Welburn *et al.* 2010; Lampson and Cheeseman 2011; Tooley and Stukenberg 2011). MT-binding affinity of yeast and human Ndc80 complexes is decreased when the Ndc80 N-terminal tail is deleted (Wei *et al.* 2005; Ciferri *et al.* 2008). Reducing the positive charge by mutating some residues in the Ndc80 N-terminal tail abolishes MT attachments (Guimaraes *et al.* 2008; Miller *et al.* 2008; Tooley *et al.* 2011). KTs are structurally intact in engineered Hec1ΔN HeLa cells; however, the cells are unable to stably bind MTs, generate tension between sister KTs, or congress chromosomes (Miller *et al.* 2008). Surprisingly, the N-terminal tail of the Ndc80 protein is dispensable for viability in *Saccharomyces cerevisiae* although a detailed phenotypic analysis has not been reported (Kemmler *et al.* 2009; Lampert *et al.* 2010).

We confirmed that the N-terminal tail of Ndc80 was non-essential by deleting amino acids 2–112 in NDC80 (hereafter referred to as *ndc80-112*; see Supporting Information, Table S1) in a plasmid, confirmed it by DNA sequencing, replaced one allele in a diploid where the deletion mutation was physically linked to URA3, and confirmed that the deletion allele was present in the diploid by PCR. We sporulated the diploid, dissected tetrads and URA3 segregated 2:2 in all

Copyright © 2012 by the Genetics Society of America

doi: 10.1534/genetics.112.143818

Manuscript received April 16, 2012; accepted for publication July 18, 2012

Supporting information is available online at <http://www.genetics.org/lookup/suppl/doi:10.1534/genetics.112.143818/-/DC1>.

¹Corresponding author: Department of Biochemistry and Molecular Genetics, University of Virginia School of Medicine, 1300 Jefferson Park Ave., Charlottesville, VA 22908-0733. E-mail: dburke@virginia.edu

Figure 1 A mitotic delay in *ndc80-112* cells. Isogenic MATa cells were synchronized with α -factor and released to the cell cycle in the absence (A) or presence (C) of 10 μ g/ml of benomyl and 25 μ g/ml carbendazim (BEN). Cells were sampled every 15 min and prepared for flow cytometry and Western blots (Yellman and Burke 2006). The untreated cells from A were analyzed by fluorescence microscopy for nuclear division (B). Serial 10-fold dilutions of saturated cultures of the indicated genotypes were spotted onto YPD plates containing 15 μ g/ml benomyl and grown at 23° for 4 days (D).

tetrads, confirming that the N-terminal tail of Ndc80 is not required for viability in yeast and that colony growth was unaffected at standard temperatures for growth (Figure S1). We measured cell cycle kinetics in synchronized wild-type and *ndc80-112* cells to determine if deleting the N-terminal tail induced a delay in mitotic timing. We synchronized cells with α -factor, released them into the cell cycle, and followed cell cycle progression by flow cytometry and Pds1 stability. The *ndc80-112* cells cycled with nearly wild-type kinetics, and we could detect a subtle increase in the number of cells with a 2C content of DNA and a subtle delay in the proteolysis of Pds1 and a slight delay in nuclear division (Figure 1, A and B). It was possible that the *ndc80-112* cells did not delay in mitosis because the N-terminal tail of Ndc80 is required for the spindle assembly checkpoint (SAC). However, *ndc80-112* cells grew as well as wild type in the presence of a sublethal concentration of benomyl unlike *mad2* cells, suggesting that the SAC was minimally impaired (Figure 1D). We synchronized wild-type and *ndc80-112* cells with α -factor and released them to the cell cycle in the presence of benomyl and monitored cell cycle progression by flow cytometry and Pds1 stability. Both wild-type and *ndc80-112* cells arrested with a 2C content of DNA, and Pds1 was stabilized (Figure 1C). We conclude that the N-terminal tail of Ndc80 is dispensable for the SAC.

The N-terminal tail of Ndc80, which is essential in human cells, may have a more limited role in yeast, which could account for the subtle mitotic delay observed in *ndc80-112* cells. We analyzed KT separation in mitotic cells expressing Mtw1-GFP to label all KTs and Spc42-DsRed to label spindle pole bodies (SPBs). There are two populations of cells that are typical (Figure 2A). One has two clearly separated lobes of GFP between the SPBs (bi-lobed), and the other has a single mass of GFP between the SPBs (uni-lobed). The bi-lobed

arrangement reflects sufficient tension between the majority of kinetochores bi-oriented on the spindle to separate the sister chromatids generating the two lobes of GFP. The uni-lobed arrangement occurs when some of the kinetochores have lost the tension between sister chromatids to produce a single mass of GFP fluorescence. We measured the proportion of cells with uni-lobed and bi-lobed GFP signals and found that *ndc80-112* cells had a larger proportion of cells with uni-lobed GFP, suggesting that the N-terminal tail is necessary to produce proper tension between sister chromatids (Figure 2A). We measured the spindle lengths as the distance between spindle poles in the uni-lobed and bi-lobed populations. There was no significant difference in spindle lengths between the uni-lobed spindles of wild-type and *ndc80-112* cells ($1.05 \pm 0.18 \mu\text{m}$ vs. $1.11 \pm 0.14 \mu\text{m}$, $n = 70$, $P = 0.22$ by two-tailed t -test). However, the spindles of bi-lobed *ndc80-112* cells were significantly longer than wild type ($1.78 \pm 0.19 \mu\text{m}$ vs. $1.65 \pm 0.17 \mu\text{m}$, $n = 70$, $P = 0.003$ by two-tailed t -test, Figure 2B). The differences probably reflect different forces exerted on the spindles in the presence or absence of the Ndc80 tail. The effects of deleting the Ndc80 tail have a measurable, albeit subtle, effect on spindle and kinetochore architecture in yeast, which could account for the delayed mitotic progression in *ndc80-112* cells.

The subtle phenotypes associated with *ndc80-112* are surprising, given that the N-terminal tail of Ndc80 is essential in human cells and is required for yeast Ndc80 to bind MTs *in vitro* (Wei *et al.* 2005; Ciferri *et al.* 2008; Miller *et al.* 2008; Tooley *et al.* 2011). One of the differences between kinetochores of yeast and human cells is the Dam1 complex that serves as a processivity factor for the Ndc80 complex, allowing Ndc80 to track the plus ends of microtubules and establish load-bearing connections (Lampert *et al.* 2010;

Figure 2 The N-terminal tail of Ndc80 is required for spindle architecture and is essential under conditions of limited Dam1. Cells expressing Mtw1-GFP and Spc42-DsRed were synchronized by growth in hydroxyurea, released to the cell cycle, and photographed after 20 min at a time just preceding nuclear division when >90% of the cells were pre-anaphase (Keyes and Burke 2009). Cells from the time point were analyzed and representative cells with bi-lobed and uni-lobed GFP fluorescence are shown and quantified. Bar, 1 μ m (A). The lengths of spindles as determined by the distance between Spc42-DsRed spots in the bi-lobed cells were rank-ordered and plotted (B). Serial 10-fold dilutions of saturated cultures of MAT cells of the indicated genotypes were spotted onto YPD plates and grown at the indicated temperatures (C). The cells were assayed for a-like fakers (ALF) as described (Warren *et al.* 2004).

Tien *et al.* 2010). We constructed double mutants between *ndc80-112* and *dam1-1* and found that the double mutant was impaired for growth compared to the single mutants when grown at a semipermissive temperature (Figure 2C). Therefore, the Ndc80 tail is essential for growth under conditions where cells are limited for Dam1 activity. There is no synthetic interaction with five other kinetochore mutants (Figure S2), and therefore the synthetic interaction with *dam1-1* does not reflect a structurally weakened kinetochore in *ndc80-112* cells. We measured the fidelity of chromosome

segregation in the strains using the established “a-like faker” assay that measures the loss of chromosome III in haploid MAT α cells (Strathern *et al.* 1981; Warren *et al.* 2004). The *ndc80-112* and *dam1-1* cells lose chromosomes at three times the rate of wild-type cells (Figure 2C). The *ndc80-112 dam1-1* cells lose chromosomes at 56 times the rate of wild-type cells, confirming that the double mutants have reduced viability due to increased chromosome loss. The microtubule architecture of the mitotic spindle in *ndc80-112 dam1-1* cells appeared normal at the semipermissive temperature (Figure S3). Together, these data suggest redundant functions for Ndc80 and Dam1 in establishing KT-MT attachments. The functional interaction between the N-terminal tail of Ndc80 and Dam1 is not at the level of recruitment of Ndc80 to the plus ends of dynamic microtubules *in vitro* (Lampert *et al.* 2010). The difference between human and yeast Ndc80 may be the combination of Dam1 and the architecture of a single MT attachment to kinetochores in yeast as opposed to the multiple MT attachments and the cooperative nature of Ndc80-MT interactions in human cells (Alushin *et al.* 2010; Tooley and Stukenberg 2011).

Acknowledgments

We thank all members of the Stukenberg and Foltz labs for reagents, equipment, and helpful discussions throughout this work. The work was supported by National Institutes of Health grant GM086502.

Literature Cited

- Alushin, G. M., V. H. Ramey, S. Pasqualato, D. A. Ball, N. Grigorieff *et al.*, 2010 The Ndc80 kinetochore complex forms oligomeric arrays along microtubules. *Nature* 467: 805–810.
- Ciferri, C., S. Pasqualato, E. Screpanti, G. Varetto, S. Santaguida *et al.*, 2008 Implications for kinetochore-microtubule attachment from the structure of an engineered Ndc80 complex. *Cell* 133: 427–439.
- Guimaraes, G. J., Y. Dong, B. F. McEwen, and J. G. Deluca, 2008 Kinetochore-microtubule attachment relies on the disordered N-terminal tail domain of Hec1. *Curr. Biol.* 18: 1778–1784.
- Kemmler, S., M. Stach, M. Knapp, J. Ortiz, J. Pfannstiel *et al.*, 2009 Mimicking Ndc80 phosphorylation triggers spindle assembly checkpoint signalling. *EMBO J.* 28: 1099–1110.
- Keyes, B. E., and D. J. Burke, 2009 Irc15 is a microtubule-associated protein that regulates microtubule dynamics in *Saccharomyces cerevisiae*. *Curr. Biol.* 19: 472–478.
- Lampert, F., P. Hornung, and S. Westermann, 2010 The Dam1 complex confers microtubule plus end-tracking activity to the Ndc80 kinetochore complex. *J. Cell Biol.* 189: 641–649.
- Lampson, M. A., and I. M. Cheeseman, 2011 Sensing centromere tension: Aurora B and the regulation of kinetochore function. *Trends Cell Biol.* 21: 133–140.
- Li, Y., J. Bachant, A. A. Alcasabas, Y. Wang, J. Qin *et al.*, 2002 The mitotic spindle is required for loading of the DASH complex onto the kinetochore. *Genes Dev.* 16: 183–197.
- McClelland, M. L., R. D. Gardner, M. J. Kallio, J. R. Daum, G. J. Gorbsky *et al.*, 2003 The highly conserved Ndc80 complex is

- required for kinetochore assembly, chromosome congression, and spindle checkpoint activity. *Genes Dev.* 17: 101–114.
- Miller, S. A., M. L. Johnson, and P. T. Stukenberg, 2008 Kinetochore attachments require an interaction between unstructured tails on microtubules and Ndc80(Hec1). *Curr. Biol.* 18: 1785–1791.
- Miranda, J. J., P. De Wulf, P. K. Sorger, and S. C. Harrison, 2005 The yeast DASH complex forms closed rings on microtubules. *Nat. Struct. Mol. Biol.* 12: 138–143.
- Strathern, J., J. Hicks, and I. Herskowitz, 1981 Control of cell type in yeast by the mating type locus. The alpha 1-alpha 2 hypothesis. *J. Mol. Biol.* 147: 357–372.
- Tien, J. F., N. T. Umbreit, D. R. Gestaut, A. D. Franck, J. Cooper *et al.*, 2010 Cooperation of the Dam1 and Ndc80 kinetochore complexes enhances microtubule coupling and is regulated by aurora B. *J. Cell Biol.* 189: 713–723.
- Tooley, J., and P. T. Stukenberg, 2011 The Ndc80 complex: integrating the kinetochore's many movements. *Chromosome Res.* 19: 377–391.
- Tooley, J. G., S. A. Miller, and P. T. Stukenberg, 2011 The Ndc80 complex uses a tripartite attachment point to couple microtubule depolymerization to chromosome movement. *Mol. Biol. Cell* 22: 1217–1226.
- Warren, C. D., D. M. Eckley, M. S. Lee, J. S. Hanna, A. Hughes *et al.*, 2004 S-phase checkpoint genes safeguard high-fidelity sister chromatid cohesion. *Mol. Biol. Cell* 15: 1724–1735.
- Wei, R. R., P. K. Sorger, and S. C. Harrison, 2005 Molecular organization of the Ndc80 complex, an essential kinetochore component. *Proc. Natl. Acad. Sci. USA* 102: 5363–5367.
- Welburn, J. P., M. Vleugel, D. Liu, J. R. Yates III, M. A. Lampson *et al.*, 2010 Aurora B phosphorylates spatially distinct targets to differentially regulate the kinetochore-microtubule interface. *Mol. Cell* 38: 383–392.
- Westermann, S., H. W. Wang, A. Avila-Sakar, D. G. Drubin, E. Nogales *et al.*, 2006 The Dam1 kinetochore ring complex moves processively on depolymerizing microtubule ends. *Nature* 440: 565–569.
- Wigge, P. A., and J. V. Kilmartin, 2001 The Ndc80p complex from *Saccharomyces cerevisiae* contains conserved centromere components and has a function in chromosome segregation. *J. Cell Biol.* 152: 349–360.
- Yellman, C. M., and D. J. Burke, 2006 The role of Cdc55 in the spindle checkpoint is through regulation of mitotic exit in *Saccharomyces cerevisiae*. *Mol. Biol. Cell* 17: 658–666.

Communicating editor: O. Cohen-Fix

GENETICS

Supporting Information

<http://www.genetics.org/lookup/suppl/doi:10.1534/genetics.112.143818/-/DC1>

A Redundant Function for the N-Terminal Tail of Ndc80 in Kinetochore–Microtubule Interaction in *Saccharomyces cerevisiae*

Pinar B. Demirel, Brice E. Keyes, Mandovi Chaterjee, Courtney E. Remington, and Daniel J. Burke

Figure S1 *ndc80-112* cells are viable at temperatures between 14 and 37 degrees C. A. DNA was extracted from cells after sporulation and tetrad dissection and subjected to PCR with primers flanking the deletion site. The results from six tetrads are shown. The white arrow indicates the deletion allele, which was completely linked to Ura⁺ in all six tetrads. B. Serial ten-fold dilutions of cells grown to saturation were spotted onto YPD plates and incubated at the indicated temperatures. There was no obvious growth phenotype associated with *ndc80-112* cells at any of the four temperatures

Figure S2 Synthetic interactions with *ndc80-112* are not a general feature of kinetochore mutants. Cells of the indicated genotypes were grown to saturation at 23 degrees and serial 10 fold dilutions were spotted onto YPD agar plates and incubated at the indicated temperatures.

Anti-tubulin

DAPI

dam1-1

dam1-1 ndc80-112

Figure S3 Indle architecture is normal as determined by indirect immunofluorescence. Cells were grown to mid-log in YM-1 plus glucose at 23 degrees before incubation at 32 degrees for three hours. Cell were fixed in 4% paraformaldehyde and processed for indirect immunofluorescence to detect microtubules with YOL 1/34 (anti-tubulin) and DNA with DAPI.

Table S1 List of Strains

3139-8-4	<i>MATa his3Δ1 leu2-3,112 ura3-52 can1 mad2::NAT can1H cyh2H gal1H</i>
3238-5	<i>MATα ade3 his4 lys2 trp1 tyr1 sup4-3 cyh2</i>
8058-3-4	<i>MATa ade2-1 bar1::lox P his3Δ1 leu2-3,112 ura3-52 trp1-289 can1Hcyh2H gal1H</i>
8058-13-2	<i>MATα ade6-1 bar1::lox P his3Δ1 leu2-3,112 ura3-52 trp1-289 can1Hcyh2H gal1H</i>
3748-3-3	<i>MATα ade2-1 bar1::lox P his3Δ1 leu2-3,112 ura3-52 trp1-289 ndc80-112:URA3 can1H cyh2H gal1H</i>
3748-3-4	<i>MATa ade2-1 bar1::lox P his3Δ1 leu2-3,112 ura3-52 trp1-289 ndc80-112:URA3 can1H cyh2H gal1H</i>
3746-10-2	<i>MATα ade2-1 his3Δ1 leu2-3,112 ura3-52 dam1-1::KanMX can1H cyh2Hgal1H</i>
3746-2-4	<i>MATα ade2-1 bar1::loxP his3Δ1 leu2-3,112 ura3-52 dam1-1::KanMXndc80-112:URA3 can1H cyh2Hgal1H</i>
3751-21-3	<i>MATα ade2-1 bar1::loxP his3Δ1 leu2-3,112 ura3-52 MTW1-GFP:URA3 SPC42- DsRed:KanMX4 can1H cyh2H gal1H</i>
3751-14-1	<i>MATa ade2-1 bar1::loxP his3Δ1 leu2-3,112 ura3-52 ndc80-112 :URA3MTW1-GFP:URA3 SPC42-DsRed:KanMX4 can1H cyh2H gal1H</i>

All strains are isogenic with A364a except 3238-5, which is isogenic with 381G