Effect of pentoxifylline in reducing oxidative stress-induced embryotoxicity¹

Xiaoyan Zhang,² Rakesh K. Sharma,^{2,3} Ashok Agarwal,² and Tommaso Falcone²

Submitted May 13, 2005; accepted July 6, 2005

Purpose: To 1) evaluate the embryotoxic effects of hydrogen peroxide on mouse embryo development and 2) examine if pentoxifylline can reverse hydrogen peroxide induced embryotoxicity.

Methods: Prospective in vitro study examining the effects of varying concentrations of hydrogen peroxide and pentoxifylline on the blastocyst development rate alone as well as in combination.

Results: A dose-dependent decrease in % BDR was seen with increasing concentrations of H₂O₂. High concentrations of hydrogen peroxide (>60 μM) were embryotoxic. Pentoxifylline (500 μM) was able to reduce the embryotoxic effect of hydrogen peroxide. Percent blastocyst development rate increased from 44% in hydrogen peroxide alone to 85% in hydrogen peroxide and pentoxifylline coinubation.

Conclusions: Pentoxifylline may be beneficial in reducing hydrogen peroxide induced embryo damage and improve IVF outcome. Patients with endometriosis-associated infertility may benefit from the use of pentoxifylline without significantly affecting embryo development.

KEY WORDS: Embryotoxicity; hydrogen peroxide; pentoxifylline.

INTRODUCTION

Reactive oxygen species (ROS) is involved in the modulation of an entire spectrum of physiological reproductive functions such as oocyte maturation, ovarian steroidogenesis, ovulation, implantation, formation of blastocyst, maintenance in pregnancy, corpus luteal function and even luteolysis (1,2). Hydrogen peroxide (H₂O₂) is a strong oxidant and a major source of ROS. Oxidative stress occurs as

a result of an imbalance between the pro-oxidants and the ability of the antioxidants to scavenge the excess ROS production. Oxidative stress in female reproduction may be a major link in the infertility puzzle and in some reproductive diseases of women such as endometriosis (1,3).

High ROS levels are associated with endometriosis, unexplained infertility and poor in vitro fertilization (IVF) outcome. Peritoneal, follicular and hydrosalpingeal fluids represent important reproductive microenvironments (3–6). Evidence show increased oxidative stress in women with endometriosis (7). There may be multiple sources of ROS in an IVF setting including the oocytes, cumulus cell mass and spermatozoa used for insemination (8). Oxidative stress can damage oocytes and cause mitochondrial alterations, embryo block, adenosine triphosphatase depletion and apoptosis (9,10).

Pentoxifylline, a methylxanthine acts as a phosphodiesterase inhibitor. It inhibits phagocytosis and

¹ Presented at the 60th Annual Meeting of the American Society for Reproductive Medicine, Philadelphia, PA, October 16–20, 2004.

² Center for Advanced Research in Human Reproduction, Infertility and Sexual Function, Department of Obstetrics-Gynecology, and Glickman Urological Institute, Cleveland Clinic Foundation, Cleveland, Ohio.

³ To whom correspondence should be addressed at Desk A19.1, Cleveland Clinic Foundation, 9500 Euclid Avenue, Cleveland, Ohio 44195, USA; e-mail: sharmar@ccf.org.

generation of toxic ROS and proteolytic enzymes. Pentoxifylline influences both the production of inflammatory mediators and the responsiveness of immunocompetent cells to inflammatory stimuli (11–14). The importance of protecting preimplantation embryos from oxidative stress is important and must be carefully monitored. The present study was designed to evaluate the embryotoxic effects of H_2O_2 and pentoxifylline alone as well as in combination and examine if pentoxifylline could reverse H_2O_2 induced embryotoxicity.

MATERIALS AND METHODS

A culture system using cryopreserved two-cell mouse embryos (Embryotech laboratories, Inc., Wilmington, MA) was used to evaluate the effects of exogenous addition of hydrogen peroxide and pentoxifylline (Sigma Chemical Co., St. Louis, MO) alone as well in combination. In brief, frozen straws were thawed and only embryos with a normal morphological appearance were used. Incubation was carried out at 37°C under a humidified atmosphere of 5% CO_2 in air. Groups of 9–10 embryos were randomly added to the inner well of a petri dish containing 1 ml of human tubal fluid (HTF, Irvine Scientific, Santa Ana, CA) without oil overlay. Embryos in all groups were monitored daily at the same time in the morning using Normarski and bright field inverted optics. The number of embryos cleaving to morula and blastocysts (early, expanding, and expanded) were recorded. A blastocyst development rate (% BDR) of over 80% was considered satisfactory for internal control.

Hydrogen peroxide was diluted with HTF to give five working concentrations of 25, 50, 60, 75 and $100\ \mu\text{M}$. A total of 326 embryos were divided into five groups and each concentration of H_2O_2 was repeated at least three times. Similarly, a total of 239 embryos were used to examine the embryotoxicity of pentoxifylline. Pentoxifylline was dissolved in HTF to give five working concentrations of 250, 500, 1000, 2000 and $4000\ \mu\text{M}$. Embryos cultured in HTF media ($n = 72$) served as the control. Once the lowest dose of H_2O_2 that was embryotoxic and the highest dose of pentoxifylline that was not embryotoxic was determined, we examined the beneficial effect of pentoxifylline in reversing H_2O_2 induced embryotoxicity. A total of 175 embryos were cultured in three groups: H_2O_2 ($60\ \mu\text{M}$) alone, pentoxifylline ($500\ \mu\text{M}$) alone and H_2O_2 ($60\ \mu\text{M}$) + pentoxifylline ($500\ \mu\text{M}$).

Comparisons between groups were performed with Fisher's exact test. A P -value of <0.05 was considered significant using the two-tailed test. Computations were performed with GraphPad In-Stat version 3.00 statistical software (GraphPad Software, Inc., San Diego, CA).

RESULTS

A total of 740 embryos were used in the study to examine the embryotoxic effects of hydrogen peroxide and pentoxifylline. A dose-dependent decrease in % BDR was seen with increasing concentrations of H_2O_2 compared with controls. A significant increase in embryotoxicity was seen at $50\ \mu\text{M}$ H_2O_2 compared with controls (% BDR: 69% versus 94% $P < 0.01$). Higher concentrations were embryotoxic and a complete block was seen at the two-cell stages with $75\ \mu\text{M}$ of H_2O_2 . Concentrations up to $500\ \mu\text{M}$ were not embryotoxic and. Compared with control, a significant decrease in % BDR (75% versus 96% $P < 0.008$) was seen at $1000\ \mu\text{M}$ of pentoxifylline. Pentoxifylline at $500\ \mu\text{M}$ was able to reduce the embryotoxic effect of H_2O_2 . A significant increase in % BDR was seen when embryos were incubated in the presence of pentoxifylline (Fig. 1). The % BDR increased from 44% in embryos cultured in the presence of H_2O_2 ($60\ \mu\text{M}$) alone to 85% in culture medium containing H_2O_2 and pentoxifylline ($P = 0.0043$).

Fig. 1. Effect of pentoxifylline (PTX: $500\ \mu\text{M}$) in combination with $60\ \mu\text{M}$ H_2O_2 in reducing the embryotoxic effect of H_2O_2 . Each point represents the mean of the readings in triplicate with 6–8 embryos in each group. The % BDR increased from 44% in H_2O_2 alone to 85% in H_2O_2 and pentoxifylline ($P = 0.0043$).

DISCUSSION

Oxidative stress has been proposed as a potential factor involved in the pathophysiology of endometriosis. Large amounts of ROS are released after chronic stimulation of peritoneal fluid macrophages in women with endometriosis. Oxidative stress-induced increase in autoantibody titers along with elevated levels of lysophosphatidyl choline have been demonstrated in the peritoneal fluid women with endometriosis (15).

During in vitro fertilization-embryo transfer (IVF-ET), the oxygen concentration is higher in in vitro culture environment compared with the in vivo conditions. Embryos and sperms may also generate ROS when they are cultured in vitro. Furthermore, when oocytes and embryos are removed from their natural environment for assisted reproduction techniques, their natural defense system is lost resulting in formation of ROS.

Reactive oxygen species in culture media affect fertilization, embryo development and clinical pregnancy rates. Elevated concentrations of ROS in day-1 culture media have been shown to be associated with lower pregnancy rates both with IVF and intracytoplasmic sperm injection (ICSI) (8). Our study demonstrates that the blastocyst development rate decreased significantly even at 60 μM H_2O_2 and higher concentrations caused a block at two-cell stage. Overproduction of ROS resulting from impaired intracellular milieu and disturbed metabolism is detrimental for the embryo (2).

Pentoxifylline is a 3',5'-nucleotidase phosphodiesterase inhibitor that protects cells from lipid peroxidation by ROS. It could be used as antioxidant in IVF. Furthermore, because of its potent immunomodulatory properties, pentoxifylline has been proposed as a treatment for endometriosis-associated infertility. Steinleitner and collaborators reported that pentoxifylline might fully reverse the infertility effects observed in rodent models of endometriosis (12,13). In our study, pentoxifylline at 500 μM significantly reduced the embryotoxic effects of H_2O_2 . This concentration of pentoxifylline is much higher than the peak plasma concentrations of pentoxifylline seen after intravenous infusion of 200 mg (about 10 μM) or 400 mg (about 4 μM) oral doses of capsule formulation (11). In this study, we did not examine the lower dose, it would be interesting to see if the lower doses of pentoxifylline are also equally effective in reducing the H_2O_2 induced embryotoxicity. Our results indicate that pentoxifylline

may be beneficial in reducing H_2O_2 induced embryo damage and improve IVF outcome. Patients with endometriosis-associated infertility may benefit from the use of pentoxifylline without significantly affecting the embryo development.

REFERENCES

1. Sharma RK, Agarwal A: Role of reactive oxygen species in gynecologic diseases. *Reprod Med Biol* 2004;3:177–199
2. Guerin P, El Mouatassim S, Menezo Y: Oxidative stress and protection against reactive oxygen species in the pre-implantation embryo and its surroundings. *Hum Reprod Update* 2001;7:175–189
3. Bedaiwy MA, Falcone T, Sharma RK, *et al.*: Prediction of endometriosis with serum and peritoneal fluid markers: A prospective controlled trial. *Hum Reprod* 2002;17:426–431
4. Attaran M, Pasqualotto E, Falcone T, *et al.*: The effect of follicular fluid reactive oxygen species on the outcome of in vitro fertilization. *Int J Fertil Womens Med* 2000;45:314–320
5. Pasqualotto EB, Agarwal A, Sharma RK, *et al.*: Effect of oxidative stress in follicular fluid on the outcome of assisted reproductive procedures. *Fertil Steril* 2004;81:973–976
6. Bedaiwy MA, Goldberg JM, Falcone T, Singh M, Nelson D, Azab H, Sharma RK: Relationship between oxidative stress and embryotoxicity of hydrosalpingeal fluid. *Hum Reprod* 2002;17:601–604
7. Szczepanska M, Kozlik J, Skrzypczak J, Mikolsajczyk M: Oxidative stress may be a piece in the endometriosis puzzle. *Fertil Steril* 2003;79:1288–1293
8. Bedaiwy MA, Falcone T, Mohamed MS, *et al.*: Differential growth of human embryos *in vitro*: Role of reactive oxygen species. *Fertil Steril* 2004;82:593–600
9. Yang HW, Hwang KJ, Kwon HC: Detection of reactive oxygen species (ROS) and apoptosis in human fragmented embryos. *Hum Reprod* 1998;13:998–1002
10. Nasr-Esfahani MH, Aitken RJ, Johnson MH: Hydrogen peroxide levels in mouse oocytes and early cleavage stage embryos developed in vitro or in vivo. *Development* 1990;109:501–507
11. Ward A, Stephen PC: Pentoxifylline. A review of its pharmacodynamic and pharmacokinetic properties, and its therapeutic efficacy. *Drugs* 1987;34:50–97
12. Steinleitner A, Lambert H, Roy S: Immunomodulation with pentoxifylline abrogates macrophage-mediated infertility in an in vivo model: A paradigm for a novel approach to the treatment of endometriosis-associated subfertility. *Fertil Steril* 1991;55:26–31
13. Steinleitner A, Lambert H, Suarez M: Immunomodulation in the treatment of endometriosis-associated subfertility: Use of pentoxifylline to reverse the inhibition of fertilization by surgically induced endometriosis in a rodent model. *Fertil Steril* 1991;55:975–979
14. Juan B, Montserrat C, Francisco F: Pentoxifylline versus placebo in the treatment of infertility associated with minimal or mild endometriosis: A pilot randomized clinical trial. *Hum Reprod* 1997;12:1046–1050
15. Shanti A, Santanam N, Morales AJ, Parthasarathy S, Murphy AA: Autoantibodies to markers of oxidative stress are elevated in women with endometriosis. *Fertil Steril* 1999;71:1115–1118