SHORT COMMUNICATIONS

TAIPEI, TAIWAN

Pregnancy After Preimplantation Genetic Diagnosis by Fluorescence In Situ Hybridization Using 18-, X-, and Y-Chromosome Probes in an Infertile Male with Mosaic Trisomy 18

Submitted: August 30, 1999

Accepted: December 27, 1999

INTRODUCTION

Infertility affects about 10% of couples, and male infertility is one of the major causes and may be on the rise in humans. Male infertility also has attracted a great deal of attention from geneticists (1,2). It is suggested by Peschka *et al.* (3) that chromosomal analysis should be performed before intracytoplasmic sperm injection (ICSI) treatment. From their series, there are 13.1% patients in 781 male infertility couples with chromosomal abnormality including one trisomy 18 mosaicism (3). In our reproduction center we routinely do chromosome analysis and Y chromosome microdeletion detection in the azoospermic or severely oligozoospermic patient when their sperm count is lower than 5×10^6 . We report a man with severe oligoasthenozoospermia whose chromosome study showed mosaic trisomy of chromosome 18. His wife went through in vitro fertilization (IVF) with ICSI. Preimplantation genetic diagnosis was done by applying fluorescence in situ hybridization using 18-, X-, and Y-chromosome probes (5). She became pregnant and the karyotype from the amniocentesis showed a normal female baby.

CASE REPORT

A 30-year-old male patient was first diagnosed to be azoospermic at another hospital. He visited our

urology clinic where semen analysis and blood chromosome and Y chromosome microdeletion were examined. Semen analysis showed azoospermia. After the sperm wash and spin down there was 0 to 1 sperm per high-power field. The blood chromosome karyotyping showed mosaic trisomy 18 in 20% of the cell lines. A Yq11 microdeletion was examined by applying two multiplex polymerase chain reaction (PCR) and each with six sets of PCR primers. The locus of each primer was listed in Table 1. After the PCR reaction, 10 μ l of the product was run on 3% agarose gel and the picture was shown (Fig. 1). No microdeletion was detected. His wife was scheduled to go through our IVF and PGD program. Before ovulation induction, down-regulation was done first by using 0.2 cc leupron subcutaneous injection on day 21 of her menstrual cycle. After down regulation, 2 ampoules of Metrodin and Pergonal were used for ovulation induction. When follicle size was around 1.6 cm, human chorionic gonadotropin (hCG) 10,000 IU was injected and oocyte retrieval was scheduled 32 h later. Twenty-seven eggs were retrieved with 25 of them at the metaphase-II phase.

ICSI was done with 7 motile and 18 immotile sperm. The total fertilization rate was 40%. The fertilization rate with motile sperm was 42.9% (3/7), and immotile sperm was 38.9% (7/18), excluding one abnormal fer-

Table I

Locus	Primer name	Size
sy95	5' DYS280-a 3' DYS280-b	303 bp
sy117	5' DYS209-a 3' DYS209-b	280 bp
sy159	5' DYZ2-a 3' DYZ2-b	213 bp
sy127	5' DYS7-a 3' DYS7-b	160 bp
DYS239	5' DYS239-a 3' DYS239-b	126 bp
sy153	5' DYS237-a 3' DYS237-b	100 bp

Table II

Locus	Primer name	Size
RPS4Y	5' RPS4Y 3' sy274	350 bp
DAZ	5' DAZ 3' sy277	306 bp
AMELY	5' AMELY 3' sy276	200 bp
DYS271	5' DYS271 3' sy81	170 bp
DYS1	5' DYS1 3' sy149	132 bp
DYS232	5' DYS232 3' sy147	100 bp

tilization (3PN). All the fertilized eggs cleaved; only one embryo went through embryo biopsy because of poor embryo quality of the rest of the embryos. The biopsy was done on the third day after oocyte retrieval when it was at the 8-cell stage. One blastomere was removed and fixed on the slide as per the procedure published by Muune' 1993 (4,5). Fluorescence in situ hybridization was done by using 18-, X-, and Y-chromosome probes and the result was XX with euploid chromosome 18. Embryo quality was compared and it seemed that embryo quality was better in the group fertilized by motile sperm. The embryos that were not transferred and unfertilized eggs were all fixed and (FISH) was done by using the same probes. One embryo had only one chromosome 18 signal with 2 X signals and one embryo had one X and one chromosome 18 signals. Both embryos had been injected with immotile sperm. The unfertilized eggs were cleaved at day 3 and FISH was done with only one chromosome 18 and one X signals. The β -hCG, checked 10 days

after the embryo was transferred, was 105 mIU. Amniocentesis was done at 16 weeks gestation, with the result of a normal female baby.

DISCUSSION

Male infertility seems to be on the rise in humans. This has attracted a great deal of attention from geneticists. It has been reported that a baby born by ICSI procedure has a higher chance of chromosomal anomaly (6). From Dr. Peschka's series there are 13.1% patients in 781 male infertility couples with chromosomal abnormality (3). These chromosomal aberrations included constitutional, low-level mosaicism of sex chromosomes, fragile sites of autosomes, and secondary structural chromosome aberrations. Combinations of different types of abnormalities also are stated. In their 34 autosomal aberration patients, there is one patient with trisomy 18 mosaicism. We report a male infertility patient with trisomy 18 mosaicism. This couple went through IVF, ICSI, and PGD and had a normal pregnancy. We thought patients scheduled for ICSI procedure should have blood chromosome study. A male infertility patient who also has a chromosome anomaly can be scheduled for IVF, ICSI, and PGD program. This way, these patients can obtain better medical care as afforded by the new reproductive technology.

REFERENCES

1. Reijo R, Lee TY, Salo P, Alagappan R, Brown LG, Rosenberg M, Rosen S, Jaffe T, Straus D, Hovatta O, Capelle A, Silber S, Page DC: Diverse spermatogenic defects in humans caused by Y chromosome deletions encompassing a novel RNA-binding protein gene. *Nature Genet* 1995;10:383-393
2. Gianaroli L, Munne S, Magli MC, Ferraretti AP: Preimplantation genetic diagnosis of aneuploidy and male infertility. *Intern J Androl* 1997;20(suppl):31-34
3. Peschka B, Leygraaf J, Van der Ven K, Montag M, Scharmann B, Schubert R, Van der Ven H, Schwanitz G: Type and frequency of chromosome aberrations in 781 couples undergoing intracytoplasmic sperm injection. *Hum Reprod* 1999;14:2257-2263
4. Munne S, Lee A, Rosenwaks Z, Grifo J, Chon J: Diagnosis of major chromosome aneuploidies in human preimplantation embryos. *Hum Reprod* 1993;8:2185-2191
5. Munne S, Sultan KM, Weier HU, Grifo JA, Cohen J, Rosenwaks Z: Assessment of numeric abnormalities of X, Y, 18, and 16 chromosomes in preimplantation human embryos before transfer. *Am J Obstet Gynecol* 1995;172:1191-1199
6. Bonduellu M, Aytoz A, Van Assche E, Devroey P, Liebaers I, Van Steirteghem A: Incidence of chromosomal aberrations in

Fig. 1.

children born after assisted reproduction through intracytoplasmic sperm injection. Hum Reprod 1998;13:781–782

Hsiao-Jui Wei,¹

Han-Sun Chiang

Wei-Min Lin

Jaw-Yuan Wen

Departments of Obstetrics and Gynecology, and Urology
Taipei Medical College Hospital, 9F, No.252
Wu-Hsing St., Taipei, Taiwan

¹ To whom correspondence should be addressed

BRISBANE, AUSTRALIA

Repeated Oocyte Maturation Block

Submitted: October 20, 1999

Accepted: January 22, 2000

INTRODUCTION

While it is common in assisted reproduction treatments for one or two oocytes in any cohort to be immature and even to fail to subsequently mature during in vitro culture, it is highly unusual for all aspirated oocytes to be immature. In 1990, Rudak *et al.* (1) reported a case in which two oocyte aspirations yielded oocytes largely arrested at the germinal vesicle stage. The same report also described two further cases involving sisters whose oocytes fertilized both normally and abnormally but failed to undergo subsequent cleavage. Hartshorne *et al.* (2) recently reported a patient in whom the oocytes from three stimulated cycles failed to mature past the metaphase-I (MI) stage. An attempt to mature those obtained from the third cycle in culture medium supplemented with gonadotropins also failed. While the oocytes from this patient frequently exhibited failure of coronal cells to expand, all other aspects of them, including size and cumulus expansion, indicated maturity.

Two cases are reported here of recognition of an oocyte maturation block when patients proceeded to intracytoplasmic sperm injection (ICSI) after failure of oocytes to fertilize with conventional in vitro fertilization (IVF).

CASE 1

History

The patient presented in 1998, aged 30, with a history of 4 years primary infertility. Her menstrual cycles were slightly irregular, averaging 31 days duration. Laparotomy revealed patent tubes with some minor pelvic adhesions and no evidence of endometriosis. A postcoital test showed numerous sperm but an absence of any motility. The husband's semen revealed normal sperm parameters. She was admitted to the assisted reproduction program with a diagnosis of unexplained infertility.

Treatment with Gamete Intrafallopian Transfer

After pituitary down-regulation from day 20 using Synarel nasal spray (Searle, Delpharm, France), ovarian stimulation was begun using recombinant follicle-stimulating hormone (FSH) (Puregon, Organon, Netherlands) 150 IU daily from day 9 to day 16. Human chorionic gonadotropin (hCG) (Pregnyl, Organon, Netherlands) 10,000 IU was administered on day 17, and laparoscopic oocyte aspiration 36 hr later yielded 18 cumulus–oocyte complexes that appeared morphologically mature. Details of endocrine levels and follicle sizes are shown in Table I. Two oocytes and 100,000 sperm were placed into one fallopian tube in the gamete intrafallopian transfer (GIFT) procedure. The remaining 16 oocytes were inseminated in vitro with 100,000 sperm from an apparently normal semen specimen 4 hr after aspiration and placed in standard embryo culture conditions using human tubal fluid (HTF) culture medium (3) supplemented with 5% patient's heat-denatured preovulatory serum. The oocytes were denuded a further 18 hr later to check fertilization. Of the 18 oocytes, 16 were found to be at the MI stage having failed to extrude the first polar body and the remaining 2 were atretic. The oocytes did not mature any further with another 24-hr incubation in vitro.

Treatment with ICSI

Following similar pituitary down-regulation, ovarian stimulation was begun more gently based on the response in the previous cycle. FSH (Puregon, Organon) 75 IU daily was administered on days 6 to 10, 100 IU on days 11 and 12, and increased to 150 IU on days 14 to 17 due to low response. Details of endocrine levels and follicle sizes are shown in Table I.

Table I. Details of Ovarian Response in the Four Cycles Reported^a

Treatment	Ovarian stimulation	Days FSH	Estradiol (pmole/Liter)	Progesterone (nmole/Liter)	EPU day	Follicle size	Oocytes aspirated
Patient 1							
GIFT	Synarel	8	8262	2.5	19	20 × 3, 19, 18 × 6,	18
	Puregon					16 × 5, 12, 9 × 2	
ICSI	Synarel	12	4814	1.8	19	18 × 4, 16 × 2, 12	7
	Puregon						
Patient 2							
GIFT	Synarel	7	2539	3.7	18	20, 18, 16 × 4,	10
	Puregon					9 × 4	
ICSI	Synarel	12	6025	3.2	19	19 × 7, 9 × 4	11
	Gonal-F						

^a Estradiol, progesterone, and follicle size on day of hCG administration.

Ovulation was triggered with 10,000 IU hCG (Pregnyl, Organon) on day 17, and transvaginal oocyte aspiration a further 36 hr later yielded seven apparently mature cumulus–oocyte complexes. When they were denuded enzymatically 2 hr later in preparation for ICSI, all were found to be at the MI stage again and of normal size. They were held in culture for 20 hr but failed to show any in vitro maturation. ICSI was performed on the MI oocytes 24 hr after aspiration but no fertilization was achieved.

CASE 2

History

The patient presented in 1997, aged 25, with a history of 5 years primary infertility. Her menstrual cycle was regular and of 28 days duration. Laparoscopy revealed normal tubes and ovaries and no endometriosis. The husband's semen showed normal sperm parameters. She was admitted to the assisted reproduction program with a diagnosis of unexplained infertility.

Treatment with GIFT

After pituitary down-regulation from day 20 using Synarel nasal spray (Searle, Delpharm, France), FSH (Puregon, Organon) 150 IU daily was administered on days 9 to 15. Details of endocrine levels and follicle sizes are shown in Table I. Ovulation was triggered with 10,000 IU hCG (Pregnyl, Organon) on day 16 and transvaginal oocyte aspiration a further 36 hr later yielded 10 cumulus–oocyte complexes, a majority showing compacted coronal cells although cumulus expansion was normal. Three oocytes and 100,000

sperm were then placed into one tube by a laparoscopic GIFT procedure. The other 7 oocytes were inseminated in vitro with 100,000 apparently good quality sperm 3 hr after aspiration and placed in standard embryo culture conditions using HTF culture medium (3) supplemented with 5% patient's heat-denatured preovulatory serum. The oocytes were denuded 14 hr later to check fertilization, revealing all to be at the MI stage having failed to extrude the first polar body. The oocytes, which were of normal size, did not mature any further with another 24 hr incubation in vitro.

Treatment with ICSI

Following pituitary down-regulation, ovarian stimulation was commenced using FSH (Gonal-F, Serono, Switzerland) 150 IU from days 6 to 10, and increased to 300 IU on days 11 to 17 due to low response. Details of endocrine levels and follicle sizes are shown in Table I. Ovulation was triggered with 10,000 IU hCG (Profasi, Serono) on day 17 and transvaginal oocyte aspiration a further 36 hr later yielded 11 cumulus–oocyte complexes again with most exhibiting compacted coronal cells but expanded cumulus. After enzymatic denudation 4 hr later in preparation for ICSI, all were found to be at the MI stage again and of normal size. A further 24 hr incubation failed to produce any in vitro maturation and no fertilization was achieved from ICSI of the oocytes at that stage.

DISCUSSION

The cases reported here seem very similar to that of Hartshorne *et al.* (2) in that the maturation of the oocytes had arrested at entry to metaphase and they failed to subsequently mature in vitro. It is difficult to

identify any marker in the treatments to indicate that the defect was due to the ovarian stimulation protocol. Follicular response to the gonadotropins was normal, the follicles grew at a normal rate, the expanded cumulus cells in both cases and coronal cells of the oocytes in one case suggested normal response to the ovulation triggering dose of hCG, and the oocytes were of a normal size. As has been noted by others however, the development of the coronal and cumulus cells correlates poorly with nuclear maturity (4). The presence of patent fallopian tubes, a regular menstrual cycle, and long primary infertility to partners with normal semen suggest that the maturation defect also is likely to be occurring in natural cycles in vivo. Further evidence that the problem is an absolute maturation block comes from evidence that normal MI oocytes from cohorts including others at the MII stage can be fertilized by ICSI, albeit at a quite reduced rate (5) and often abnormally (6). In the cases reported here the intracytoplasmic injection of sperm failed to produce any fertilization.

Both couples were made aware of the potential complications of ICSI on MI oocytes but elected to proceed with it after allowing the oocytes a period on in vitro incubation that has been shown to benefit ICSI fertilization of such oocytes (5). It was our intention in both cases that any oocytes fertilizing as a result of the ICSI and continuing cleavage to the eight-cell stage be screened for numerical chromosomal anomalies by fluorescent in situ hybridization before their transfer was considered.

Unfortunately for couples with this rare problem it usually takes a second treatment cycle to provide more conclusive evidence that an absolute oocyte maturation block exists. Potential mechanisms have been postulated but we are far from being able to correct the problem. Hypoxanthine potentially is involved (7). It has been isolated from follicular fluid, has been added to culture medium to inhibit germinal vesicle breakdown in oocytes from several species, and produces a block at the two-cell stage of mouse embryo development in vitro. In both cases reported here the couples declined any further investigatory cycles to study their oocytes and have transferred to donor oocyte programs.

REFERENCES

1. Rudak E, Dor J, Kimchi M, Goldman B, Levran D, Mashiach S: Anomalies of human oocytes from infertile women undergoing treatment by in vitro fertilization. *Fertil Steril* 1990;54:292–296
2. Hartshorne G, Montgomery S, Klentzeris L: A case of failed oocyte maturation in vivo and in vitro. *Fertil Steril* 1999;71:567–570
3. Quinn P, Kerin JF, Warnes GM: Improved pregnancy rate in human in vitro fertilization with the use of a medium based on the composition of human tubal fluid. *Fertil Steril* 1985;44:493–498
4. Rattanachaiyanont M, Leader A, Leveille M-C: Lack of correlation between oocyte–corona–cumulus complex morphology and nuclear maturity in oocytes collected in stimulated cycles for intracytoplasmic sperm injection. *Fertil Steril* 1999;71:937–940
5. Strassburger D, Komarovsky D, Bern O, Kasterstein E, Raziel A, Freidler S, Schachter M, Ron-El R: *Hum Reprod* 1999; Abstract Book 1; Abstract O-011: 6
6. Rybouchkin A, Van der Elst J, Dhont M: Intracytoplasmic sperm injection into immature human oocytes. *Hum Reprod* 1999; Abstract Book 1; Abstract O-012: 7
7. Downs SM: Purine control of mouse oocyte maturation: Evidence that nonmetabolized hypoxanthine maintains meiotic arrest. *Mol Reprod Dev* 1993;35:82–94

Keith L. Harrison¹

Deborah A. Sherrin

J. Douglas Keeping

Queensland Fertility Group

225 Wickham Terrace

Brisbane 4000, Australia

¹ To whom correspondence should be addressed.

HYOGO AND TOCHIGI, JAPAN

Successful Pregnancies in a Case of Retrograde Ejaculation Associated with Tethered Spinal Cord Syndrome

INTRODUCTION

Retrograde ejaculation is an uncommon cause of male infertility but the most common cause of an absent ejaculate. The incidence of retrograde ejaculation has been reported as 0.4% to less than 2.0% (1–3). The patient complains of aspermia while the orgasmic sensation is preserved (4). In some cases, retrograde ejaculation may appear as an incomplete form that may lead to them being diagnosed as severe oligozoospermia (3,5,6). The diagnosis of retrograde ejaculation is confirmed when spermatozoa are present in postcoital or postmasturbatory specimens of urine.

A variety of physiological or neurological abnormalities have been postulated to cause retrograde ejacula-

tion. Any alteration in the neurological stimulus involving semen emission and bladder neck competence may affect a man's normal antegrade ejaculation, leading to retrograde ejaculation. Yavetz *et al.* (2) reviewed the etiologies among 91 infertile patients with retrograde ejaculation collected from 27 reports. Nine had congenital abnormalities, while others had acquired causes including neurogenic, traumatic, and idiopathic problems.

Here we report a rare case of retrograde ejaculation associated with tethered spinal cord syndrome, which is caused by a lipoma confluent with the spinal cord. The first pregnancy was achieved by the intrauterine insemination (IUI) using sperm obtained from the bladder, and subsequently a healthy female infant was born at term. The second pregnancy was established by intracytoplasmic sperm injection (ICSI) using testicular sperm because obstruction probably caused by the inflammation at the level of bilateral epididymis was diagnosed after the initial pregnancy. The course of the second pregnancy also was uneventful and a healthy female infant was born at term.

CASE REPORT

A 27-year-old man was referred to our clinic because of infertility resulting from aspermia. The patient was sexually active and able to have erections and sexual intercourse. As the patient had bladder dysfunction from childhood, he was familiar with self-catheterization. Congenital tethered spinal cord syndrome had been diagnosed when he was 21 years old. A computerized tomography (CT) (not shown) was carried out and revealed a lipoma confluent with the spinal cord. A myelography (Fig. 1) was performed and the tethered spinal cord syndrome was diagnosed. His wife was 23 years old and had no abnormal findings in the endocrinological and immunological examinations as well as in a hysterosalpingography. The diagnosis of retrograde ejaculation was given by the presence of spermatozoa in the postmasturbatory urine.

To evaluate semen characteristics, the patient inserted a catheter to empty the bladder before masturbation. The second self-catheterization was done to collect sperm immediately after masturbation. Semen examination including the detection of antisperm antibodies as well as the sperm function tests such as the hemizona assay (HZA) (7), the hypoosmotic swelling test (HOST) (8), and the Penetrak test (9) were performed using the sample obtained from the bladder. As shown in Table I, the semen characteristics were

Fig. 1. A myelography of the patient showing that the spinal cord is tethered to the level of L5, leading to a diagnosis of the tethered spinal cord syndrome.

normal except for the sperm motility (30% with forward progression), according to the criteria of the World Health Organization (10). The HZA and the HOST tested normal. The value of penetration assessed by the Penetrak test showed a slightly unsatisfactory result (29 mm/90 min). The sperm collected from the patient had neither sperm-agglutinating nor sperm-immobilizing antibody. Moreover, a medical treatment with imipramine hydrochloride did not lead to a restoration of antegrade ejaculation. These findings indicated a less invasive treatment, such as IUI, could be the initial treatment strategy for this couple.

To minimize major problems associated with the acidity, osmolarity, and urine contamination, which might cause less sperm-fertilizing ability due to sperm immobilization and disruption of the sperm cell membrane (11), the patient emptied the bladder, then collected semen directly into a wide-mouthed sterile

Table I. Initial Evaluation of the Sperm Collected from the Patient's Bladder

Semen characteristics ^a		(normal range)
Volume	2.5 ml	2.0 $\leq$
Sperm concentration	22 $\times 10^6$ /ml	20 $\times 10^6 \leq$
Total sperm count	55 $\times 10^6$ /ejaculate	40 $\times 10^6 \leq$
Motility	30% with forward progression	50 $\leq$
Morphology	80% with normal forms	30 $\leq$
Sperm function tests		
Hemizona index ^b	80	36 $\leq$
Total swollen sperm ^c	54%	50 $\leq$
Penetration value ^d	29 mm/90 min	30 $\leq$
Antisperm antibody tests		
Sperm agglutination test	Negative	Negative
Sperm immobilization test	Negative	Negative

^a The normal criteria was cited from the text of WHO (10).

^b Hemizona index was assessed by the hemizona assay (7).

^c Total swollen sperm was assessed by the hypo-osmotic swelling test (8).

^d Penetration value was assessed by the Penetrak test (9).

container filled with a 12 ml volume of sperm washing medium (Irvine Scientific, USA) by catheterization soon after masturbation. Sperm suspension was immediately centrifuged at $300 \times g$ for 5 min in a test tube and the pellet was collected, resuspended using the same medium, and washed twice. A 0.5 ml volume of sperm washing medium was added on the pellet and mixed. Following 1 hr of sperm incubation at 37°C in air, a treatment with IUI was performed at the time of ovulation. The total volume inseminated, the sperm concentration and the sperm motility were 0.8 ml, 10×10^6 /ml, and 78%, respectively at the 4th IUI treatment cycle when the patient conceived. The pregnancy course was uneventful. In August 1993, a normal female infant weighing 3048 g was born transvaginally at 41 weeks' gestation.

In April 1995, the treatment with IUI using the same procedure was started because the couple wished for another pregnancy. Following three unsuccessful treatment cycles with IUI, the patient's semen quality worsened. Severe oligoasthenozoospermia with a bacterial contamination in the seminal plasma was diagnosed. The use of antibiotics in combination with IUI treatment was continued for another five cycles. However, the IUI treatment was discontinued because an obstructive azoospermia was diagnosed in November 1997. After a consultation, the couple wished to proceed to surgical retrieval of the sperm.

The patient's partner was stimulated using a combination of a gonadotropin-releasing hormone (GnRH) agonist started in the luteal phase (suppression protocol) followed by gonadotropins as previously reported (12–14). Because the sperm could not be obtained by microsurgical epididymal sperm aspiration (MESA), the samples were retrieved from testicular tissue by

the tissue-excision method, which minimized the risk of a testicular hematoma. Under spinal anesthesia, the scrotum was opened by making a small incision, then further incision was made in the tunica albuginea. A portion of extruding testicular tissue was incised and isolated by surgical scissors. The isolated tissue was transferred to a test tube containing modified human tubal fluid (mHTF) (Irvine Scientific, USA). The sperm were isolated from the testicular tissue by mincing with a surgical blade and passing through a mesh (FALCON 2350, Becton Dickinson, USA). The isolated sperm were suspended in the same medium, centrifuged at $300 \times g$ for 5 min, and the pellet was collected for ICSI. The procedure of ICSI was followed as we previously reported (15–17). Eleven metaphase-II oocytes were injected with the immotile testicular sperm. Three of them fertilized, cleaved, and were transferred. A single pregnancy was achieved and the pregnancy course was uneventful. In June 1999, a normal female infant weighing 3190 g was born transvaginally at 40 week's gestation.

DISCUSSION

In this case report, the first pregnancy in a case of retrograde ejaculation associated with the tethered spinal cord syndrome was described. To our knowledge, this is also the first demonstration of ejaculation disorder complicated in a case of tethered spinal cord syndrome. The tethered spinal cord syndrome occurs commonly in children, causing pain, neurological deficits, and bowel and bladder dysfunction (18,19). It is commonly believed that surgical spinal cord untethering in children is likely to prevent neurological dete-

rioration (19,20). It also is reported that the older children and the adults with occult spinal dysraphism are more likely to present with irreversible urological and neurological findings than the younger children (20). In our case, a lipoma confluent with the spinal cord had been undiagnosed until he was 21 years old because the only problem had been bladder dysfunction. The incidence of a lipoma, which causes the tethered spinal cord syndrome, has been reported as 14 to 28% (18–20). It is commonly believed that the children who have a congenital tethered spinal cord syndrome benefit from surgical untethering because it prevents neurological deterioration (19,20). However, controversy has surrounded the issue of surgery in newly diagnosed adult patients. Early diagnosis should be done, because the older children and the adults with occult spinal dysraphism were more likely to present with irreversible urological and neurological findings than the younger children (20). After a consultation, our patient wished to be treated without any surgical treatments, because the disease then was not so severe.

Although the assisted reproductive technology, such as in vitro fertilization (IVF) and ICSI, is a powerful treatment option for the patients with retrograde ejaculation, it is recommended that initially less invasive and more convenient method be applied for treatment (2,21). The medical treatment for the correction of retrograde ejaculation is based either on increasing sympathetic tone at the bladder neck or on decreasing the parasympathetic activity. One example is a selective and long-acting α -adrenergic stimulator, imipramine hydrochloride, which produced antegrade ejaculation for the patients with retrograde ejaculation (22) and successful pregnancies were achieved (3,23). However, results of the administration of imipramine hydrochloride to this patient were unsuccessful. Then, the patient was treated with IUI using the sperm collected from the bladder, since the sperm quality, including motile function and fertilizing ability, was fairly good (as shown in Table I). There are several methods to recover the sperm from the bladder in the patients with retrograde ejaculation. First, is the invasive method in which the urine is replaced with a few milliliters of isotonic buffer solution by catheterization before masturbation. After the patient masturbates, the catheter is reintroduced and the specimen is withdrawn, and his spouse is inseminated with the sperm obtained (24). The second technique is noninvasive, compared with the Hotchkiss's method, in which the patient is given sodium bicarbonate to adjust the urine pH and osmolarity prior to sperm recovery (25). In this case, the former method was simplified and used because the patient was familiar with self-catheterization. A

successful pregnancy was achieved in the fourth IUI treatment cycle. Although the patient was repeatedly treated, another pregnancy was not established by the IUI treatment because repeated urethral infection caused inflammatory obstruction at the level of the epididymis.

For the treatment of the patients with retrograde ejaculation, the assisted reproductive technologies including IVF (3,26–28), gamete intrafallopian tubal transfer (GIFT) (29), pronuclear stage tubal transfer (PROST) (30), and ICSI (31,32) were introduced. In this case, obstructive azoospermia was diagnosed before proceeding to use these treatments using the sperm recovered from the bladder. An ICSI treatment using the sperm recovered by testicular sperm extraction (TESE) was applied and a pregnancy was achieved at the first attempt of the TESE–ICSI treatment. Thus, some of the patients with retrograde ejaculation preserving normal sperm motility and fertilizing ability can be treated with IUI using the sperm collected from the bladder. However, IVF or ICSI treatment to achieve a pregnancy in the cases of retrograde ejaculation when the patients have poor sperm motility and fertilizing ability should be considered. Moreover, the patients with retrograde ejaculation can be aided by ICSI using the sperm retrieved from the epididymis or the testis when severe urethral infection leads obstructive azoospermia.

ACKNOWLEDGMENTS

We thank Akiko Hasegawa, PhD, and Yukari Hamada, BS, for their technical assistance.

REFERENCES

1. Dubin L, Amelar RD: Etiologic factors in 1294 consecutive cases of male infertility. *Fertil Steril* 1971;22:469–474
2. Yavetz H, Yogev L, Hauser R, Lessing JB, Paz G, Homonnai ZT: Retrograde ejaculation. *Hum Reprod* 1994;9:381–386
3. Okada H, Fujioka H, Tatsumi N, Kanzaki M, Inaba Y, Fujisawa M, Gohji K, Arakawa S, Kamidono A: Treatment of patients with retrograde ejaculation in the era of modern assisted reproductive technology. *J Urol* 1998;159:848–850
4. Sandler B: Idiopathic retrograde ejaculation. *Fertil Steril* 1979;32:474–475
5. Keisermann WM, Dubin L, Amelar RD: A new type of retrograde ejaculation: Report of three cases. *Fertil Steril* 1974;25:1071–1072
6. Thomas AJ: Ejaculatory dysfunction. *Fertil Steril* 1983;39:445–454
7. Burkman LJ, Kruger TF, Coddington CC: The hemizona assay (HZA): Development of a diagnostic test for the binding of

- human spermatozoa to human hemizona pellucida to predict fertilization potential. *Fertil Steril* 1988;49:688–697
8. Jeyendran RS, Van der Ven HH, Perez-Pelaez M, Crabo BG, Zaneveld LJ: Development of an assay to assess the functional integrity of the human sperm membrane and its relationship to the other semen characteristics. *J Reprod Fertil* 1984;70:219–228
 9. Shutte B: Penetration ability of human spermatozoa into standardized bovine cervical mucus (Penetrak) in patients with normal and pathological semen samples. *Andrologia* 1987;19:217–224
 10. World Health Organization: WHO laboratory manual for the examination of human semen and sperm-cervical mucus interaction, 3rd ed. Cambridge University Press, Cambridge, England, 1992
 11. Hershlag A, Schiff SF, DeCherney AH: Clinical review: retrograde ejaculation. *Hum Reprod* 1991;6:255–258
 12. Shibahara H, Mitsuo M, Ikeda Y, Shigeta M, Koyama K: Effects of sperm immobilizing antibodies on pregnancy outcome in infertile women treated with IVF-ET. *Am J Reprod Immunol* 1996;36:96–100
 13. Shibahara H, Funabiki M, Shiotani T, Ikeda Y, Koyama K: A case of primary ovarian pregnancy after in-vitro fertilization and embryo transfer. *J Assist Reprod Gen* 1997;14:63–64
 14. Shibahara H, Naito S, Hasegawa A, Mitsuo M, Shigeta M, Koyama K: Evaluation of sperm fertilizing ability using the Sperm Quality Analyzer. *Int J Androl* 1997;20:112–117
 15. Shibahara H, Mitsuo M, Inoue M, Hasegawa A, Shigeta M, Koyama K: Relationship between human in-vitro fertilization and intracytoplasmic sperm injection and the zona-free hamster egg penetration test. *Hum Reprod* 1998;13:1928–1932
 16. Shibahara H, Hamada Y, Hasegawa A, Wakimoto E, Toji H, Shigeta M, Koyama K: Relationship between sperm motility index assessed by the Sperm Quality Analyzer and the outcome of intracytoplasmic sperm injection. *J Assist Reprod Gen* 1999; in press
 17. Shibahara H, Hamada Y, Hasegawa A, Toji H, Shigeta M, Yoshimoto T, Shima H, Koyama K: Correlation between the motility of frozen-thawed epididymal spermatozoa and the outcome of intracytoplasmic sperm injection. *Int J Androl* 1999; in press
 18. Anderson FM: Occult spinal dysraphism: a series of 73 cases. *Pediatrics* 1975;55:826–835
 19. Hoffman HJ, Hendrick EB, Humphreys RP: The tethered spinal cord: Its protean manifestations, diagnosis and surgical correction. *Childs Brain* 1976;2:145–155
 20. Satar N, Bauer SB, Shefner J, Kelly MD, Durbey MM: The effects of delayed diagnosis and treatments in patients with an occult spinal dysraphism. *J Urol* 1995;154:754–758
 21. Ranieri DM: Is IVF/ICSI always the first step to treat couples with infertility due to retrograde ejaculation? *Fertil Steril* 1998;69:1160–1161
 22. Brooks ME, Berzin M, Braf Z: Treatment of retrograde ejaculation with imipramine. *Urology* 1980;15:353–355
 23. Eppel SM, Berzin M: Pregnancy following treatment of retrograde ejaculation with clomipramine hydrochloride. *S Afr Med J* 1984;66:889–891
 24. Hotchkiss R, Pinto A, Kleegman S: Artificial insemination with semen recovered from the bladder. *Fertil Steril* 1955;6:37–40
 25. Urry RL, Middleton RG, McGavin S: A simple and effective technique for increasing pregnancy rate in couples with retrograde ejaculation. *Fertil Steril* 1986;46:1124–1127
 26. Bosman E, Fourie FLR, van der Merwe JV: Successful pregnancy despite retrograde ejaculation. A case report. *S Afr Med J* 1990;77:368–369
 27. Hirsch AV, Mills C, Tan SL, Bekir J, Rainsbury P: Case report: Pregnancy using spermatozoa aspirated from the vas deferens in a patient with ejaculatory failure due to spinal injury. *Hum Reprod* 1993;8:89–90
 28. Tay JJ, Whitehead A, Joyce AD, Rutherford AJ: Twin pregnancy following IVF treatment using frozen-thawed retrograde ejaculated sperm. *J Assist Reprod Gen* 1996;13:731–732
 29. Vernon M, Wilson E, Muse K, Estes S, Curry T: Successful pregnancies from men with retrograde ejaculation with the use of washed sperm and gamete intrafallopian tube transfer (GIFT). *Fertil Steril* 1988;50:822–824
 30. Hulme VA, Stander FSH, Kruger TF, van der Merwe JP, Windt ML: Retrograde ejaculation and pronuclear stage tubal transfer. *Arch Androl* 1991;26:21–24
 31. Gerris J, Van Royen E, Mangel-Schots K, Joostens M, de Vits A: Pregnancy after intracytoplasmic sperm injection of metaphase II oocytes with spermatozoa from a man with complete retrograde ejaculation. *Hum Reprod* 1994;9:1293–1296
 32. Jimenez C, Grizard G, Pouly J-L, Boucher D: Birth after combination of cryopreservation of sperm recovered from urine and intracytoplasmic sperm injection in a case of complete retrograde ejaculation. *Fertil Steril* 1997;68:542–544

Hiroaki Shibahara
Hideaki Toji
Minoru Shigeta
Testuro Yoshimoto
Hiroki Shima
Koji Koyama¹

Department of Obstetrics and Gynecology, and
 Department of Urology
 Hyogo College of Medicine, 1-1
 Mukogawa-cho, Nishinomiya
 Hyogo 663-8501, Japan
 and

Department of Obstetrics and Gynecology
 Jichi Medical School
 3311-1 Yakushiji, Minamikawachi-machi, Kawachi-gun
 Tochigi 329-0498, Japan

¹ To whom correspondence should be addressed.