Published in final edited form as:

Neurobiol Dis. 2013 September ; 0: 28–37. doi:10.1016/j.nbd.2012.05.022.

Environmental and pharmacological modulation of cellular plasticity: role in the pathophysiology and treatment of depression

Kristie T. Ota and Ronald S. Duman

Division of Molecular Psychiatry, Abraham Ribicoff Research Facilities, Connecticut Mental Health Center, Yale University School of Medicine, New Haven, Connecticut 06508

Abstract

Atrophy of neurons and gross structural alterations of limbic brain regions, including the prefrontal cortex (PFC) and hippocampus, have been reported in brain imaging and postmortem studies of depressed patients. Preclinical findings have suggested that prolonged negative stress can induce changes comparable to those seen in major depressive disorder (MDD), through dendritic retraction and decreased spine density in PFC and hippocampal CA3 pyramidal neurons. Interestingly, recent studies have suggested that environmental and pharmacological manipulations, including antidepressant medication, exercise, and diet, can block or even reverse many of the molecular changes induced by stress, providing a clear link between these factors and susceptibility to MDD. In this review, we will discuss the environmental and pharmacological factors, as well as the contribution of genetic polymorphisms, involved in the regulation of neuronal morphology and plasticity in MDD and preclinical stress models. In particular, we will highlight the pro-depressive changes incurred by stress and the reversal of these changes by antidepressants, exercise, and diet.

Keywords

Major depressive disorder; Stress; Animal models; Antidepressants; Exercise; Omega-3 fatty acids; Early life stress

Introduction

Major depressive disorder (MDD) is a recurring psychiatric illness that affects up to 17% of the American population, and can be severely debilitating, affecting sleep, work, social relationships, and appetite (Kessler et al., 2005). Both clinical and pre-clinical studies have revealed that prolonged stress can lead to depressive symptoms that are associated with atrophy of neurons and related behavioral alterations in limbic brain areas, most notably the prefrontal cortex (PFC) and hippocampus. For example, postmortem findings show cell atrophy and decreased volume of the PFC in MDD patients (Drevets et al., 1997; Rajkowska et al., 1999). Further, a recent meta-analysis of imaging studies revealed that lower gray matter volumes are seen in cortical limbic structures (dorsal frontomedial and paracingulate

© 2012 Elsevier Inc. All rights reserved.

Correspondence: Ronald S. Duman, 34 Park Street, New Haven, CT 06508. ronald.duman@yale.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

cortex) of MDD patients, in parallel with increases in glucose metabolism in related cortical regions (subgenual and pregenual anterior cingulate) (Sacher et al., 2011). Consistent with this, morphological and morphometric analyses of the hippocampus in depressed patients show reduced volume, gray matter, and neuropil (Bremner et al., 2000; Campbell et al., 2004; McKinnon et al., 2009; Sheline et al., 1996; Stockmeier et al., 2004). The amygdala is another limbic structure that undergoes structural alterations, although in this case amygdala neurons undergo a hypertrophy in response to repeated stress that could contribute to anxiety and certain emotional symptoms of depression (see Roozendaal et al. (2009) for review of the role of the amygdala in stress).

Understanding the mechanisms by which these structural and morphological changes arise could provide insight into the biological basis of MDD. Importantly, recent work has shown that these changes are not permanent and that these structural alterations can be altered bi-directionally through environmental and pharmacological influences. Here, we provide an overview of the morphological changes in rodent models of stress and MDD, and the influence of environmental, pharmacological, and genetic factors. This work provides a framework for understanding the impact of lifetime exposure to environmental factors in the context of genetic susceptibility or resilience to depression and will lead to updated recommendations for healthy emotional preventative lifestyles and new therapeutic interventions.

Structural and Neurotrophic Deficits Caused by Chronic Stress and Depression

Stress, defined as an alteration of normal homeostasis (Selye, 1936), is a strong environmental factor that can have a wide-range of effects on multiple organ systems, including depression. Consistent with this, chronic stress is commonly used as an animal model of depression. Preclinical studies demonstrate that chronic stress causes dendritic retraction and decreased spine density, in PFC pyramidal neurons (layer II/III and layer V) and hippocampal CA3 pyramidal neurons following repeated stress (Izquierdo et al., 2006; McEwen, 1999; Radley et al., 2006; Radley et al., 2004; Shansky et al., 2009; Sousa et al., 2000). Reductions of spine density and length are observed at both the distal and proximal segments of the apical tuft, and can be measured by reduction in expression of synaptic proteins (Li et al., 2011). These cellular alterations may contribute to the reduced volume of PFC and hippocampus reported in brain imaging studies, and to the cognitive and emotional deficits observed in depressed patients (Drevets et al., 1997; McEwen, 2007; Sacher et al., 2011; Sheline et al., 2003).

Chronic stress is one of the better rodent models of depression, resulting in structural alterations as well as behavioral anhedonia, a core symptom of depression (Willner, 1997). Anhedonia and behavioral despair, typically assessed in the forced swim test, are also reversed by antidepressant treatment, providing a level of construct validity. However, it is difficult to interpret the meaning of behavioral despair in these models and to model other aspects of depression that are evaluated by verbal assessment in humans. In addition, the heterogeneous nature of MDD makes it difficult to accurately model all aspects of depression at behavioral, as well as molecular and cellular levels. Nevertheless, when used with an understanding of the limitations, the chronic stress models can provide powerful information on the underlying molecular and cellular determinants of depression and the circuitry controlling complex stress-related behaviors.

One mechanism that may mediate the cellular alterations observed following chronic stress is through altered expression of neurotrophic factors. Neurotrophic factors are important for growth and survival during neuronal development but continue to play a critical role in the

survival and function of neurons in the adult brain (McAllister, 2002). Here we discuss some of the factors best-characterized for their roles in stress and depression, including brain derived neurotrophic factor (BDNF), vascular endothelial growth factor (VEGF), VGF (non-acronym), and insulin-like growth factor (IGF)-1.

Stress Decreases Neurotrophic Factor Expression and Signaling

BDNF—Perhaps the most studied neurotrophic factor in stress and MDD is BDNF (Castrén et al., 2007; Duman and Monteggia, 2006; Monteggia, 2007). BDNF levels are decreased in cortical and hippocampal regions of postmortem brains of depressed subjects relative to controls (Chen et al., 2001; Dwivedi et al., 2003; Karege et al., 2005). Interestingly, levels of BDNF in blood correlate with that in brain tissue, and are decreased in depressed patients relative to controls (Bocchio-Chiavetto et al., 2010; Brunoni et al., 2008; Karege et al., 2005; Klein et al., 2011; Sen et al., 2008; Shimizu et al., 2003). However, BDNF is also expressed in many peripheral tissues, and the source of BDNF that contributes to altered levels in blood of MDD patients, as well as the underlying mechanisms, requires further investigation (Schmidt et al., 2011).

Further supporting a role of BDNF in MDD is evidence that this neurotrophic factor is decreased in animal models of chronic stress (Duman, 2004b; Murakami et al., 2005). Moreover, applications of recombinant BDNF (i.e., i.c.v., intra-hippocampus, or intra-midbrain) have antidepressant and anxiolytic effects in behavioral models. These effects include increased mobility in the forced swim test, increased escape in the learned helplessness test, increased sucrose consumption in the chronic unpredictable stress (CUS) paradigm, decreased latency in the novelty induced hypophagia test, and increased time spent in the open arms of an elevated plus maze (Schmidt and Duman, 2010; Shirayama et al., 2002; Siuciak et al., 1997).

In contrast, deletion of BDNF blocks the behavioral effects of antidepressants in several of these models, including the forced swim test and learned helplessness model (Duman and Monteggia, 2006; Monteggia et al., 2004). Consistent with this, constitutive heterozygous (+/−) deletion of BDNF (homozygous deletion mutants are lethal) results in neuronal atrophy in the hippocampus and PFC (Chen et al., 2006; Magariños et al., 2011), which may contribute to blockade of antidepressant responses. Interestingly, most studies have shown that the deletion of BDNF alone is not sufficient to cause depressive behavior in rodents, but instead increases susceptibility to depressive effects (Duman et al., 2007; Martinowich et al., 2007). A recent study, however, has shown that short hairpin RNA (shRNA) knockdown of BDNF in the hippocampus is sufficient to cause depressive behaviors, suggesting that perhaps BDNF plays a specific regional role in the etiology of depression (Taliaz et al., 2010). The discrepancy between this and previous approaches that do not target a specific brain region could be due to selective knockdown in the hippocampus, as BDNF in the mesolimbic dopamine circuit produces opposing depressive actions (Krishnan and Nestler, 2008). Taken together, these preclinical studies demonstrate a requirement for BDNF in the actions of antidepressants and implicate BDNF deficits as a risk factor that contributes to neuronal atrophy and is associated with depressive symptoms.

The expression of BDNF is induced by activation of the cyclic AMP (cAMP) and Ca²⁺/calmodulin (CAM) signaling cascades, activation of cAMP- and CAM-dependent protein kinases (PKA and CAMK, respectively), and the cAMP response element binding protein (CREB) (Greer and Greenberg, 2008; Nestler et al., 2002). BDNF also has numerous downstream molecular effectors (Figure 1). BDNF activates TrkB, which can stimulate the mitogen-activated protein kinase (MAPK) and phosphatidylinositol 3-kinase (PI3K)-Akt signaling pathways. Based on the significant role of BDNF in depression and treatment response, it is not surprising that MAPK signaling is similarly required for the behavioral

effects of antidepressant action (Duman et al., 2007). Consistent with this, recent studies have shown that MAPK signaling is decreased in postmortem brains of depressed subjects and that MAPK phosphatase-1 (MKP-1), which negatively regulates the MAPK signaling pathway, is increased in the hippocampus of depressed subjects (Duric et al., 2010; Dwivedi et al., 2006; Dwivedi et al., 2007). Induction of MKP-1 using a viral expression approach produces stress-like behaviors in the absence of stress in rodents, while MKP-1 deletion creates a state resilience to chronic stress exposure (Duric et al., 2010). Taken together, these studies elucidate some of the downstream signaling mechanisms by which BDNF can underlie depression behaviors, although additional studies are needed to further elucidate the complex signaling pathways that contribute to the actions of BDNF.

Interestingly, a human functional genetic variant of BDNF, the Val66Met single nucleotide polymorphism (SNP), has provided further evidence for a role for BDNF in MDD. The BDNF Met polymorphism results in decreased activity-dependent release of BDNF by decreasing the processing and targeting of mature BDNF to dendrites and synapses (Chen et al., 2004; Chiaruttini et al., 2009; Egan et al., 2003). Carriers of the BDNF Met allele have decreased hippocampal volume and display cognitive deficits compared to the Val allele (Bueller et al., 2006; Egan et al., 2003; Hariri et al., 2003; Pezawas et al., 2004; Szeszko et al., 2005). Although there is no direct association of the BDNF Met allele with depression (Verhagen et al., 2010), there are reports that early life stress or trauma increase the vulnerability to depressive phenotypes in BDNF Met carriers (Gatt et al., 2009) (Figure 2).

Studies of the Met allele in rodents provide similar evidence for structural and functional alterations. Mice with a knock in of the BDNF Met allele exhibit atrophy of pyramidal neurons in the hippocampus and PFC (Chen et al., 2006; Liu et al., 2011). BDNF Met knock in mice also display decreased density of spines, decreased number of large diameter “mushroom” or mature spines, and a reduction in excitatory postsynaptic currents in the PFC (Liu et al., 2011). Similar reductions in dendrite length and branching have been observed in constitutive BDNF heterozygous mutant mice that express one half the normal complement of BDNF (Chen et al., 2006; Magariños et al., 2011). These effects of the BDNF Met allele, as well as constitutive BDNF heterozygous deletion mutant mice, are reminiscent of the atrophy of neurons in the PFC and hippocampus caused by chronic stress.

VEGF—Although VEGF was first characterized in and is best known for its effects on endothelial cell proliferation and angiogenesis, it is also expressed in brain and influences neurite outgrowth and maturation during development (Fournier and Duman, 2011). Studies in humans suggest that a VEGF polymorphism, VEGF2578C/A, which leads to lower circulating levels of VEGF (Lambrechts et al., 2003), is linked to treatment-resistant depression (Viikki et al., 2010). In parallel, CUS in rodents decreases expression of VEGF in the granule cell layer and hilar regions of the hippocampus, providing evidence that disruption of this factor could also contribute to the structural and behavioral deficits caused by stress (Bergström et al., 2008; Heine et al., 2005).

Stress and Neurogenesis

One of the functional consequences of alterations in neurotrophic factors is regulation of neurogenesis. Neurogenesis occurs during early postnatal development, but also continues in certain neurogenic zones in the adult, notably the dentate gyrus of hippocampus and subventricular zone. Different types of stress, including CUS, intruder stress, predator stress, and footshock stress, all decrease the number of newborn neurons in the granule cell layer of the adult hippocampus (Drew and Hen, 2007; Duman, 2004a; McEwen, 1999). Interestingly, the loss of new neurons is not sufficient to produce depressive behaviors (Drew and Hen, 2007; Sahay and Hen, 2007), but a recent study demonstrates that new neurons are

necessary for proper control of endocrine and behavioral responses to stress (Snyder et al., 2011). Consistent with the effects of stress on BDNF and its role in depression, there is strong evidence that BDNF also contributes to the regulation of neurogenesis. For example, central or peripheral BDNF administration increases adult hippocampal neurogenesis (Schmidt and Duman, 2010). In contrast, conditional deletion of TrkB in hippocampal neural progenitor cells leads to impaired proliferation and neurogenesis (Li et al., 2008). Similarly, BDNF +/- mice have decreased basal proliferation (Lee et al., 2002a), and both BDNF +/- and dominant negative TrkB over expressing mice show impaired long-term survival of newborn neurons (Sairanen et al., 2005).

Taken together, these findings demonstrate that chronic stress serves as a valid model for depression-related alterations: many of the same structural and cellular changes are observed in depression and chronic stress, and there is evidence that the same trophic factors and signaling pathways may be involved in humans and animal models. Although we have focused on the role of neurotrophic factors, there are a number of other factors and systems that have also been implicated in depression that will not be discussed due to space limitations. This includes recent work demonstrating alterations of the GABA neurotransmitter system in depressed patients (see Sanacora (2010) for review).

Early Life Stress and Maternal Care

In addition to stress in adulthood, early postnatal or adolescent stress, as well as maternal care, can have major consequences on neuronal function and behavior in the adult. Studies have shown that individuals with a history of childhood abuse are more likely to develop depression or other stress-related illnesses (e.g., PTSD) in adulthood (Felitti et al., 1998; Heim and Binder, 2011; Mullen et al., 1996; Wilkinson and Goodyer, 2011). In support of this, preclinical studies have shown that exposure to early life stress, such as prolonged daily maternal separation, are associated with increased emotionality and stress-responsiveness (Heim and Nemeroff, 1999; Meaney, 2001). At the neural level, these effects are associated with alterations in the activity of the HPA axis. Maternal separation stress is associated with decreases in the levels of glucocorticoid receptors, which bind the stress hormone corticosterone and provides negative feedback on further activity of the HPA axis. These changes persist into adulthood and are associated with increased corticosterone release and changes in emotional behavior (Anisman et al., 1998).

BDNF expression can also be altered by early life stress or variations on maternal care (Figure 2). Maternal separation (24 hr) on postnatal day 9 in rats can lead to reduced BDNF (Roceri et al., 2004), while high levels of maternal care, measured by licking and grooming behavior, increase hippocampal BDNF (Liu et al., 2000). These alterations in BDNF expression can ultimately affect synaptic development and behavior in the adult.

Further, studies have found that cell proliferation and number of immature neurons are decreased in the dentate gyrus of adult rats that have undergone maternal separation as pups (Mirescu et al., 2004). This effect can be localized to the ventral, but not dorsal, hippocampus, the former being associated with circuitry underlying anxiety (Hulshof et al., 2011). Moreover, the reduction in cell proliferation can be reversed by decreasing corticosterone levels, supporting the hypothesis that early life stress causes hypersensitivity to glucocorticoids and decreases structural plasticity in adults (Mirescu et al., 2004). Animals subjected to maternal separation also show increased anxiety- and depression-like behaviors in adulthood (George et al., 2010; Lee et al., 2007; Wei et al., 2010). As an example, rat pups that have been subjected to prolonged maternal separation exhibit altered emotional responses, including an unwillingness to explore an open field, feed in a novel environment, or interact socially (Caldji et al., 2000).

Interestingly, recent studies have identified genetic polymorphisms that can increase the susceptibility to depression in the human population. This helps to account for the fact that although child abuse is a major risk factor for the development of depression during the lifetime, not all abused children become depressed (Charney and Manji, 2004; Heim and Nemeroff, 2001). For example, in a cross-sectional study of individuals with lifetime MDD, patients that have the BDNF Val66Met polymorphism show a greater susceptibility to depression later in life following childhood abuse, compared to those with intact BDNF release (Elzinga et al., 2011) (Figure 2).

Another genetic risk factor is the serotonin transporter gene, 5-HTTLPR, polymorphism. This polymorphism confers susceptibility to depression in individuals subjected to early life stress (Caspi et al., 2003; Eley et al., 2004; Kaufman et al., 2004; Kendler, 2005; Risch et al., 2009; Uher et al., 2011). In addition, there is a three-way interaction of the 5-HTTLPR and the BDNF Val66Met polymorphism with childhood adversity (Grabe et al., 2012; Kaufman et al., 2006; Nederhof et al., 2010; Wichers et al., 2008), suggesting that the BDNF and 5-HT systems work in concert to influence susceptibility to mood disorders. Possible interactions of these systems have been demonstrated in pharmacological studies, showing that BDNF expression is up-regulated by chronic administration of a selective serotonin reuptake inhibitor (SSRI) (Chen et al., 2001).

Collectively, these findings suggest that exposure to early life stress can lead to long-term effects on neuroplasticity in neural circuits that control emotion, which, in turn, can have a profound influence on susceptibility to depression in adulthood.

Antidepressants, Neurotrophic Factors, and Neurogenesis

Pharmacological antidepressants, including SSRI and norepinephrine selective reuptake inhibitors, monoamine oxidase inhibitors, and atypical antidepressants, have many effects that are the opposite to those of stress and depression that have been discussed thus far. Clinical studies demonstrate that levels of BDNF and new cell birth are increased in the hippocampus of those patients treated with antidepressants at the time of death (Boldrini et al., 2009; Chen et al., 2001). Similarly, serum BDNF levels (Gervasoni et al., 2005) and hippocampal volume (Sheline et al., 2003; Vermetten et al., 2003) are partially restored in depressed patients receiving antidepressant treatment.

These findings have been corroborated in preclinical studies. Antidepressant administration in rats increases the expression of BDNF in the hippocampus and PFC (Baj et al., 2012; Duman, 2004b; Krishnan and Nestler, 2008; Monteggia, 2007; Nibuya et al., 1995; Nibuya et al., 1996) and increases neurogenesis in the adult hippocampus (Duman, 2004b; Duman and Monteggia, 2006; Samuels and Hen, 2011). Up-regulation of BDNF, as well as induction of neurogenesis, is seen after chronic, but not acute, antidepressant administration, consistent with the time course for the therapeutic action of these agents (Duman, 2005).

In addition, VEGF and IGF- 1, growth factors known to increase neurogenesis in the adult hippocampus (Duman, 2004b; Sairanen et al., 2005), produce antidepressant effects in behavioral models when infused into the brain (Duman et al., 2009; Warner-Schmidt and Duman, 2007). In contrast, blockade of VEGF signaling prevents the behavioral actions of antidepressants (Duman et al., 2008; Warner-Schmidt and Duman, 2007). Blockade of neurogenesis blocks the behavioral actions of antidepressants in certain behavioral models (David et al., 2009; Santarelli et al., 2003), suggesting that increased neurogenesis could contribute to the antidepressant effects of trophic factors. Further, deletion of TrkB blocks antidepressant induction of neurogenesis (Li et al., 2008), supporting a role for BDNF or a related neurotrophic factor in these actions.

The requirement for chronic administration of traditional antidepressants suggests that alterations of intracellular signaling pathways and long-term gene expression may underlie the mechanisms of antidepressant action. Chronic, but not acute, administration of both 5-HT and norepinephrine selective reuptake inhibitors increases cAMP-CREB signaling, which subsequently increases the expression of BDNF (Duman and Monteggia, 2006; Nibuya et al., 1996). In addition, MAPK signaling is required for the behavioral actions of traditional antidepressants, whereby blockade of MAPK signaling prevents behavioral antidepressant action (Duman et al., 2007). The glycogen synthase kinase-3 (GSK-3) signaling pathway, which leads to activation of transcription factors promoting cell cycle arrest and apoptosis, has also been linked to mood disorders and is inhibited by the mood stabilizer, lithium, suggesting its role in the regulation of depressive behaviors (Voleti and Duman, 2011). This is just a partial list of the signaling pathways that have been implicated in the actions of traditional antidepressant treatments (Krishnan and Nestler, 2008; Martinowich et al., 2007; Voleti and Duman, 2011).

Rapid Acting Antidepressant: NMDA Receptor Antagonists

Traditional antidepressants take 6–8 weeks to produce a therapeutic response, but recent clinical studies have reported that ketamine, an NMDA receptor antagonist, produces antidepressant effects within hours (Berman et al., 2000; Liebrecht et al., 2007; Zarate et al., 2006). Further, the rapid antidepressant actions of ketamine are observed in treatment-resistant patients (i.e., failed two or more typical antidepressants), addressing another major unmet need (improved therapeutic efficacy). These findings represent a major advance for the treatment of depression, identifying a rapid, efficacious therapeutic agent with a completely novel mechanism of action.

Preclinical studies demonstrate that ketamine produces a rapid induction of synaptic protein synthesis, spine density and maturation (Li et al., 2010). These effects are mediated by activation of the mammalian target of rapamycin (mTOR) signaling system (Li et al., 2010), which controls synaptic protein synthesis in dendrites (Hoeffer and Klann, 2010). The rapid synaptic and behavioral actions of ketamine are blocked by rapamycin, a selective inhibitor of mTOR signaling (Koike et al., 2011; Li et al., 2010). In addition, a single dose of ketamine can reverse the reduction in spine density and function caused by chronic stress exposure in a rapamycin-sensitive manner (Li et al., 2011). A possible role for mTOR in depression is supported by a recent postmortem study reports that mTOR signaling is decreased in the PFC of depressed subjects (Jernigan et al., 2011).

The induction of mTOR and synaptogenesis are thought to occur via increased glutamate-AMPA transmission leading to release of BDNF and related downstream signaling (Hoeffer and Klann, 2010; Jourdi et al., 2009; Slipczuk et al., 2009; Takei et al., 2004) (Figure 3). Notably, the synaptic and behavioral actions of ketamine are blocked in BDNF conditional deletion mutant mice (Autry et al., 2011) and in BDNF Met knock in mice (Liu et al., 2011). Although Autry et al. (2011) did not observe a role for mTOR, this may be explained by methodological differences (behavior in the forced swim test was examined 30 min after ketamine instead of 2 hrs or longer, when the therapeutic response is observed in depressed patients, see Zarate et al., (2006). The 30 min time point is also problematic as this is when levels of glutamate are increased and the psychotomimetic effects of ketamine are observed (Berman et al., 2000; Moghaddam et al., 1997; Zarate et al., 2006).

Additional evidence has suggested that GSK-3 may also play a role in mediating the effects of ketamine. Specifically, Beurel et al. (2011) found that ketamine increases the phosphorylation of GSK-3, and that blockade of this phosphorylation prevents the action of ketamine. Similar to the activation of mTOR signaling, induction of phospho-GSK-3 may be

mediated by Akt, which is a negative regulator of GSK-3 and is activated by ketamine (Beurel et al., 2011; Duman et al., 2011).

Due to the abuse potential of ketamine, recent work has begun to explore other targets that may have fast-acting antidepressant actions similar to ketamine. For example, inhibition of presynaptic metabotropic glutamate receptor 2/3 (mGluR2/3) or activation of metabotropic glutamate receptor 7 (mGluR7), both of which regulate glutamate release and subsequently the postsynaptic AMPA receptors, can similarly produce antidepressant effects (Bradley et al., 2012; Dwyer et al., 2011; Koike et al., 2011). Moreover, the effects of mGluR2/3 antagonists are also dependent on mTOR signaling, providing further evidence of the potential to produce rapid antidepressant effects (Dwyer et al., 2011; Koike et al., 2011).

Studies of the actions of typical and novel rapid acting antidepressants have provided insight into the cellular and molecular mechanisms underlying these agents and to depression, and further validates the chronic stress model. These studies also demonstrate how typical antidepressant medications provide a pharmacological approach to alleviate depressive symptoms, albeit with limitations, some of which may be addressed with novel ketamine-like agents.

Exercise, Neurotrophic Factors, and Neurogenesis

The stress effects discussed thus far have focused on uncontrollable stressors, be it chronic stress during adulthood or maternal separation during early life, which are perceived as negative or aversive. However, it has become increasingly clear that different types of stress result in different outcomes. In addition to these negative stressors, recent studies have examined the influence of controllable stress. Here, we define stress as “controllable” if the person or animal is able to terminate the stressor at any given time. The differences between uncontrollable and controllable stress have been described for decades, starting with the learned helplessness experiments. In this paradigm, one rat can behaviorally terminate tail shocks (controllable stress) while another yoked rat can not (uncontrollable stress), with each rat receiving identical shocks (Seligman and Beagley, 1975). In rats that received uncontrollable, but not controllable stress, impaired escape learning and exaggerated fear conditioning are observed (Maier and Watkins, 2005).

One particular type of controllable stress is voluntary exercise. Voluntary exercise is considered to be a type of stress due to activation of the sympathetic nervous system, leading to epinephrine production and regulation of hypothalamic-pituitary-adrenal (HPA) axis and increased glucocorticoid production (Droste et al., 2007; Makatsori et al., 2003). Interestingly, exercise exerts downstream effects that are in the opposite direction from that of the uncontrollable stressors described above. For example, in clinical studies, exercise has been found to be beneficial for reducing depressive symptoms in mild to moderately depressed patients (Babyak et al., 2000; Dunn et al., 2005; Herring et al., 2012).

Preclinical studies have extended this research to further elucidate the molecular and cellular mechanisms underlying the effect of exercise on depression. Wheel running is most often used as a form of voluntary exercise in animal studies (Stranahan et al., 2008). Exercise is a positive reinforcer for rats: they will bar press for access to a running wheel and can even develop conditioned place preference to environments previously paired to a running wheel (Belke and Wagner, 2005; Iversen, 1993). Using the wheel-running exercise paradigm, preclinical studies have also reported an antidepressant effect of exercise. For example, rodents given chronic free access to a running wheel show antidepressant-like effects in learned helplessness, forced swim test, and tail suspension test relative to sedentary controls (Bjørnebekk et al., 2005; Dishman et al., 1997; Duman et al., 2008; Greenwood et al., 2005; Greenwood et al., 2003).

Exercise may alter the same neurotrophic factors and signaling pathways as antidepressant drug treatment. Chronic exercise increases BDNF expression and phosphorylated MAPK in the hippocampus and cerebral cortex (Adlard and Cotman, 2004; Chen and Russo-Neustadt, 2009; Duman et al., 2008; Russo-Neustadt et al., 1999; Russo-Neustadt et al., 2000), similar to the actions of antidepressants (Baj et al., 2012). Further, the antidepressant effects of exercise in the forced swim test are abolished in mice heterozygous for BDNF deletion (Duman et al., 2008). Consistent with this, blockade of MAPK using PD184161 prevents the antidepressant actions of exercise following forced swim test in wild-type mice (Duman et al., 2008). The antidepressant effects of exercise may depend on post-translational processing of BDNF. Specifically, Sartori et al. (2011) found that voluntary exercise on a running wheel increases levels of mature BDNF protein levels and increases levels of mRNA of the proBDNF proteolytic cleavage-related genes p11 and tPA, with no differences in precursor or truncated forms of BDNF.

Further, voluntary running can even *prevent* acute and chronic stress-induced alterations in depressive behavior and decreases in BDNF (Adlard and Cotman, 2004; Zheng et al., 2006). Exercise protects against decreases in hippocampal glucocorticoid receptors in response to CUS (Zheng et al., 2006) and prevents decreases in BDNF levels in immobilization stress-exposed animals (Adlard and Cotman, 2004). It is thought that these exercise-induced alterations in BDNF exert their actions, at least in part, through increased neurogenesis (Erickson et al., 2011; Lafenêtre et al., 2010), as exercise increases neurogenesis in the granule cell layer of the adult hippocampus (van Praag et al., 1999).

Exercise also up regulates other neurotrophic factors that are increased by antidepressant treatment and have been shown to have antidepressant effects in rodent models, including IGF-1 (Trejo et al., 2001). IGF-1 uptake in the hippocampus is stimulated by exercise (Trejo et al., 2001), and peripheral blockade of IGF-I blocks the exercise-induced antidepressant effects on both hippocampal neurogenesis and in the forced swim test. These data suggest that IGF-I is required for the antidepressant effects of exercise (Duman et al., 2009; Trejo et al., 2001; Trejo et al., 2008). There is evidence that IGF-1 differentially regulates neurogenesis depending on the differentiation state of new neurons, suggesting that the response of proliferating precursors and post mitotic immature neurons to exercise is dependent on IGF-1 (Llorens-Martín et al., 2010).

VEGF is also thought to play a role in the mediation of the antidepressant effects of exercise. Blockade of peripheral VEGF prevents exercise-induced antidepressant effects, including behavioral and neurogenic responses, as well as alterations in blood vessel density in the hippocampus (Fabel et al., 2003; Kiuchi et al., 2012). Early studies were unable to find alterations in VEGF expression in the hippocampus following running in mice, concluding that VEGF was induced in the peripheral vasculature and transported into the brain where it produces central effects, including increased hippocampal neurogenesis (Fabel et al., 2003). However, a more recent study using a VEGF-luciferase reporter mouse found that exercise increases VEGF transcription, mRNA and protein levels in the hippocampus (Tang et al., 2010).

VGF (nonacronym) is a nerve growth factor that is regulated in the hippocampus by exercise at both the mRNA and protein levels (Hunsberger et al., 2007). Originally studied for its roles in energy metabolism (Hahm et al., 1999) and synaptic plasticity (Alder et al., 2003), a recent finding that VGF is also involved in mood regulation has opened up a new avenue of research on this peptide. Administration of a VGF-derived peptide into the brain produces antidepressant effects in mice, and heterozygous VGF deletion mutant mice do not show the exercise-induced antidepressant responses that are observed in wild-type mice (Hunsberger et al., 2007). VGF also promotes proliferation of hippocampal neurons (Thakker-Varia et al.,

2007), providing a potential cellular mechanism for the antidepressant-like effects of VGF and exercise. Taken together, these data suggest that VGF also contributes to the antidepressant actions of exercise.

Recent data have also suggested a role for endocannabinoids in mediating the antidepressant effects of exercise (Gorzalka and Hill, 2011; Sparling et al., 2003). In humans, acute exercise increases plasma endocannabinoid levels (Sparling et al., 2003), and preclinical studies report that chronic exercise increases levels of endocannabinoid and the cannabinoid 1 receptor in the hippocampus (Hill et al., 2010; Wolf et al., 2010). Endocannabinoid signaling can also modulate exercise-induced neurogenesis (Hill et al., 2010; Wolf et al., 2010), demonstrating a downstream cellular mechanism that could underlie the antidepressant actions of this interesting neuronal signaling system.

Exercise also influences synaptic plasticity and structure. Long-term potentiation (LTP) is increased in the dentate gyrus both *in vitro* and *in vivo* in rodents given access to running (Farmer et al., 2004; van Praag et al., 1999). Further, exercise increases spine density in hippocampal area CA1, the dentate gyrus, and entorhinal cortex layer III (Redila and Christie, 2006; Stranahan et al., 2007), and accelerates neuronal maturation (Zhao et al., 2006).

Taken together, these results suggest that functional antidepressant responses that result from exercise occur via a combination of neurotrophic/growth factor systems and related signaling pathway alterations, and that subsequent increases in hippocampal neurogenesis and alterations in synaptic plasticity and structure may underlie these effects.

Diet, Neurotrophic and Neurogenic responses

Dietary Restriction

Dietary restriction was first discovered to increase lifespan in a wide range of organisms (Roth et al., 2000; Weindruch et al., 1979), and since then has been studied for other beneficial effects, including reduction in aging-associated illnesses and improved response to stress (Mattson, 2005; Mattson et al., 2004b). In the brain, caloric restriction has similarly beneficial effects, including increased BDNF expression and increased hippocampal neurogenesis via increased survival of newborn neurons (Duan et al., 2001; Lee et al., 2002b; Mattson et al., 2004a). However, these effects are dependent on age and duration of food restriction. One study found that adolescent rats that are food-restricted for 5 weeks starting at postnatal day 28 show significantly *increased* depressive and anxiety-like behaviors and aberrant serotonergic activity (Jahng et al., 2007). Similarly, another study has suggested that cyclic caloric restriction combined with periods of palatable foods (i.e. binge eating) results in aberrant monoamine activity in limbic regions and increases levels of pro-depressive and anxiety behaviors (Chandler-Laney et al., 2007). Thus, dietary restriction can play both a positive and negative role in depressive behaviors, depending on the developmental stage and context of restriction, possibly through alterations of BDNF and other signaling pathways.

Omega-3 Polyunsaturated Fatty Acids

Omega-3 polyunsaturated fatty acids are the most widely used non-vitamin supplement in the United States (Barnes et al., 2008), and recent epidemiological and clinical studies have suggested that they may have antidepressant actions. In the nervous system, one out of every three fatty acids belongs to the omega-3 fatty acids group, which make up an essential component of the CNS membrane phospholipid-acyl chains (Bourre et al., 1991). Omega-3 fatty acids are often ingested as docosahexaenoic acid (DHA) and eicosapentenoic acid (EPA), which are long-chain polyunsaturated fatty acids found in fish and marine sources.

Interest in the antidepressant actions of omega-3 fatty acids was first sparked when a cross-national study found that there was a correlation between fish consumption and lower annual prevalence of MDD (Hibbeln, 1998). Follow-up studies have found that depressed patients have significantly lower red blood cell omega-3 levels in erythrocyte membranes and plasma as compared to non-depressed patients (Freeman, 2006), and there appears to be a correlation between omega-3 fatty acid intake and corticolimbic brain matter volume (Conklin et al., 2007).

Although clinical studies have also suggested a role for omega-3 fatty acids in depression, results have been somewhat mixed (Grenyer et al., 2007; Hakkarainen et al., 2004; Lespérance et al., 2011; Marangell et al., 2003; Mischoulon and Fava, 2000; Nemets et al., 2002; Rees et al., 2008; Rocha Araujo et al., 2010; Su et al., 2003). Some of the variability may be attributable to methodological differences, including the type of omega-3 fatty acids and dosing schedule, as one meta-analysis found that EPA, but not DHA, may play a role in antidepressant action (Martins, 2009). Additionally, in most of the studies, the omega-3 fatty acids were used in conjunction with antidepressant treatment, and thus it remains unclear whether omega-3 fatty acids are effective independently. A recent study suggests that omega-3 fatty acid supplementation started concurrently with SSRI treatment can enhance the SSRI's effectiveness as compared to placebo supplement (Gertsik et al., 2012). However, it remains to be seen whether supplementation is effective in all or only in a subpopulation of patients with a low baseline level of omega-3 fatty acids, and if supplements have independent antidepressant effects.

Preclinical work has proposed several mechanisms by which omega-3 fatty acids may play an antidepressant role, many of which are similar to the effects of pharmacological antidepressants. Consistent with the clinical data, omega-3 fatty acid deprivation in rats for 15 weeks results in a pro-depressive phenotype, as measured by increased immobility time on the forced swim test (DeMar et al., 2006) and decreased BDNF levels in the frontal cortex (Rao et al., 2007). Further, CREB protein DNA-binding activity and nuclear phosphorylated MAPK levels are also reduced with omega-3 deprivation, possibly underlying the reduction of BDNF levels (Rao et al., 2007). Conversely, omega-3 fatty acid supplementation results in an antidepressant effect in forced swim test and increased BDNF levels in the cortex and hippocampus (Vines et al., 2011). Similarly, in rat primary cortical astrocytes, omega-3 supplementation increases BDNF levels (Rao et al., 2007). Other studies have shown that omega-3 fatty acid supplementation can actually reverse the effect of chronic restraint stress on plasma corticosterone levels (Ferraz et al., 2011).

At a cellular level, omega-3 fatty acids may oppose the actions of stress by increasing neurite growth and dendritic arborization. Both *in vitro* and *in vivo* studies have shown that low levels of DHA result in decreased neurite length and dendritic arborization (Calderon and Kim, 2004; Dagai et al., 2009), and that this effect can be rescued by DHA supplementation (Calderon and Kim, 2004). Further, administration of omega-3 fatty acids increases hippocampal volume, synaptophysin expression, and newborn cells in rodents, effects that are similar to the actions of antidepressants (Venna et al., 2009).

Although we have focused on the effects of omega-3 fatty acids on neurotrophic factor levels and dendrite arborization, another potential mechanism of antidepressant action is via regulation of proinflammatory cytokines, which are reported to be elevated in certain depressed patient populations (James and Cleland, 1997). Moreover, despite the evidence suggesting a role of omega-3 fatty acids in modulating depressive behaviors, it is important to keep in mind that these findings have been mixed, in both clinical and preclinical studies (Carlezon et al., 2005; Huang et al., 2008; Naliwaiko et al., 2004). Additional clinical and

preclinical studies are required to further elucidate the effects of this and other dietary supplements on cellular and behavioral determinants of stress and depression.

Summary and Conclusions

It has become increasingly clear that changes observed in MDD patients are the result of a number of environmental factors. Here, we have highlighted behavioral, cellular and morphological changes in models of MDD that are influenced by genetic and environmental factors, as well as pharmacological agents. Through the action of BDNF and other neurotrophic factors, genetic polymorphisms and stress (early and late life) can lead to neuronal atrophy and decreased neurogenesis, while maternal care, exercise, diet, and antidepressants can produce the opposite effects (Figure 2). Importantly, this evidence suggests that exercise and diet, two factors that can be controlled throughout life, can alter fundamental aspects of neuronal signaling and neurogenesis that are required for mental health. Although further studies will be necessary to obtain a comprehensive understanding of the mechanisms by which the environment can influence depression and mental health, the findings presented here demonstrate the importance of counteracting the inevitable exposure to stress in everyday life. One effective approach for improved mental health may include implementing policies that make recommendations to promote exercise and proper diet. Taken together, this review has presented an array of evidence demonstrating that environment can play an important role in altering neuroplasticity related to depressive behaviors, and suggests that understanding the complex interactions between stress, environment, and depression, can contribute to better treatments for MDD and other stress-related mood disorders.

References

- Adlard PA, Cotman CW. Voluntary exercise protects against stress-induced decreases in brain-derived neurotrophic factor protein expression. *Neuroscience*. 2004; 124:985–92. [PubMed: 15026138]
- Alder J, et al. Brain-derived neurotrophic factor-induced gene expression reveals novel actions of VGF in hippocampal synaptic plasticity. *J Neurosci*. 2003; 23:10800–8. [PubMed: 14645472]
- Anisman H, et al. Do early-life events permanently alter behavioral and hormonal responses to stressors? *Int J Dev Neurosci*. 1998; 16:149–64. [PubMed: 9785112]
- Autry AE, et al. NMDA receptor blockade at rest triggers rapid behavioural antidepressant responses. *Nature*. 2011:1–7.
- Babyak M, et al. Exercise treatment for major depression: maintenance of therapeutic benefit at 10 months. *Psychosom Med*. 2000; 62:633–8. [PubMed: 11020092]
- Baj G, et al. Physical Exercise and Antidepressants Enhance BDNF Targeting in Hippocampal CA3 Dendrites: Further Evidence of a Spatial Code for BDNF Splice Variants. *Neuropsychopharmacology* : official publication of the American College of Neuropsychopharmacology. 2012
- Barnes PM, et al. Complementary and alternative medicine use among adults and children: United States, 2007. *Natl Health Stat Report*. 2008:1–23. [PubMed: 19361005]
- Belke TW, Wagner JP. The reinforcing property and the rewarding aftereffect of wheel running in rats: a combination of two paradigms. *Behav Processes*. 2005; 68:165–72. [PubMed: 15686826]
- Bergström A, et al. Stress sensitivity and resilience in the chronic mild stress rat model of depression; an in situ hybridization study. *Brain Res*. 2008; 1196:41–52. [PubMed: 18234161]
- Berman RM, et al. Antidepressant effects of ketamine in depressed patients. *Biol Psychiatry*. 2000; 47:351–4. [PubMed: 10686270]
- Beurel E, et al. Inhibition of glycogen synthase kinase-3 is necessary for the rapid antidepressant effect of ketamine in mice. *Mol Psychiatry*. 2011; 16:1068–70. [PubMed: 21502951]
- Bjørnebekk A, et al. The antidepressant effect of running is associated with increased hippocampal cell proliferation. *Int J Neuropsychopharmacol*. 2005; 8:357–68. [PubMed: 15769301]

- Bocchio-Chiavetto L, et al. Serum and plasma BDNF levels in major depression: a replication study and meta-analyses. *World J Biol Psychiatry*. 2010; 11:763–73. [PubMed: 20334574]
- Boldrini M, et al. Antidepressants increase neural progenitor cells in the human hippocampus. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology*. 2009; 34:2376–89. [PubMed: 19606083]
- Bourre JM, et al. Essentiality of omega 3 fatty acids for brain structure and function. *World Rev Nutr Diet*. 1991; 66:103–17. [PubMed: 2053331]
- Bradley SR, et al. The mGluR7 allosteric agonist AMN082 produces antidepressant-like effects by modulating glutamatergic signaling. *Pharmacol Biochem Behav*. 2012; 101:35–40. [PubMed: 22138407]
- Bremner JD, et al. Hippocampal volume reduction in major depression. *Am J Psychiatry*. 2000; 157:115–8. [PubMed: 10618023]
- Brunoni AR, et al. A systematic review and meta-analysis of clinical studies on major depression and BDNF levels: implications for the role of neuroplasticity in depression. *Int J Neuropsychopharmacol*. 2008; 11:169–80. [PubMed: 18752720]
- Bueller JA, et al. BDNF Val66Met allele is associated with reduced hippocampal volume in healthy subjects. *Biol Psychiatry*. 2006; 59:812–5. [PubMed: 16442082]
- Calderon F, Kim H-Y. Docosahexaenoic acid promotes neurite growth in hippocampal neurons. *J Neurochem*. 2004; 90:979–88. [PubMed: 15287904]
- Caldji C, et al. The effects of early rearing environment on the development of GABAA and central benzodiazepine receptor levels and novelty-induced fearfulness in the rat. *Neuropsychopharmacology*. 2000; 22:219–29. [PubMed: 10693149]
- Campbell S, et al. Lower hippocampal volume in patients suffering from depression: a meta-analysis. *Am J Psychiatry*. 2004; 161:598–607. [PubMed: 15056502]
- Carlezon WA, et al. Antidepressant-like effects of uridine and omega-3 fatty acids are potentiated by combined treatment in rats. *Biol Psychiatry*. 2005; 57:343–50. [PubMed: 15705349]
- Caspi A, et al. Influence of life stress on depression: moderation by a polymorphism in the 5-HTT gene. *Science*. 2003; 301:386–9. [PubMed: 12869766]
- Castrén E, et al. Role of neurotrophic factors in depression. *Curr Opin Pharmacol*. 2007; 7:18–21. [PubMed: 17049922]
- Chandler-Laney PC, et al. A history of caloric restriction induces neurochemical and behavioral changes in rats consistent with models of depression. *Pharmacol Biochem Behav*. 2007; 87:104–14. [PubMed: 17490740]
- Charney DS, Manji HK. Life stress, genes, and depression: multiple pathways lead to increased risk and new opportunities for intervention. *Sci STKE*. 2004; 2004:re5. [PubMed: 15039492]
- Chen B, et al. Increased hippocampal BDNF immunoreactivity in subjects treated with antidepressant medication. *Biol Psychiatry*. 2001; 50:260–5. [PubMed: 11522260]
- Chen MJ, Russo-Neustadt AA. Running exercise-induced up-regulation of hippocampal brain-derived neurotrophic factor is CREB-dependent. *Hippocampus*. 2009; 19:962–72. [PubMed: 19294650]
- Chen Z-Y, et al. Genetic variant BDNF (Val66Met) polymorphism alters anxiety-related behavior. *Science*. 2006; 314:140–3. [PubMed: 17023662]
- Chen Z-Y, et al. Variant brain-derived neurotrophic factor (BDNF) (Met66) alters the intracellular trafficking and activity-dependent secretion of wild-type BDNF in neurosecretory cells and cortical neurons. *J Neurosci*. 2004; 24:4401–11. [PubMed: 15128854]
- Chiaruttini C, et al. Dendritic trafficking of BDNF mRNA is mediated by translin and blocked by the G196A (Val66Met) mutation. *Proc Natl Acad Sci USA*. 2009; 106:16481–6. [PubMed: 19805324]
- Conklin SM, et al. Long-chain omega-3 fatty acid intake is associated positively with corticolimbic gray matter volume in healthy adults. *Neurosci Lett*. 2007; 421:209–12. [PubMed: 17574755]
- Dagai L, et al. Docosahexaenoic acid significantly stimulates immediate early response genes and neurite outgrowth. *Neurochem Res*. 2009; 34:867–75. [PubMed: 18781386]
- David DJ, et al. Neurogenesis-Dependent and -Independent Effects of Fluoxetine in an Animal Model of Anxiety/Depression. *Neuron*. 2009; 62:479–493. [PubMed: 19477151]

- DeMar JC, et al. One generation of n-3 polyunsaturated fatty acid deprivation increases depression and aggression test scores in rats. *J Lipid Res.* 2006; 47:172–80. [PubMed: 16210728]
- Dishman RK, et al. Activity wheel running reduces escape latency and alters brain monoamine levels after footshock. *Brain Res Bull.* 1997; 42:399–406. [PubMed: 9092882]
- Drevets WC, et al. Subgenual prefrontal cortex abnormalities in mood disorders. *Nature.* 1997; 386:824–7. [PubMed: 9126739]
- Drew MR, Hen R. Adult hippocampal neurogenesis as target for the treatment of depression. *CNS Neurol Disord Drug Targets.* 2007; 6:205–18. [PubMed: 17511617]
- Droste SK, et al. Voluntary exercise impacts on the rat hypothalamic-pituitary-adrenocortical axis mainly at the adrenal level. *Neuroendocrinology.* 2007; 86:26–37. [PubMed: 17595533]
- Duan W, et al. Brain-derived neurotrophic factor mediates an excitoprotective effect of dietary restriction in mice. *J Neurochem.* 2001; 76:619–26. [PubMed: 11208925]
- Duman CH, et al. A role for MAP kinase signaling in behavioral models of depression and antidepressant treatment. *Biol Psychiatry.* 2007; 61:661–70. [PubMed: 16945347]
- Duman CH, et al. Voluntary exercise produces antidepressant and anxiolytic behavioral effects in mice. *Brain Res.* 2008; 1199:148–58. [PubMed: 18267317]
- Duman CH, et al. Peripheral insulin-like growth factor-I produces antidepressant-like behavior and contributes to the effect of exercise. *Behav Brain Res.* 2009; 198:366–71. [PubMed: 19056428]
- Duman RS. Depression: a case of neuronal life and death? *Biol Psychiatry.* 2004a; 56:140–5. [PubMed: 15271581]
- Duman RS. Role of neurotrophic factors in the etiology and treatment of mood disorders. *Neuromolecular Med.* 2004b; 5:11–25. [PubMed: 15001809]
- Duman RS. Neurotrophic factors and regulation of mood: role of exercise, diet and metabolism. *Neurobiol Aging.* 2005; 26(Suppl 1):88–93. [PubMed: 16226350]
- Duman RS, et al. Signaling pathways underlying the rapid antidepressant actions of ketamine. *Neuropharmacology.* 2011
- Duman RS, Monteggia LM. A neurotrophic model for stress-related mood disorders. *Biol Psychiatry.* 2006; 59:1116–27. [PubMed: 16631126]
- Duman RS, Voleti B. Signaling pathways underlying the pathophysiology and treatment of depression: novel mechanisms for rapid-acting agents. *Trends Neurosci.* 2012; 35:47–56. [PubMed: 22217452]
- Dunn AL, et al. Exercise treatment for depression: efficacy and dose response. *Am J Prev Med.* 2005; 28:1–8. [PubMed: 15626549]
- Duric V, et al. A negative regulator of MAP kinase causes depressive behavior. *Nat Med.* 2010; 16:1328–32. [PubMed: 20953200]
- Dwivedi Y, et al. ERK MAP kinase signaling in post-mortem brain of suicide subjects: differential regulation of upstream Raf kinases Raf-1 and B-Raf. *Mol Psychiatry.* 2006; 11:86–98. [PubMed: 16172610]
- Dwivedi Y, et al. Altered gene expression of brain-derived neurotrophic factor and receptor tyrosine kinase B in postmortem brain of suicide subjects. *Arch Gen Psychiatry.* 2003; 60:804–15. [PubMed: 12912764]
- Dwivedi Y, et al. Aberrant extracellular signal-regulated kinase (ERK) 5 signaling in hippocampus of suicide subjects. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology.* 2007; 32:2338–50. [PubMed: 17342168]
- Dwyer JM, et al. mTOR activation is required for the antidepressant effects of mGluR2/3 blockade. *Int J Neuropsychopharmacol.* 2011:1–6.
- Egan MF, et al. The BDNF val66met polymorphism affects activity-dependent secretion of BDNF and human memory and hippocampal function. *Cell.* 2003; 112:257–69. [PubMed: 12553913]
- Eley TC, et al. Gene-environment interaction analysis of serotonin system markers with adolescent depression. *Mol Psychiatry.* 2004; 9:908–15. [PubMed: 15241435]
- Elzinga BM, et al. The impact of childhood abuse and recent stress on serum brain-derived neurotrophic factor and the moderating role of BDNF Val66Met. *Psychopharmacology (Berl).* 2011; 214:319–28. [PubMed: 20703451]

- Erickson KI, et al. Exercise training increases size of hippocampus and improves memory. *Proc Natl Acad Sci USA*. 2011; 108:3017–22. [PubMed: 21282661]
- Fabel K, et al. VEGF is necessary for exercise-induced adult hippocampal neurogenesis. *Eur J Neurosci*. 2003; 18:2803–12. [PubMed: 14656329]
- Farmer J, et al. Effects of voluntary exercise on synaptic plasticity and gene expression in the dentate gyrus of adult male Sprague-Dawley rats in vivo. *Neuroscience*. 2004; 124:71–9. [PubMed: 14960340]
- Felitti VJ, et al. Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults. The Adverse Childhood Experiences (ACE) Study. *Am J Prev Med*. 1998; 14:245–58. [PubMed: 9635069]
- Ferraz AC, et al. Chronic ω -3 fatty acids supplementation promotes beneficial effects on anxiety, cognitive and depressive-like behaviors in rats subjected to a restraint stress protocol. *Behav Brain Res*. 2011; 219:116–22. [PubMed: 21192985]
- Fournier NM, Duman RS. Role of vascular endothelial growth factor in adult hippocampal neurogenesis: Implications for the pathophysiology and treatment of depression. *Behav Brain Res*. 2011
- Freeman MP. Omega-3 fatty acids and perinatal depression: a review of the literature and recommendations for future research. *Prostaglandins Leukot Essent Fatty Acids*. 2006; 75:291–7. [PubMed: 16930971]
- Gatt JM, et al. Interactions between BDNF Val66Met polymorphism and early life stress predict brain and arousal pathways to syndromal depression and anxiety. *Mol Psychiatry*. 2009; 14:681–95. [PubMed: 19153574]
- George ED, et al. Maternal separation with early weaning: a novel mouse model of early life neglect. *BMC Neurosci*. 2010; 11:123. [PubMed: 20920223]
- Gertsik L, et al. Omega-3 fatty acid augmentation of citalopram treatment for patients with major depressive disorder. *J Clin Psychopharmacol*. 2012; 32:61–4. [PubMed: 22198441]
- Gervasoni N, et al. Partial normalization of serum brain-derived neurotrophic factor in remitted patients after a major depressive episode. *Neuropsychobiology*. 2005; 51:234–8. [PubMed: 15897674]
- Gorzalka BB, Hill MN. Putative role of endocannabinoid signaling in the etiology of depression and actions of antidepressants. *Prog Neuropsychopharmacol Biol Psychiatry*. 2011; 35:1575–85. [PubMed: 21111017]
- Grabe HJ, et al. Genetic epistasis between the brain-derived neurotrophic factor Val66Met polymorphism and the 5-HTT promoter polymorphism moderates the susceptibility to depressive disorders after childhood abuse. *Prog Neuropsychopharmacol Biol Psychiatry*. 2012; 36:264–70. [PubMed: 21996278]
- Greenwood BN, et al. The consequences of uncontrollable stress are sensitive to duration of prior wheel running. *Brain Res*. 2005; 1033:164–78. [PubMed: 15694921]
- Greenwood BN, et al. Freewheel running prevents learned helplessness/behavioral depression: role of dorsal raphe serotonergic neurons. *J Neurosci*. 2003; 23:2889–98. [PubMed: 12684476]
- Greer PL, Greenberg ME. From synapse to nucleus: calcium-dependent gene transcription in the control of synapse development and function. *Neuron*. 2008; 59:846–60. [PubMed: 18817726]
- Grenyer BFS, et al. Fish oil supplementation in the treatment of major depression: a randomised double-blind placebo-controlled trial. *Prog Neuropsychopharmacol Biol Psychiatry*. 2007; 31:1393–6. [PubMed: 17659823]
- Hahm S, et al. Targeted deletion of the Vgf gene indicates that the encoded secretory peptide precursor plays a novel role in the regulation of energy balance. *Neuron*. 1999; 23:537–48. [PubMed: 10433265]
- Hakkarainen R, et al. Is low dietary intake of omega-3 fatty acids associated with depression? *Am J Psychiatry*. 2004; 161:567–9. [PubMed: 14992986]
- Hariri AR, et al. Brain-derived neurotrophic factor val66met polymorphism affects human memory-related hippocampal activity and predicts memory performance. *J Neurosci*. 2003; 23:6690–4. [PubMed: 12890761]

- Heim C, Binder EB. Current research trends in early life stress and depression: Review of human studies on sensitive periods, gene-environment interactions, and epigenetics. *Exp Neurol*. 2011
- Heim C, Nemeroff CB. The impact of early adverse experiences on brain systems involved in the pathophysiology of anxiety and affective disorders. *Biol Psychiatry*. 1999; 46:1509–22. [PubMed: 10599479]
- Heim C, Nemeroff CB. The role of childhood trauma in the neurobiology of mood and anxiety disorders: preclinical and clinical studies. *Biol Psychiatry*. 2001; 49:1023–39. [PubMed: 11430844]
- Heine VM, et al. Chronic stress in the adult dentate gyrus reduces cell proliferation near the vasculature and VEGF and Flk-1 protein expression. *Eur J Neurosci*. 2005; 21:1304–14. [PubMed: 15813940]
- Herring MP, et al. Effect of exercise training on depressive symptoms among patients with a chronic illness: a systematic review and meta-analysis of randomized controlled trials. *Arch Intern Med*. 2012; 172:101–11. [PubMed: 22271118]
- Hibbeln JR. Fish consumption and major depression. *Lancet*. 1998; 351:1213. [PubMed: 9643729]
- Hill MN, et al. Endogenous cannabinoid signaling is required for voluntary exercise-induced enhancement of progenitor cell proliferation in the hippocampus. *Hippocampus*. 2010; 20:513–23. [PubMed: 19489006]
- Hoeffler CA, Klann E. mTOR signaling: at the crossroads of plasticity, memory and disease. *Trends Neurosci*. 2010; 33:67–75. [PubMed: 19963289]
- Huang S-Y, et al. Omega-3 fatty acids on the forced-swimming test. *J Psychiatr Res*. 2008; 42:58–63. [PubMed: 17070845]
- Hulshof HJ, et al. Maternal separation decreases adult hippocampal cell proliferation and impairs cognitive performance but has little effect on stress sensitivity and anxiety in adult Wistar rats. *Behav Brain Res*. 2011; 216:552–60. [PubMed: 20816703]
- Hunsberger JG, et al. Antidepressant actions of the exercise-regulated gene VGF. *Nat Med*. 2007; 13:1476–82. [PubMed: 18059283]
- Iversen IH. Techniques for establishing schedules with wheel running as reinforcement in rats. *J Exp Anal Behav*. 1993; 60:219–38. [PubMed: 8354968]
- Izquierdo A, et al. Brief uncontrollable stress causes dendritic retraction in infralimbic cortex and resistance to fear extinction in mice. *J Neurosci*. 2006; 26:5733–8. [PubMed: 16723530]
- Jahng JW, et al. Chronic food restriction in young rats results in depression- and anxiety-like behaviors with decreased expression of serotonin reuptake transporter. *Brain Res*. 2007; 1150:100–7. [PubMed: 17383614]
- James MJ, Cleland LG. Dietary n-3 fatty acids and therapy for rheumatoid arthritis. *Semin Arthritis Rheum*. 1997; 27:85–97. [PubMed: 9355207]
- Jernigan CS, et al. The mTOR signaling pathway in the prefrontal cortex is compromised in major depressive disorder. *Prog Neuropsychopharmacol Biol Psychiatry*. 2011; 35:1774–9. [PubMed: 21635931]
- Jourdi H, et al. BDNF mediates the neuroprotective effects of positive AMPA receptor modulators against MPP⁺-induced toxicity in cultured hippocampal and mesencephalic slices. *Neuropharmacology*. 2009; 56:876–85. [PubMed: 19371576]
- Karege F, et al. Low brain-derived neurotrophic factor (BDNF) levels in serum of depressed patients probably results from lowered platelet BDNF release unrelated to platelet reactivity. *Biol Psychiatry*. 2005; 57:1068–72. [PubMed: 15860348]
- Kaufman J, et al. Brain-derived neurotrophic factor-5-HTTLPR gene interactions and environmental modifiers of depression in children. *Biol Psychiatry*. 2006; 59:673–80. [PubMed: 16458264]
- Kaufman J, et al. Social supports and serotonin transporter gene moderate depression in maltreated children. *Proc Natl Acad Sci USA*. 2004; 101:17316–21. [PubMed: 15563601]
- Kendler KS. “A gene for...”: the nature of gene action in psychiatric disorders. *Am J Psychiatry*. 2005; 162:1243–52. [PubMed: 15994704]
- Kessler RC, et al. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry*. 2005; 62:617–27. [PubMed: 15939839]

- Kiuchi T, et al. Regular exercise cures depression-like behavior via VEGF-Flk-1 signaling in chronically stressed mice. *Neuroscience*. 2012; 207:208–17. [PubMed: 22306286]
- Klein AB, et al. Blood BDNF concentrations reflect brain-tissue BDNF levels across species. *Int J Neuropsychopharmacol*. 2011; 14:347–53. [PubMed: 20604989]
- Koike H, et al. Involvement of the mammalian target of rapamycin signaling in the antidepressant-like effect of group II metabotropic glutamate receptor antagonists. *Neuropharmacology*. 2011; 61:1419–23. [PubMed: 21903115]
- Krishnan V, Nestler EJ. The molecular neurobiology of depression. *Nature*. 2008; 455:894–902. [PubMed: 18923511]
- Lafenêtre P, et al. Exercise can rescue recognition memory impairment in a model with reduced adult hippocampal neurogenesis. *Front Behav Neurosci*. 2010; 3:34. [PubMed: 20204139]
- Lambrechts D, et al. VEGF is a modifier of amyotrophic lateral sclerosis in mice and humans and protects motoneurons against ischemic death. *Nature Genetics*. 2003; 34:383–94. [PubMed: 12847526]
- Lee J, et al. Evidence that brain-derived neurotrophic factor is required for basal neurogenesis and mediates, in part, the enhancement of neurogenesis by dietary restriction in the hippocampus of adult mice. *J Neurochem*. 2002a; 82:1367–75. [PubMed: 12354284]
- Lee J, et al. Dietary restriction enhances neurotrophin expression and neurogenesis in the hippocampus of adult mice. *J Neurochem*. 2002b; 80:539–47. [PubMed: 11905999]
- Lee J-H, et al. Depressive behaviors and decreased expression of serotonin reuptake transporter in rats that experienced neonatal maternal separation. *Neurosci Res*. 2007; 58:32–9. [PubMed: 17298851]
- Lespérance F, et al. The efficacy of omega-3 supplementation for major depression: a randomized controlled trial. *J Clin Psychiatry*. 2011; 72:1054–62. [PubMed: 20584525]
- Li N, et al. mTOR-dependent synapse formation underlies the rapid antidepressant effects of NMDA antagonists. *Science*. 2010; 329:959–64. [PubMed: 20724638]
- Li N, et al. Glutamate N-methyl-D-aspartate receptor antagonists rapidly reverse behavioral and synaptic deficits caused by chronic stress exposure. *Biol Psychiatry*. 2011; 69:754–61. [PubMed: 21292242]
- Li Y, et al. TrkB regulates hippocampal neurogenesis and governs sensitivity to antidepressive treatment. *Neuron*. 2008; 59:399–412. [PubMed: 18701066]
- Liebrenz M, et al. Intravenous ketamine therapy in a patient with a treatment-resistant major depression. *Swiss Med Wkly*. 2007; 137:234–6. [PubMed: 17525879]
- Liu D, et al. Maternal care, hippocampal synaptogenesis and cognitive development in rats. *Nat Neurosci*. 2000; 3:799–806. [PubMed: 10903573]
- Liu R-J, et al. Brain-Derived Neurotrophic Factor Val66Met Allele Impairs Basal and Ketamine-Stimulated Synaptogenesis in Prefrontal Cortex. *Biol Psychiatry*. 2011
- Llorens-Martín M, et al. Exercise modulates insulin-like growth factor 1-dependent and -independent effects on adult hippocampal neurogenesis and behaviour. *Mol Cell Neurosci*. 2010; 44:109–17. [PubMed: 20206269]
- Magariños AM, et al. Effect of brain-derived neurotrophic factor haploinsufficiency on stress-induced remodeling of hippocampal neurons. *Hippocampus*. 2011; 21:253–64. [PubMed: 20095008]
- Maier SF, Watkins LR. Stressor controllability and learned helplessness: the roles of the dorsal raphe nucleus, serotonin, and corticotropin-releasing factor. *Neurosci Biobehav Rev*. 2005; 29:829–41. [PubMed: 15893820]
- Makatsori A, et al. Voluntary wheel running modulates glutamate receptor subunit gene expression and stress hormone release in Lewis rats. *Psychoneuroendocrinology*. 2003; 28:702–14. [PubMed: 12727136]
- Marangell LB, et al. A double-blind, placebo-controlled study of the omega-3 fatty acid docosahexaenoic acid in the treatment of major depression. *Am J Psychiatry*. 2003; 160:996–8. [PubMed: 12727707]
- Martinowich K, et al. New insights into BDNF function in depression and anxiety. *Nat Neurosci*. 2007; 10:1089–93. [PubMed: 17726474]

- Martins JG. EPA but not DHA appears to be responsible for the efficacy of omega-3 long chain polyunsaturated fatty acid supplementation in depression: evidence from a meta-analysis of randomized controlled trials. *J Am Coll Nutr.* 2009; 28:525–42. [PubMed: 20439549]
- Mattson MP. Energy intake, meal frequency, and health: a neurobiological perspective. *Annu Rev Nutr.* 2005; 25:237–60. [PubMed: 16011467]
- Mattson MP, et al. BDNF and 5-HT: a dynamic duo in age-related neuronal plasticity and neurodegenerative disorders. *Trends Neurosci.* 2004a; 27:589–94. [PubMed: 15374669]
- Mattson MP, et al. A neural signaling triumvirate that influences ageing and age-related disease: insulin/IGF-1, BDNF and serotonin. *Ageing Res Rev.* 2004b; 3:445–64. [PubMed: 15541711]
- McAllister AK. Spatially restricted actions of BDNF. *Neuron.* 2002; 36:549–50. [PubMed: 12441043]
- McEwen BS. Stress and hippocampal plasticity. *Annu Rev Neurosci.* 1999; 22:105–22. [PubMed: 10202533]
- McEwen BS. Physiology and neurobiology of stress and adaptation: central role of the brain. *Physiological Reviews.* 2007; 87:873–904. [PubMed: 17615391]
- McKinnon MC, et al. A meta-analysis examining clinical predictors of hippocampal volume in patients with major depressive disorder. *J Psychiatry Neurosci.* 2009; 34:41–54. [PubMed: 19125212]
- Meaney MJ. Maternal care, gene expression, and the transmission of individual differences in stress reactivity across generations. *Annu Rev Neurosci.* 2001; 24:1161–92. [PubMed: 11520931]
- Mirescu C, et al. Early life experience alters response of adult neurogenesis to stress. *Nat Neurosci.* 2004; 7:841–6. [PubMed: 15273691]
- Mischoulon D, Fava M. Docosahexanoic acid and omega-3 fatty acids in depression. *Psychiatr Clin North Am.* 2000; 23:785–94. [PubMed: 11147247]
- Moghaddam B, et al. Activation of glutamatergic neurotransmission by ketamine: a novel step in the pathway from NMDA receptor blockade to dopaminergic and cognitive disruptions associated with the prefrontal cortex. *J Neurosci.* 1997; 17:2921–7. [PubMed: 9092613]
- Monteggia LM. Elucidating the role of brain-derived neurotrophic factor in the brain. *Am J Psychiatry.* 2007; 164:1790. [PubMed: 18056232]
- Monteggia LM, et al. Essential role of brain-derived neurotrophic factor in adult hippocampal function. *Proc Natl Acad Sci USA.* 2004; 101:10827–32. [PubMed: 15249684]
- Mullen PE, et al. The long-term impact of the physical, emotional, and sexual abuse of children: a community study. *Child Abuse Negl.* 1996; 20:7–21. [PubMed: 8640429]
- Murakami S, et al. Chronic stress, as well as acute stress, reduces BDNF mRNA expression in the rat hippocampus but less robustly. *Neurosci Res.* 2005; 53:129–39. [PubMed: 16024125]
- Naliwaiko K, et al. Effects of fish oil on the central nervous system: a new potential antidepressant? *Nutr Neurosci.* 2004; 7:91–9. [PubMed: 15279495]
- Nederhof E, et al. Interaction between childhood adversity, brain-derived neurotrophic factor val/met and serotonin transporter promoter polymorphism on depression: the TRAILS study. *Biol Psychiatry.* 2010; 68:209–12. [PubMed: 20553751]
- Nemets B, et al. Addition of omega-3 fatty acid to maintenance medication treatment for recurrent unipolar depressive disorder. *Am J Psychiatry.* 2002; 159:477–9. [PubMed: 11870016]
- Nestler EJ, et al. Preclinical models: status of basic research in depression. *Biol Psychiatry.* 2002; 52:503–28. [PubMed: 12361666]
- Nibuya M, et al. Regulation of BDNF and trkB mRNA in rat brain by chronic electroconvulsive seizure and antidepressant drug treatments. *J Neurosci.* 1995; 15:7539–47. [PubMed: 7472505]
- Nibuya M, et al. Chronic antidepressant administration increases the expression of cAMP response element binding protein (CREB) in rat hippocampus. *J Neurosci.* 1996; 16:2365–72. [PubMed: 8601816]
- Pezawas L, et al. The brain-derived neurotrophic factor val66met polymorphism and variation in human cortical morphology. *J Neurosci.* 2004; 24:10099–102. [PubMed: 15537879]
- Radley JJ, et al. Repeated stress induces dendritic spine loss in the rat medial prefrontal cortex. *Cereb Cortex.* 2006; 16:313–20. [PubMed: 15901656]

- Radley JJ, et al. Chronic behavioral stress induces apical dendritic reorganization in pyramidal neurons of the medial prefrontal cortex. *Neuroscience*. 2004; 125:1–6. [PubMed: 15051139]
- Rajkowska G, et al. Morphometric evidence for neuronal and glial prefrontal cell pathology in major depression. *Biol Psychiatry*. 1999; 45:1085–98. [PubMed: 10331101]
- Rao JS, et al. n-3 polyunsaturated fatty acid deprivation in rats decreases frontal cortex BDNF via a p38 MAPK-dependent mechanism. *Mol Psychiatry*. 2007; 12:36–46. [PubMed: 16983391]
- Redila VA, Christie BR. Exercise-induced changes in dendritic structure and complexity in the adult hippocampal dentate gyrus. *Neuroscience*. 2006; 137:1299–307. [PubMed: 16338077]
- Rees A-M, et al. Omega-3 fatty acids as a treatment for perinatal depression: randomized double-blind placebo-controlled trial. *Aust N Z J Psychiatry*. 2008; 42:199–205. [PubMed: 18247194]
- Risch N, et al. Interaction between the serotonin transporter gene (5-HTTLPR), stressful life events, and risk of depression: a meta-analysis. *JAMA*. 2009; 301:2462–71. [PubMed: 19531786]
- Roceri M, et al. Postnatal repeated maternal deprivation produces age-dependent changes of brain-derived neurotrophic factor expression in selected rat brain regions. *Biol Psychiatry*. 2004; 55:708–14. [PubMed: 15038999]
- Rocha Araujo DM, et al. What is the effectiveness of the use of polyunsaturated fatty acid omega-3 in the treatment of depression? *Expert Rev Neurother*. 2010; 10:1117–29. [PubMed: 20586692]
- Roozendaal B, et al. Stress, memory and the amygdala. *Nat Rev Neurosci*. 2009; 10:423–33. [PubMed: 19469026]
- Roth GS, et al. Effects of reduced energy intake on the biology of aging: the primate model. *Eur J Clin Nutr*. 2000; 54(Suppl 3):S15–20. [PubMed: 11041070]
- Russo-Neustadt A, et al. Exercise, antidepressant medications, and enhanced brain derived neurotrophic factor expression. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology*. 1999; 21:679–82. [PubMed: 10516964]
- Russo-Neustadt AA, et al. Physical activity and antidepressant treatment potentiate the expression of specific brain-derived neurotrophic factor transcripts in the rat hippocampus. *Neuroscience*. 2000; 101:305–12. [PubMed: 11074154]
- Sacher J, et al. Mapping the depressed brain: A meta-analysis of structural and functional alterations in major depressive disorder. *Journal of affective disorders*. 2011
- Sahay A, Hen R. Adult hippocampal neurogenesis in depression. *Nat Neurosci*. 2007; 10:1110–5. [PubMed: 17726477]
- Sairanen M, et al. Brain-derived neurotrophic factor and antidepressant drugs have different but coordinated effects on neuronal turnover, proliferation, and survival in the adult dentate gyrus. *J Neurosci*. 2005; 25:1089–94. [PubMed: 15689544]
- Samuels BA, Hen R. Neurogenesis and affective disorders. *Eur J Neurosci*. 2011; 33:1152–9. [PubMed: 21395859]
- Sanacora G. Cortical inhibition, gamma-aminobutyric acid, and major depression: there is plenty of smoke but is there fire? *Biol Psychiatry*. 2010; 67:397–8. [PubMed: 20159143]
- Santarelli L, et al. Requirement of hippocampal neurogenesis for the behavioral effects of antidepressants. *Science*. 2003; 301:805–9. [PubMed: 12907793]
- Sartori CR, et al. The antidepressive effect of the physical exercise correlates with increased levels of mature BDNF, and proBDNF proteolytic cleavage-related genes, p11 and tPA. *Neuroscience*. 2011; 180:9–18. [PubMed: 21371535]
- Schmidt HD, Duman RS. Peripheral BDNF produces antidepressant-like effects in cellular and behavioral models. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology*. 2010; 35:2378–91. [PubMed: 20686454]
- Schmidt HD, et al. Functional Biomarkers of Depression: Diagnosis, Treatment, and Pathophysiology. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology*. 2011
- Seligman ME, Beagley G. Learned helplessness in the rat. *J Comp Physiol Psychol*. 1975; 88:534–41. [PubMed: 1150935]
- Selye H. A syndrome produced by diverse nocuous agents. *Nature*. 1936:138.

- Sen S, et al. Serum brain-derived neurotrophic factor, depression, and antidepressant medications: meta-analyses and implications. *Biol Psychiatry*. 2008; 64:527–32. [PubMed: 18571629]
- Shansky RM, et al. Stress-induced dendritic remodeling in the prefrontal cortex is circuit specific. *Cereb Cortex*. 2009; 19:2479–84. [PubMed: 19193712]
- Sheline YI, et al. Untreated depression and hippocampal volume loss. *Am J Psychiatry*. 2003; 160:1516–8. [PubMed: 12900317]
- Sheline YI, et al. Hippocampal atrophy in recurrent major depression. *Proc Natl Acad Sci USA*. 1996; 93:3908–13. [PubMed: 8632988]
- Shimizu E, et al. Alterations of serum levels of brain-derived neurotrophic factor (BDNF) in depressed patients with or without antidepressants. *Biol Psychiatry*. 2003; 54:70–5. [PubMed: 12842310]
- Shirayama Y, et al. Brain-derived neurotrophic factor produces antidepressant effects in behavioral models of depression. *J Neurosci*. 2002; 22:3251–61. [PubMed: 11943826]
- Siuciak JA, et al. Antidepressant-like effect of brain-derived neurotrophic factor (BDNF). *Pharmacol Biochem Behav*. 1997; 56:131–7. [PubMed: 8981620]
- Slipczuk L, et al. BDNF activates mTOR to regulate GluR1 expression required for memory formation. *PLoS ONE*. 2009; 4:e6007. [PubMed: 19547753]
- Snyder JS, et al. Adult hippocampal neurogenesis buffers stress responses and depressive behaviour. *Nature*. 2011; 476:458–61. [PubMed: 21814201]
- Sousa N, et al. Reorganization of the morphology of hippocampal neurites and synapses after stress-induced damage correlates with behavioral improvement. *Neuroscience*. 2000; 97:253–66. [PubMed: 10799757]
- Sparling PB, et al. Exercise activates the endocannabinoid system. *Neuroreport*. 2003; 14:2209–11. [PubMed: 14625449]
- Stockmeier CA, et al. Cellular changes in the postmortem hippocampus in major depression. *Biol Psychiatry*. 2004; 56:640–50. [PubMed: 15522247]
- Stranahan AM, et al. Running induces widespread structural alterations in the hippocampus and entorhinal cortex. *Hippocampus*. 2007; 17:1017–22. [PubMed: 17636549]
- Stranahan AM, et al. Central mechanisms of HPA axis regulation by voluntary exercise. *Neuromolecular Med*. 2008; 10:118–27. [PubMed: 18273712]
- Su K-P, et al. Omega-3 fatty acids in major depressive disorder. A preliminary double-blind, placebo-controlled trial. *Eur Neuropsychopharmacol*. 2003; 13:267–71. [PubMed: 12888186]
- Szeszko PR, et al. Brain-derived neurotrophic factor val66met polymorphism and volume of the hippocampal formation. *Mol Psychiatry*. 2005; 10:631–6. [PubMed: 15768049]
- Takei N, et al. Brain-derived neurotrophic factor induces mammalian target of rapamycin-dependent local activation of translation machinery and protein synthesis in neuronal dendrites. *J Neurosci*. 2004; 24:9760–9. [PubMed: 15525761]
- Taliaz D, et al. Knockdown of brain-derived neurotrophic factor in specific brain sites precipitates behaviors associated with depression and reduces neurogenesis. *Mol Psychiatry*. 2010; 15:80–92. [PubMed: 19621014]
- Tang K, et al. Exercise-induced VEGF transcriptional activation in brain, lung and skeletal muscle. *Respir Physiol Neurobiol*. 2010; 170:16–22. [PubMed: 19853064]
- Thakker-Varia S, et al. The neuropeptide VGF produces antidepressant-like behavioral effects and enhances proliferation in the hippocampus. *J Neurosci*. 2007; 27:12156–67. [PubMed: 17989282]
- Trejo JL, et al. Circulating insulin-like growth factor I mediates exercise-induced increases in the number of new neurons in the adult hippocampus. *J Neurosci*. 2001; 21:1628–34. [PubMed: 11222653]
- Trejo JL, et al. The effects of exercise on spatial learning and anxiety-like behavior are mediated by an IGF-I-dependent mechanism related to hippocampal neurogenesis. *Mol Cell Neurosci*. 2008; 37:402–11. [PubMed: 18086533]
- Uher R, et al. Serotonin transporter gene moderates childhood maltreatment's effects on persistent but not single-episode depression: replications and implications for resolving inconsistent results. *Journal of affective disorders*. 2011; 135:56–65. [PubMed: 21439648]

- van Praag H, et al. Running increases cell proliferation and neurogenesis in the adult mouse dentate gyrus. *Nat Neurosci.* 1999; 2:266–70. [PubMed: 10195220]
- Venna VR, et al. PUFA induce antidepressant-like effects in parallel to structural and molecular changes in the hippocampus. *Psychoneuroendocrinology.* 2009; 34:199–211. [PubMed: 18848400]
- Verhagen M, et al. Meta-analysis of the BDNF Val66Met polymorphism in major depressive disorder: effects of gender and ethnicity. *Mol Psychiatry.* 2010; 15:260–71. [PubMed: 18852698]
- Vermetten E, et al. Long-term treatment with paroxetine increases verbal declarative memory and hippocampal volume in posttraumatic stress disorder. *Biol Psychiatry.* 2003; 54:693–702. [PubMed: 14512209]
- Viikki M, et al. Vascular endothelial growth factor (VEGF) polymorphism is associated with treatment resistant depression. *Neurosci Lett.* 2010; 477:105–8. [PubMed: 20417690]
- Vines A, et al. The role of 5-HT(1A) receptors in fish oil-mediated increased BDNF expression in the rat hippocampus and cortex: A possible antidepressant mechanism. *Neuropharmacology.* 2011
- Voleti B, Duman RS. The Roles of Neurotrophic Factor and Wnt Signaling in Depression. *Clinical pharmacology and therapeutics.* 2011
- Warner-Schmidt JL, Duman RS. VEGF is an essential mediator of the neurogenic and behavioral actions of antidepressants. *Proc Natl Acad Sci USA.* 2007; 104:4647–52. [PubMed: 17360578]
- Wei L, et al. Early life stress increases anxiety-like behavior in Balbc mice despite a compensatory increase in levels of postnatal maternal care. *Horm Behav.* 2010; 57:396–404. [PubMed: 20096699]
- Weindruch RH, et al. Influence of controlled dietary restriction on immunologic function and aging. *Fed Proc.* 1979; 38:2007–16. [PubMed: 437143]
- Wichers M, et al. The BDNF Val(66)Met x 5-HTTLPR x child adversity interaction and depressive symptoms: An attempt at replication. *Am J Med Genet B Neuropsychiatr Genet.* 2008; 147B: 120–3. [PubMed: 17579366]
- Wilkinson PO, Goodyer IM. Childhood adversity and allostatic overload of the hypothalamic-pituitary-adrenal axis: a vulnerability model for depressive disorders. *Dev Psychopathol.* 2011; 23:1017–37. [PubMed: 22018079]
- Willner P. Validity, reliability and utility of the chronic mild stress model of depression: a 10-year review and evaluation. *Psychopharmacology (Berl).* 1997; 134:319–29. [PubMed: 9452163]
- Wolf SA, et al. Cannabinoid receptor CB1 mediates baseline and activity-induced survival of new neurons in adult hippocampal neurogenesis. *Cell Commun Signal.* 2010; 8:12. [PubMed: 20565726]
- Zarate CA, et al. A randomized trial of an N-methyl-D-aspartate antagonist in treatment-resistant major depression. *Arch Gen Psychiatry.* 2006; 63:856–64. [PubMed: 16894061]
- Zhao C, et al. Distinct morphological stages of dentate granule neuron maturation in the adult mouse hippocampus. *J Neurosci.* 2006; 26:3–11. [PubMed: 16399667]
- Zheng H, et al. Beneficial effects of exercise and its molecular mechanisms on depression in rats. *Behav Brain Res.* 2006; 168:47–55. [PubMed: 16290283]

Highlights

1. Reviews the literature on environmental and pharmacological modulation of MDD.
2. MDD is associated with neuronal atrophy and structural alterations.
3. Chronic stress provides a model for depression.
4. Pharmacological antidepressants, exercise, and diet can reverse changes incurred by stress and MDD.

Figure 1.

BDNF signaling pathways and opposing effects of stress, depression, exercise and antidepressants. BDNF binds to the TrkB receptor, resulting in autophosphorylation of tyrosine residues. These sites then interact to activate the MAPK, CaMKII, and PI3K signaling cascades. These pathways ultimately regulate multiple cellular systems, including neuronal survival, growth, and neuroplasticity, which can lead to alterations in antidepressant behavioral responses. Stress and depression decrease activation of these pathways, while antidepressants and exercise increase activation of these pathways. Abbreviations: BDNF, brain derived neurotrophic factor; TrkB, tropomyosin-related kinase B; MAPK, mitogen-activated protein kinase; CaMKII, calmodulin-dependent protein kinase; PI3K, phosphatidylinositol 3-kinase; Grb2, growth factor receptor-bound protein 2; GAB1, Grb2-associated binder-1; SOS, son of sevenless; Shc, Src homology 2-domain containing adaptor protein; PLCγ phospholipase C-γ; ERK, extracellular signal-regulated kinase; MEK, MAP/ERK kinase; MKP-1, MAP kinase phosphatase 1; PDK1, phosphoinositide dependent kinase 1; Akt, serine threonine kinase or protein kinase B; IP3, inositol triphosphate. Figure adapted from Duman and Voleti (2012).

Figure 2.

Life timeline of genetic and environmental modulation of healthy emotional state or depression by BDNF and other neurotrophic factors. Genetic risk factors and stress can exacerbate depressive symptoms, while antidepressants, exercise, and diet may ameliorate depressive symptoms, in part, through the action of neurotrophic factors. This up- or down-regulation of BDNF by these environmental factors can shift the balance from depressed to healthy, and vice versa. Abbreviations: BDNF, brain derived neurotrophic factor.

Figure 3.

Pathways mediating the regulation of the mTOR signaling cascade by novel rapid-acting antidepressants. Ketamine, an NMDA receptor antagonist, rapidly stimulates glutamate transmission and AMPA receptor activation, leading to activity-dependent release of BDNF. This leads to phosphorylation of ERK and Akt, subsequently regulating mTOR signaling. This can lead, through regulation of P70S6K and 4E-BP1, to increased translation and increased synaptic proteins, which contributes to increased synaptogenesis and antidepressant behaviors. Abbreviations: BDNF, brain derived neurotrophic factor; TrkB, tropomyosin-related kinase B; mTOR, mammalian target of rapamycin; NMDA, N-methyl-D-aspartic acid; AMPA, alpha-amino-3-hydroxy-5-methyl-4-isoxazole-propionic acid; ERK, extracellular signal-regulated kinase; Akt, serine threonine kinase or protein kinase B; P70S6K, P70 ribosomal protein S6 kinase; 4E-BP1, eukaryotic translation initiation factor 4E-binding protein 1; TSC, tuberous sclerosis complex. Figure adapted from Duman and Voleti (2012).