Published in final edited form as:

Psychoneuroendocrinology. 2012 December ; 37(12): 2042–2046. doi:10.1016/j.psyneuen.2012.04.004.

Neuroendocrine Predictors of Emotional and Behavioral Adjustment in Boys: Longitudinal Follow-Up of a Community Sample

Audrey R. Tyrka^{1,2}, Janet K Lee¹, Julia A. Graber³, Ashley M. Clement¹, Megan M. Kelly^{4,5}, Laura DeRose⁶, Michelle P. Warren⁷, and Jeanne Brooks-Gunn^{7,8}

¹Mood Disorders Research Program and Laboratory for Clinical Neuroscience, Butler Hospital, Providence, RI

²Department of Psychiatry and Human Behavior, Warren Alpert Medical School of Brown University, Providence, RI

³Department of Psychology, University of Florida, Gainesville, FL

⁴Department of Psychiatry, University of Massachusetts Medical School, Worcester, MA

⁵Edith Nourse Rogers Memorial Veterans Hospital, Bedford, MA

⁶Department of Developmental Psychology, Adelphi University, Garden City, NY

⁷College of Physicians & Surgeons, Columbia University, NY, NY

⁸Teachers College, Columbia University, NY, NY

Abstract

Background—Dysregulation of the hypothalamic-pituitary-adrenal (HPA) axis has been observed in adults and children with mood and anxiety disorders and is thought to be involved in the pathogenesis of these disorders. We recently studied a diverse community sample of boys and found associations of behavioral problems, including symptoms of depression and anxiety, with basal and stress-induced cortisol concentrations. Here we examine cortisol-emotional/behavioral associations at a two-year follow-up and test whether initial cortisol is predictive of worsening of emotional/behavioral problems two years later.

Method—Seventy-eight 10–14 year-old boys and their mothers completed a battery of psychosocial assessments, provided morning and afternoon saliva samples, and participated in a home visit involving mildly stressful tasks and saliva collection for cortisol assay during a two-year follow-up assessment.

© 2012 Elsevier Ltd. All rights reserved.

Address Correspondence to: Audrey R. Tyrka, M.D., Ph.D., Butler Hospital, 345 Blackstone Blvd., Providence, RI 02906. TEL: (401) 455-6520. FAX: (401) 455-6534. Audrey_Tyrka@Brown.edu.

Disclosure Statement

The authors report no conflicts of interest.

Contributions

Drs. Tyrka, Graber, Kelly, Warren, and Brooks-Gunn designed the study and participated in manuscript preparation. Drs. Tyrka and Kelly, Ms. Lee and Ms. Clement participated in data analysis and manuscript preparation. Dr. Warren performed the cortisol assays. Dr. DeRose participated in data collection and manuscript preparation.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Results—Consistent with the findings from our time 1 assessment, greater declines in cortisol across the home-visit challenge task were significantly associated with internalizing and externalizing behaviors as well as attention problems and social problems at the two-year follow-up. In addition, morning and afternoon cortisol concentrations at the initial assessment were significant positive predictors of the later development of child depressive symptoms at follow-up after controlling for initial depressive symptoms.

Conclusion—These findings demonstrate that children in the community with internalizing and externalizing behavior problems have altered patterns of HPA axis stress reactivity. In addition, our prospective findings suggest that elevated cortisol concentrations may influence the later development of emotional/behavioral problems in boys.

Keywords

children; adolescents; boys; depression; cortisol; anxiety; HPA axis; stress reactivity; longitudinal; prospective

Introduction

Excessive activation of the hypothalamic-pituitary-adrenal (HPA) axis is seen in a substantial proportion of adults and children with depressive and anxiety disorders (Guerry and Hastings, 2011; Pervanidou, 2008). Youth with subsyndromal symptoms of depression and anxiety (internalizing symptoms) also show increased basal and provoked cortisol concentrations in several studies, whereas externalizing behaviors have been linked to blunted cortisol concentrations and responses in some reports. Other studies have documented elevated cortisol concentrations with externalizing behaviors, possibly due to the co-occurrence of internalizing and externalizing behaviors (Tyrka et al., 2010).

HPA axis dysfunction may lead to behavioral symptoms through effects on glucocorticoid receptors (GRs), which are widely distributed in limbic brain regions involved in the regulation of stress responses, cognition, affect, and behavior. Animal studies show that excessive or prolonged activation of GRs results in neurostructural changes in these regions (McEwen, 2007). Therefore, longitudinal studies of the association between cortisol activity and the later development of psychopathology are of great interest. Statistically controlling for the contemporaneous association between cortisol and emotional/behavioral problems at the initial assessment is necessary to show that cortisol is linked to the *later* development of emotional/behavioral problems.

Only a few prior investigations have conducted such prospective analyses, but the findings are suggestive of an effect of cortisol on the later development of emotional/behavior problems. In a study of 4.5 year-old boys, higher afternoon cortisol predicted the development of greater internalizing behaviors and social wariness in Kindergarten, after controlling for initial behavior (Smider et al., 2002). Elevated morning saliva cortisol in a sample of 13 year-olds predicted depressive symptoms at age 16, after controlling for initial depressive symptoms (Halligan et al., 2007). In a 6-month follow-up study of 7–17 year-old clinic-referred children, cortisol reactivity to a parent-child conflict discussion task was inversely predictive of later internalizing and externalizing behaviors, but positively predictive of anxiety disorders (Granger et al., 1996). Finally, in adolescents at-risk for psychopathology, high morning cortisol predicted onset of a major depressive episode in the following 12 months (Goodyer et al., 2000).

While most prior work on this topic has involved white, middle-class samples of preschool or adolescent-aged boys and girls, we recently found, in a study of boys, aged 8–11 from diverse backgrounds, that internalizing problems were linked to higher basal afternoon

cortisol concentrations (Tyrka et al., 2010). Additionally, a greater decline in cortisol concentrations across a home-visit challenge task was seen in boys with internalizing behaviors and social problems, as well as attention and thought problems. Here we report a two-year follow-up assessment in this sample to determine whether these associations persist. In addition, we hypothesized a prospective relationship between initial cortisol concentrations and later increases in emotional/behavioral problems at the two-year follow-up.

Methods

Participants

Boys aged 8–11 ($M=9.10$, $SD=0.73$) were recruited from several New York City area public schools and followed two years later (Range=1.5–2.5 years) when they were age 10–14 ($M=12.3$, $SD=0.72$). One hundred two of these participants provided data for analyses of the association between cortisol measures and emotion/behavior at the time 1 assessment and were thus included in our initial report on this topic (Tyrka et al., 2010). The current study includes seventy-eight boys with follow-up cortisol and emotional/behavioral measures, including 61 who were included in our initial report (Tyrka et al., 2010), and an additional 17 who participated in the study initially but did not have complete cortisol and emotional/behavioral data at the initial assessment for inclusion in the prior report. The prospective analyses included 69 of these boys who had both the follow-up cortisol and emotional/behavioral measures as well as the initial emotional/behavioral measure that was controlled in the analysis.

The follow-up sample did not differ from the initial cohort at time 1 (T1) on the demographic measures (Table 1). Those who did not participate at follow-up had higher self-reported depressive symptoms at T1 ($p<.05$), but did not differ on any of the other emotional/behavioral measures.

Procedures

Methodological details are described in Tyrka et al., 2010. Briefly, boys and their mothers participated in a home visit stress paradigm involving four challenging tasks with saliva sampling at baseline and after each task. These visits took place between 4:00 and 8:00 PM ($M=4:49$ PM, $SD=2:03$). In addition, mothers and boys completed questionnaires and collected saliva samples at home on two consecutive days: a morning saliva sample immediately upon awakening and before brushing teeth or consuming anything, and an afternoon sample upon return from school. Cortisol concentrations from the two sampling days were averaged for AM and PM samples. Procedures used at T1 and the follow-up assessment were identical except for the use of an age-appropriate cognitive task during the home visit: Object Assembly task from the WISC (Wechsler, 1991) at T1, and the Mazes task (Wechsler, 1974) at follow-up. In addition, while salivettes were used to collect saliva during home visits, boys provided free spit using a straw at follow-up for morning and afternoon samples. Cortisol values using these two methods are nearly identical (Sarstedt, 2007). Procedures were approved by the IRB of Teachers College at Columbia University.

Cortisol assay—Saliva cortisol (Table 2) was assayed in duplicate using radioimmunoassay procedures outlined by Kirschbaum and colleagues (Diagnostic Products Company), with a lower detection limit of 0.02 $\mu\text{g/dl}$, and intra- and inter-assay coefficients of variation less than 5% and 3%, respectively.

Measures

Child Report of Depressive and Anxiety Symptoms—Boys completed the Children's Depression Inventory (CDI; Kovacs, 1985), ($\alpha = .81$ and $\alpha = .85$ for T1 and follow-up assessments, respectively) and the anxiety subscale of the Hopkins Symptom Checklist (HSCL; Rickels et al., 1976) ($\alpha = .79$ and $\alpha = .90$).

Parent Report of Behavior Problems—To assess behavior problems, we used the Internalizing ($\alpha = .84$ and $\alpha = .88$ at T1 and follow-up assessments, respectively) and externalizing ($\alpha = .88$ and $\alpha = .87$) subscales of the widely-used parent-report Child Behavior Checklist (CBCL; Achenbach, 2001). We also included the CBCL subscales social problems, thought problems, and attention problems ($\alpha = .70$, $.65$, and $.79$ at T1, and $\alpha = .66$, $.53$, and $.79$, at the follow-up assessment, respectively).

Statistical Analyses

Data were analyzed using SPSS for Mac version 19, all tests were two-tailed, and alpha was set to .05. For cortisol measures, mean morning and afternoon cortisol concentrations were computed from samples collected on two consecutive days. Outliers, three standard deviations from the mean, were identified and Winsorized to equal the next highest value. For home visit cortisol, at both assessments the first saliva sample yielded the highest cortisol concentration, suggesting that anticipatory adrenocortical activation and degree of subsequent decline in activation was indicative of HPA reactivity in this context. Therefore, a home visit change score was calculated by subtracting the value of the last cortisol sample from the first cortisol sample.

Time of sample was controlled in all analyses to account for the diurnal variation in cortisol concentrations. Correlation coefficients and analysis of variance were computed to determine whether any of the emotional/behavioral measures were correlated with the following potential covariates: age, race (Black, Hispanic and White non-Hispanic), socioeconomic status (SES, assessed with the Combined Family Hollingshead Score) and pubertal development. For the prospective analyses, these tests assessed associations between the potential covariates reported at T1 with emotional/behavioral measures at follow-up. In both concurrent and prospective analyses, SES was identified as a significant covariate for the CBCL externalizing scale. In addition, Tanner stage was correlated with CBCL social problems in the concurrent analysis only. These demographic variables were included as covariates in the relevant analyses. Finally, given the mixed findings for externalizing behaviors in prior studies and that internalizing and externalizing behaviors covary ($r = .57$ and $r = .59$, at T1 and follow-up assessments, respectively), we controlled for internalizing scores in any analyses with significant findings for externalizing behaviors.

Partial correlations, controlling for time of cortisol sample and significant covariates, tested for associations between the follow-up cortisol and emotional/behavioral measures. For the prospective hypotheses, general linear models were used to test for associations between T1 cortisol measures and the follow-up emotional/behavioral measures, controlling for T1 time of sample, the T1 emotional/behavioral measure, and any other significant covariates.

Results

Concurrent Analyses of Cortisol and Emotional/Behavioral Measures at Follow-Up

Home Visit Cortisol Change Score—The follow-up home visit cortisol change score was significantly associated with CBCL internalizing behaviors ($r(72) = .29$, $p = .011$), externalizing behaviors ($r(71) = .31$, $p = .007$), social problems ($r(72) = .32$, $p = .006$), and attention problems ($r(72) = .26$, $p = .027$), and there was a trend for thought problems ($r(72) = .$

20, $p=.09$). For each of these subscales, higher levels of behavior problems were associated with greater declines in saliva cortisol during the home visit. The cortisol change score was not significantly correlated with anxiety symptoms or CDI score. Externalizing behaviors were no longer significant after controlling for internalizing behaviors ($r(70)=.18$, $p=.14$).

Morning and Afternoon Cortisol—Partial correlations testing for associations of behavior and symptoms with morning or afternoon cortisol concentrations at the follow-up were not significant. There was a trend-level negative association between externalizing behaviors and morning cortisol concentrations ($r(73)=-.21$, $p=.07$), which remained after controlling for Internalizing behaviors ($r(72)=-.26$, $p=.052$).

Prospective Analysis of T1 Cortisol Measures and Behavior/Emotion at Follow-up

Home Visit Cortisol Change Score—There were no significant effects of T1 home visit cortisol on the later development of emotional/behavioral problems. There were trend level effects of T1 cortisol change score on follow-up anxiety symptoms ($F(1, 57)=3.9$, $p=.053$) and thought problems ($F(1, 63)=3.6$, $p=.061$), after controlling for T1 symptoms and time of sample.

Morning and Afternoon Cortisol—Both T1 morning cortisol and afternoon cortisol were positive predictors of CDI score at follow-up, controlling for T1 CDI score and time of sample [$F(1, 56)=5.2$, $p=.027$, and $F(1, 58)=4.2$, $p=.045$, respectively]. None of the other emotional/behavioral measures was predicted by T1 morning or afternoon cortisol concentrations.

Discussion

Our concurrent findings of associations between change in home visit cortisol and internalizing, externalizing, social, and attention problems are consistent with our prior findings for internalizing and social problems when boys were 8–11 years of age. Our prospective analyses showed that both T1 morning and afternoon cortisol concentrations predicted later depressive symptoms (and greater declines in home visit cortisol at T1 predicted the development of thought problems and anxiety at trend level). These findings extend previous work on this topic to a diverse community sample of young adolescent boys. Taken together, these findings raise the possibility that elevated cortisol concentrations may lead to changes in brain regions implicated in mood, anxiety, and cognition.

Although basal cortisol concentrations predicted increases in CDI score (and we previously found concurrent associations between basal cortisol and behavior at T1, Tyrka et al., 2010), we did not find concurrent associations at follow-up between behavior and morning or afternoon basal cortisol concentrations, possibly due to power limitations. The present sample is limited by the modest size due to attrition as well as missing data for some who did participate in the follow-up assessment. Although our follow-up sample did not differ from those who did not continue with respect to most emotional/behavioral measures or the demographic characteristics, they did have lower self-reported depressive symptoms on the CDI at T1 and this may have influenced our findings.

It is important to note that we did not assess effects of adverse experiences which are linked to altered adrenocortical responding as well as behavioral problems (Cicchetti et al., 2010). We also acknowledge that the home visit cortisol measure did not reflect a rise in cortisol, but rather different degrees of decline across the home visit, so that it remains unclear whether this represents a classic stress-induced cortisol response. In addition, this is a non-clinical community sample with emotional/behavioral problems and cortisol concentrations in the normative range. Only a few participants were above the clinical threshold for

depressive symptoms on the CDI, however, several participants met the clinical thresholds for internalizing and externalizing at the initial assessment (n=10 and 10, respectively) and at follow-up (n=16 and 13, respectively). The findings for internalizing and externalizing remained unchanged using this categorical variable in logistic regression models (data not shown). Whether these findings reflect processes related to clinical conditions remains to be determined, but it should be noted that the literature on cortisol in children and youth with major depression has been inconsistent (Guerry and Hastings, 2011).

In conclusion, our results confirm and extend findings from the few existing studies on the prospective link between cortisol levels and the later development of internalizing problems in youth. Given the literature on the effects of glucocorticoids on brain regions implicated in cognition, mood, and anxiety, our findings are consistent with the possibility that elevated cortisol concentrations could be causally implicated in the development of these symptoms.

Acknowledgments

*Role of the Funding Source

Supported by NIMH Grants K23 MH067947 (ART) and R01 MH083704-01 (ART) and MH56557 (JBG).

We thank the staff of the Boys Health and Development Project and the boys and their families who devoted their time and energy to this study.

References

1. Achenbach, TM. Child Behavior Checklist for Ages 6 to 18. Burlington: University of Vermont Research Center for Children, Youth, & Families; 2001.
2. Cicchetti D, Rogosch FA, Gunnar MR, Toth SL. The differential impacts of early physical and sexual abuse and internalizing problems on daytime cortisol rhythm in school-aged children. *Child Dev.* 2010; 81(1):252–69. [PubMed: 20331666]
3. Goodyer IM, Herbert J, Tamplin A, Altham PM. Recent life events, cortisol, dehydroepiandrosterone and the onset of major depression in high-risk adolescents. *Br J Psychiatry.* 2000; 177:499–504. [PubMed: 11102323]
4. Granger DA, Weisz JR, McCracken JT, Ikeda SC, Douglas P. Reciprocal influences among adrenocortical activation, psychosocial processes, and the behavioral adjustment of clinic-referred children. *Child Dev.* 1996; 67(6):3250–62. [PubMed: 9071780]
5. Guerry JD, Hastings PD. In search of HPA axis dysregulation in child and adolescent depression. *Clin Child Fam Psychol Rev.* 2011; 14(2):135–60. [PubMed: 21290178]
6. Halligan SL, Murray L, Martins C, Cooper PJ. Maternal depression and psychiatric outcomes in adolescent offspring: a 13-year longitudinal study. *J Affect Disord.* 2007; 97(1–3):145–54. [PubMed: 16863660]
7. Kovacs M. The Children's Depression, Inventory (CDI). *Psychopharmacol Bull.* 1985; 21(4):995–8. [PubMed: 4089116]
8. McEwen BS. Physiology and neurobiology of stress and adaptation: central role of the brain. *Physiol Rev.* 2007; 87(3):873–904. [PubMed: 17615391]
9. Pervanidou P. Biology of post-traumatic stress disorder in childhood and adolescence. *J Neuroendocrinol.* 2008; 20(5):632–8. [PubMed: 18363804]
10. Rickels K, Garcia CR, Lipman RS, Derogatis LR, Fisher EL. The Hopkins Symptom Checklist. Assessing emotional distress in obstetric-gynecologic practice. *Prim Care.* 1976; 3(4):751–64. [PubMed: 1051525]
11. Sarstedt. Cortisol-Salivette Saliva Collection Device Evaluation Report. 2007.
12. Smider NA, Essex MJ, Kalin NH, Buss KA, Klein MH, Davidson RJ, Goldsmith HH. Salivary cortisol as a predictor of socioemotional adjustment during kindergarten: a prospective study. *Child Dev.* 2002; 73(1):75–92. [PubMed: 14717245]

13. Tyrka AR, Kelly MM, Graber JA, DeRose L, Lee JK, Warren MP, Brooks-Gunn J. Behavioral adjustment in a community sample of boys: links with basal and stress-induced salivary cortisol concentrations. *Psychoneuroendocrinology*. 2010; 35(8):1167–77. [PubMed: 20223598]
14. Wechsler, D. The Wechsler Intelligence Scale for Children. 3. New York: The Psychological Corporation; 1991.
15. Wechsler, D. Wechsler Intelligence Scale for Children Revised. New York: The Psychological Corporation; 1974.

\$watermark-text

\$watermark-text

\$watermark-text

Table 1

Demographics T1 and Follow-up

Demographics (Range)	Initial <i>M (SD)</i>	Follow-Up <i>M (SD)</i>
Age (8–11 initial, 10–14 follow up)	9.13 (<i>0.74</i>)	12.28 (<i>0.69</i>)
SES: Combined family Hollingshead (8–66 T1, 4–63 follow up)	36.50 (<i>13.89</i>)	30.12 (<i>16.86</i>)
Puberty: Tanner Stage	One: 77 (77%) Two: 19 (18.6%)	One: 4 (4.6%) Two: 21 (24.1%) Three: 31 (35.6%) Four: 28 (32.2%) Five: 3 (3.4%)
Race	Black: 46 (45%) White: 33 (32%) Hispanic: 23 (23%)	White: 32 (41%) Black: 29 (37.2 %) Hispanic: 17 (21.8%)

Table 2

Behavioral Measures and Cortisol at Time 1 and Follow-Up.

Behavioral Measures	Initial (Range)	Follow-Up (Range)
<u>Parent Report</u> (t-scores)		
CBCL Internalizing Symptoms	37.6–76.7	39.3–79.2
CBCL Externalizing Symptoms	38.6–77.9	39.4–78.0
CBCL Social Problems	40.4–77.1	41.5–79.3
CBCL Thought Problems	41.8–81.5	43.1–79.5
CBCL Attention Problems	39.2–77.1	39.3–78.1
<u>Child Report</u>		
Child Depression Inventory (CDI), (0–52)	5.7 (4.6)	6.2 (5.5)
HSCL Anxiety Subscale, (1–5)	1.5 (0.6)	1.4 (0.5)
<u>Cortisol Concentrations</u>		
AM Cortisol (μg/dl)	0.413 (0.278)	0.358 (0.191)
PM Cortisol (μg/ul)	0.103 (0.101)	0.118 (0.097)
Home Visit Sample 1 (μg/dl)	0.092 (0.074)	0.115 (0.105)
Home Visit Sample 2 (μg/dl)	0.060 (0.037)	0.084 (0.062)
Home Visit Sample 3 (μg/dl)	0.059 (0.045)	0.076 (0.061)
Home Visit Sample 4 (μg/dl)	0.055 (0.044)	0.066 (0.050)
Home Visit Sample 5 (μg/dl)	0.062 (0.074)	0.060 (0.046)
Home Visit Δ Score (μg/dl) (HV 1–HV 5)	0.030 (0.061)	0.050 (0.058)

Note. CBCL is the Child Behavior Checklist; HSCL refers to the Hopkins Symptom Checklist.