Published in final edited form as:

Cancer Causes Control. 2012 May ; 23(5): 709–716. doi:10.1007/s10552-012-9940-y.

Obesity and colorectal cancer screening among black and white adults

Sarah S Cohen¹, Harvey J Murff², Lisa B Signorello^{1,2}, and William J Blot^{1,2}

¹International Epidemiology Institute, Rockville, MD

²Division of Epidemiology, Department of Medicine, Vanderbilt Epidemiology Center, Vanderbilt University School of Medicine; and the Vanderbilt-Ingram Cancer Center, Nashville, TN

Abstract

Purpose—To examine whether body mass index is associated with reduced colorectal cancer (CRC) screening in a large population of black and white adults.

Methods—Cross-sectional data collected at baseline for 9,547 black males, 14,515 black females, 3,519 white males, and 7,245 white females age 50–79 enrolled in the Southern Community Cohort Study from 2002–2009 were used to examine odds ratios (OR) with 95% confidence intervals (CI) for use of colonoscopy or sigmoidoscopy in relation to body mass index (BMI) categories (<18.5, 18.5–24.9 (referent), 25–29.9, 30–34.9, 35–39.9, and 40+ (extreme obesity), kg/m²) using logistic regression controlling for age, education, income, health insurance status, last physician visit, cigarette smoking, and alcohol consumption.

Results—Increased BMI was not associated with reduced CRC screening among whites (OR [95% CI] for BMI 40 = 1.02 [0.71–1.46] for white males and 0.99 [0.83–1.19] for white females), and odds of CRC screening were increased with high BMI among blacks (OR [95% CI] for BMI 40 = 1.34 [1.03–1.74] for black males and 1.13 [0.98–1.29] for black females). Extreme obesity was associated with reduced odds of CRC screening only among white women in subgroup analyses limited to those with health insurance or income < \$25,000/year.

Conclusions—Elevated BMI was not a deterrent to CRC screening overall in this population. In light of low overall screening rates for colorectal cancer nationally, efforts to increase screening in all individuals should remain the focus of public health initiatives.

Keywords

Colonoscopy; sigmoidoscopy; obesity; body mass index; race; epidemiology

INTRODUCTION

Colorectal cancer (CRC) is the third most common cancer in the United States with approximately 141,210 new cases and 49,380 deaths expected in 2011 (1). CRC screening is recommended for all adults over the age of 50 in order to reduce CRC incidence through the removal of precancerous lesions and to reduce mortality with early detection (2, 3). Obesity has been consistently linked to increased incidence and mortality of CRC (4, 5), and with the rapidly increasing prevalence of obesity in recent decades (6, 7), screening is of particular importance to prevent substantial morbidity and mortality from this common cancer. However, CRC screening rates are low with only about half of adults over the age of 50

meeting CRC screening guidelines (8). There is evidence to indicate that cancer screening may be lower in overweight or obese individuals and, further, some studies indicate that this association may be modified by race or gender (9–12). However data are limited and have been inconsistent for CRC screening specifically (9), and most studies have not been able to adequately compare associations between obesity and CRC screening across race and gender groups. In this analysis, we examined the association between body mass index (BMI) and CRC screening in the Southern Community Cohort Study which includes large numbers of males and females over a wide range of age and body sizes, and further, includes black and white participants of similar, and often low, socioeconomic status.

METHODS

Study population and data collection

The Southern Community Cohort Study (SCCS) is an ongoing prospective cohort study assessing risks of and disparities in cancer and other chronic diseases and their health impact (13, 14). Study enrollment took place at 71 participating community health centers (CHCs) located in twelve southeastern states (Alabama, Arkansas, Florida, Georgia, Kentucky, Louisiana, Mississippi, North Carolina, South Carolina, Tennessee, Virginia, and West Virginia). CHCs are institutions that provide health and preventative services primarily to low income and uninsured persons (15). Potential study participants were CHC patients, friends or family members who were accompanying patients, and users of CHC pharmacies. Participants were recruited to join the study in-person at the CHC by a trained study interviewer. SCCS eligibility requirements included being age 40–79 years, English-speaking, and not having been under treatment for cancer in the past 12 months. Nearly 73,000 participants were recruited via CHCs from 2002 to 2009. The study was approved by Institutional Review Boards at Vanderbilt University and Meharry Medical College.

Participants were administered an in-person baseline interview by a trained interviewer at enrollment. The computer-assisted personal interview contained questions about demographic, medical, familial, lifestyle and other participant characteristics (a copy of the questionnaire is available on the study website www.southerncommunitystudy.org). For participants enrolled from 2002 until September 2007, current weight and height were obtained by self-report. For 24% of these participants, measured height and weight were abstracted from medical records for validation purposes. Starting in October 2007, current height and weight were measured by study interviewers using a SECA 703 digital scale and height rod.

All participants were asked “Have you ever had a colonoscopy?” and “Have you ever had a sigmoidoscopy?” A sigmoidoscopy was described as “[a] short tube inserted into the rectum while you are awake and un-sedated to look for colon or rectal cancer” and a colonoscopy was described as “[a] long tube inserted into the rectum after you are sedated or put to sleep to look for colon or rectal cancer.” Participants who answered yes to these questions were then asked when their last colonoscopy or sigmoidoscopy took place.

Statistical methods

Current CRC screening guidelines recommend that screening begin at age 50 (2); therefore this analysis was limited to the 37,731 SCCS participants who were age 50 years or older at the time of enrollment. Study exclusions included 1,613 participants who reported their race as other than White or Black/African-American; 388 participants with missing anthropometric values; 200 participants reporting a prior diagnosis of either colon or rectal cancer; and 704 participants with missing values for sigmoidoscopy or colonoscopy.

screening. 34,826 participants age 50–79 at enrollment (9,547 black males, 14,515 black females, 3,519 white males, and 7,245 white females) were included in the final analysis.

The primary outcome for analysis was a dichotomous measure of up-to-date colorectal cancer screening by either sigmoidoscopy in the past five years or colonoscopy in the past ten years (hereafter called “CRC screening”). Sigmoidoscopy alone and colonoscopy alone were also examined as outcome measures. BMI was calculated as $[\text{weight (kg)}]/[\text{height (m)}^2]$ and categorized using standard cut-offs (16) as underweight (BMI < 18.5 kg/m²), healthy weight (18.5–24.9), overweight (25.0–29.9), obesity class I (30.0–34.5), obesity class II (35.0–39.9) and obesity class III or extreme obesity (40.0+).

Odds ratios (OR) with 95% confidence intervals (95% CI) for up-to-date CRC screening associated with categories of BMI were estimated using logistic regression. Participants of healthy weight (BMI 18.5–24.9 kg/m²) were used as the referent category. Potential confounders were determined from the literature and included age, educational attainment, annual household income, health insurance coverage, current cigarette smoking status, alcohol consumption, time since last doctor’s visit, and family history of colon or rectal cancer (categorized as shown in Table 1). Physical activity (obtained from a physical activity questionnaire at the baseline interview and including light, moderate, and vigorous occupational and household activity plus moderate and vigorous sports activity, measured as MET-hours/day) and dietary intake (obtained from an 89-item food frequency questionnaire at the baseline interview, measured as total energy intake in kcal/day) were also examined as potential confounders but their inclusion in the multivariate models did not alter the odds ratios between BMI and CRC screening in a meaningful way. Therefore all models presented here do not include adjustment for physical activity or dietary intake.

Logistic regression models were examined separately for males and females. Effect modification by race was evaluated by comparing models with and without interaction terms for race*BMI using the likelihood ratio test (LRT). The LRT was significant ($p < 0.1$, determined *a priori*) for males ($p = 0.03$) and females ($p = 0.004$). Thus models were stratified by race as well as by gender for this report.

In the SCCS, the prevalence of obesity differed significantly by health insurance status (χ^2 $p < 0.0001$). Because CRC screening is also strongly associated with insurance coverage (8), we conducted analyses limited to the 22,220 (64%) individuals with health insurance coverage to reduce the potential for residual confounding. For similar reasons, analyses stratified by categories of annual household income (<\$15,000/year, \$15,000–\$24,999/year, and \$25,000/year) were also conducted within the race and sex strata. The same set of covariates was used in these subgroup analyses as in the main race-sex stratified models.

To assess potential effects of the dependent nature of the data for participants who enrolled in the SCCS at the same CHC, models that used generalized estimating equations (GEE) to produce standard errors and test statistics were compared to models that did not use GEE. In the models that used GEE, an exchangeable working correlation structure was assumed, meaning that a common covariance was assumed among all participants enrolled in a single CHC. The standard errors produced in the models using GEE were negligibly different from those produced in models without the use of GEE, and therefore the results presented are from models that did not use GEE.

All analyses were conducted using SAS/STAT software Version 9.2 of the SAS System for Windows (SAS Institute Inc., Cary, NC).

RESULTS

Overall, the SCCS population is of relatively low socioeconomic status with 61% of participants reporting a household income less than \$15,000 per year (63% of blacks and 55% of whites) and approximately one-third of the participants reporting less than a high school education (38% of blacks and 29% of whites). Overweight and obesity were both common in the study population with the highest prevalence among black females (86% overweight and 60% obese) and the lowest in black males (64% overweight and 29% obese). Black participants were on average younger than white participants at enrollment (mean age=57.0 v. 59.0 for males and 58.3 v. 59.3 for females).

Crude CRC screening prevalences were 30% in black males, 37% in black females, 36% in white males, and 44% in white females. Age-adjusted up-to-date CRC screening prevalences are shown in Table 1 by race, sex, and covariate categories. Higher screening prevalence was seen among older participants, and those with a college education, a household income of \$50,000 or more, a recent physician visit, and a family history of colorectal cancer while lower screening prevalence was observed among those individuals reporting no health insurance, current smokers, and those reporting more than 2 alcohol drinks per day.

A dichotomous measure of obesity was not associated with decreased CRC in any of the four sex-race groups, with the odds of screening slightly higher among those with BMI over 30 kg/m² versus 18.5–29.9 kg/m² (Table 2). In race and sex-stratified multivariate logistic regression models examining categories of BMI, obesity classes I, II, and III were all associated with higher odds of CRC screening among black males compared with those of a healthy weight. A similar trend was observed among black women although the magnitude of the association was smaller and the confidence intervals slightly overlapped the null. Among whites males, being overweight and obese was generally associated with higher odds of CRC screening although not among men with extreme obesity. No association was observed between overweight or obesity and CRC screening among white women. Models including sigmoidoscopy or colonoscopy as individual outcome measures produced very similar results (data not shown).

Analyses limited to SCCS participants reporting any health insurance coverage showed a similar picture to the entire study population for males of both races (data not shown), but the trends for females, while not statistically significant, were somewhat different than among the SCCS population as a whole. Among insured black females, the modest trend towards increased CRC screening with increasing body size was attenuated; neither overweight nor obesity classes I, II, or III were associated with CRC screening in this group (OR [95% CI]=1.04 [0.90–1.20], 0.99 [0.86–1.14], 1.05 [0.89–1.22], and 1.03 [0.88–1.21], respectively). In contrast, among white women, those in obesity classes II and III had lower odds of CRC screening than healthy weight women (0.89 [0.72–1.10] and 0.88 [0.71–1.10]). While household income was strongly positively associated with CRC screening rates overall (Table 1), within strata of household income (<\$15,000, \$15,000–24,999, and \$25,000+), CRC screening rates across BMI categories were similar to the overall population for most race and sex groups (data not shown). White women had a notably different pattern, however, with a CRC screening prevalence among women with household income of \$25,000+ of 43% in women with extreme obesity compared with 57% in healthy weight women; these differences in screening prevalence were not seen among white women with household incomes of <\$15,000 year (42% versus 38% for extreme obesity versus healthy weight) or \$15,000–24,999 (40% versus 39%). In income-stratified multivariate logistic regression models, income was an effect modifier of the BMI-CRC screening relationship only among white women. For white women with income <\$15,000 and \$15,000–24,999 per year, extreme obesity was not associated with CRC screening

(OR=1.14 [0.80–1.61] and 1.20 [0.80–1.80], respectively) but among women with income of \$25,000 or more, extreme obesity was associated with reduced odds of being up-to-date for CRC screening (OR=0.66 [0.44–1.00]). Similar trends by income category were seen for obesity classes I and II among white women (data not shown).

DISCUSSION

In this large population of generally low-income black and white adults, overall we found that BMI was not associated with CRC screening among whites, and, contrary to our initial hypothesis, that increased body size was associated with small increases in screening odds among black men and women. Previous reports examining CRC screening in relation to body size have been inconsistent. Using 2001 Behavioral Risk Factor Surveillance System (BRFSS) data, obese women were found to be less likely to have received a sigmoidoscopy within the past five years while obese males were more likely to have done so (17) while among the 130,000 individuals (97% white) in the Cancer Prevention Study (CPS) II Nutrition Cohort, lower odds of ever having a colonoscopy or sigmoidoscopy were found among overweight and obese males and females (18). Another study reported that overweight and obese males and females were both slightly more likely to have received CRC screening compared to their healthy weight counterparts (19), and two found no association between CRC screening and overweight or obesity (20, 21). None of these studies presented results specifically among African-Americans. Two relatively small studies have focused solely on African American populations with one reporting non-statistically significant increases in CRC screening among obese African Americans (22), and the other reporting decreased CRC screening among obese black women but not obese black men (12).

Studies examining associations between obesity and CRC screening separately by both race and gender are rare, a gap in the literature that this report uniquely fills. In the few studies that have presented separate results by race and gender, it appears that white women may be the most susceptible to lower screening in relation to larger body size. Using 2005 National Health Interview Study (NHIS) data for 4,430 white and 690 black women, Leone et al. observed lower odds of colonoscopy among obese white women (OR for BMI ≥ 30 v. < 30 = 0.66 [0.50–0.85]) and higher odds among obese black women (OR=1.30 [0.83–2.96] (23). With our much larger population of black and white women in the SCCS, we classified women into obese versus non-obese categories, as well as finer categories of BMI. When we grouped women by dichotomous obesity (BMI ≥ 30 versus 18.5–29.9 kg/m²), the OR of 1.05 [0.97–1.14] for obese black women was similar to the OR for of 1.06 [0.96–1.18] for obese white women. However, comparing BMI ≥ 40 vs 18.5–24.9 kg/m², the OR were 1.13 [0.98–1.29] for black and 0.99 [0.83–1.19] for white women. In addition, among the subset of white women with income $>$ \$25,000, the OR for BMI ≥ 40 vs 18.5–24.9 kg/m² was 0.66 [0.44–1.00], while no parallel decline was seen among black women (OR=1.05 [0.73–1.49]). Hence both the SCCS and NHIS data suggest that racial differences may exist, with lower CRC screening associated with obesity only among moderate to upper income white women.

These differences by race may reflect issues related to negative body image, an issue that has been shown to be more pervasive among white women than black women (24). If white women are more self-conscious about their weight, this may prevent them from receiving certain medical procedures, particularly those that are more invasive such as sigmoidoscopy or colonoscopy. Reasons for the observed difference in screening rates between obese and non-obese individuals in higher but not lower-income women are unclear. It may be that among women of the lowest means, simply meeting the basic needs of food, shelter, and medical care when absolutely necessary is the sole focus while body size is a stronger

influence among women with greater means because basic needs have been met. This is an area deserving of further inquiry.

The overall differences observed between the NHIS and CPS populations and the SCCS could likely be related to SES differences. The CPS study population is of higher educational attainment and has more health-conscious behaviors than the general population, and the NHIS is designed to be nationally representative. In contrast, the SCCS has a much higher proportion of low income, low educational attainment, and uninsured individuals than either the NHIS or CPS.

Among black men, we observed higher CRC screening rates among overweight and obese SCCS participants. It has been reported that overweight and obese individuals utilize health care more often than those of healthy weight and thus have more opportunities to discuss cancer screening with their physician (25). It is also possible that medical practitioners are recommending CRC screening more often to patients who are overweight because excess weight is a known risk factor for CRC. However, we did not observe increased CRC screening among overweight white participants and there is not reason to expect that increased medical visits or screening recommendations for obese individuals would be limited to blacks.

One limitation of this analysis was that we were able only to examine CRC screening by sigmoidoscopy or colonoscopy as information on fecal occult blood testing (FOBT) or barium enema was not obtained as part of the SCCS baseline survey. There is some evidence to indicate that FOBT is used more commonly among individuals of low-income, low education, and without health insurance (26); notably, though, the proportion of screening exams done by FOBT is relatively low and has been dropping in recent years (26). In studies that have examined endoscopy alone versus FOBT and endoscopy together in relation to BMI, associations have generally been similar for the two screening measures; ORs for endoscopy alone versus FOBT or endoscopy by categories of BMI were very similar in an analysis of 2000 NHIS data (21) as were adjusted CRC screening rates using BRFSS data (27). In contrast, using 2005 NHIS data, Leone et al. reported an association between obesity and colonoscopy but not FOBT (23). On balance, the relatively low proportion of CRC screening exams done by FOBT coupled with multiple reports that showed similar associations between obesity and CRC screening regardless of screening modality indicate that the results we found for endoscopy are likely reflective of patterns for all CRC screening.

A further limitation is that both the use of sigmoidoscopy and colonoscopy as well as body size were obtained by self-report. However, self-reported endoscopy by either sigmoidoscopy or colonoscopy had high sensitivity (>96%) and specificity (>92%) when compared with health plan records in at least one study (28). As for self-reported body size, while a recent review indicates that among women, height tends to be over-reported and weight under-reported (29), data from the 1999–2004 National Health and Examination Survey show that despite errors in self-report, BMI categories based on self-reported values still generally demonstrate good agreement with BMI categories from measured values (30). Additionally, within the SCCS, BMI values calculated from self-reported height and weight were very highly correlated with BMI values calculated from medical record data overall (Pearson correlation coefficient > 0.95) as well as across strata of race and BMI, indicating that the self-reported values are generally of good quality.

Finally, it should be noted that the SCCS population, due to its unique recruitment through Community Health Centers and the resulting high proportion of low income individuals, is not reflective of the socioeconomic or race distributions of the entire US population. Thus,

differences in effects compared with other studies, particular those that are nationally representative such as the NHS, may be due to the socioeconomic status of the underlying study populations. However, this study design does not detract from our ability to make within-cohort comparisons between race groups, and is in fact a major strength of this study because residual confounding due to differences in socioeconomic status between race groups is minimized by design, and additionally because low-income individuals have traditionally been understudied.

In summary, in this large population of black and white adults, being overweight or obese does not appear to play a deterring role in the receipt of CRC screening in any race or sex group, except perhaps among the subset of upper income white women. Given that only half of adults over the age of 50 are currently being screened for colorectal cancer, efforts to increase screening rates in all individuals should remain the focus of public health initiatives.

Acknowledgments

The Southern Community Cohort Study is funded by grant R01 CA92447 from the National Cancer Institute including special allocations from the American Recovery and Reinvestment Act (3R01 CA092447-08S1).

References

1. Siegel R, Ward E, Brawley O, Jemal A. Cancer statistics, 2011: the impact of eliminating socioeconomic and racial disparities on premature cancer deaths. *CA Cancer J Clin.* 2011; 61(4): 212–36. Epub 2011/06/21. [PubMed: 21685461]
2. Levin B, Lieberman DA, McFarland B, Smith RA, Brooks D, Andrews KS, et al. Screening and surveillance for the early detection of colorectal cancer and adenomatous polyps, 2008: a joint guideline from the American Cancer Society, the US Multi-Society Task Force on Colorectal Cancer, and the American College of Radiology. *CA Cancer J Clin.* 2008; 58(3):130–60. [PubMed: 18322143]
3. U.S. Preventive Services Task Force. Screening for colorectal cancer: recommendation statement. Rockville, MD: US Department of Health and Human Services, Agency for Healthcare Research and Quality; 2008.
4. Calle EE, Thun MJ. Obesity and cancer. *Oncogene.* 2004; 23(38):6365–78. [PubMed: 15322511]
5. IARC. IARC Handbooks of Cancer Prevention. Weight Control and Physical Activity. International Agency for Research on Cancer; Lyon: 2002.
6. Flegal KM, Carroll MD, Ogden CL, Curtin LR. Prevalence and trends in obesity among US adults, 1999–2008. *Jama.* 2010; 303(3):235–41. [PubMed: 20071471]
7. Flegal KM, Carroll MD, Ogden CL, Johnson CL. Prevalence and trends in obesity among US adults, 1999–2000. *Jama.* 2002; 288(14):1723–7. [PubMed: 12365955]
8. American Cancer Society. Cancer Prevention & Early Detection Facts & Figures 2010. Atlanta: American Cancer Society; 2010.
9. Cohen SS, Palmieri RT, Nyante SJ, Koralek DO, Kim S, Bradshaw P, et al. Obesity and screening for breast, cervical, and colorectal cancer in women: a review. *Cancer.* 2008; 112(9):1892–904. [PubMed: 18361400]
10. Maruthur NM, Bolen S, Brancati FL, Clark JM. Obesity and mammography: a systematic review and meta-analysis. *J Gen Intern Med.* 2009; 24(5):665–77. Epub 2009/03/12. [PubMed: 19277790]
11. Maruthur NM, Bolen SD, Brancati FL, Clark JM. The association of obesity and cervical cancer screening: a systematic review and meta-analysis. *Obesity (Silver Spring, Md.* 2009; 17(2):375–81. Epub 2008/11/11.
12. James AS, Leone L, Katz ML, McNeill LH, Campbell MK. Multiple health behaviors among overweight, class I obese, and class II obese persons. *Ethn Dis.* 2008; 18(2):157–62. Epub 2008/05/30. [PubMed: 18507267]

13. Signorello LB, Hargreaves MK, Blot WJ. The Southern Community Cohort Study: investigating health disparities. *J Health Care Poor Underserved*. 2010; 21(1 Suppl):26–37. [PubMed: 20173283]
14. Signorello LB, Hargreaves MK, Steinwandel MD, Zheng W, Cai Q, Schlundt DG, et al. Southern community cohort study: establishing a cohort to investigate health disparities. *J Natl Med Assoc*. 2005; 97(7):972–9. [PubMed: 16080667]
15. Hargreaves, MK.; Arnold, C.; Blot, WJ. Community health centers: Their role in the treatment of minorities and in health disparities research. In: Satcher, D.; Pamies, R., editors. *Multicultural Medicine and Health Disparities*. New York: McGraw-Hill; 2006. p. 485-94.
16. World Health Organization. Report of a WHO Consultation on Obesity. Geneva: WHO; 1997. Obesity: Preventing and Managing the Global Epidemic.
17. Heo M, Allison DB, Fontaine KR. Overweight, obesity, and colorectal cancer screening: disparity between men and women. *BMC Public Health*. 2004; 4:53. [PubMed: 15533259]
18. Chao A, Connell CJ, Cokkinides V, Jacobs EJ, Calle EE, Thun MJ. Under use of screening sigmoidoscopy and colonoscopy in a large cohort of US adults. *Am J Public Health*. 2004; 94(10):1775–81. [PubMed: 15451749]
19. Yancy WS Jr, McDuffie JR, Stechuchak KM, Olsen MK, Oddone EZ, Kinsinger LS, et al. Obesity and receipt of clinical preventive services in veterans. *Obesity* (Silver Spring, Md. 2010; 18(9): 1827–35.
20. Vlahov D, Ahern J, Vazquez T, Johnson S, Philips LA, Nash D, et al. Racial/ethnic differences in screening for colon cancer: report from the New York Cancer Project. *Ethn Dis*. 2005; 15(1):76–83. [PubMed: 15720052]
21. Wee CC, McCarthy EP, Phillips RS. Factors associated with colon cancer screening: the role of patient factors and physician counseling. *Prev Med*. 2005; 41(1):23–9. [PubMed: 15916989]
22. Satia JA, Galanko JA. Demographic, behavioral, psychosocial, and dietary correlates of cancer screening in African Americans. *J Health Care Poor Underserved*. 2007; 18(4 Suppl):146–64. Epub 2007/12/11. [PubMed: 18065857]
23. Leone LA, Campbell MK, Satia JA, Bowling JM, Pignone MP. Race moderates the relationship between obesity and colorectal cancer screening in women. *Cancer Causes Control*. 21(3):373–85. [PubMed: 19941158]
24. Fitzgibbon ML, Blackman LR, Avellone ME. The relationship between body image discrepancy and body mass index across ethnic groups. *Obes Res*. 2000; 8(8):582–9. [PubMed: 11156434]
25. Bertakis KD, Azari R. Obesity and the use of health care services. *Obes Res*. 2005; 13(2):372–9. [PubMed: 15800296]
26. Klabunde CN, Cronin KA, Breen N, Waldron WR, Ambs AH, Nadel MR. Trends in colorectal cancer test use among vulnerable populations in the United States. *Cancer Epidemiol Biomarkers Prev*. 2011; 20(8):1611–21. Epub 2011/06/10. [PubMed: 21653643]
27. Rosen AB, Schneider EC. Colorectal cancer screening disparities related to obesity and gender. *J Gen Intern Med*. 2004; 19(4):332–8. [PubMed: 15061742]
28. Baier M, Calonge N, Cutter G, McClatchey M, Schoentgen S, Hines S, et al. Validity of self-reported colorectal cancer screening behavior. *Cancer Epidemiol Biomarkers Prev*. 2000; 9(2): 229–32. [PubMed: 10698488]
29. Gorber SC, Tremblay M, Moher D, Gorber B. A comparison of direct vs. self-report measures for assessing height, weight and body mass index: a systematic review. *Obes Rev*. 2007; 8(4):307–26. [PubMed: 17578381]
30. Craig BM, Adams AK. Accuracy of body mass index categories based on self-reported height and weight among women in the United States. *Matern Child Health J*. 2009; 13(4):489–96. [PubMed: 18607705]

Up-to-date colorectal cancer screening (colonoscopy within past 10 years or Sigmoidoscopy within past 5 years) in relation to body mass index (BMI) and potential confounders of the relationship between BMI and screening among black and white participants in the Southern Community Cohort Study, 2002–2009.

Table 1

	Black males (N=9,547)			Black females (N=14,515)			White males (N=3,519)			White females (N=7,245)		
	N	Age-adjusted % screened	% screened	N	Age-adjusted % screened	% screened	N	Age-adjusted % screened	% screened	N	Age-adjusted % screened	% screened
BMI (kg/m ²)												
<18.5	154	32%		115	27%		35	23%		116	40%	
18.5–24.9	3,364	32%		1,975	38%		913	37%		1,545	48%	
25–29.9	3,331	40%		3,807	42%		1,248	46%		1,973	48%	
30–34.9	1,712	45%		3,817	43%		785	48%		1,728	52%	
35–39.9	676	46%		2,525	44%		338	49%		953	48%	
40 +	310	49%		2,276	46%		200	44%		930	49%	
Age at interview												
50–59	6,854	24%		9,330	32%		2,076	28%		4,130	37%	
60–69	2,133	42%		3,838	45%		1,068	44%		2,361	50%	
70–79	560	54%		1,347	52%		375	61%		754	59%	
Education												
< 9 years	1,384	35%		1,641	39%		490	39%		721	44%	
9–11 years	2,524	34%		3,642	40%		563	39%		1,358	47%	
High school ^a	3,414	40%		5,348	42%		1,274	42%		2,934	48%	
Some college/Junior college	1,481	45%		2,553	45%		706	44%		1,397	49%	
College graduate and beyond	742	56%		1,326	53%		486	57%		834	59%	
Household income												
<\$15,000	6,107	35%		9,098	40%		1,973	38%		3,955	46%	
\$15,000–\$24,999	1,951	42%		3,162	42%		710	41%		1,492	46%	
\$25,000–\$49,999	1,002	50%		1,577	52%		493	55%		1,025	57%	
\$50,000+	390	66%		508	64%		308	60%		671	65%	
Health insurance												
Any private	1,661	47%		3,234	48%		648	59%		2,001	56%	
Medicare only	1,221	44%		2,031	45%		591	44%		1,003	53%	
Medicaid only	1,234	37%		2,157	40%		318	42%		739	47%	

	Black males (N=9,547)			Black females (N=14,515)			White males (N=3,519)			White females (N=7,245)		
	N	Age-adjusted % screened	% screened	N	Age-adjusted % screened	% screened	N	Age-adjusted % screened	% screened	N	Age-adjusted % screened	% screened
Other (includes military)	1,536	46%		2,197	49%		661	48%		988	55%	
No health insurance	3,878	24%		4,877	29%		1,296	27%		2,506	32%	
Smoking status												
Current smoker	5,042	31%		3,732	36%		1,467	34%		2,150	43%	
Former smoker	2,508	47%		3,453	45%		1,316	50%		2,084	52%	
Never smoked	1,992	41%		7,317	43%		734	46%		3,006	49%	
Alcohol consumption (drinks/day)												
None	3,528	46%		9,382	43%		1,686	46%		4,699	49%	
1-2	3,500	36%		4,273	41%		1,238	44%		2,270	49%	
> 2	2,486	27%		767	32%		582	33%		244	42%	
Months since last physician visit ^b												
< 1	2,638	45%		4,534	47%		1,146	50%		2,303	52%	
1-3	4,009	41%		7,062	43%		1,443	46%		3,437	50%	
4-6	800	39%		1,225	38%		310	38%		645	48%	
7-12	1,078	27%		1,084	33%		284	37%		526	37%	
13-24	377	21%		253	22%		133	22%		153	33%	
>24	587	11%		299	19%		194	20%		165	16%	
Family history of colorectal cancer ^c												
Yes	530	45%		1,090	53%		291	53%		687	65%	
No	9,017	39%		13,425	42%		3,228	43%		6,558	47%	

^a Includes those with a GED and those who attended vocational school.

^b Visit to a doctor or other medical person

^c Family history includes birth mother, birth father, any full sisters, or any full brothers

Table 2

Odds ratios from race and gender-stratified multivariate logistic regression models examining associations between colorectal cancer screening (colonoscopy within past 10 years or sigmoidoscopy within past 5 years) and body mass index.

	Black Males		Black Females		White Males		White Females	
	OR	95% CI	OR	95% CI	OR	95% CI	OR	95% CI
BMI (kg/m ²)								
<18.5	1.10	(0.74 – 1.64)	0.64	(0.40 – 1.02)	0.60	(0.24 – 1.50)	0.75	(0.49 – 1.14)
18.5–24.9	1.0	REF	1.0	REF	1.0	REF	1.0	REF
25–29.9	1.21	(1.07 – 1.37)	1.04	(0.92 – 1.18)	1.29	(1.06 – 1.59)	0.97	(0.84 – 1.12)
30–34.9	1.32	(1.14 – 1.52)	1.08	(0.95 – 1.22)	1.25	(1.00 – 1.57)	1.08	(0.93 – 1.25)
35–39.9	1.24	(1.02 – 1.51)	1.09	(0.96 – 1.25)	1.33	(0.99 – 1.77)	0.99	(0.83 – 1.19)
40+	1.34	(1.03 – 1.74)	1.13	(0.98 – 1.29)	1.02	(0.71 – 1.46)	0.99	(0.83 – 1.19)
Obese ^a								
No	1.0	REF	1.0	REF	1.0	REF	1.0	REF
Yes	1.16	(1.04–1.29)	1.06	(0.98–1.14)	1.04	(0.89–1.23)	1.06	(0.96–1.17)

Adjusted for age (3 categories), education (5 categories), income (4 categories), health insurance (5 categories), cigarette smoking (3 categories), alcohol consumption (3 categories), last doctor's visit (6 categories), and family history of colon/rectal cancer (Y/N)

^aObesity defined as Yes (BMI ≥ 30 kg/m²) versus No (BMI 18.5–29.9 kg/m²)