Published in final edited form as:

J Genet. 2009 December ; 88(4): 399–407.

RPGR-containing protein complexes in syndromic and non-syndromic retinal degeneration due to ciliary dysfunction

Carlos A. Murga-Zamalloa¹, Anand Swaroop², and Hemant Khanna^{1,*}

¹Department of Ophthalmology and Visual Sciences, University of Michigan, Ann Arbor, MI 48105, USA

²Neurobiology-Neurodegeneration and Repair laboratory (N-NRL), National Eye Institute, National Institutes of Health, Bethesda, MD 20892, USA

Abstract

Dysfunction of primary cilia due to mutations in cilia-centrosomal proteins is associated with pleiotropic disorders. The primary (or sensory) cilium of photoreceptors mediates polarized trafficking of proteins for efficient phototransduction. Retinitis pigmentosa GTPase regulator (RPGR) is a cilia-centrosomal protein mutated in >70% of X-linked RP cases and 10%–20% of simplex RP males. Accumulating evidence indicates that RPGR may facilitate the orchestration of multiple ciliary protein complexes. Disruption of these complexes due to mutations in component proteins is an underlying cause of associated photoreceptor degeneration. Here, we highlight the recent developments in understanding the mechanism of cilia-dependent photoreceptor degeneration due to mutations in RPGR and RPGR-interacting proteins in severe genetic diseases, including retinitis pigmentosa, Leber congenital amaurosis (LCA), Joubert syndrome, and Senior–Loken syndrome, and explore the physiological relevance of photoreceptor ciliary protein complexes.

Keywords

primary cilia; centrosome; transition zone; ciliopathies; photoreceptor; retinal degeneration; retina; RPGR; RP2; CEP290; RPGRIP1L; NPHP

Introduction

The cilium is an extension of the cell membrane formed by nucleation of microtubules. The primary cilium is present in almost all cell types and has diverse functions (Pazour and Witman 2003; Davenport and Yoder 2005; Scholey and Anderson 2006); it contains a central core structure (the axoneme), which comprises of nine outer doublet microtubules with no central microtubule pair (9+0) (Pazour and Witman 2003). The ciliary subunits are assembled at the basal body or mother centriole of post-mitotic cells. This process involves coordinated action of centrosomal proteins and small GTPases that control the switch between cytokinesis and ciliogenesis (Doxsey 2001; Spektor *et al.* 2007; Tsang *et al.* 2008). During ciliogenesis, protein complexes are transported distally for growth of the axoneme using an elaborate mechanism called intraflagellar transport (IFT) (Rosenbaum *et al.* 1999). According to the current model of IFT, protein and membrane cargo are transported

bidirectionally along the axoneme by coordinated action of kinesin (anterograde; KIF family members) and dynein (retrograde) motors (Besharse *et al.* 2003; Follit *et al.* 2006).

Given their near-ubiquitous presence, cilia are involved in diverse cellular processes, including establishment of left–right asymmetry, sonic hedgehog signalling, mechanosensation, olfaction, chemosensation, and photo-transduction (Gerdes *et al.* 2009). Commensurate with this, defects in primary cilia result in severe developmental and lethal disorders, such as altered embryonic patterning, renal cystic diseases, mental retardation and photoreceptor degeneration (Gerdes *et al.* 2009).

Photoreceptor sensory cilium

Photoreceptor inner segment (having the metabolic machinery) and the outer segment (membranous disks containing phototransduction proteins) are linked by a connecting cilium, which is a modified primary cilium (Young 1968) (figure 1). Approximately 10% of outer segments are turned over each day, with new discs being formed proximally and shed distally. About 2000 opsin molecules are transported per minute to maintain the function/integrity of each rod outer segment (Besharse 1986); these molecules are synthesized in the inner segment, sorted at the post-Golgi vesicles and transported to the base of the connecting cilium, where they probably associate with transport proteins for trafficking to the outer segment (Chuang and Sung 1998; Deretic *et al.* 1998). The polarized post-Golgi trafficking and docking of rhodopsin at the basal bodies involve the activity of small GTPases, including Rab8 and ARF4 (Deretic *et al.* 1995; Moritz *et al.* 2001) and is also probably mediated by a FYVE-domain containing protein SARA, phosphatidylinositol 1-phosphate (PI3P), and syntaxin-3 (Chuang *et al.* 2007; Mazelova *et al.* 2009). Perturbation of rhodopsin transport leads to RP (Sung *et al.* 1994; Colley *et al.* 1995).

The transition zone (TZ) of photoreceptor sensory cilium serves as a ‘transport corridor’ for bidirectional trafficking of macromolecular complexes along the microtubule network (figure 1). IFT particle proteins, including Tg737/Polaris/IFT88, localize at the basal body and the axoneme of photoreceptor cilium (Pazour *et al.* 2002). Disruption of IFT in *Kif3a* conditional knockout mice (Marszalek *et al.* 2000) and in *Tg737^{orpk}* mice (Pazour *et al.* 2002) results in opsin accumulation in inner segments and consequently photoreceptor degeneration. The phototransduction proteins, transducin, arrestin, and recoverin undergo lightdependent reversible translocation between outer and inner segments (Sokolov *et al.* 2002; Strissel *et al.* 2005, 2006). Transport of arrestin is, at least in part, mediated by simple diffusion and anchoring through protein–protein interactions (Nair *et al.* 2005).

Photoreceptor ciliopathies

Retinal degeneration due to ciliary dysfunction appears as part of a spectrum of diseases, where one end are some forms of Leber congenital amaurosis (LCA) (MIM 204000), characterized at birth or during early childhood. On the other side of the spectrum resides RP (MIM 268000), wherein a majority of patients exhibit early signs of night blindness in the early stages (due to rod photoreceptor dysfunction) progressing to decreased visual fields and culminating in complete blindness usually in the later stages of life. Depending upon the gene, age of onset phenotype can vary. RP is inherited in autosomal dominant (~30% of cases), autosomal recessive (~20%) as well as X-linked manner (Daiger *et al.* 2007).

X-linked RP (XLRP)

XLRP is one of the severe forms of RP associated with considerable clinical and genetic heterogeneity and accounts for 10%–20% of inherited nonsyndromic RP (Fishman 1978; Jay 1982; Breuer *et al.* 2002). Most affected males exhibit early-onset visual symptoms with

night blindness in the first decade and rapid progression towards blindness by age 40 (Bird 1975; Fishman *et al.* 1988; Daiger *et al.* 2007; Shu *et al.* 2007). Heterozygous carrier females can show electroretinographic (ERG) abnormalities and tapetal-like reflex (Bird 1975; Fishman *et al.* 1986; Cideciyan and Jacobson 1994; Sieving 1995). Some XLRP patients have abnormal sperm phenotype (Hunter *et al.* 1988) or hearing defects (Zito *et al.* 2003; Iannaccone *et al.* 2004). To date, six genetic loci have been identified of which, two causative genes *RP2* and *RP3* (*RPGR*) have been cloned (Wright *et al.* 1991; McGuire *et al.* 1995; Fujita *et al.* 1996; Meindl *et al.* 1996; Roepman *et al.* 1996; Gieser *et al.* 1998; Schwahn *et al.* 1998; Hardcastle *et al.* 2000; Melamud *et al.* 2006).

RP2

Mutations in *RP2* account for approximately 10% of XLRP (Hardcastle *et al.* 1999; Mears *et al.* 1999; Sharon *et al.* 2000, 2003; Breuer *et al.* 2002). The *RP2* gene encodes a putative protein of 350 amino acids (Schwahn *et al.* 1998; Chapple *et al.* 2000). The crystal structure of the RP2 protein reveals an amino-terminal β -helix that is structurally and functionally homologous to the tubulin-specific chaperone, cofactor C; most disease-causing missense mutations are present in this domain (Bartolini *et al.* 2002; Grayson *et al.* 2002; Kuhnel *et al.* 2006). RP2 appears to be targeted predominantly to the plasma membrane (Chapple *et al.* 2000; Grayson *et al.* 2002). It interacts with ADP-ribosylation factor-like 3 (ARL3) (Grayson *et al.* 2002; Kuhnel *et al.* 2006), a microtubule-associated small GTP-binding protein (Kahn *et al.* 2005), which localizes to the connecting cilium of photoreceptors (Grayson *et al.* 2002; Schrick *et al.* 2006).

Retinitis pigmentosa GTPase regulator (RPGR)

Mutations in the retinitis pigmentosa GTPase regulator (*RPGR*) gene are associated with more than 70% of the patients with X-linked retinitis pigmentosa (XLRP) and 10%–20% of simplex RP males (Meindl *et al.* 1996; Roepman *et al.* 1996; Breuer *et al.* 2002; Vervoort and Wright 2002; Sharon *et al.* 2003). These data indicate that RPGR mutations are one of the most common causes of retinal degeneration in addition to rhodopsin. In addition, some patients with mutations in RPGR may also display syndromic features like primary cilia dyskinesia or hearing loss (van Dorp *et al.* 1992; Iannaccone *et al.* 2003, 2004; Koenekoop *et al.* 2003; Zito *et al.* 2003; Moore *et al.* 2006). RPGR is predominantly located at the primary cilium of photoreceptors and multiple different isoforms are recognized (Hong *et al.* 2003; Khanna *et al.* 2005; He *et al.* 2008); however two major ones are recognized: RPGR^{ORF15} and RPGR^{EX1–19} (Vervoort *et al.* 2000; Breuer *et al.* 2002).

Expression and localization of RPGR protein isoforms

Complex-splicing patterns are reported for *RPGR* though the physiological relevance of these transcripts is unclear (Yan *et al.* 1998; Kirschner *et al.* 1999; Vervoort *et al.* 2000; Hong and Li 2002; Ferreira 2005). Purine-rich exon splicing enhancers in ORF15 may also modify the efficiency of splicing (Hong and Li 2002). Multiple immunoreactive bands are observed using isoform-specific RPGR antibodies (Yan *et al.* 1998; Hong and Li 2002; Mavlyutov *et al.* 2002; He *et al.* 2008; Khanna *et al.* 2005; Otto *et al.* 2005; Shu *et al.* 2005; Chang *et al.* 2006). The constitutive RPGR^{EX1–19} isoform is isoprenylated and localizes to golgi in transfected cells (Yan *et al.* 1998). The amino-terminal region of RPGR encoded by exons 1–15 encompasses a common RCC1-like domain (Meindl *et al.* 1996; Vervoort *et al.* 2000) (figure 2). Although RCC1 is a guanine nucleotide exchange factor (GEF) for Ran GTPases (Meindl *et al.* 1996; Renault *et al.* 2001; Sazer *et al.* 2005), no such activity has yet been reported for RPGR. The RPGR^{ORF15} isoforms that include a C-terminal acidic domain rich in Glu–Gly repeats (Vervoort *et al.* 2000) are reported to have distinct subcellular localizations (Mavlyutov *et al.* 2002; Hong *et al.* 2003). In addition, multiple predicted

homology domains can also be detected in the primary structure of the RPGR protein. RPGR^{ORF15} localizes predominantly to the cilia and basal bodies of both mouse and human photoreceptors though some additional labelling can be found in the inner and outer segments (Khanna *et al.* 2005; Shu *et al.* 2005). In proliferating cells, centrosomes and midbody are labelled with anti-RPGR antibodies (Chang *et al.* 2006; Shu *et al.* 2005). Our analysis of RPGR using isoform-specific antibodies revealed that the RPGR^{EX1-19} and RPGR^{ORF15} isoforms localize to overlapping as well as distinct subcellular compartments in the retina (He *et al.* 2008). Moreover, the RPGR^{EX1-19} isoforms exist in overlapping protein complexes with RPGR^{ORF15}.

Animal models of RPGR

A *Rpgr*-knockout (ko) mouse with deletion of exons 4–6 of *Rpgr* is reported to show late-onset cone-rod degeneration (Hong *et al.* 2000); however, this *Rpgr*-ko mouse is not a complete null and expresses some RPGR^{ORF15} isoforms that localize to the photoreceptor cilium (Khanna *et al.* 2005). The phenotype of this mouse can be partially rescued by an ORF15-variant (Hong *et al.* 2005). However, the same variant in transgenic mice may result in dominant gain-of-function phenotype exhibiting rapid disease progression (Hong *et al.* 2004). Attempts to generate a complete *Rpgr* null mutation in mice showing no RPGR isoforms have so far been futile. RPGR^{ORF15} frameshift mutations have been identified in two naturally occurring canine mutants; the XLPRA2 dog exhibits relatively rapid photoreceptor degeneration and severe ERG abnormalities, whereas the XLPRA1 mutant shows a milder phenotype (Zhang *et al.* 2002; Beltran *et al.* 2006).

RPGR-interactome in photoreceptors

Until 2005, two RPGR-interacting proteins were identified by Y2H analysis using the RLD bait: RPGRIP1, which is localized to the sensory cilium and mutated in patients with LCA (Boylan and Wright 2000; Dryja *et al.* 2001; Hong *et al.* 2001); and PDE6d, a prenyl-binding protein involved in the retrieval of PDE from rod outer segment membranes by interacting with the GTPase Rab13 (Linari *et al.* 1999; Zhang *et al.* 2004). Two chromosome-associated proteins, SMC1 and SMC3 and a centrosomal protein, nucleophosmin (NPM) (Shu *et al.* 2005) directly interact with RPGR (Khanna *et al.* 2005; Hirano 2006). Later IP studies identified intraflagellar transport protein, Tg737/Polaris/IFT88, which is involved in ciliogenesis and photoreceptor outer segment maintenance in mice (Pazour *et al.* 2002; Davenport and Yoder 2005) and with several microtubule transport proteins, including cytoplasmic dynein and kinesin-II as part of RPGR containing protein complexes in the retina (Khanna *et al.* 2005). Studies on the analysis of selected RPGR-interacting ciliary disease proteins are described below.

RPGR-interacting ciliary disease proteins

RPGRIP1

The RPGR interacting protein 1 (RPGRIP1) was first described as an interactor of RPGR after screening a bovine retina library by yeast two-hybrid analysis (Boylan and Wright 2000; Roepman *et al.* 2000). Initial analysis showed that RPGR and RPGRIP1 can co-localize from the connecting cilium towards the outer segment of bovine and human photoreceptors, and disease associated mutations in *RPGR* can disrupt this interaction. Subsequent studies revealed that mutations in RPGRIP1 are associated with LCA (Dryja *et al.* 2001; Gerber *et al.* 2001). RPGRIP1 is predominantly localized at the connecting cilium of mouse photoreceptors (Hong *et al.* 2001), and the *Rpgrip1* ko mice features early onset retinal degeneration (Zhao *et al.* 2003). Analysis of the *Rpgrip1*^{-/-} mice revealed redistribution of RPGR in the inner segments as opposed to the sensory cilium of

photoreceptors, suggesting that RPGRIP1 functions as an anchoring protein for RPGR (Hong *et al.* 2001; Zhao *et al.* 2003).

NPHP5

Nephronophthisis (NPHP) is characterized by progressive loss of renal function when presented with RP is termed Senior–Loken syndrome (SLSN). Nine NPHP genes (NPHP1–9) have been identified; all are implicated in ciliary function (Hildebrandt and Otto 2005; Chang *et al.* 2006; Khanna *et al.* 2009). All patients with *NPHP5* mutations develop RP (Otto *et al.* 2005). We have shown that NPHP5 localizes to photoreceptor cilium and interacts with RPGR, highlighting their functional interaction in ciliary transport (Otto *et al.* 2005).

CEP290/NPHP6

CEP290 is a cilia-centrosomal protein that was initially identified as a tumour antigen (Guo *et al.* 2004). Centrosomes are the major microtubule-organizing centre for the cell (Doxsey *et al.* 2005) and mutations in centrosomal proteins are associated with syndromic diseases like Alstrom syndrome (MIM 203800) or complex pathological processes like cancer (Andersen *et al.* 2003; Badano *et al.* 2005). CEP290 is not the exception; mutations in the *CEP290/NPHP6* are associated with syndromic disorders, including Joubert syndrome, Meckel–Gruber syndrome, and BBS (Sayer *et al.* 2006; Valente *et al.* 2006; Baala *et al.* 2007; Leitch *et al.* 2008). We showed that a hypomorphic allele of the *Cep290* gene in the *rd16* (retinal degeneration 16) mouse model of autosomal recessive early-onset retinal degeneration and olfactory dysfunction (Chang *et al.* 2006). The mutant CEP290 protein exhibits increased association with RPGR and likely results in the mislocalization of RPGR in photoreceptors (Chang *et al.* 2006).

Based on these studies, den Hollander *et al.* and others screened LCA patients for mutations in the *CEP290* gene and reported predicted hypomorphic alleles as a frequent cause of LCA (den Hollander *et al.* 2006; Helou *et al.* 2007; Perrault *et al.* 2007). We later showed that hypomorphic *CEP290* alleles are also associated with olfactory dysfunction in patients (McEwen *et al.* 2007). Taken together, we suggest that loss of function mutations in CEP290 can cause pleiotropic and developmental disorders while a moderately functional CEP290 protein primarily results in sensory deficits.

RPGRIP1L/NPHP8

Mutations in RPGR interacting protein 1-like (RP-GRIP1L/NPHP8) are associated with Joubert syndrome, Meckel–Gruber syndrome and Bardet–Biedl syndrome (Arts *et al.* 2007; Delous *et al.* 2007; Wolf *et al.* 2007; Brancati *et al.* 2008; Doherty *et al.* 2009). RPGRIP1L regulates sonic hedgehog signalling pathway and mediates left–right asymmetry as well as limb patterning. Interestingly, a missense variant of RPGRIP1L and A229T is frequently associated with retinal degeneration in ciliopathy patients (Khanna *et al.* 2009). RPGRIP1L can physically interact with RPGR, and the A229T in RPGRIP1L variation severely compromises this interaction (Khanna *et al.* 2009). These data suggest that variations in NPHP8 can act as modifier alleles that affect the penetrance and expressivity of the retinopathy phenotype, likely through its interaction with RPGR.

Conclusions and future directions

Over 200 different RPGR mutations have so far been reported in patients with X-linked retinopathies of diverse clinical phenotypes. A vast majority of these are nonsense mutations or deletions/insertions resulting in frameshift, which are predicted to cause premature truncation of the RPGR^{ORF15} protein. While mutations in the ORF15 exon are generally

associated with a milder disease, mutations in RPGR exons 1–14 result in a more severe disease. Initially, most human RPGR mutations were hypothesized to have a null phenotype in males; however, wide variations in clinical phenotype of males and carrier females, and complexities associated with RPGR transcripts and protein isoforms strongly indicate that several disease alleles may in fact be hypomorphs (with partial function).

What is the role of RPGR as part of distinct multiprotein complexes at the cilium and how do mutations in RPGR cause photoreceptor degeneration? Though basic components associated with ciliary transport have been discovered (Rosenbaum *et al.* 1999; Rosenbaum 2002), the mechanisms of cargo sorting and assembly of protein complexes, and their regulation by signalling pathways in photoreceptors have not been elucidated. Periodic turnover of photoreceptor outer segments demands efficient functioning of the ciliary transport process (Besharse *et al.* 2003). We hypothesize that RPGR facilitates the assembly of transport protein complexes by interacting with distinct ciliary—basal body—centrosome (CBC) proteins, and that RPGR's localization in the photoreceptor cilia is necessary for efficient intersegmental transport (figure 3).

Owing to the associated clinical heterogeneity, a detailed genotype–phenotype correlation analysis is critical to understand the progression and pathogenesis of RPGR-associated disease. These studies will benefit from characterization of additional animal model systems representing RPGR mutations as well as identification of components of the RPGR-interactome in photoreceptors. For example, characterization of photoreceptor dysfunction and degeneration in knock-in mouse mutants of *Rpgr* can assist in understanding associated disease pathogenesis. In addition, functional analysis of the disease-causing mutations of RPGR, such as the effect on RPGR localization, integrity of the interactome and effect on cilia-dependent development are required to delineate the mechanism of heterogenic phenotype observed in patients. These investigations should assist in designing rational therapeutic paradigms for XLRP as well as associated ciliary disorders.

Acknowledgments

This work is supported by National Eye Institute intramural funds and grants from the National Institutes of Health (EY007961), Foundation Fighting Blindness (FFB), and Midwest Eye Banks and Transplantation Center.

References

- Andersen JS, Wilkinson CJ, Mayor T, Mortensen P, Nigg EA, Mann M. Proteomic characterization of the human centrosome by protein correlation profiling. *Nature*. 2003; 426:570–574. [PubMed: 14654843]
- Arts HH, Doherty D, van Beersum SE, Parisi MA, Letteboer SJ, Gordon NT, et al. Mutations in the gene encoding the basal body protein RPGRIP1L, a nephrocystin-4 interactor, cause Joubert syndrome. *Nat. Genet.* 2007; 39:882–888. [PubMed: 17558407]
- Baala L, Audollent S, Martinovic J, Ozilou C, Babron MC, Sivanandamoorthy S, et al. Pleiotropic effects of CEP290 (NPHP6) mutations extend to Meckel syndrome. *Am. J. Hum. Genet.* 2007; 81:170–179. [PubMed: 17564974]
- Badano JL, Teslovich TM, Katsanis N. The centrosome in human genetic disease. *Nat. Rev. Genet.* 2005; 6:194–205. [PubMed: 15738963]
- Bartolini F, Bhamidipati A, Thomas S, Schwahn U, Lewis SA, Cowan NJ. Functional overlap between retinitis pigmentosa 2 protein and the tubulin-specific chaperone cofactor C. *J. Biol. Chem.* 2002; 277:14629–14634. [PubMed: 11847227]
- Beltran WA, Hammond P, Acland GM, Aguirre GD. A frameshift mutation in RPGR exon ORF15 causes photoreceptor degeneration and inner retina remodeling in a model of X-linked retinitis pigmentosa. *Invest. Ophthalmol. Vis. Sci.* 2006; 47:1669–1681. [PubMed: 16565408]

- Besharse, JC. The retina: a model for cell biological studies part I. New York, USA: Academic Press; 1986.
- Besharse JC, Baker SA, Luby-Phelps K, Pazour GJ. Photoreceptor intersegmental transport and retinal degeneration: a conserved pathway common to motile and sensory cilia. *Adv. Exp. Med. Biol.* 2003; 533:157–164. [PubMed: 15180260]
- Bird AC. X-linked retinitis pigmentosa. *BrJ. Ophthalmol.* 1975; 59:177–199.
- Boylan JP, Wright AF. Identification of a novel protein interacting with RPGR. *Hum. Mol. Genet.* 2000; 9:2085–2093. [PubMed: 10958647]
- Brancati F, Travaglini L, Zablocka D, Boltshauser E, Accorsi P, Montagna G, et al. RPGRIP1L mutations are mainly associated with the cerebello-renal phenotype of Joubert syndrome-related disorders. *Clin. Genet.* 2008; 74:164–170. [PubMed: 18565097]
- Breuer DK, Yashar BM, Filippova E, Hiriyanna S, Lyons RH, Mears AJ, et al. A comprehensive mutation analysis of RP2 and RPGR in a North American cohort of families with X-linked retinitis pigmentosa. *Am. J. Hum. Genet.* 2002; 70:1545–1554. [PubMed: 11992260]
- Chang B, Khanna H, Hawes N, Jimeno D, He S, Lillo C, et al. In-frame deletion in a novel centrosomal/ciliary protein CEP290/NPHP6 perturbs its interaction with RPGR and results in early-onset retinal degeneration in the rd16 mouse. *Hum. Mol. Genet.* 2006; 15:1847–1857. [PubMed: 16632484]
- Chapple JP, Hardcastle AJ, Grayson C, Spackman LA, Willison KR, Cheetham ME. Mutations in the N-terminus of the X-linked retinitis pigmentosa protein RP2 interfere with the normal targeting of the protein to the plasma membrane. *Hum. Mol. Genet.* 2000; 9:1919–1926. [PubMed: 10942419]
- Chuang JZ, Sung CH. The cytoplasmic tail of rhodopsin acts as a novel apical sorting signal in polarized MDCK cells. *J. Cell Biol.* 1998; 142:1245–1256. [PubMed: 9732285]
- Chuang JZ, Zhao Y, Sung CH. SARA-regulated vesicular targeting underlies formation of the light-sensing organelle in mammalian rods. *Cell.* 2007; 130:535–547. [PubMed: 17693260]
- Cideciyan AV, Jacobson SG. Image analysis of the tapetal-like reflex in carriers of X-linked retinitis pigmentosa. *Invest. Ophthalmol. Vis. Sci.* 1994; 35:3812–3824. [PubMed: 7928178]
- Colley NJ, Cassill JA, Baker EK, Zuker CS. Defective intracellular transport is the molecular basis of rhodopsindependent dominant retinal degeneration. *Proc. Natl. Acad. Sci. USA.* 1995; 92:3070–3074. [PubMed: 7708777]
- Daiger SP, Bowne SJ, Sullivan LS. Perspective on genes and mutations causing retinitis pigmentosa. *Arch. Ophthalmol.* 2007; 125:151–158. [PubMed: 17296890]
- Davenport JR, Yoder BK. An incredible decade for the primary cilium: a look at a once-forgotten organelle. *Am. J. Physiol. Renal. Physiol.* 2005; 289:F1159–F1169. [PubMed: 16275743]
- Delous M, Baala L, Salomon R, Laclef C, Vierkotten J, Tory K, et al. The ciliary gene RPGRIP1L is mutated in cerebellooculo-renal syndrome (Joubert syndrome type B) and Meckel syndrome. *Nat. Genet.* 2007; 39:875–881. [PubMed: 17558409]
- den Hollander AI, Koenekoop RK, Yzer S, Lopez I, Arends ML, Voeselek KE, et al. Mutations in the CEP290 (NPHP6) gene are a frequent cause of Leber congenital amaurosis. *Am. J. Hum. Genet.* 2006; 79:556–561. [PubMed: 16909394]
- Deretic D, Huber LA, Ransom N, Mancini M, Simons K, Papermaster DS. rab8 in retinal photoreceptors may participate in rhodopsin transport and in rod outer segment disk morphogenesis. *J. Cell Sci.* 1995; 108:215–224. [PubMed: 7738098]
- Deretic D, Schmerl S, Hargrave PA, Arendt A, McDowell JH. Regulation of sorting and post-Golgi trafficking of rhodopsin by its C-terminal sequence QVS(A)PA. *Proc. Natl. Acad. Sci. USA.* 1998; 95:10620–10625. [PubMed: 9724753]
- Doherty D, Parisi MA, Finn LS, Gunay-Aygun M, Al-Mateen M, Bates D, et al. Mutations in 3 genes (MKS3, CC2D2A and RPGRIP1L) cause COACH syndrome (Joubert syndrome with congenital hepatic fibrosis). *J. Med. Genet.* 2009
- Doxsey S. Re-evaluating centrosome function. *Nat. Rev. Mol. Cell Biol.* 2001; 2:688–698. [PubMed: 11533726]
- Doxsey S, McCollum D, Theurkauf W. Centrosomes in cellular regulation. *Annu. Rev. Cell Dev. Biol.* 2005; 21:411–434. [PubMed: 16212501]

- Dryja TP, Adams SM, Grimsby JL, McGee TL, Hong DH, Li T, et al. Null RPGRIP1 alleles in patients with Leber congenital amaurosis. *Am. J. Hum. Genet.* 2001; 68:1295–1298. [PubMed: 11283794]
- Ferreira PA. Insights into X-linked retinitis pigmentosa type 3, allied diseases and underlying pathomechanisms. *Hum. Mol. Genet.* 2005; 14:R259–R267. [PubMed: 16244324]
- Fishman GA. Retinitis pigmentosa. Genetic percentages. *Arch. Ophthalmol.* 1978; 96:822–826. [PubMed: 655919]
- Fishman GA, Weinberg AB, McMahon TT. X-linked recessive retinitis pigmentosa. Clinical characteristics of carriers. *Arch. Ophthalmol.* 1986; 104:1329–1335. [PubMed: 3753283]
- Fishman GA, Farber MD, Derlacki DJ. X-linked retinitis pigmentosa. Profile of clinical findings. *Arch. Ophthalmol.* 1988; 106:369–375. [PubMed: 3257866]
- Follit JA, Tuft RA, Fogarty KE, Pazour GJ. The intraflagellar transport protein IFT20 is associated with the Golgi complex and is required for cilia assembly. *Mol. Biol. Cell.* 2006; 17:3781–3792. [PubMed: 16775004]
- Fujita R, Bingham E, Forsythe P, McHenry C, Aita V, Navia BA, et al. A recombination outside the BB deletion refines the location of the X linked retinitis pigmentosa locus RP3. *Am. J. Hum. Genet.* 1996; 59:152–158. [PubMed: 8659520]
- Gerber S, Perrault I, Hanein S, Barbet F, Ducrocq D, Ghazi I, et al. Complete exon-intron structure of the RPGR-interacting protein (RPGRIP1) gene allows the identification of mutations underlying Leber congenital amaurosis. *Eur J. Hum. Genet.* 2001; 9:561–571.
- Gerdes JM, Davis EE, Katsanis N. The vertebrate primary cilium in development, homeostasis, and disease. *Cell.* 2009; 137:32–45. [PubMed: 19345185]
- Gieser L, Fujita R, Goring HH, Ott J, Hoffman DR, Cideciyan AV, et al. A novel locus (RP24) for X-linked retinitis pigmentosa maps to Xq26-27. *Am. J. Hum. Genet.* 1998; 63:1439–1447. [PubMed: 9792872]
- Grayson C, Bartolini F, Chapple JP, Willison KR, Bhamidipati A, Lewis SA, et al. Localization in the human retina of the X-linked retinitis pigmentosa protein RP2, its homologue cofactor C and the RP2 interacting protein Arl3. *Hum. Mol. Genet.* 2002; 11:3065–3074. [PubMed: 12417528]
- Guo J, Jin G, Meng L, Ma H, Nie D, Wu J, et al. Subcellular localization of tumor-associated antigen 3H11Ag. *Biochem. Biophys. Res. Commun.* 2004; 324:922–930. [PubMed: 15474516]
- Hardcastle AJ, Thiselton DL, Van Maldergem L, Saha BK, Jay M, Plant C, et al. Mutations in the RP2 gene cause disease in 10% of families with familial X-linked retinitis pigmentosa assessed in this study. *Am. J. Hum. Genet.* 1999; 64:1210–1215. [PubMed: 10090907]
- Hardcastle AJ, Thiselton DL, Zito I, Ebenezer N, Mah TS, Gorin MB, Bhattacharya SS. Evidence for a new locus for X-linked retinitis pigmentosa (RP23). *Invest. Ophthalmol. Vis. Sci.* 2000; 41:2080–2086. [PubMed: 10892847]
- He S, Parapuram SK, Hurd TW, Behnam B, Margolis B, Swaroop A, Khanna H. Retinitis pigmentosa GTPase regulator (RPGR) protein isoforms in mammalian retina: Insights into X-linked retinitis pigmentosa and associated ciliopathies. *Vision Res.* 2008; 48:366–376. [PubMed: 17904189]
- Helou J, Otto EA, Attanasio M, Allen SJ, Parisi MA, Glass I, et al. Mutation analysis of NPHP6/CEP290 in patients with Joubert syndrome and Senior-Loken syndrome. *J. Med. Genet.* 2007; 44:657–663. [PubMed: 17617513]
- Hildebrandt F, Otto E. Cilia and centrosomes: a unifying pathogenic concept for cystic kidney disease? *Nat. Rev. Genet.* 2005; 6:928–940. [PubMed: 16341073]
- Hirano T. At the heart of the chromosome: SMC proteins in action. *Nat. Rev. Mol. Cell Biol.* 2006; 7:311–322. [PubMed: 16633335]
- Hong DH, Li T. Complex expression pattern of RPGR reveals a role for purine-rich exonic splicing enhancers. *Invest. Ophthalmol. Vis. Sci.* 2002; 43:3373–3382. [PubMed: 12407146]
- Hong DH, Pawlyk BS, Shang J, Sandberg MA, Berson EL, Li T. A retinitis pigmentosa GTPase regulator (RPGR)-deficient mouse model for X-linked retinitis pigmentosa (RP3). *Proc. Natl. Acad. Sci. USA.* 2000; 97:3649–3654. [PubMed: 10725384]
- Hong DH, Yue G, Adamian M, Li T. Retinitis pigmentosa GTPase regulator (RPGR)-interacting protein is stably associated with the photoreceptor ciliary axoneme and anchors RPGR to the connecting cilium. *J. Biol. Chem.* 2001; 276:12091–12099. [PubMed: 11104772]

- Hong DH, Pawlyk B, Sokolov M, Strissel KJ, Yang J, Tulloch B, et al. RPGR isoforms in photoreceptor connecting cilia and the transitional zone of motile cilia. *Invest. Ophthalmol. Vis. Sci.* 2003; 44:2413–2421. [PubMed: 12766038]
- Hong DH, Pawlyk BS, Adamian M, Li T. Dominant, gain-of-function mutant produced by truncation of RPGR. *Invest. Ophthalmol. Vis. Sci.* 2004; 45:36–41. [PubMed: 14691151]
- Hong DH, Pawlyk BS, Adamian M, Sandberg MA, Li T. A single, abbreviated RPGR-ORF15 variant reconstitutes RPGR function *in vivo*. *Invest. Ophthalmol. Vis. Sci.* 2005; 46:435–441. [PubMed: 15671266]
- Hunter DG, Fishman GA, Kretzer FL. Abnormal axonemes in X-linked retinitis pigmentosa. *Arch. Ophthalmol.* 1988; 106:362–368. [PubMed: 3345154]
- Iannaccone A, Breuer DK, Wang XF, Kuo SF, Normando EM, Filippova E, et al. Clinical and immunohistochemical evidence for an X linked retinitis pigmentosa syndrome with recurrent infections and hearing loss in association with an RPGR mutation. *J. Med. Genet.* 2003; 40:e118. [PubMed: 14627685]
- Iannaccone A, Wang X, Jablonski MM, Kuo SF, Baldi A, Cosgrove D, et al. Increasing evidence for syndromic phenotypes associated with RPGR mutations. *Am. J. Ophthalmol.* 2004; 137:785–786. author reply 786. [PubMed: 15059739]
- Jay M. Figures and fantasies: the frequencies of the different genetic forms of retinitis pigmentosa. *Birth Defects, Orig. Artic. Ser.* 1982; 18:167–173. [PubMed: 7171752]
- Kahn RA, Volpicelli-Daley L, Bowzard B, Shrivastava-Ranjan P, Li Y, Zhou C, Cunningham L. Arf family GTPases: roles in membrane traffic and microtubule dynamics. *Biochem. Soc. Trans.* 2005; 33:1269–1272. [PubMed: 16246095]
- Khanna H, Hurd TW, Lillo C, Shu X, Parapuram SK, He S, et al. RPGR-ORF15, which is mutated in retinitis pigmentosa, associates with SMC1, SMC3, and microtubule transport proteins. *J. Biol. Chem.* 2005; 280:33580–33587. [PubMed: 16043481]
- Khanna H, Davis EE, Murga-Zamalloa CA, Estrada-Cuzcano A, Lopez I, den Hollander AI, et al. A common allele in RPGRIP1L is a modifier of retinal degeneration in ciliopathies. *Nat. Genet.* 2009; 41:739–745. [PubMed: 19430481]
- Kirschner R, Rosenberg T, Schultz-Heienbrok R, Lenzner S, Feil S, Roepman R, et al. RPGR transcription studies in mouse and human tissues reveal a retina-specific isoform that is disrupted in a patient with X-linked retinitis pigmentosa. *Hum. Mol. Genet.* 1999; 8:1571–1578. [PubMed: 10401007]
- Koenekoop RK, Loyer M, Hand CK, Al Mahdi H, Dembinska O, Beneish R, et al. Novel RPGR mutations with distinct retinitis pigmentosa phenotypes in French–Canadian families. *Am. J. Ophthalmol.* 2003; 136:678–687. [PubMed: 14516808]
- Kuhnel K, Veltel S, Schlichting I, Wittinghofer A. Crystal structure of the human retinitis pigmentosa 2 protein and its interaction with Arl3. *Structure.* 2006; 14:367–378. [PubMed: 16472755]
- Leitch CC, Zaghoul NA, Davis EE, Stoetzel C, Diaz-Font A, Rix S, et al. Hypomorphic mutations in syndromic encephalocele genes are associated with Bardet–Biedl syndrome. *Nat. Genet.* 2008; 40:443–448. [PubMed: 18327255]
- Linari M, Ueffing M, Manson F, Wright A, Meitinger T, Becker J. The retinitis pigmentosa GTPase regulator, RPGR, interacts with the delta subunit of rod cyclic GMP phosphodiesterase. *Proc. Natl. Acad. Sci. USA.* 1999; 96:1315–1320. [PubMed: 9990021]
- Marszalek JR, Liu X, Roberts EA, Chui D, Marth JD, Williams DS, Goldstein LS. Genetic evidence for selective transport of opsin and arrestin by kinesin-II in mammalian photoreceptors. *Cell.* 2000; 102:175–187. [PubMed: 10943838]
- Mavlyutov TA, Zhao H, Ferreira PA. Species-specific subcellular localization of RPGR and RPGRIP isoforms: implications for the phenotypic variability of congenital retinopathies among species. *Hum. Mol. Genet.* 2002; 11:1899–1907. [PubMed: 12140192]
- Mazelova J, Ransom N, Astuto-Gribble L, Wilson MC, Deretic D. Syntaxin 3 and SNAP-25 pairing, regulated by omega-3 docosahexaenoic acid, controls the delivery of rhodopsin for the biogenesis of cilia-derived sensory organelles, the rod outer segments. *J. Cell Sci.* 2009; 122:2003–2013. [PubMed: 19454479]

- McEwen DP, Koenekoop RK, Khanna H, Jenkins PM, Lopez I, Swaroop A, Martens JR. Hypomorphic CEP290/NPHP6 mutations result in anosmia caused by the selective loss of G proteins in cilia of olfactory sensory neurons. *Proc. Natl. Acad. Sci. USA*. 2007; 104:15917–15922. [PubMed: 17898177]
- McGuire RE, Sullivan LS, Blanton SH, Church MW, Heckenlively JR, Daiger SP. X-linked dominant cone-rod degeneration: linkage mapping of a new locus for retinitis pigmentosa (RP 15) to Xp22.13- p22.11. *Am. J. Hum. Genet.* 1995; 57:87–94. [PubMed: 7611300]
- Mears AJ, Gieser L, Yan D, Chen C, Fahrner S, Hiriyan S, et al. Protein-truncation mutations in the RP2 gene in a North American cohort of families with X-linked retinitis pigmentosa. *Am. J. Hum. Genet.* 1999; 64:897–900. [PubMed: 10053026]
- Meindl A, Dry K, Herrmann K, Manson F, Ciccodicola A, Edgar A, et al. A gene (RPGR) with homology to the RCC1 guanine nucleotide exchange factor is mutated in X-linked retinitis pigmentosa (RP3). *Nat. Genet.* 1996; 13:35–42. [PubMed: 8673101]
- Melamud A, Shen GQ, Chung D, Xi Q, Simpson E, Li L, et al. Mapping a new genetic locus for X linked retinitis pigmentosa to Xq28. *J. Med. Genet.* 2006; 43:e27. [PubMed: 16740911]
- Moore A, Escudier E, Roger G, Tamalet A, Pelosse B, Marlin S, et al. RPGR is mutated in patients with a complex X linked phenotype combining primary ciliary dyskinesia and retinitis pigmentosa. *J. Med. Genet.* 2006; 43:326–333. [PubMed: 16055928]
- Moritz OL, Tam BM, Hurd LL, Peranen J, Deretic D, Papermaster DS. Mutant rab8 Impairs docking and fusion of rhodopsin-bearing post-Golgi membranes and causes cell death of transgenic *Xenopus* rods. *Mol. Biol. Cell.* 2001; 12:2341–2351. [PubMed: 11514620]
- Nair KS, Hanson SM, Mendez A, Gurevich EV, Kennedy MJ, Shestopalov VI, et al. Light-dependent redistribution of arrestin in vertebrate rods is an energy-independent process governed by protein-protein interactions. *Neuron.* 2005; 46:555–567. [PubMed: 15944125]
- Otto EA, Loeys B, Khanna H, Hellemans J, Sudbrak R, Fan S, et al. Nephrocystin-5, a ciliary IQ domain protein, is mutated in Senior-Loken syndrome and interacts with RPGR and calmodulin. *Nat. Genet.* 2005; 37:282–288. [PubMed: 15723066]
- Pazour GJ, Witman GB. The vertebrate primary cilium is a sensory organelle. *Curr. Opin. Cell Biol.* 2003; 15:105–110. [PubMed: 12517711]
- Pazour GJ, Baker SA, Deane JA, Cole DG, Dickert BL, Rosenbaum JL, et al. The intraflagellar transport protein, IFT88, is essential for vertebrate photoreceptor assembly and maintenance. *J. Cell Biol.* 2002; 157:103–113. [PubMed: 11916979]
- Perrault I, Delphin N, Hanein S, Gerber S, Dufier JL, Roche O, et al. Spectrum of NPHP6/CEP290 mutations in Leber congenital amaurosis and delineation of the associated phenotype. *Hum. Mutat.* 2007; 28:416. [PubMed: 17345604]
- Renault L, Kuhlmann J, Henkel A, Wittinghofer A. Structural basis for guanine nucleotide exchange on Ran by the regulator of chromosome condensation (RCC1). *Cell.* 2001; 105:245–255. [PubMed: 11336674]
- Roepman R, van Duijnhoven G, Rosenberg T, Pinckers AJ, Bleeker-Wagemakers LM, Bergen AA, et al. Positional cloning of the gene for X-linked retinitis pigmentosa 3: homology with the guanine-nucleotide-exchange factor RCC1. *Hum. Mol. Genet.* 1996; 5:1035–1041. [PubMed: 8817343]
- Roepman R, Bernoud-Hubac N, Schick DE, Maugeri A, Berger W, Ropers HH, et al. The retinitis pigmentosa GTPase regulator (RPGR) interacts with novel transport-like proteins in the outer segments of rod photoreceptors. *Hum. Mol. Genet.* 2000; 9:2095–2105. [PubMed: 10958648]
- Rosenbaum J. Intraflagellar transport. *Curr. Biol.* 2002; 12:R125. [PubMed: 11864582]
- Rosenbaum JL, Cole DG, Diener DR. Intraflagellar transport: the eyes have it. *J. Cell Biol.* 1999; 144:385–388. [PubMed: 9971734]
- Sayer JA, Otto EA, O'Toole JF, Nurnberg G, Kennedy MA, Becker C, et al. The centrosomal protein nephrocystin-6 is mutated in Joubert syndrome and activates transcription factor ATF4. *Nat. Genet.* 2006; 38:674–681. [PubMed: 16682973]
- Sazer S. The view from Awaji island: past, present, and future of RCC1 and the Ran GTPase system. *Dev. Cell.* 2005; 9:729–733. [PubMed: 16363093]
- Scholey JM, Anderson KV. Intraflagellar transport and cilium-based signaling. *Cell.* 2006; 125:439–442. [PubMed: 16678091]

- Schrick JJ, Vogel P, Abuin A, Hampton B, Rice DS. ADP-ribosylation factor-like 3 is involved in kidney and photoreceptor development. *Am. J. Pathol.* 2006; 168:1288–1298. [PubMed: 16565502]
- Schwahn U, Lenzner S, Dong J, Feil S, Hinzmann B, van Duijnhoven G, et al. Positional cloning of the gene for X-linked retinitis pigmentosa 2. *Nat. Genet.* 1998; 19:327–332. [PubMed: 9697692]
- Sharon D, Bruns GA, McGee TL, Sandberg MA, Berson EL, Dryja TP. X-linked retinitis pigmentosa: mutation spectrum of the RPGR and RP2 genes and correlation with visual function. *Invest. Ophthalmol. Vis. Sci.* 2000; 41:2712–2721. [PubMed: 10937588]
- Sharon D, Sandberg MA, Rabe VW, Stillberger M, Dryja TP, Berson EL. RP2 and RPGR mutations and clinical correlations in patients with X-linked retinitis pigmentosa. *Am. J. Hum. Genet.* 2003; 73:1131–1146. [PubMed: 14564670]
- Shu X, Fry AM, Tulloch B, Manson FD, Crabb JW, Khanna H, et al. RPGR ORF15 isoform co-localizes with RPGRIP1 at centrioles and basal bodies and interacts with nucleophosmin. *Hum. Mol. Genet.* 2005; 14:1183–1197. [PubMed: 15772089]
- Shu X, Black GC, Rice JM, Hart-Holden N, Jones A, O'Grady A, et al. RPGR mutation analysis and disease: an update. *Hum. Mutat.* 2007; 28:322–328. [PubMed: 17195164]
- Sieving PA. Diagnostic issues with inherited retinal and macular dystrophies. *Semin. Ophthalmol.* 1995; 10:279–294. [PubMed: 10160215]
- Sokolov M, Lyubarsky AL, Strissel KJ, Savchenko AB, Govardovskii VI, Pugh EN Jr, Arshavsky VY. Massive light-driven translocation of transducin between the two major compartments of rod cells: a novel mechanism of light adaptation. *Neuron.* 2002; 34:95–106. [PubMed: 11931744]
- Spektor A, Tsang WY, Khoo D, Dynlacht BD. Cep97 and CP110 suppress a cilia assembly program. *Cell.* 2007; 130:678–690. [PubMed: 17719545]
- Strissel KJ, Lishko PV, Trieu LH, Kennedy MJ, Hurley JB, Arshavsky VY. Recoverin undergoes light-dependent intracellular translocation in rod photoreceptors. *J. Biol. Chem.* 2005; 280:29250–29255. [PubMed: 15961391]
- Strissel KJ, Sokolov M, Trieu LH, Arshavsky VY. Arrestin translocation is induced at a critical threshold of visual signaling and is superstoichiometric to bleached rhodopsin. *J. Neurosci.* 2006; 26:1146–1153. [PubMed: 16436601]
- Sung CH, Makino C, Baylor D, Nathans J. A rhodopsin gene mutation responsible for autosomal dominant retinitis pigmentosa results in a protein that is defective in localization to the photoreceptor outer segment. *J. Neurosci.* 1994; 14:5818–5833. [PubMed: 7523628]
- Tsang WY, Bossard C, Khanna H, Peranen J, Swaroop A, Malhotra V, Dynlacht BD. CP110 suppresses primary cilia formation through its interaction with CEP290, a protein deficient in human ciliary disease. *Dev. Cell.* 2008; 15:187–197. [PubMed: 18694559]
- Valente EM, Silhavy JL, Brancati F, Barrano G, Krishnaswami SR, Castori M, et al. Mutations in CEP290, which encodes a centrosomal protein, cause pleiotropic forms of Joubert syndrome. *Nat. Genet.* 2006; 38:623–625. [PubMed: 16682970]
- van Dorp DB, Wright AF, Carothers AD, Bleeker-Wagemakers EM. A family with RP3 type of X-linked retinitis pigmentosa: an association with ciliary abnormalities. *Hum. Genet.* 1992; 88:331–334. [PubMed: 1733835]
- Vervoort R, Wright AF. Mutations of RPGR in X-linked retinitis pigmentosa (RP3). *Hum. Mutat.* 2002; 19:486–500. [PubMed: 11968081]
- Vervoort R, Lennon A, Bird AC, Tulloch B, Axton R, Miano MG, et al. Mutational hot spot within a new RPGR exon in X-linked retinitis pigmentosa. *Nat. Genet.* 2000; 25:462–466. [PubMed: 10932196]
- Wolf MT, Saunier S, O'Toole JF, Wanner N, Groshong T, Attanasio M, et al. Mutational analysis of the RPGRIP1L gene in patients with Joubert syndrome and nephronophthisis. *Kidney Int.* 2007; 72:1520–1526. [PubMed: 17960139]
- Wright AF, Bhattacharya SS, Aldred MA, Jay M, Carothers AD, Thomas NS, et al. Genetic localisation of the RP2 type of X linked retinitis pigmentosa in a large kindred. *J. Med. Genet.* 1991; 28:453–457. [PubMed: 1895315]

- Yan D, Swain PK, Breuer D, Tucker RM, Wu W, Fujita R, et al. Biochemical characterization and subcellular localization of the mouse retinitis pigmentosa GTPase regulator (mRpg). *J. Biol. Chem.* 1998; 273:19656–19663. [PubMed: 9677393]
- Young RW. Passage of newly formed protein through the connecting cilium of retina rods in the frog. *J. Ultrastruct. Res.* 1968; 23:462–473. [PubMed: 5692302]
- Zhang Q, Acland GM, Wu WX, Johnson JL, Pearce-Kelling S, Tulloch B, et al. Different RPGR exon ORF15 mutations in Canids provide insights into photoreceptor cell degeneration. *Hum. Mol. Genet.* 2002; 11:993–1003. [PubMed: 11978759]
- Zhang H, Liu XH, Zhang K, Chen CK, Frederick JM, Prestwich GD, Baehr W. Photoreceptor cGMP phosphodiesterase delta subunit (PDEdelta) functions as a prenyl-binding protein. *J. Biol. Chem.* 2004; 279:407–413. [PubMed: 14561760]
- Zhao Y, Hong DH, Pawlyk B, Yue G, Adamian M, Grynberg M, et al. The retinitis pigmentosa GTPase regulator (RPGR)-interacting protein: subserving RPGR function and participating in disk morphogenesis. *Proc. Natl. Acad. Sci. USA.* 2003; 100:3965–3970. [PubMed: 12651948]
- Zito I, Downes SM, Patel RJ, Cheetham ME, Ebenezer ND, Jenkins SA, et al. RPGR mutation associated with retinitis pigmentosa, impaired hearing, and sinorespiratory infections. *J. Med. Genet.* 2003; 40:609–615. [PubMed: 12920075]

Figure 1.

(A) Schematic of a rod photoreceptor cell showing the sensory cilium axoneme (Ax); TZ, transition zone; R, the rootlet; IS, inner segment; OS, outer segment connected by the TZ; N, nucleus. (B) Immunofluorescence image of photoreceptor layer of mouse retina stained with anti-acetylated α -tubulin antibody, a ciliary marker (green); ONL, outer nuclear layer.

HUMAN RPGR^{1-ORF15}

Figure 2. Schematic representation of the predicted domain organization of human RPGR protein. P-Loop, ATP/GTP binding loop; RLD, RCC1-like domain; AP-sorting, adaptor protein sorting domain; Glu-Gly, glutamic acid and glycine rich domain; CC, coiled-coil domain.

Figure 3. Schematic representation of the RPGR and ciliarybasal bodycentrosomal (CBC) protein complexes in photoreceptors; binary interactions and macromolecular protein complexes facilitate microtubule-based ciliary transport. Dashed lines indicate interactions as part of complexes.