Published in final edited form as:

J Abnorm Psychol. 2012 February ; 121(1): 119–128. doi:10.1037/a0023938.

Visuospatial Attention in Schizophrenia: Deficits in Broad Monitoring

Britta Hahn,
University of Maryland School of Medicine

Benjamin M. Robinson,
University of Maryland School of Medicine

Alexander N. Harvey,
University of Maryland School of Medicine

Samuel T. Kaiser,
University of Maryland School of Medicine

Carly J. Leonard,
University of California, Davis

Steven J. Luck, and
University of California, Davis

James M. Gold
University of Maryland School of Medicine

Abstract

Although selective attention is thought to be impaired in people with schizophrenia (PSZ), prior research has found no deficit in the ability to select 1 location and withdraw attention from another. PSZ and healthy control subjects (HCS) performed a stimulus detection task in which 1, 2, or all 4 peripheral target locations were cued. When 1 or 2 locations were cued, both PSZ and HCS responded faster when the target appeared at a cued than uncued location. However, increases in the number of validly cued locations had much more deleterious effects on performance for PSZ than HCS, especially for targets of low contrast whose detection was more dependent on attention. PSZ also responded more slowly in trials with 4 cued locations relative to trials with 1 or 2 invalidly cued locations. Thus, visuospatial attention deficits in schizophrenia arise when broad monitoring is required rather than when attention must be focused narrowly.

Correspondence concerning this article should be addressed to Britta Hahn, Maryland Psychiatric Research Center, P.O. box 21247, Baltimore, MD 21228, bhahn@mprc.umaryland.edu.

Britta Hahn, Benjamin M. Robinson, Alexander N. Harvey, Samuel T. Kaiser, and James M. Gold: University of Maryland School of Medicine, Maryland Psychiatric Research Center, P.O. box 21247, Baltimore, MD 21228

Carly J. Leonard, and Steven J. Luck: University of California, Davis, Center for Mind & Brain and Department of Psychology, Davis, CA 95618

Publisher's Disclaimer: The following manuscript is the final accepted manuscript. It has not been subjected to the final copyediting, fact-checking, and proofreading required for formal publication. It is not the definitive, publisher-authenticated version. The American Psychological Association and its Council of Editors disclaim any responsibility or liabilities for errors or omissions of this manuscript version, any version derived from this manuscript by NIH, or other third parties. The published version is available at www.apa.org/pubs/journals/abn

Introduction

Neurocognitive deficits in schizophrenia best predict long-term functional disease outcome (Green, 1996; Green, Kern, & Heaton, 2004). These deficits are manifold but circumscribed, affecting numerous distinct mechanisms but sparing others (Gold, Hahn, Strauss, & Waltz, 2009). Efforts to develop effective pharmacotherapy for these symptoms rely on a mapping, characterization, and reduction of their complexity to a finite number of underlying problems (Marder, & Fenton, 2004; Carter, & Barch, 2007). A target of particular interest is selective attention, which has been frequently considered to be the root of impairments across a wide variety of cognitive tasks.

The visuospatial selective attention domain has been studied extensively in people with schizophrenia (PSZ), using variants of the Posner orienting paradigm (Posner, 1980), in which a cue directs attention either voluntarily or involuntarily to one of two possible target locations. Although many of these studies focused on examining possible lateralized abnormalities (Bustillo et al., 1997; Carter, Robertson, Chaderjian, Celaya, & Nordahl, 1992; Carter, Robertson, Chaderjian, O'Shara-Celaya, & Nordahl, 1994; Gold et al., 1992; Liotti, Dazzi, & Umiltà, 1993; Maruff, Hay, Malone, & Currie, 1995; Posner, Early, Reiman, Pardo, & Dhawan, 1988; Sapir, Henik, Dobrusin, & Hochman, 2001; Strauss, Novakovic, Tien, Bylsma, & Pearlson, 1991; Wigal, Swanson, & Potkin, 1997), they also provide more general clues about visuospatial selective attention mechanisms in schizophrenia. Notably, collapsed across hemifields, the reaction time (RT) difference between trials with a valid cue (i.e. one that correctly predicts the location of an upcoming target) and an invalid cue (i.e. one that directs attention to a location where the target does not appear) is no smaller in PSZ than in HCS. This is true for all of the above studies except Posner et al. A deficit in the ability to select one location and withdraw attention from another would have manifested itself in slower RT in valid trials and faster RT in invalid trials, and thus would have resulted in a smaller validity effect. PSZ displayed no such selection deficit, which is surprising given the hypothesized dysfunction of selective attention (Nuechterlein, & Dawson, 1984; Luck, & Gold, 2008). However, there is a suggestion of other, unexpected abnormalities.

A replicated finding is that the RT benefit of trials with a valid cue relative to a neutral condition that does not provide information about where to attend tends to be larger in PSZ than in HCS (Gold et al., 1992; Liotti et al., 1993; Bustillo et al., 1997; Sapir et al., 2001). A recent study confirmed this phenomenon in a direct assessment using optimized task conditions (Spencer et al., 2011). Thus, against all expectations, PSZ seemingly use the cue information more efficiently to orient attention in space. Alternatively, PSZ may be impaired on the neutral trials, with a reduced ability to spread attention widely and to maintain a broad focus of attention. The resulting disproportionate slowing in the neutral condition would result in the appearance of greater RT benefits of valid cues and reduced RT costs of invalid cues relative to the neutral condition. Indeed, the RT cost of invalid cues tends to be reduced in PSZ relative to HCS, although only in studies employing a no-cue neutral condition in which the target is not preceded by any signal (Liotti et al.; Nestor et al., 1992; Oie, Rund, & Sundet, 1998). When a double-cue neutral condition has been used, with peripheral cues at both locations, the RT cost on invalid relative to neutral trials tends to be larger in PSZ (Carter et al., 1992, 1994; Bustillo et al.; Daban et al., 2004). A possible explanation, given that PSZ do not appear to derive greater alerting effects from cues (Daban et al.; Gouzoulis-Mayfrank et al. 2007), is that the physical onset of the double-cue automatically directed the attentional focus to the two possible target locations in a bottom-up manner. Thus, PSZ may derive greater benefit than HCS from attention being spatially directed by peripheral cues to both locations prior to target onset, and conversely, the state of not having external signals guide attention to any specific locations may create

disproportional impairment. That is, PSZ may have a deficit in attending broadly on the basis of endogenous attentional control mechanisms.

The current study directly tested the hypothesis that PSZ have difficulty distributing attention broadly under endogenous control. We employed a visuospatial attention paradigm in which a central cue predicted the location of a peripheral target stimulus. One, two, or all four possible target locations could be cued simultaneously, manipulating the degree to which narrow focusing versus broad spatial monitoring was required. The target usually appeared at a cued location (valid trials), but occasionally at an uncued location (invalid trials), thus allowing us to simultaneously assess the ability to spread attention across varying numbers of locations and the ability to focus attention (by comparing performance for valid versus invalid trials). We predicted that stimulus detection deficits in PSZ relative to HCS would be particularly pronounced when all four locations were cued, necessitating a broad and diffuse attentional focus. If deficits are specific to broad monitoring rather than disengaging and shifting attention, performance of PSZ on these trials should be impaired relative to both valid and invalid predictive cue trials, but the difference between valid and invalid trials should be equivalent in PSZ and HCS. We also manipulated target contrast, predicting that the differences between PSZ and HCS would be larger for low-contrast targets because high-contrast targets evoke automatic detection mechanisms that are less influenced by the top-down distribution of spatial attention (Hawkins, Shafto, & Richardson, 1988). Because we manipulated target contrast, and PSZ sometimes exhibit contrast sensitivity impairments (reviewed by Javitt, 2009), we included a perceptual control task to ensure that the observed differences between PSZ and HCS were not a result of low-level sensory mechanisms.

Methods

Participants

Twenty-nine clinically stable, medicated outpatients meeting Diagnostic and Statistical Manual of Mental Disorders-IV (DSM-IV; American Psychiatric Association, 1994) criteria for schizophrenia (N=13 paranoid, 7 undifferentiated, 2 residual, 1 disorganized) or schizoaffective disorder (N=6), and 26 matched HCS participated. Demographic and clinical information is summarized in Table 1. Diagnosis was established using a best estimate approach in which information from a Structured Clinical Interview for DSM-IV (SCID) was combined with a review of medical records at a consensus diagnosis meeting chaired by one of the authors (JMG). All patients were receiving antipsychotic medication at time of testing; 4 were treated with first-generation antipsychotics, 23 with second-generation antipsychotics, and 2 with both. Fifteen patients additionally received mood stabilizing medication, 5 a benzodiazepine and 3 benztropine, an antiparkinsonian medication. A set of analyses of medication effects will be presented at the end of the Results section. Only patients whose medication had not changed in the preceding four weeks were enrolled. Control participants were recruited from the community via random digit dialing and word of mouth and were not taking any psychotropic medication. None of the control participants had a current Axis 1 or 2 diagnosis, as established by a SCID, and no self-reported family history of psychosis. Two controls had a history of Major Depression, now in full remission. Groups did not differ in age [$t(53)=0.2$, $P>0.8$], parental education [$t(51)=1.13$, $P>0.2$], sex (Chi-square $P>0.7$) or ethnicity (Chi-square $P>0.3$).

However, PSZ had fewer years of education than HCS [$t(53)=2.88$, $P<0.01$]. All participants provided informed consent for a protocol approved by the University of Maryland School of Medicine Institutional Review Board. Before participants signed the consent form, the investigator reviewed its content with them and answered any questions. For PSZ, basic

understanding of study demands and risks was formally evaluated in the presence of a third-party witness.

Neuropsychological Testing

Participants completed the Wechsler Abbreviated Scale of Intelligence (WASI; Wechsler, 1999), the Wide Range Achievement Test (WRAT 4; Wilkinson, & Robertson, 2006), the Wechsler Test of Adult Reading (WTAR; Wechsler, 2001), and the MATRICS battery (Nuechterlein, & Green, 2006). These tests were usually given on a separate day from the main experiment. PSZ scored lower than HCS on the WASI ($P=0.004$) and MATRICS battery ($P<0.001$), but there were no group differences on the WRAT 4 ($P>0.8$) or WTAR ($P>0.6$), suggesting similar premorbid functioning (Table 1).

Equipment

Tasks were completed in a dimly illuminated room on a 17" CRT monitor with a 60 Hz refresh rate. Eye-tracking was performed throughout both tasks to monitor central fixation, using an EyeLink 1000 eye-tracking system (SR Research Ltd., Mississauga, Ontario) operating at 2000 Hz. The eye-tracker consisted of an infrared light source and video camera, providing an image of the participant's right eye. Participants rested their heads on a chin and forehead rest at 70 cm viewing distance from the monitor.

Stimuli and Tasks

Spatial Attentional Resource Allocation Task (SARAT)—The SARAT has been previously described and validated as a tool for manipulating the size of the attentional focus in space (Hahn, Ross, & Stein, 2006). The task was slightly modified from the original version. Participants were required to keep their eyes fixated on a central circle containing a fixation cross (trials with eye movements were eliminated) and to detect a target signal at any of four peripheral locations marked by placeholders (Figure 1). With eyes directed at the center of the fixation cross, the center of the target location placeholders was positioned at an eccentricity of 12.5° . The diameter of the central circle was 3.0° and that of each of the placeholders was 1.5° . The central circle and placeholders, black against white, formed a background that remained on display throughout the task. A target consisted of one of the placeholders filling with a checkerboard of grey and white squares of 3×3 pixels each, yielding a spatial frequency of ~ 5.4 cycles/ $^\circ$ (note that these squares are too small to be discernible in Figure 1). The luminance of the white checks equaled that of the white background. Two target contrasts were tested. The contrast of the grey checks was 80% for the high-contrast targets and 20% for the low-contrast targets. Contrast was calculated as $(\text{white luminance} - \text{luminance of grey checks}) / \text{white luminance}$ (as measured with a J17 LumaColor photometer, Tektronix, Beaverton, OR). Upon detecting a target, participants pressed a button with their dominant index finger as quickly as possible.

A trial began once continuous central fixation was maintained for 500 ms. A cue then appeared in the central circle. Target presentation followed after a variable stimulus-onset-asynchrony of 400, 700, 1000 or 1300 ms. The target was visible for 500 ms, and the cue remained on display until 500 ms after target offset. The cue consisted of one, two or four quarters of the fixation circle turning black, indicating that the subsequent target was likely to appear in one of the corresponding quadrants of the display. When two quadrants were cued, they were always adjoining (both top, both bottom, both left, or both right).

The number of cued locations is related to the predictability of the target location. Fewer cued locations provide more precise information about the target location, allowing for a narrower and more intense attentional focus at the cued location(s) (Hahn et al., 2006). Conversely, increasing the number of cued locations increases spatial uncertainty and the

need to monitor broadly. The cue provided invalid information on 20% of the trials in which one or two locations were cued.

The cue was not followed by a target on 9.7% of trials, presented unpredictably, to discourage anticipatory responding to the cue. These cue-only trials were identical to the other trials, except that no target was presented during the 500-ms target interval. False alarms during these trials averaged $1.3 \pm 3.8\%$ in HCS and $7.8 \pm 25\%$ in PSZ [$t(53)=1.28$, NS]. All trials were followed by a 1500-ms intertrial interval, during which only the task background was presented. In total, there were 336 valid trials (56 x 1/2/4 cued locations x high/low target contrast), 56 invalid trials (14 x 1/2 cued locations x high/low contrast) and 42 cue-only trials (14 x 1/2/4 cued locations), tested over 14 blocks interspersed by rest periods. All trial types were randomized over every two consecutive blocks. The task took approximately one hour to complete.

Perceptual Control Task—To test whether possible group differences in perceptual sensitivity could explain the results we obtained, we invited all participants of the main experiment back to perform a contrast sensitivity measure that used the SARAT stimuli but had minimal attentional requirements and manipulated target contrast across a wide range of values (Figure 1). A standard 2-interval forced choice procedure was used to avoid response bias effects (Macmillan, & Creelman, 1991), and the method of constant stimuli was used rather than an adaptive staircase to avoid confounding sensitivity with lapses of attention. Originally, contrast levels of 4, 8, 16, 32 and 64% were tested. After 5 HCS and 9 PSZ had completed the task, a 12% contrast condition was added for the remaining participants. Each trial began with a 500-ms fixation period, followed by the onset of a one-location cue. 1000 ms after cue onset, a brief, clearly audibly tone was presented, and 1500 ms after the first tone, a different, easily differentiable tone was presented. A 500-ms target stimulus came on at the cued location simultaneously with one of the two tones. Thus, participants could always predict exactly and with absolute certainty where (at the cued location) and when (coinciding with one of the two tones) the target would appear. The central cue disappeared 1500 ms after the onset of the second tone. At the end of each trial, participants made an unspeeded forced choice indicating whether the target coincided with the first or second tone. Thirty trials of each contrast were tested, divided over 5 runs, each followed by a rest period. Contrast levels were randomized within each run. The task took approximately 30 min to complete. Twenty-five PSZ and 22 HCS completed it; the others were lost to follow-up. Two PSZ performed at chance across all contrast levels and clearly had difficulty following instructions. Their data were excluded from analysis of the control task, resulting in $N=23$ PSZ. Their SARAT data were not excluded because there was no indication that they had problems understanding this simple stimulus detection task.

Data analysis

SARAT—Trials with RT below 200 ms or above 2500 ms were considered outliers and excluded from analyses (0.13% and 0.14% of trials in PSZ and HCS, respectively). RTs were expressed as means and omission errors as the percentage of trials in which a target was presented but no response was made, as in previous studies employing this paradigm (Hahn et al., 2006; Hahn, Ross, & Stein, 2007; Hahn, Ross, Yang, et al., 2007). Trials with fixations (defined as stationary eye-position for 10 ms) outside a central circle measuring 4° of visual angle in diameter that occurred between cue onset and target offset were excluded from analyses of RT and omission errors. The percentage of trials with such eye movements was also analyzed. Eye-tracking data were lost for one HCS, for whom all trials were included in analyses of RT and omission errors. Exclusion of this HCS did not change any of the results.

Performance indices were analyzed by 3-factor mixed-model ANOVAs with Group (PSZ, HCS) as a between-subject factor and the number of cued locations (NumCuedLoc; 1,2,4) and Target contrast (high, low) as within-subject factors. One ANOVA included valid trials with 1 or 2 cued locations and nonpredictive trials (4 cued locations) and a separate ANOVA included invalid trials with 1 or 2 cued locations and nonpredictive trials. To compare the validity effect (invalid vs. valid trial performance) between groups, RT and omission errors in trials with 1 or 2 cued locations were also submitted to a 4-factor ANOVA that included both valid and invalid trials (Group x Validity x NumCuedLoc x Target contrast).

Perceptual Control Task—The percentage of trials in which the participant correctly reported which interval contained the target was analyzed using a 2-factor ANOVA with Group as a between-subjects factor and Target contrast as a within-subject factor. Trials with fixations outside the central fixation area between cue onset and the end of the second potential target interval were excluded from analysis.

Results

SARAT

Reaction Time

Valid versus nonpredictive trials: First we consider trials with 1, 2, or 4 cued locations on which the target appeared at a cued location. PSZ responded more slowly than HCS overall (Figure 2), leading to a significant main effect of Group [$F(1,53)=6.79$, $P<0.02$] in 3-factor ANOVA. Both groups responded more slowly to low- than high-contrast targets, as supported by a main effect of Target contrast [$F(1,53)=101.1$, $P<0.001$]. RTs increased monotonically with greater spatial uncertainty, leading to a main effect of NumCuedLoc [$F(2,106)=108.4$, $P<0.001$]. This uncertainty-dependent slowing was of approximately twice the magnitude in PSZ as that in HCS, as supported by a significant interaction of Group with NumCuedLoc [$F(2,106)=11.2$, $P<0.001$]. The average RT slowing from trials with 1 to trials with 4 cued locations was 49 ms in HCS vs. 94 ms in PSZ [$t(53)=3.79$, $P<0.01$; Cohen's $d=0.63$].

The three-way interaction (Group x NumCuedLoc x Target contrast) was also significant [$F(2,106)=4.05$, $P=0.02$]. This reflected a larger effect of the number of cued locations for low-contrast than high-contrast targets in PSZ but not in HCS. Supporting this, separate 2-factor ANOVAs in PSZ and HCS yielded a significant interaction of NumCuedLoc with Target contrast only in PSZ [$F(2,56)=4.23$, $P=0.02$]. In PSZ, the RT slowing from trials with 1 to trials with 4 cued locations averaged 81.4 ms for high-contrast targets and 106 ms for low-contrast targets [$t(28)=2.18$, $P<0.038$; paired samples t -test]. In HCS, this RT slowing was almost identical between the target contrasts (49.9 vs. 48.1 ms; NS).¹ The slowing did not correlate with scores on any of the neuropsychological measures in either PSZ or HCS for either target intensity.

Invalid versus nonpredictive trials: Next, we compared invalid cue trials with 1 and 2 cued locations and nonpredictive cue trials (4 cued locations). In HCS, RTs were approximately the same for all cue types. Remarkably, PSZ were actually slower on the nonpredictive than invalid cue trials. These differences yielded a significant interaction of

¹Disproportionate slowing with more cued locations in PSZ was not restricted to trials with 4 cued locations but became gradually more pronounced with an increasing number of cued locations. This is substantiated by the finding that, for low-contrast targets, PSZ and HCS differed significantly in their degree of slowing not only for the RT difference between trials with 1 and 4 cued locations [$t(53)=3.70$, $P<0.001$, independent samples t -test], but also for the RT difference between trials with 1 and 2 cued locations [$t(53)=2.73$, $P<0.01$].

NumCuedLoc (1,2,4) with Group [$F(2,106)=5.44$, $P=0.006$] in 3-factor ANOVA. One-factor ANOVAs confirmed an effect of NumCuedLoc in PSZ [$F(2,56)=8.87$, $P<0.001$] but not in HCS [$F(2,50)<1$]. Interactions involving Target contrast were not significant. Thus, PSZ but not HCS were impaired when the cue directed them to spread attention across four locations compared to when the cue directed them to focus on an incorrect location.

Valid versus invalid trials: Both groups displayed slower RT on invalid than valid cue trials with 1 or 2 cued locations, and this was confirmed by a significant main effect of Validity [$F(1,53)=25.7$, $P<0.001$] in a 4-factor ANOVA [Group x Validity x NumCuedLoc (1,2) x Target contrast]. There were no significant interactions involving Validity and Group [Validity x Group: $F(1,53)=1.1$, $P>0.3$], and the Validity effect was significant ($P<0.001$) in both HCS (41 ms, averaged over NumCuedLoc and Target contrast) and PSZ (45 ms). The effect size of the group difference in the validity effect was Cohen's $d=0.13$. Thus, as in many prior studies, we found no evidence of an impairment in the ability of PSZ to focus attention onto 1 or 2 cued locations.

% Omission Errors²

Valid versus nonpredictive trials: Both PSZ and HCS made more omission errors with low- than high-contrast targets (Figure 3). However, PSZ but not HCS showed an increase in omission errors with greater spatial unpredictability for the low-contrast targets. This was supported by a significant 3-way interaction [$F(2,106)=4.55$, $P=0.013$]. Separate 2-factor ANOVAs in PSZ and HCS confirmed a significant NumCuedLoc x Target contrast interaction in PSZ [$F(2,56)=5.82$, $P=0.005$], which was absent in HCS [$F(2,50)<1$]. In follow-up 1-factor ANOVAs, the effect of NumCuedLoc in PSZ was significant for low-contrast [$F(2,56)=3.84$, $P=0.027$] but not high-contrast targets [$F(2,56)<1$]. The increase in omission errors from trials with 1 to trials with 4 cued locations for low-contrast targets was of moderate effect size ($d=0.58$). It did not correlate with scores on any of the neuropsychological measures.

Invalid versus nonpredictive trials: PSZ made a disproportionately large number of omission errors on both nonpredictive and invalid cue trials with low-contrast targets. This was substantiated by a Group x Target contrast interaction [$F(1,53)=7.37$, $P<0.01$] in a 3-factor ANOVA. Follow-up t-tests confirmed a significant group difference for low-contrast [$t(53)=2.93$, $P=0.005$] but not high-contrast targets [$t(53)=1.43$, NS]. In PSZ, the omission rate was numerically greater on nonpredictive trials than on invalid trials with a single cued location, whereas HCS showed the opposite pattern. However, effects involving NumCuedLoc were not significant.

Valid versus invalid trials: Both groups made more omission errors on invalid than valid trials, as confirmed by a significant main effect of Validity [$F(1,53)=5.38$, $P<0.05$] in a 4-factor ANOVA. There were no significant interactions involving Validity and Group (Validity x Group: $P=0.15$). This provides additional evidence that PSZ are unimpaired at directing attention toward some locations and withdrawing attention from others.

²This measure failed Shapiro-Wilk tests for normality in both HCS ($P<0.001$) and PSZ ($P=0.05$). Data transformation could not remedy this. Skewness (HCS: 2.49, PSZ: 0.71) was mainly due to a few large outlier values. Skewness was substantially reduced (HCS: 0.569, PSZ: 0.585) by excluding the 2 HCS and the 3 PSZ with the largest percentage of omission errors, which also aided normality (Shapiro-Wilk test: $P=0.12$ in HCS, $P=0.15$ in PSZ). After excluding these subjects from the ANOVA of valid trials, the critical 3-way interaction was still significant ($P<0.045$).

Percentage of Trials with Eye Movements

Valid versus nonpredictive trials: Trials with eye movements away from the fixation point occurring between cue onset and target offset were more numerous in PSZ than HCS (Figure 4). Their number increased with spatial unpredictability in PSZ but not in HCS, especially for low-contrast targets. This led to a significant 3-way interaction [$F(2,104)=6.93$, $P=0.001$]. In HCS, the percentage of trials with eye movements did not depend on NumCuedLoc or Target contrast, as confirmed by an absence of main effects or interaction in a 2-factor ANOVA. In PSZ, however, the NumCuedLoc x Target contrast interaction was significant [$F(2,56)=5.54$, $P=0.006$]. In 1-factor ANOVAs, the effect of NumCuedLoc was significant in PSZ for low-contrast targets [$F(2,56)=15.3$, $P<0.001$] but not high-contrast targets [$F(2,56)=1.80$, NS], indicating that spatial uncertainty increased eye movements particularly for trials with low-contrast targets (Cohen's $d=0.64$ for the difference between 1 and 4 cued locations for low-intensity targets).

Invalid versus nonpredictive trials, and valid versus invalid trials: Invalid trials were not analyzed because the percentage of trials with eye movements was typically zero due to the low total number of invalid cue trials.

Effects of medication status—Additional 3-factor ANOVAs were performed comparing PSZ who received a given medication with all other PSZ. Each ANOVA included NumCuedLoc (1, 2, 4) and Target contrast as within-subject factors, but one ANOVA included Benzodiazepine, one Mood stabilizer, one Clozapine, and one Typical antipsychotic (present, absent) as a between-subject factor. As above, separate ANOVAs were performed for valid and non-predictive and for invalid and nonpredictive trials. The only significant interaction involving Benzodiazepine was with NumCuedLoc on valid trial RT [$F(2,54)=3.44$, $P<0.05$]. This effect was driven by two of the five patients treated with benzodiazepines displaying RT differences of >170 ms between trials with 1 and 4 cued locations. When repeating the original Group x NumCuedLoc x Target contrast ANOVA without these 5 patients, the same main effects and interactions were observed. The only significant interaction involving Mood stabilizer was with Target contrast on the percentage of trials with eye movements [$F(1,27)=4.36$, $P<0.05$]. In PSZ not receiving mood stabilizers, trials with eye movements were more numerous for low- than high-contrast targets [13 vs. 10%; $t(13)=3.15$, $P<0.01$], but in PSZ treated with mood stabilizers trials with eye movements were almost identical between high- and low-contrast targets (13% in each case). When repeating the original Group x NumCuedLoc x Target contrast ANOVA without these 15 patients, the same results were obtained. There were no interactions involving Clozapine or Typical antipsychotic.

Finally, for PSZ, we performed Pearson correlations of haloperidol equivalents (Andreasen, Pressler, Nopoulos, Miller, & Ho, 2010) with the difference between valid trials with 1 and 4 cued locations in the three dependent variables (RT, omission errors, trials with eye-movements). There were no significant correlations for trials with low-contrast or high-contrast targets.

Perceptual Control Task

The purpose of this task was to test whether the difficulty of PSZ in detecting the low-contrast targets under conditions of spatial uncertainty and the specific psychophysical parameters of the SARAT could have been an artifact of poorer perceptual sensitivity rather than a consequence of an impaired ability to distribute attention broadly. Reduced contrast sensitivity in PSZ to our task stimuli would be indicated by impaired discrimination performance relative to HCS especially for trials with low contrast levels. As can be seen from Figure 5, performance accuracy was almost identical between PSZ and HCS, and this

was supported by the absence of a Group main effect [$F(1,43)<1$] or a Group x Target contrast interaction [$F(4,172)<1$] in a 2-factor ANOVA. When including the 12% data point and limiting ANOVA to participants for whom this data point was available ($N=14$ PSZ, $N=17$ HCS), the same results were obtained [Group: $F(1,29)<1$; Group x Target contrast: $F(5,145)<1$]. Accuracy did not differ between groups at any contrast level in independent-samples t-tests ($P>0.3$ in each case). Thus, we conclude that although impaired contrast sensitivity has been observed in PSZ under different stimulus and task conditions (reviewed by Javitt, 2009), our sample of PSZ had no perceptual deficit relative to HCS in detecting the low-contrast stimuli under the present conditions. It is extremely unlikely that any difference in perceptual sensitivity that could not be detected when tested directly was responsible for the performance patterns in the SARAT.

Discussion

The present findings resolve the apparent discrepancy between the widespread belief that schizophrenia involves impaired attentional selection and the repeated finding that PSZ exhibit normal and even superior visuospatial attentional cuing effects (Gold et al., 2009; Spencer et al., 2011). Specifically, the present study confirms the observation that PSZ are unimpaired at focusing attention on one location and withdrawing attention from others, but it demonstrates a substantial impairment in the ability of PSZ to distribute attention broadly.

On valid trials, both groups displayed slower responding when the target location became more uncertain, but this effect was substantially more pronounced for PSZ than HCS. If this was due to better attentional selection with more precise cueing, larger performance costs would have been observed on invalid cue trials. Instead, the performance difference between valid and invalid trials when one or two locations were cued did not differ between PSZ and HCS, as observed in previous studies (Gold et al., 2009). Furthermore, PSZ but not HCS actually responded more slowly on nonpredictive than invalid trials. Thus, attending broadly is actually more deleterious to performance in PSZ than focusing attention away from the location of the upcoming target. The results indicate that the observed performance pattern arose because PSZ were disproportionately impaired in nonpredictive cue trials that required monitoring all four possible target locations.

Impairment with more spatial uncertainty was particularly pronounced in trials with low-contrast targets in PSZ. First, RT slowing with spatial uncertainty was greater with low- than high-contrast targets in PSZ but not HCS. Second, omission errors in trials with low-contrast targets increased with greater spatial uncertainty in PSZ but not HCS. Third, on trials with low-contrast targets, PSZ made more eye movements away from central fixation when the cue was nonpredictive. Thus, spatial unpredictability combined with low physical target salience created the largest performance impairment in PSZ. However, our perceptual control experiment indicated that, under the current task conditions, PSZ and HCS did not differ in contrast sensitivity to the targets per se; that is, PSZ had no problems detecting the peripheral low-contrast stimuli when attentional demands were minimized. The disparity between this finding and studies that did identify contrast sensitivity reductions in PSZ may be due to differences in stimulus properties (ours are likely to be processed by the magnocellular; Javitt, 2009), in patient populations (reduced contrast sensitivity may be associated with negative symptoms; Slaghuys, 2004; Keri, Kiss, Kelemen, Benedek, & Janka, 2005), in medication status (the current sample received mostly atypical neuroleptics, while reduced contrast sensitivity may be specific to PSZ medicated with typical antipsychotics; Chen et al. 2003), or in the measurement method (by using the method of constant stimuli rather than an adaptive staircase we rule out differences resulting from nonspecific factors such as lapses of attention). Importantly, without drawing any more

general conclusions about perceptual abnormalities in schizophrenia, we can say that the current stimulus detection deficit observed when multiple locations were cued was clearly attentional and not sensory in nature.

Why would a broad monitoring deficit be moderated by target contrast? The greater sensitivity of the low-contrast targets to the deleterious effects of a broad attentional state may be conceptualized under limited resource capacity models of attention (e.g. Kahneman, 1973). Broad monitoring all four locations may be more effortful for PSZ than HCS and exceed the available processing resources, thus leading to a suboptimal attentional state. Bottom-up orienting may have aided detection of high-contrast targets even when PSZ were in a suboptimal attentional state (Hawkins et al., 1988). Detection of low-contrast targets, however, is more dependent on spatial attention. When spatial uncertainty is high, the size of the attentional window must be expanded to effectively monitor all possible target locations (Eriksen, & Yeh, 1985; Belopolsky, Zwaan, Theeuwes, & Kramer, 2007), and this spreading of the benefits of attentional processing across the visual field enables the detection of even low-contrast targets. Thus, the patient deficit may be described as suboptimal maintenance of a wide attentional window. Indeed, it has recently been suggested that PSZ have a narrowed “attentional spotlight” and insufficient attentional resources to maintain a wide visual span (Elahipanah, Christensen, & Reingold, 2010).

Psychophysical models suggest that inattention slows the rate of perceptual information acquisition (Luck, & Vecera, 2002; Palmer, 1998) and thus increases the time required to reach detection threshold. For low-contrast stimuli, the detection threshold may never be reached on some trials, leading to omission errors. Thus, a failure of adopting a broader attentional window with more spatial uncertainty can explain both the increased RTs for low- and high-contrast targets and the increased rate of omission errors for low-contrast targets in PSZ. This failure, quantified as the difference in RT or omission errors between trials with 1 and 4 cued locations in PSZ, did not correlate with any of the neuropsychological indices collected, including IQ, MATRICS domains, WTAR and WRAT. Thus, at this point in time, there is no obvious clue about the degree to which a broad monitoring deficit may relate to other aspects of cognitive dysfunction described in PSZ. Interpreted broadly and generalized beyond the visuospatial domain, a narrowed attentional focus, or inability to spread attention broadly, may limit the ability to process multiple inputs or perceive multiple concomitant possibilities. We speculate that this type of processing limitation may translate into a reduced ability to consider multiple alternatives and may underlie the reduced cognitive flexibility described in PSZ (e.g. Elliot, McKenna, Robbins & Sahakian, 1995). Clearly, the current findings do not suffice to support such generalization, but they may lay the foundation for future work.

The eye movement data suggest that PSZ may have tried to overcome their difficulty in efficiently spreading attention across the visual field by resorting to a serial focusing of the target locations once they failed to detect a target. PSZ made more fixations outside the central fixation area on nonpredictive cue trials, in particular those with low-contrast targets. These eye movements must have been exploratory rather than being triggered by the physical target onset because exogenous triggering would have yielded more eye movements for high-contrast targets. Instead, PSZ appeared to initiate a large portion of eye movements upon guessing that they missed the target, or in an effort to verify the occurrence of a low-contrast target. Thus, RT, omission errors, and central fixation performance provided converging evidence for a deficit in the ability of PSZ to monitor broadly and maintain a wide attentional window. Based on the pattern of results obtained previously using the Posner paradigm (see Introduction), we suggest that this impairment is a result of dysfunction in top-down attentional control rather than an inability to distribute attention

widely as a result of bottom-up orienting. Future studies may test this possibility by employing peripheral cues.

An fMRI study of the SARAT in healthy adults found that activity in the rostral anterior cingulate and posterior cingulate cortex was predictive of trial-by-trial RT, but only on nonpredictive cue trials that require broad monitoring (Hahn, Ross, & Stein, 2007). These areas are central hubs of the default network of resting brain function, which other studies have shown to be dysfunctional in PSZ (Pomarol-Clotet et al., 2008; Whitfield-Gabrieli et al. 2009). The combination of these previous results suggests that the broad monitoring deficit of PSZ may be caused by dysfunction of the default network. Although the default network is usually associated with inward focusing and task-independent thought processes (Raichle et al., 2001), some studies suggested that this network is also involved in maintaining a broad attentional state of diffuse “watchfulness” toward the external environment (Gilbert, Dumontheil, Simons, Frith, & Burgess, 2007; Gilbert, Simons, Frith, & Burgess, 2006; Hahn, Ross, & Stein, 2007). This “sentinel function” (Buckner, Andrews-Hanna, & Schacter, 2008) is thought of as a safety mode entered when perceptual processing resources are not directed to specific external stimuli. We suggest that a “sentinel dysfunction” may underlie the impaired performance of PSZ when broad attention is required.

There are several potential limitations that should be considered. At this point in time, it is unclear to what degree the identified deficit in spreading attention widely generalizes beyond the spatial domain. Additionally, it is unknown whether it reflects the number of discrete possible target locations or simply the size of the area over which attention has to be spread. Thus, future experiments should address, for example, whether the same results are obtained if there is greater positional uncertainty about target locations. Another limitation is that the current PSZ sample represents a largely stable, medicated outpatient population, and future studies will need to establish to what degree the present findings generalize across disease stages or states. Furthermore, future studies should expand on the present findings by differentiating between different symptom profiles and by more expansive analyses of potential medication effects than possible within the current sample. The current study provides the first clear evidence that visuospatial attention in schizophrenia is marked by a deficit in maintaining a wide attentional window rather than in focusing narrowly. This suboptimal attentional state appears to increase the threshold of physical target salience necessary to trigger target detection, and it may be reflective of an impaired sentinel function of the default network.

Acknowledgments

This research was supported by a grant from the National Institute of Mental Health (grant number MH065034 to J.M.G. and S.J.L.). We thank Valerie M. Beck for help with setting up the eye-tracking procedure, Jacqueline Kiwanuka, Sharon August and Leeka Hubzin for their assistance with the conduct of the study, Dr. Robert McMahon for statistical advice, and all volunteers participating in this study.

References

- Addington D, Addington J, Maticka-Tyndale E, Joyce J. Reliability and validity of a depression rating scale for schizophrenics. *Schizophrenia Research*. 1992; 6:201–208. [PubMed: 1571313]
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 4. Washington, DC: Author; 1994.
- Andreasen, NC. *The Scale for the Assessment of Negative Symptoms (SANS)*. Iowa City, IA: University of Iowa; 1984.

- Andreasen NC, Pressler M, Nopoulos P, Miller D, Ho BC. Antipsychotic dose equivalents and dose-years: a standardized method for comparing exposure to different drugs. *Biological Psychiatry*. 2010; 67:255–262. [PubMed: 19897178]
- Belopolsky AV, Zwaan L, Theeuwes J, Kramer AF. The size of an attentional window modulates attentional capture by color singletons. *Psychonomic Bulletin & Review*. 2007; 14:934–938. [PubMed: 18087962]
- Buckner RL, Andrews-Hanna JR, Schacter DL. The brain's default network: anatomy, function, and relevance to disease. *Annals of the New York Academy of Sciences*. 2008; 1124:1–38. [PubMed: 18400922]
- Bustillo JR, Thaker G, Buchanan RW, Moran M, Kirkpatrick B, Carpenter WT Jr. Visual information-processing impairments in deficit and nondeficit schizophrenia. *American Journal of Psychiatry*. 1997; 154:647–654. [PubMed: 9137120]
- Carter CS, Barch DM. Cognitive neuroscience-based approaches to measuring and improving treatment effects on cognition in schizophrenia: the CNTRICS initiative. *Schizophrenia Bulletin*. 2007; 33:1131–1137. [PubMed: 17630405]
- Carter CS, Robertson LC, Chaderjian MR, Celaya LJ, Nordahl TE. Attentional asymmetry in schizophrenia: controlled and automatic processes. *Biological Psychiatry*. 1992; 31:909–918. [PubMed: 1637931]
- Carter CS, Robertson LC, Chaderjian MR, O'Shara-Celaya L, Nordahl TE. Attentional asymmetry in schizophrenia: the role of illness subtype and symptomatology. *Progress in Neuropsychopharmacology and Biological Psychiatry*. 1994; 18:661–683.
- Daban C, Krebs MO, Bourdel MC, Willard D, Loo H, Olie JP, et al. Effects of atypical neuroleptics on alertness and visual orienting in stabilized schizophrenic patients: a preliminary study. *International Journal of Neuropsychopharmacology*. 2004; 7:255–263. [PubMed: 15050039]
- Elahipanah A, Christensen BK, Reingold EM. Visual search performance among persons with schizophrenia as a function of target eccentricity. *Neuropsychology*. 2010; 24:192–198. [PubMed: 20230113]
- Elliot R, McKenna PJ, Robbins TW, Sahakian BJ. Neuropsychological evidence for frontostriatal dysfunction in schizophrenia. *Psychological Medicine*. 1995; 25:619–30. [PubMed: 7480441]
- Eriksen CW, Yeh YY. Allocation of attention in the visual field. *Journal of Experimental Psychology: Human Perception and Performance*. 1985; 11:583–597. [PubMed: 2932532]
- Gilbert SJ, Dumontheil I, Simons JS, Frith CD, Burgess PW. Comment on "Wandering minds: the default network and stimulus-independent thought". *Science*. 2007; 317:43. [PubMed: 17615325]
- Gilbert SJ, Simons JS, Frith CD, Burgess PW. Performance-related activity in medial rostral prefrontal cortex (area 10) during low-demand tasks. *Journal of Experimental Psychology: Human Perception and Performance*. 2006; 32:45–58. [PubMed: 16478325]
- Gold JM, Hahn B, Strauss GP, Waltz JA. Turning it upside down: areas of preserved cognitive function in schizophrenia. *Neuropsychology Review*. 2009; 19:294–311. [PubMed: 19452280]
- Gold JM, Randolph C, Coppola R, Carpenter CJ, Goldberg TE, Weinberger DR. Visual orienting in schizophrenia. *Schizophrenia Research*. 1992; 7:203–209. [PubMed: 1390400]
- Gouzoulis-Mayfrank E, Balke M, Hajsamou S, Ruhrmann S, Schultze-Lutter F, Daumann J, et al. Orienting of attention in unmedicated patients with schizophrenia, prodromal subjects and healthy relatives. *Schizophrenia Research*. 2007; 97:35–42. [PubMed: 17869065]
- Green MF. What are the functional consequences of neurocognitive deficits in schizophrenia? *American Journal of Psychiatry*. 1996; 153:321–330. [PubMed: 8610818]
- Green MF, Kern RS, Heaton RK. Longitudinal studies of cognition and functional outcome in schizophrenia: implications for MATRICS. *Schizophrenia Research*. 2004; 72:41–51. [PubMed: 15531406]
- Hahn B, Ross TJ, Stein EA. Neuroanatomical dissociation between bottom-up and top-down processes of visuospatial selective attention. *Neuroimage*. 2006; 32:842–853. [PubMed: 16757180]
- Hahn B, Ross TJ, Stein EA. Cingulate activation increases dynamically with response speed under stimulus unpredictability. *Cerebral Cortex*. 2007; 17:1664–1671. [PubMed: 16963517]

- Hahn B, Ross TJ, Yang Y, Kim I, Huestis MA, Stein EA. Nicotine enhances visuospatial attention by deactivating areas of the resting brain default network. *Journal of Neuroscience*. 2007; 27:3477–3489. [PubMed: 17392464]
- Hawk AB, Carpenter WT, Strauss JS. Diagnostic criteria and five-year outcome in schizophrenia: a report from the International Pilot Study of Schizophrenia. *Archives of General Psychiatry*. 1975; 32:343–347. [PubMed: 1115575]
- Hawkins HL, Shafto MG, Richardson K. Effects of target luminance and cue validity on the latency of visual detection. *Perception & Psychophysics*. 1988; 44:484–492. [PubMed: 3226899]
- Javitt DC. When doors of perception close: bottom-up models of disrupted cognition in schizophrenia. *Annual Reviews of Clinical Psychology*. 2009; 5:249–275.
- Kahneman, D. *Attention and effort*. Englewood Cliffs, NJ: Prentice-Hall; 1973.
- Liotti M, Dazzi S, Umiltà C. Deficits of the automatic orienting of attention in schizophrenic patients. *Journal of Psychiatric Research*. 1993; 27:119–130. [PubMed: 8515384]
- Luck SJ, Gold JM. The construct of attention in schizophrenia. *Biological Psychiatry*. 2008; 64:34–39. [PubMed: 18374901]
- Luck, SJ.; Vecera, SP. Attention. In: Pashler, H.; Yantis, S., editors. *Stevens' Handbook of Experimental Psychology: Vol. 1. Sensation and Perception*. 3. New York: Wiley; 2002. p. 235–286.
- Macmillan, NA.; Creelman, CD. *Detection Theory: A User's Guide*. New York: Cambridge University Press; 1991.
- Marder SR, Fenton W. Measurement and Treatment Research to Improve Cognition in Schizophrenia: NIMH MATRICS initiative to support the development of agents for improving cognition in schizophrenia. *Schizophrenia Research*. 2004; 72:5–9. [PubMed: 15531402]
- Maruff P, Hay D, Malone V, Currie J. Asymmetries in the covert orienting of visual spatial attention in schizophrenia. *Neuropsychologia*. 1995; 33:1205–1223. [PubMed: 8552225]
- Nestor PG, Faux SF, McCarley RW, Penhune V, Shenton ME, Pollak S, et al. Attentional cues in chronic schizophrenia: abnormal disengagement of attention. *Journal of Abnormal Psychology*. 1992; 101:682–689. [PubMed: 1430608]
- Nuechterlein KH, Dawson ME. Information processing and attentional functioning in the developmental course of schizophrenic disorders. *Schizophrenia Bulletin*. 1984; 10:160–203. [PubMed: 6729409]
- Nuechterlein, KH.; Green, MF. *MATRICS Consensus Cognitive Battery, Manual*. Los Angeles, CA: MATRICS Assessment Inc; 2006.
- Oie M, Rund BR, Sundet K. Covert visual attention in patients with early-onset schizophrenia. *Schizophrenia Research*. 1998; 34:195–205. [PubMed: 9850986]
- Overall JE, Gorman DR. The Brief Psychiatric Rating Scale. *Psychological Reports*. 1962; 10:799–812.
- Palmer, J. Attentional effects in visual search: Relating search accuracy and search time. In: Wright, RD., editor. *Visual attention*. Vol. 8. New York: Oxford University Press; 1998. p. 348–388.
- Pomarol-Clotet E, Salvador R, Sarro S, Gomar J, Vila F, Martinez A, et al. Failure to deactivate in the prefrontal cortex in schizophrenia: dysfunction of the default mode network? *Psychological Medicine*. 2008; 38:1185–1193. [PubMed: 18507885]
- Posner MI. Orienting of attention. *Quarterly Journal of Experimental Psychology*. 1980; 32:3–25. [PubMed: 7367577]
- Posner MI, Early TS, Reiman E, Pardo PJ, Dhawan M. Asymmetries in hemispheric control of attention in schizophrenia. *Archives of General Psychiatry*. 1988; 45:814–821. [PubMed: 3415424]
- Raichle ME, MacLeod AM, Snyder AZ, Powers WJ, Gusnard DA, Shulman GL. A default mode of brain function. *Proceedings of the National Academy of Sciences of the United States of America*. 2001; 98:676–682. [PubMed: 11209064]
- Sapir A, Henik A, Dobrusin M, Hochman EY. Attentional asymmetry in schizophrenia: disengagement and inhibition of return deficits. *Neuropsychology*. 2001; 15:361–370. [PubMed: 11499991]

- Spencer KM, Nestor PG, Valdman O, Niznikiewicz MA, Shenton ME, McCarley RW. Enhanced facilitation of spatial attention in schizophrenia. *Neuropsychology*. 2011; 25:76–85. [PubMed: 20919764]
- Strauss ME, Novakovic T, Tien AY, Bylsma F, Pearlson GD. Disengagement of attention in schizophrenia. *Psychiatry Research*. 1991; 37:139–146. [PubMed: 1876627]
- Wechsler, D. Wechsler Abbreviated Scale of Intelligence (WASI). San Antonio, TX: The Psychological Corporation; 1999.
- Wechsler, D. Wechsler Test of Adult Reading (WTAR). San Antonio, TX: The Psychological Corporation; 2001.
- Whitfield-Gabrieli S, Thermenos HW, Milanovic S, Tsuang MT, Faraone SV, McCarley RW, et al. Hyperactivity and hyperconnectivity of the default network in schizophrenia and in first-degree relatives of persons with schizophrenia. *Proceedings of the National Academy of Sciences of the United States of America*. 2009; 106:1279–1284.
- Wigal SB, Swanson JM, Potkin SG. Lateralized attentional deficits in drug-free and medicated schizophrenic patients. *Neuropsychologia*. 1997; 35:1519–1525. [PubMed: 9460721]
- Wilkinson, GS.; Robertson, GJ. Wide Range Achievement Test (WRAT) 4. Lutz, FL: Psychological Assessment Resources; 2006.

Figure 1.

Trial examples of the Spatial Attentional Resource Allocation Task (SARAT, top) and the perceptual control task (bottom). For the SARAT, one, two, or all four locations were cued and subjects responded upon detecting the target. For the perceptual control task, one location was always cued. Target onset coincided with either one of two briefly presented tones ("BEEP" or "BOOP"), and subjects made a forced choice as to with which tone the target coincided.

Figure 2.

Reaction times of healthy control subjects (HCS) and people with schizophrenia (PSZ) in the SARAT. The graph compares the averages ($\pm$ SEM) of trials with high-contrast targets (“high”) and low-contrast targets (“low”), and trials with 1, 2 or 4 validly cued target locations and 1 or 2 invalidly cued locations. (*) $P=0.07$, *** $P<0.001$ in Tukey tests comparing RT, averaged over high- and low-contrast targets, between trials with 1, 2 or 4 validly cued locations, and between trials with 1 and 2 invalidly cued locations and trials with 4 cued locations.

Figure 3.
The percentage of omission errors (averages $\pm$ SEM) of HCS and PSZ in the SARAT.

Figure 4.

The percentage of trials with eye movement in HCS and PSZ in the SARAT. The graph compares trials with high-contrast targets (“high”) and low-contrast targets (“low”) with 1, 2 or 4 validly cued locations. ** $P < 0.01$, *** $P < 0.001$ in Tukey test comparing only trials with low-contrast targets because a cue effect was identified only for these trials.

Figure 5.

Response accuracy (averages $\pm$ SEM) in the perceptual control task for HCS (N=22) and PSZ (N=23). The 12% contrast level data points include only 17 HCS and 14 PSZ.

Table 1Group Demographics (mean $\pm$ stdev)

	Patients	Controls
Age	41.4 $\pm$ 9.8 (range 22–53)	41.9 $\pm$ 9.0 (range 23–54)
Male : Female	16 : 13	13 : 13
AA : C : A : AI ^a	10 : 16 : 2 : 1	12 : 14 : 0 : 0
Education (years)	13.1 $\pm$ 2.3	14.7 $\pm$ 1.8 *
Parental education ^b	14.3 $\pm$ 3.3 ^c	13.5 $\pm$ 1.9
WASI	101.5 $\pm$ 13.5	112.0 $\pm$ 11.9 *
MATRICES total score	33.8 $\pm$ 15.3	48.9 $\pm$ 10.5 **
WRAT 4 standard score	98.9 $\pm$ 14.2	99.4 $\pm$ 12.7
WTAR standard score	101.9 $\pm$ 16.8	103.5 $\pm$ 12.2
BPRS	36.2 $\pm$ 7.4 (range 24–53)	
SANS	33.8 $\pm$ 12.9 (range 4–57)	
LOFS	20.5 $\pm$ 6.3 (range 10–34)	
Calgary Depression Scale	2.8 $\pm$ 2.6 (range 0–9)	

^a AA = African American; C = Caucasian; A = Asian; AI = American Indian

^b average over mother's and father's years of education

^c data unavailable for 2 subjects

* P<0.01,

** P<0.001;

significant difference between PSZ and HCS in independent samples t-test

BPRS = Brief Psychiatric Rating Scale (Overall, & Gorman, 1962)

SANS = Scale for the Assessment of Negative Symptoms (Andreasen, 1984)

LOFS = Level Of Functioning Scale (Hawk, Carpenter, & Strauss, 1975)

Calgary Depression Scale (Addington, Addington, Maticka-Tyndale, & Joyce, 1992)