Published in final edited form as:

Prog Neurobiol. 2012 February ; 96(2): 208–219. doi:10.1016/j.pneurobio.2012.01.004.

Unmasking the mysteries of the habenula in pain and analgesia

L. Shelton^{a,b}, L. Becerra^{a,b,c,d}, and D. Borsook^{a,b,c,d,*}

^aCenter for Pain and the Brain, Harvard Medical School, United States

^bChildren's Hospital Boston, Harvard Medical School, United States

^cMcLean Hospital, Harvard Medical School, United States

^dMassachusetts General Hospital, Harvard Medical School, United States

Abstract

The habenula is a small bilateral structure in the posterior–medial aspect of the dorsal thalamus that has been implicated in a remarkably wide range of behaviors including olfaction, ingestion, mating, endocrine and reward function, pain and analgesia. Afferent connections from forebrain structures send inputs to the lateral and medial habenula where efferents are mainly projected to brainstem regions that include well-known pain modulatory regions such as the periaqueductal gray and raphe nuclei. A convergence of preclinical data implicates the region in multiple behaviors that may be considered part of the pain experience including a putative role in pain modulation, affective, and motivational processes. The habenula seems to play a role as an evaluator, acting as a major point of convergence where external stimuli is received, evaluated, and redirected for motivation of appropriate behavioral response. Here, we review the role of the habenula in pain and analgesia, consider its potential role in chronic pain, and review more recent clinical and functional imaging data of the habenula from animals and humans. Even though the habenula is a small brain structure, advances in structural and functional imaging in humans should allow for further advancement of our understanding of its role in pain and analgesia.

Keywords

Habenula; Pain; Imaging; fMRI; Analgesia

1. Introduction

The habenula is a small bilateral structure in the posterior–medial aspect of the dorsal thalamus, located on the floor of the third ventricle above the pineal complex (together known as the epithalamus). It has been implicated in a remarkably wide range of behaviors including olfaction, ingestion, mating, endocrine, reward function, addiction and pain (Hikosaka, 2010). Given the broad involvement across numerous functions and behaviors, the region seems to have a basic but vital role as an evaluator, and acts as a major point of convergence where external stimuli is received, evaluated, and redirected for motivation of appropriate behavioral response.

In this review, we address the role of the habenula in five main sections: (1) Anatomy of Habenula and Known Connectivity with Afferent and Modulatory Pain Systems – here, we

report on the known anatomy of the structure; (2) *The Habenula in Pain and Analgesia* – we provide a summary of preclinical data suggesting a major role for the structure in pain and analgesia; (3) *Potential Role of the Habenula in Chronic Pain* – given the pivotal role in a number of putative processes that are part of the chronic pain experience (e.g., depression) we integrate data on habenula functions that are salient to chronic pain; (4) *Therapeutic Targets and the Habenula* – is a section that focuses on pharmacological modulation of the structure; and (5) in the section, Functional Imaging – Potential to Further Understanding the Habenula's Role in Pain and Analgesia, we look to the future in utilizing such approaches in the clinical domain in chronic pain.

2. The habenula and afferent and efferent connections

The connectivity of the habenula is complex (Herkenham and Nauta, 1979; Sutherland, 1982; Bianco and Wilson, 2009). The habenula is present across species, being larger relative to the whole brain in subhuman mammals. As shown in Fig. 1, it is a small bilateral structure in the dorsal-posterior thalamus, and sits astride the inferior lateral ventricles and above the pineal complex. It is divided into medial and lateral components that have different afferent and efferent connections that underlie differences in their function. Afferent and efferent pathways to and from the medial and lateral habenula show a segregation of functional connectivity. Fig. 2 shows connectivity of the medial (MHb) and lateral (LHb) habenula from other central brain and spinal cord regions. The habenula seems to be organized into multiple channels conveying parallel streams of information to the contralateral habenula, midbrain, and brainstem (Kim, 2009). In addition, inter and intrahabenular connections have been defined. The connections between the habenula nuclei are asymmetrical; only the medial habenula sends projection to the lateral habenula (Kim and Chang, 2005) while the commissural projection arises primarily in the lateral nucleus (Kim, 2009).

Pain inputs to the habenula include direct and indirect (via structures such as the lateral hypothalamus) afferents from the spinal cord. The lateral hypothalamus is involved in pain modulation (Dafny et al., 1996). Of the known afferent inputs that may be involved in pain transmission, the lateral hypothalamus projects to the habenula complex (Herkenham and Nauta, 1977; Parent et al., 1981). Stimulation of the habenula significantly modulates lateral hypothalamus noxious-evoked activity. The lateral hypothalamus receives afferents from deep and superficial lamina in the spinal cord (Burstein et al., 1990). Others have also reported direct spinal-habenula connections in the cat (Craig, 2003), where spinal lamina I injections of the anterograde tracer Phaseolus vulgaris leucoagglutinin (PHA-L) terminate in the lateral habenula. Within the structure, the lateral habenula (LHb) may be more sensitive to the inhibitory effects of μ opioid receptor agonist, morphine, on nociceptive stimuli (Wu et al., 2005), supporting a more prominent role of the lateral habenula in pain processing. Dense μ opioid receptor levels are described in the habenula (Mansour et al., 1987) while delta and kappa receptors are less dense in the region.

Other indirect effects of pain may take place through multiple inputs to the habenula from CNS structures such as the frontal cortex and nucleus accumbens (NAc) (see Fig. 2). Of these, the inputs from limbic forebrain structures, well known to be involved in pain processing (Casey, 1999), include the NAc (Becerra and Borsook, 2008), and the basal ganglia (Chudler and Dong, 1995; Borsook et al., 2010).

As shown in Fig. 2, the medial habenula (MHb) projects predominantly to the interpeduncular nucleus in the midbrain (Herkenham and Nauta, 1979) while the lateral habenula has projections to the ventral tegmental area (VTA) and the substantia nigra pars compacta (SNc). The two latter structures contain dopaminergic neurons (Bromberg-Martin

et al., 2010). The lateral habenula also has projections to the dorsal and median raphe nuclei that contain serotonergic neurons (Ferraro et al., 1996; Alenina et al., 2006). Prior studies initially suggested that periaqueductal gray (PAG), nucleus accumbens, and habenula might constitute a unidirectional loop to play their roles in pain modulation (Ma and Han, 1991). More recent works indicate that intrahabenular naloxone injections inhibit analgesia produced by morphine injected into the PAG. In addition, intra-accumbens injections of naloxone diminish the analgesic effects of intrahabenular morphine injections (Ma et al., 1992), suggesting that the habenula acts as a relay in descending pain modulation from the nucleus accumbens to the PAG (Yu and Han, 1990). Acetylcholinergic projections from the MHb to the IPN are, also, well described (Fasolo et al., 1992).

The projections to the midbrain regions such as the PAG and raphe are important in habenula-mediated analgesia. As noted, electrical stimulation or morphine injection into the habenula produces analgesia. The PAG has a well described role in descending modulation of pain (Behbehani, 1995; Ren and Dubner, 2002) and the dorsal raphe, which sends afferents directly to the spinal cord via the raphe magnus in the medulla, also contribute through serotonergic mediated pain modulatory processing (Wang and Nakai, 1994). Both serotonergic and dopaminergic systems have important roles in pain processing – including pain modulation (serotonin) (Fields, 2004; Benarroch, 2008) and reward (dopamine) (Wood, 2008). Finally, the interpeduncular nucleus receives a major output from the medial habenula and modulates cells in the raphe magnus (Hentall and Budhrani, 1990) in a manner that may enhance a pain response (Meszaros et al., 1985).

The habenula maintains an array of cross-talk between networks through different neurotransmitter systems. For example, main outputs of the LHb are to the midbrain *serotonergic system* where it targets the dorsal raphe and median raphe, the *dopaminergic system* where it targets the ventral tegmental area (VTA), and the *cholinergic system*, that targets the lateral dorsal tegmental nucleus (Sutherland, 1982; Klemm, 2004).

3. The habenula in pain and analgesia

Convergent preclinical data have implicated the habenula in numerous behaviors including pain and analgesia. Direct afferent inputs to the structure have been reported to originate in lamina I of the dorsal horn and trigeminal nucleus (Craig, 2003). Other inputs, notably from the hypothalamus (a structure also involved in pain processing as described earlier), have afferent inputs to the habenula (Goto et al., 2005). Thus, the structure is directly connected to pain processing through these pathways. Other pathways (see section above) have also been reported that may also convey information related to pain and analgesia. For example, anatomical and electrical studies show connections to the midbrain central gray and serotonergic raphe nuclei (Ferraro et al., 1996), both involved in pain modulation.

Further evidence can be found from activation observed in electrophysiology or electrostimulation studies. Neurons in the lateral habenula respond to noxious stimuli (but not non-noxious stimuli) that may be excitatory or inhibitory depending on the intensity of the stimulus (Gao et al., 1996). Electrical stimulation of the habenula produces analgesia in a formalin test (Cohen and Melzack, 1986) and is reversible when naloxone is directly injected into the habenula (Mahieux and Benabid, 1987). Other markers of activity include neuronal markers and functional imaging. Using c-fos immunohistochemistry as a marker of neuronal activity shows increases in neuronal stains in the lateral habenula in experimental pain (Smith et al., 1997; Lehner et al., 2004). In a chronic model of pain (compression of the dorsal root ganglion) fos expression has been reported to be transiently increased in the habenula (Sun et al., 2002). In a measure of functional activity utilizing 2-deoxyglucose, diabetes-induced pain with streptozotocin (STZ) in rats, habenula activation was decreased

in response to evoked pain inducing stimuli compared with control rats (Paulson et al., 2007), suggesting that habenula activation may represent part of a modulatory pain system. Further details on imaging the habenula are noted in Section 5.

Preclinical data have also evaluated the effects of analgesics on habenula function. For example, morphine injections into the habenula, a region containing high levels of opioid receptors (Neal et al., 1999), result in analgesia (Cohen and Melzack, 1985). In addition, heroin decreases habenula glucose utilization supporting a role of opioid mediated responses of neurons (Martin et al., 1997). Opioids, including morphine, may have a dual modulation of glutamatergic inputs in the structure to produce these changes.

Taken together, the data support a role of the habenula in pain processing. As noted in the section on anatomy of the structure, it is situated to receive information from forebrain structures and the dorsal diencephalic conduction (DDC) system – one of two major pathways that interconnect the limbic forebrain and sites in the mid- and hindbrain (Bianco and Wilson, 2009). As such, the emotional components of pain may interact with pain modulatory systems through this structure. This is perhaps highly salient given emotional changes in chronic pain (including cognitive and reward) are clearly part of what makes chronic pain difficult for patients.

4. Habenula and behaviors associated with chronic pain

Chronic pain is associated with a number of behaviors and complex interactions including sensory, emotional, and cognitive alterations. Given the involvement of the habenula in pain (see Section 2), research related to behaviors that are considered part of overall pain behavior including analgesia suggests a potential integrative role for this structure. As noted above, the structure may be an important interactive pathway linking the limbic forebrain and brainstem nuclei to the midbrain. Chronic pain is associated with a number of alterations including a deficient reward state due to changes in reward–aversion processing; stress; in co-morbid conditions such as anxiety, depression or addiction; and in more complex processes such as defensive behaviors and prediction errors. It is perhaps highly salient that stimulation of the habenula may reverse some of these behaviors including depression (Sartorius et al., 2010) or potentially addiction (Luigjes et al., 2011).

4.1. Reward and aversion

The habenula has been implicated in reward (Hong and Hikosaka, 2008; Hikosaka, 2010). Inhibitory input from the lateral habenula (LHb) shows major influence in determining the reward-related activity of dopamine neurons (Matsumoto and Hikosaka, 2007). A projection from the globus pallidus to the lateral habenula is one pathway where reward prediction processing is activated in the habenula by pallidal inputs (Hong and Hikosaka, 2008). Given the connections of the habenula to dopaminergic regions in the VTA and SNc, the structure likely plays a role in reward-related signals. Analgesia may be a process driving these reward related activities (Schweinhardt et al., 2009). Consistent with a putative role of the habenula in negative reinforcement or responses to aversive stimuli (such as pain), stimulation of the lateral habenula results in a reward decrease as measured by sucrose self administration (Friedman et al., 2011).

Pain and addiction are considered to be at the opposite ends of the reward–aversion spectrum (Elman et al., 2011). A putative role of the structure in addiction has been supported by data relating to increased activity during drug-seeking behaviors (Zhang et al., 2005), and controlling nicotine intake (Fowler et al., 2011). Furthermore, attenuation of dopamine sensitization in the accumbens is observed following treatment with 18-Methoxycoronaridine (18-MC), an iboga alkaloid congener (Taraschenko et al., 2007).

Behaviorally 18-MC inhibits morphine self-administration in rats (Glick et al., 2006) through actions on the medial habenula. The issue of addiction and pain are complex, and it is a major focus of research as to whether pain enhances addiction potential given that chronic pain may be considered a 'reward deficit state' (Elman et al., 2011) while opioid addicts have altered pain processing (Ren et al., 2009) and purported to have opioid induced hyperalgesia (Fishbain et al., 2011).

4.2. Stress and defensive behaviors

Pain is a stressor, and the habenula, as understood from its relationship with aversive events, acts as a behavioral mediator to stressors. Implications of pain as a stressor are particularly prominent if chronic or repeated (e.g., in migraine). Studies of the medial habenula (MHb) have shown involvement in neuroendocrine and immunological responses to various kinds of stress (Blake et al., 1990). Measures of c-fos expression in primates suggests that the habenula maintains prolonged activation under traumatic stress and appeared to play a role in the process of long-term stress (Kazi et al., 2004). Such data support a function in long-term behavioral alterations after uncontrollable stress (Amat et al., 2001). Habenular influence over other behavioral functions, also appear to be enhanced with higher induced stress levels. For example, habenular lesions in rat shock avoidance experiments showed little differences to sham operated controls, until stress levels were heightened by more demanding tasks, or higher shock levels (Thornton and Bradbury, 1989). Thus, the habenula seems to maintain a role in promoting decision-making and survival driven reactions (Hikosaka, 2010) to aversive stressors that would include pain.

An extension of fear, stress, and anxiety are the resultant defensive behaviors associated with them. Serotonergic systems have been implicated in defensive behaviors through the lateral habenula (LHb that may differ depended on the 5-HT receptor subtype involved (Pobbe and Zangrossi, 2010)). Habenula lesions eliminate 5-HT increases in the dorsal raphe to inescapable shocks (Amat et al., 2001). Rats subject to swim tests after habenular lesions did not utilize escape behaviors which was not enhanced by an antidepressant nomifensine (despite promoting the escape behavior in control animals) (Thornton et al., 1985). It seems habenula-signaled adaptive behaviors to fear, anxiety, or stress further promote follow-up performance such as escape, removal, or avoidance suggesting that the habenula regulates defensive response to fear and anxiety-related inputs (Pobbe and Zangrossi, 2008). The structure seems to play a significant role in defensive behaviors including present and future avoidance. In the context of chronic pain, fear avoidance is a major problem (Rainville et al., 2011).

4.3. Fear and anxiety

Fear and anxiety also have a significant role in both acute and chronic pain (Ploghaus et al., 1999; den Hollander et al., 2010). The expectation of the pain modulates (usually increases) the pain sensation as well as playing a role in anxiety and phobia (Ploghaus et al., 2003). These interactions are a part of the important adaptive behaviors that lead to the avoidance of pain-causing agents, where fear and anxiety are the emotional mitigators to those behaviors. The habenula link to anxiety has been shown in lesioning studies where habenula efferent pathway damage resulted in enhanced levels of anxiety behaviors in rats (Murphy et al., 1996). The authors postulate that the effects on anxiety related to either a disruption of lateral habenular (LHb) projections to dopaminergic neurons in the ventral tegmentum or to regions high in benzodiazepine receptors. Further support is found in maternity studies; sensory inputs from infants or pups modulate anxiety during the postpartum period (Smith and Lonstein, 2008). Zebrafish studies have reported anxiety regulation by the habenula (Jesuthasan, 2011) and habenula efferents in controlling experience-dependent fear response (Agetsuma et al., 2010). Specifically, the dorsal habenula-interpeduncular nucleus-griseum

central pathway is critical for the modification of response in an experience-dependent manner. Some have considered the habenula as a “switching board for selection of behavioral strategy to cope with fear and anxiety” (Okamoto et al., 2011). As such it may potentially control a variety of emotional behaviors in chronic pain.

4.4. Depression

Pain and depression are frequently co-morbid (Borsook et al., 2007). Functional, chemical and anatomical changes from chronic pain can, in time, alter the sensory system, emotional and motivational systems and result in comorbidity such as depression. These changes, in turn, will affect the reward/aversion process and are plausible contributors to psychiatric disorders that have been seen to develop in chronic pain sufferers (Borsook et al., 2007). Consistently, many therapies for chronic pain and depression overlap (Borsook et al., 2007). Major depressive disorder (MDD), in general, is associated with dysfunction of the central serotonergic system. The principal source of serotonin is in the raphe nuclei in the midbrain. The influence of serotonin (5-HT) manipulation on emotional state and depression patients has been studied through tryptophan manipulation (Morris et al., 1999; Roiser et al., 2009; Evers et al., 2010). Following tryptophan depletion, habenula blood flow increases significantly in remitted major depressive disorder patients when subject to emotional stimuli (Roiser et al., 2009). The result suggests that in addition to a consistent habenular response to negative feedback, it further shows a specific sensitivity with depressive patients and might play a role in the pathogenesis of MDD. Magnetic imaging studies of the habenula volume in MDD and bipolar disorder suggest that a volume reduction might contribute to the risk of developing or the result of such affective disorders (Savitz et al., 2011). In treatment resistant depression, trials of deep brain stimulation (DBS) have been used in different regions including the major afferent bundle of the lateral habenula, which was reportedly successful in a therapy resistant patient (Sartorius et al., 2010).

4.5. Higher level functioning – cognition and prediction errors

A habenular role in cognition can be synthesized from behaviors following dysfunction. Symptoms of altered memory and attention in rats follow from habenula lesions (Lecourtier et al., 2004). These deficits have improved and in some cases declined over repeated testing, a plausible scenario for the habenula having multiple roles in cognition. In human populations, schizophrenia patients report deficits in utilizing feedback to promote appropriate problem-solving skills and learning. This is consistent with the habenular activity and dopamine damper to negative feedback as a way to guide follow-up responses. In human studies alteration of habenula activity is reported in schizophrenia patients that have altered processing to negative outcomes (Shepard et al., 2006). Lacking this activity through a limited functioning habenula seems to promote feedback-processing deficits in schizophrenia patients (Shepard et al., 2006).

Prediction error is a theoretical signal promoting learning by encoding midbrain dopamine neurons to inform the rest of the brain of expectancy and outcome mismatch (Schultz, 2010). The relationship between habenula and error evaluation has been studied in both animals (Shepard et al., 2006; Ji and Shepard, 2007; Matsumoto and Hikosaka, 2007) and humans (Ullsperger and von Cramon, 2003). It is thought that mesencephalic dopaminergic neurons are inhibited by the habenula through glutamatergic afferent input onto GABAergic neurons in the VTA (Omelchenko et al., 2009; Brinschwitz et al., 2010) and SN (Christoph et al., 1986). When expected rewards are not received or negative feedback is solicited (Christoph et al., 1986), the habenula fires through the emotional system to reinforce behavior avoidance. Supporting data are found from electro-stimulation of habenula where there is a substantial 80–95% dopamine decrease (Ji and Shepard, 2007), and in lesion studies where dopamine levels show an increase (Nishikawa et al., 1986). Detection likely

occurs from dopamine levels in the rostral cingulate motor area (rCMA) and reward values are associated with the neural pathway controlling the dopamine system, the pallidus-habenula-dopamine pathway, relying on a combination of stimulus reward pairing and abstract inference as the mechanism for determination (Bromberg-Martin et al., 2010). It seems plausible that neuronal pathways excited during pain stimulation would similarly be filtered through a negative reward analysis system. Regions involved in prediction errors for aversive events have been reported, but the habenula was not (Ploghaus et al., 2000). However, the relationship of pain and reward circuitry, noted in a report from our group (Becerra et al., 2001) remains an area of intense interest and involves structures including the nucleus accumbens. More recent human studies (see below – imaging the habenula) have evaluated negative prediction errors (Salas et al., 2010). The issue of error detection and chronic pain may be considered in light of interoceptive behaviors (Ray and Slobounov, 2002) that may be altered in chronic pain as a result of alteration in anterior insula and cingulate dysfunction (Baliki et al., 2006; Coen et al., 2009).

4.6. A model for habenula's interactions in chronic pain

Conceptually, it is useful to consider functional features of the habenula in the context of interactions between its afferent and efferent connections which, based on preclinical data, suggest some form of functional loop. Fig. 3 is a model of alterations in habenula functional loops in chronic pain. As noted in Fig. 2, afferent inputs to the lateral and medial habenula send efferents to brainstem regions. These in turn feed back onto many of the afferent pathways. These can be divided into analgesia/hyperalgesia, sensory-motor, stress, and reward/aversion habenula loops. Each loop or circuit relates to basic functions of the habenula that are altered in acute and chronic pain (see Fig. 3). As noted in the figure, in the chronic pain condition there are increases or decreases in these functional circuits (e.g., reward loop diminishes and stress circuits increase). The model, derived from preclinical data showing increased activation (electrophysiology, fos) in acute pain, and increased metabolism in chronic pain (see Fig. 3) is expressed in terms of expected functional activity in the habenula. Clearly, while the structure is not the epicenter of altered function in chronic pain (since pain is a distributed function), it would seem that it could play a significant role in sensory, emotional and modulatory processing of pain.

5. Analgesics and the habenula

Early preclinical studies indicate that the habenula is involved in analgesia through direct electrical stimulation (Cohen and Melzack, 1993). While many brain structures are involved in analgesia (e.g., PAG and cingulate), there is growing evidence that the habenula, too, plays a role in analgesia through direct effects on neurotransmitter systems present in the structure or secondary effects, say for example, by enhancing frontal (cortical and subcortical) inputs into a structure where treatments may alter these inputs and drugs may enhance brainstem processes well known to play a role in chronic pain. Overall, the structure is modulated or directly modulates systems or pathways involved in pain processing including opioridergic, serotonergic, dopaminergic, and noradrenergic.

5.1. Opioidergic systems

Opioids are present in the habenula. Indeed, some of the highest levels of μ receptor RNA in the brain have been reported in the medial habenula (Zastawny et al., 1994; Bunzow et al., 1995). Morphine acts directly on habenula neurons to produce analgesia (Zhou et al., 1981; Cohen and Melzack, 1985). Such information clearly implicates an analgesic effect that is observed when morphine is directly injected into other brain structures including the cingulate cortex (LaGraze et al., 2006) and the periaqueductal gray (Carstens et al., 1988; Bernal et al., 2007). While it can inhibit and facilitate excitatory synaptic transmission

(Hashimoto et al., 2009), how it affects analgesia through direct action on the habenula is unclear, but assumed to be through activation of brainstem modulatory systems. In regard to the latter, such a drive could potentially act on facilitatory or inhibitory systems through serotonergic systems as has been shown in animal models of chronic pain (Dogrul et al., 2009). In addition, a complex interaction between opioids and dopaminergic systems may disrupt pathways such as efferent systems that include the habenulo-interpeduncular pathway that may modulate a dopaminergic mesolimbic pathway. Levels of other opioids including nociception in the habenula (Schulz et al., 1996; Florin et al., 2000) suggest alternate functions of opioid induced neural processing. Furthermore, at least what may be deduced from other drugs of abuse is that chronic use results in degeneration in axons from the lateral habenula to midbrain (including the SN, VTA and raphe) (Ellison, 2002). Thus opioids may modulate habenula function and if given chronically may alter habenula-brainstem connectivity.

5.2. Dopaminergic systems

Dopamine is involved in reward processing (Wise and Rompre, 1989; Schultz, 2007) and also in pain and analgesia (Wood, 2008; Lapirot et al., 2011; Sarkis et al., 2011). Given the projections of the lateral habenula to dopaminergic nuclei (Goncalves et al., 2011), some of these inputs are inhibitory on midbrain dopaminergic cells (Balcita-Pedicino et al., 2011); specifically glutamatergic habenula neurons act on inhibitory GABAergic cells (Brinschwitz et al., 2010). Consistent findings show electrical stimulation of the LHb results in inhibition of neural activity of dopamine neurons (Matsumoto and Hikosaka, 2007). The effects of dopaminergic drugs on the habenula have also been evaluated. The dopaminergic agonist apomorphine produces excitation of individual LHb neurons (specifically in the lateral division of the LHb) (Kowski et al., 2009). There is also a complex regulation of habenula involvement in dopamine and reward: neurons in the LHb are known to encode reward-related signals that are opposite dopamine neurons; thus, LHb neurons are inhibited by reward and sensory stimuli predicting reward and excited by reward omission (Matsumoto, 2009). In addition, the habenula affects prefrontal and NAc dopamine release (Lecourtier et al., 2008). Thus, the habenula is intricately involved in reward processing thorough control of dopaminergic neurons in the forebrain and brainstem (Matsumoto and Hikosaka, 2007; Matsumoto, 2009; Bromberg-Martin and Hikosaka, 2011).

5.3. Noradrenergic systems

Peripheral (superior cervical ganglionectomy) and central (striatal) lesions result in increases in norepinephrine in the habenula (Gottesfeld, 1983). The former produced increases in the LHB and the latter in both the LHb and MHb (Gottesfeld, 1983). Brain NA systems are activated by stress (Morilak et al., 2005). High levels of serotonin receptors subtypes (5-HT₅) are reported to be present in the medial habenula of wild type (Waeber et al., 1998). Taken together, the effects of SNRIs in the treatment of chronic pain (Wright et al., 2011) may reflect these drug effects on the habenula.

5.4. Serotonergic systems

Both human and animal (Meng et al., 2011) models of habenula stimulation seem to reverse depression, implicating the serotonergic systems. Supporting data have shown excitatory amino acids in the lateral habenula-dorsal raphe pathway (see Fig. 2) are involved in the regulation of striatal 5-HT release (Kalen et al., 1989). In addition, fluoxetine, a selective serotonin reuptake inhibitor (SSRI) reduced metabolism in the habenula (Shumake et al., 2010). The potential mechanism of this may be through local serotonergic receptors, notably (5HT-B) (Kinsey et al., 2001), 5-HT_{2C} (Pompeiano et al., 1994) and 5-HT₅ which is similar to the 5HT_{1D} family (Plassat et al., 1992). Thus pharmacological agents that are effective in chronic pain may also contribute to their analgesic action via habenula serotonergic systems.

5.5. Other habenula neurochemical systems that may be related to pain processing

A number of other neurochemical systems are implicated in habenula function. These include the NMDA receptor antagonists which decrease activity in the habenula (Weissman et al., 1989; Eintrei et al., 1999); calcitonin gene related peptide CGRP (Skofitsch and Jacobowitz, 1985) as is substance P (Neckers et al., 1979) which is present in high levels in rat habenula. However, the involvement and distribution of these peptides is complex; for example, NK-1 and NK-3 receptor agonists excite habenula neurons and electrophysiological recording suggest that these receptors are found on different populations of neurons in medial habenula (Norris et al., 1993; Waeber et al., 1998). A cholinergic pathway from the medial habenula to the IPN has been defined (Fasolo et al., 1992). A putative role in pain is unclear but may relate to addictive propensity for nicotine (Fowler et al., 2011) and it is well documented that chronic pain and smoking are frequently comorbid.

5.6. Habenula and anesthesia

In addition to analgesics, the structure may also contribute to the underlying effects of anesthetics. Anesthesia results in loss of sensation with or without loss of consciousness and may also affect pain systems. Interestingly, a number of reports show the involvement of the habenula in anesthesia. For example, the anesthetic althesin a steroid anesthetic produced increased metabolic activity in the habenula that is inhibited by lesioning inputs from the stria terminals (McQueen et al., 1984). Thiopental also increased habenula activation in rats exposed to a noxious stimulus and the authors postulated that it may have a role in pain processing (Archer et al., 1995). So while a specific role is not understood, the habenula's interaction with anesthesia seems to relate to its role in analgesia.

5.7. Habenula deep pain stimulation – potential therapy for chronic pain

Both animal and human reports of lateral habenula stimulation provide a potential future therapeutic avenue for this approach in chronic pain patients refractive to other treatments. In studies on depression, deep brain stimulation (DBS) has been used in animal models of depression (Meng et al., 2011). In a report of a depressed patient refractory to other treatments DBS (see Fig. 4) reportedly produced remission (Sartorius et al., 2010). In addition, DBS of the lateral habenula also resulted in decreased drug or reward seeking behavior (thought to be through increases of glutamatergic drive to the VTA) (Friedman et al., 2010, 2011). Given the overlap of neural circuits in depression, addiction and pain (Borsook et al., 2007; Elman et al., 2011), the issue of whether DBS of the habenula produces pain relief in chronic pain models needs to be studied, not only to better understand the role of the structure in pain processing but also as a potential therapy.

Few studies have evaluated analgesic responses of the habenula in chronic pain models. It is likely that drugs that target the habenula may provide a useful model for chronic pain because of its integrative role in pain and analgesia. Perhaps, rather than a role in pain sensitivity or intensity, it is more likely to play a role in chronic pain conditions including migraine. Thus, it may be an ideal “light-house” or indicator of the potential effects of analgesics on chronic pain. This is particularly salient given new insights such as activation of inhibition of the rostro-ventro-medial medulla in inhibiting neuropathic pain (De Felice et al., 2011) and the habenula's modulation of these regions as well as issues related to tonic pain (King et al., 2009).

6. Functional imaging of the habenula

As noted in the Introduction, functional markers of habenula to pain activity include measures of neuronal markers (e.g., c-fos), electrophysiology (Andersen and Dafney, 1983;

Hua et al., 1998), and metabolic markers (Paulson et al., 2005, 2007). More recently the use of imaging techniques such as functional magnetic resonance imaging (fMRI) has been applied in humans in non-pain paradigms.

6.1. Functional imaging of the habenula in animals

Unlike most neuroimaging of pain, the animal data have preceded the human data. Pain and analgesic functional imaging studies have been performed in animals (Borsook and Becerra, 2011) but few have reported habenula activation. Studies employing imaging measures in pain models suggest prominent activation in the habenula (Fig. 5A). Using a marker of neuronal activity (99m)Tc-HMPAO, two studies report activation in the habenula. In a model of diabetes-induced neuropathic pain, decreased activation (compared with control animals) was seen in animals shown to have hypersensitivity to pain (Paulson et al., 2007). The conclusion of this study was that the habenula contributed to modulation (inhibition of pain) since the midbrain displayed a similar decrease in activity (Fig. 5B, top row) in contrast to activation in other regions particularly those encoding pain intensity (somatosensory cortex, ventrobasal thalamic nuclei). In an earlier study from the same group, spinal cord damage produced similar changes in the habenula (Paulson et al., 2005) (Fig. 5B, bottom row).

6.2. Functional imaging of the habenula in humans

While functional imaging techniques have advanced in humans with higher field systems (3 T and 7 T for fMRI), there are challenges imaging the habenula because it is a small structure, approximately 3 mm × 3 mm × 3 mm in humans. Current fMRI imaging protocols acquire images with a resolution of 3–5 mm. This is barely large enough to cover the habenula with one voxel. Accordingly, it is necessary to optimize current protocols to focus in the area. However, defining its location in imaging studies is made somewhat easier by its alignment to the third ventricle as well as its proximity to the pineal gland in the horizontal plane. A few functional imaging studies have reported activation in the habenula using either fMRI (Ullsperger and von Cramon, 2003; Roiser et al., 2009), or positron emission tomography (PET) (Morris et al., 1999; Sartorius et al., 2010). In a study by Morris et al., the effect of altering tryptophan (a precursor of serotonin) levels in patients with affective disorders was evaluated. Since the habenula projects to the raphe nucleus, the study took advantage of its ability to influence the activity of serotonergic neurons in the raphe nuclei and modulate release of cerebral serotonin. Subjects performed simple tasks (verbal fluency and word repetition) under these conditions during PET scans. Results show that the extent of mood change following tryptophan depletion is related to the covariation of activity between two serotonergic sub-cortical nuclei, the habenula and dorsal raphe (an increase drive to these areas with less tryptophan). Highly correlated activity between these structures was present only in subjects with high post-depletion depression ratings (Morris et al., 1999).

In an fMRI study monitoring habenular function for error and reward prediction, data were acquired during informative and non-informative feedback stimuli amid a task involving subject decisions (Ullsperger and von Cramon, 2003). The aim of the study was to determine specific regions of brain activity relating to negative and positive feedback. The expectation was a larger hemodynamic activity of habenular region on errors with negative feedback and a lower hemodynamic activity on correct responses with positive feedback. Positive feedback raised hemodynamic activity in the ventral striatum (nucleus accumbens) and the putamen, negative feedback activated the rCMA, the inferior anterior insula, and the epithalamus (habenular complex). Functional connectivity analysis has shown the direction of interaction between amygdala and VTA/SN, determining the habenula as the mediator during error detection (Ide and Li, 2011). A review of the literature suggests habenula

activation may be present some human fMRI reports on brain activation by pain although not specifically reported (Fig. 5C). Based on the animal literature of increased activation in acute and chronic pain, we provide a summary of expected activation levels within the structure for fMRI activation for acute and chronic pain (Fig. 5) and to be reversed following analgesic administration (not shown).

In a habenula targeted deep brain stimulation (DBS) in a patient who had suffered from treatment resistant depression, localization of neurological changes over time were made by co-registering PET T1 and T2 and subtracting for a difference image. This image and a post-operation CT image were both co-registered to a high resolution T1 MRT scan with the same transformation parameters (Sartorius et al., 2010). Metabolic effects can be seen in the fluorodeoxyglucose positron emission tomography images in Fig. 4. Although a small structure, these and other fMRI studies (D'Ardenne et al., 2008; Li et al., 2008) thus define that habenula can be imaged in humans and that the approaches may have important implications for pain and analgesia (viz., error and reward, depression, etc.) because of the putative role of the structure.

Under separate fMRI studies where pain itself was the research topic, activation that is likely to overlap the habenula can be seen in many of the resulting images. While determination of activation for such a small region cannot visually be determined with 100% accuracy, the data are nonetheless convincing and warrants further exploration with more robust imaging methods. A habenular response to pain could help devise a much-sought quantitative measure for the pain experience of an individual, removing subjectivity that can interfere with diagnostic and pain treatment. It has the potential to lead to a determinant that distinguishes an injury or disease that will lead to chronic pain. It would seem further understanding of the function in humans through neuroimaging would provide exciting new information on brain systems involved in pain and analgesia.

7. Conclusions

The habenula is involved in numerous behavioral functions that are implicated in pain processing (viz., cognition, addition, aversion, learned helplessness, and reward, Hikosaka, 2007; Matsumoto and Hikosaka, 2007; Frahm et al., 2011; Friedman et al., 2011; Paul et al., 2011; Li et al., 2011; Paul et al., 2011). Through various loops, habenula outputs modulate the inputs from forebrain structures and thus the structure may modulate the interactions between frontal brain regions involved in cognitive, reward, and various brainstem functions including pain modulation. With respect to the latter, these outputs to brainstem regions modulate sensory gating (Ellison, 1994), protective behaviors (Pobbe and Zangrossi, 2010) and endogenous tone in response to pain and may be thus construed as involved in stress evasion (Hikosaka, 2010). However, the role of the habenula in chronic pain in the human condition is yet to be determined. Given the insights of this structure on conditions that are seen with chronic pain including addiction (Fowler et al., 2011) and depression (Sartorius et al., 2007), abnormal functioning in the habenula may have a significant role in the chronic pain state. In this chronic pain, abnormal function of habenula may contribute to pain behavior through dysfunction of “intrinsic central process that coordinates various selective functions (including perceptual, visceral, and reinforcement processes) into a global dysfunctional state” (Ikemoto, 2010). The use of neuroimaging should provide new insights of the habenula in the human chronic pain condition.

Acknowledgments

This work was supported by a grant from NIH (1R01NS065051) and a K24 Mentoring grant (5K24NS64050).

Abbreviations

MHb	medial habenula
LHb	lateral habenula
NAc	nucleus accumbens
VTa	ventral tegmental area
PAG	periaqueductal gray
fMRI	functional magnetic resonance

References

- Agetsuma M, Aizawa H, Aoki T, Nakayama R, Takahoko M, Goto M, Sassa T, Amo R, Shiraki T, Kawakami K, Hosoya T, Higashijima S, Okamoto H. The habenula is crucial for experience-dependent modification of fear responses in zebrafish. *Nat. Neurosci.* 2010; 13:1354–1356. [PubMed: 20935642]
- Alenina N, Bashammakh S, Bader M. Specification and differentiation of serotonergic neurons. *Stem Cell Rev.* 2006; 2:5–10. [PubMed: 17142880]
- Amat J, Sparks PD, Matus-Amat P, Griggs J, Watkins LR, Maier SF. The role of the habenular complex in the elevation of dorsal raphe nucleus serotonin and the changes in the behavioral responses produced by uncontrollable stress. *Brain Res.* 2001; 917:118–126. [PubMed: 11602236]
- Andersen ERB, Dafney N. Electrophysiological evidence of concurrent dorsal raphe input to caudate, septum, habenula, thalamus hippocampus, cerebellum and olfactory bulb. *Int. J. Neurosci.* 1983; 18:107–115. [PubMed: 6302024]
- Archer DP, Froelich J, McHugh M, Pappius HM. Local cerebral glucose utilization in stimulated rats sedated with thiopental. *Anesthesiology.* 1995; 83:160–168. [PubMed: 7604994]
- Balcita-Pedicino JJ, Omelchenko N, Bell R, Sesack SR. The inhibitory influence of the lateral habenula on midbrain dopamine cells: ultrastructural evidence for indirect mediation via the rostromedial mesopontine tegmental nucleus. *J. Comp. Neurol.* 2011; 519:1143–1164. [PubMed: 21344406]
- Baliki MN, Chialvo DR, Geha PY, Levy RM, Harden RN, Parrish TB, Apkarian AV. Chronic pain and the emotional brain: specific brain activity associated with spontaneous fluctuations of intensity of chronic back pain. *J. Neurosci.* 2006; 26:12165–12173. [PubMed: 17122041]
- Becerra L, Borsook D. Signal valence in the nucleus accumbens to pain onset and offset. *Eur. J. Pain.* 2008; 12:866–869. [PubMed: 18226937]
- Becerra L, Breiter HC, Wise R, Gonzalez RG, Borsook D. Reward circuitry activation by noxious thermal stimuli. *Neuron.* 2001; 32:927–946. [PubMed: 11738036]
- Behbehani MM. Functional characteristics of the midbrain periaqueductal gray. *Prog. Neurobiol.* 1995; 46:575–605. [PubMed: 8545545]
- Benarroch EE. Descending monoaminergic pain modulation: bidirectional control and clinical relevance. *Neurology.* 2008; 71:217–221. [PubMed: 18625968]
- Bernal SA, Morgan MM, Craft RM. PAG mu opioid receptor activation underlies sex differences in morphine antinociception. *Behav. Brain Res.* 2007; 177:126–133. [PubMed: 17118467]
- Bianco IH, Wilson SW. The habenular nuclei: a conserved asymmetric relay station in the vertebrate brain. *Philos. Trans. R. Soc. Lond. B: Biol. Sci.* 2009; 364:1005–1020. [PubMed: 19064356]
- Blake MJ, Nowak TS Jr, Holbrook NJ. In vivo hyperthermia induces expression of HSP70 mRNA in brain regions controlling the neuroendocrine response to stress. *Brain Res. Mol. Brain Res.* 1990; 8:89–92. [PubMed: 2166205]
- Borsook D, Becerra L. CNS animal fMRI in pain and analgesia. *Neurosci. Biobehav. Rev.* 2011; 35:1125–1143. [PubMed: 21126534]

- Borsook D, Becerra L, Carlezon WA Jr, Shaw M, Renshaw P, Elman I, Levine J. Reward-aversion circuitry in analgesia and pain: implications for psychiatric disorders. *Eur. J. Pain.* 2007; 11:7–20. [PubMed: 16495096]
- Borsook D, Upadhyay J, Chudler EH, Becerra L. A key role of the basal ganglia in pain and analgesia—insights gained through human functional imaging. *Mol. Pain.* 2010; 6:27. [PubMed: 20465845]
- Brinshawitz K, Dittgen A, Madai VI, Lommel R, Geisler S, Veh RW. Glutamatergic axons from the lateral habenula mainly terminate on GABAergic neurons of the ventral midbrain. *Neuroscience.* 2010; 168:463–476. [PubMed: 20353812]
- Bromberg-Martin ES, Hikosaka O. Lateral habenula neurons signal errors in the prediction of reward information. *Nat. Neurosci.* 2011; 14:1209–1216. [PubMed: 21857659]
- Bromberg-Martin ES, Matsumoto M, Hikosaka O. Dopamine in motivational control: rewarding, aversive, and alerting. *Neuron.* 2010; 68:815–834. [PubMed: 21144997]
- Bunzow JR, Zhang G, Bouvier C, Saez C, Ronnekleiv OK, Kelly MJ, Grandy DK. Characterization and distribution of a cloned rat mu-opioid receptor. *J. Neurochem.* 1995; 64:14–24. [PubMed: 7798908]
- Burstein R, Cliffer KD, Giesler GJ Jr. Cells of origin of the spinothalamic tract in the rat. *J. Comp. Neurol.* 1990; 291:329–344. [PubMed: 2298937]
- Carstens E, Stelzer B, Zimmermann M. Microinjections of glutamate or morphine at coincident midbrain sites have different effects on nociceptive dorsal horn neurons in the rat. *Neurosci. Lett.* 1988; 95:185–191. [PubMed: 2906415]
- Casey KL. Forebrain mechanisms of nociception and pain: analysis through imaging. *Proc. Natl. Acad. Sci. U.S.A.* 1999; 96:7668–7674. [PubMed: 10393878]
- Christoph GR, Leonzio RJ, Wilcox KS. Stimulation of the lateral habenula inhibits dopamine-containing neurons in the substantia nigra and ventral tegmental area of the rat. *J. Neurosci.* 1986; 6:613–619. [PubMed: 3958786]
- Chudler EH, Dong WK. The role of the basal ganglia in nociception and pain. *Pain.* 1995; 60:3–38. [PubMed: 7715939]
- Coen SJ, Yaguez L, Aziz Q, Mitterschiffthaler MT, Brammer M, Williams SC, Gregory LJ. Negative mood affects brain processing of visceral sensation. *Gastroenterology.* 2009; 137:253–261. 261 e251-252. [PubMed: 19582887]
- Coghill RC, McHaffie JG, Yen YF. Neural correlates of interindividual differences in the subjective experience of pain. *Proc. Natl. Acad. Sci. U.S.A.* 2003; 100:8538–8542. [PubMed: 12824463]
- Cohen SR, Melzack R. Morphine injected into the habenula and dorsal posteromedial thalamus produces analgesia in the formalin test. *Brain Res.* 1985; 359:131–139. [PubMed: 4075141]
- Cohen SR, Melzack R. Habenular stimulation produces analgesia in the formalin test. *Neurosci. Lett.* 1986; 70:165–169. [PubMed: 3774213]
- Cohen SR, Melzack R. The habenula and pain: repeated electrical stimulation produces prolonged analgesia but lesions have no effect on formalin pain or morphine analgesia. *Behav. Brain Res.* 1993; 54:171–178. [PubMed: 8391825]
- Craig AD. Distribution of trigeminothalamic and spinothalamic lamina I terminations in the cat. *Somatosens. Mot. Res.* 2003; 20:209–222. [PubMed: 14675960]
- D'Ardenne K, McClure SM, Nystrom LE, Cohen JD. BOLD responses reflecting dopaminergic signals in the human ventral tegmental area. *Science.* 2008; 319:1264–1267. [PubMed: 18309087]
- Dafny N, Dong WQ, Prieto-Gomez C, Reyes-Vazquez C, Stanford J, Qiao JT. Lateral hypothalamus: site involved in pain modulation. *Neuroscience.* 1996; 70:449–460. [PubMed: 8848153]
- De Felice M, Sanoja R, Wang R, Vera-Portocarrero L, Oyarzo J, King T, Ossipov MH, Vanderah TW, Lai J, Dussor GO, Fields HL, Price TJ, Porreca F. Engagement of descending inhibition from the rostral ventromedial medulla protects against chronic neuropathic pain. *Pain.* 2011; 152:2701–2709. [PubMed: 21745713]
- den Hollander M, de Jong JR, Volders S, Goossens ME, Smeets RJ, Vlaeyen JW. Fear reduction in patients with chronic pain: a learning theory perspective. *Expert Rev. Neurother.* 2010; 10:1733–1745. [PubMed: 20977330]

- Dogrul A, Ossipov MH, Porreca F. Differential mediation of descending pain facilitation and inhibition by spinal 5HT-3 and 5HT-7 receptors. *Brain Res.* 2009; 1280:52–59. [PubMed: 19427839]
- Eintrei C, Sokoloff L, Smith CB. Effects of diazepam and ketamine administered individually or in combination on regional rates of glucose utilization in rat brain. *Br. J. Anaesth.* 1999; 82:596–602. [PubMed: 10472230]
- Ellison G. Stimulant-induced psychosis, the dopamine theory of schizophrenia, and the habenula. *Brain Res. Brain Res. Rev.* 1994; 19:223–239. [PubMed: 7914793]
- Ellison G. Neural degeneration following chronic stimulant abuse reveals a weak link in brain, fasciculus retroflexus, implying the loss of forebrain control circuitry. *Eur. Neuropsychopharmacol.* 2002; 12:287–297. [PubMed: 12126867]
- Elman I, Zubieta JK, Borsook D. The missing p in psychiatric training: why it is important to teach pain to psychiatrists. *Arch. Gen. Psychiatry.* 2011; 68:12–20. [PubMed: 21199962]
- Evers EA, Sambeth A, Ramaekers JG, Riedel WJ, van der Veen FM. The effects of acute tryptophan depletion on brain activation during cognition and emotional processing in healthy volunteers. *Curr. Pharm. Des.* 2010; 16:1998–2011. [PubMed: 20370668]
- Fasolo A, Virgili M, Panzica GC, Contestabile A. Immunohistochemistry and neurochemistry of the habenulo-interpeduncular connection after partial developmental depletion of habenular cholinergic neurons in the rat. *Exp. Brain Res.* 1992; 90:297–301. [PubMed: 1383020]
- Ferraro G, Montalbano ME, Sardo P, La Grutta V. Lateral habenular influence on dorsal raphe neurons. *Brain Res. Bull.* 1996; 41:47–52. [PubMed: 8883915]
- Fields H. State-dependent opioid control of pain. *Nat. Rev. Neurosci.* 2004; 5:565–575. [PubMed: 15208698]
- Fishbain DA, Lewis JE, Gao J. Are psychoactive substance (opioid)-dependent chronic pain patients hyperalgesic? *Pain Pract.* 2011; 11:337–343. [PubMed: 21199316]
- Florin S, Meunier J, Costentin J. Autoradiographic localization of [3H]nociceptin binding sites in the rat brain. *Brain Res.* 2000; 880:11–16. [PubMed: 11032985]
- Fowler CD, Lu Q, Johnson PM, Marks MJ, Kenny PJ. Habenular alpha5 nicotinic receptor subunit signalling controls nicotine intake. *Nature.* 2011; 471:597–601. [PubMed: 21278726]
- Frahm S, Slimak MA, Ferrarese L, Santos-Torres J, Antolin-Fontes B, Auer S, Filkin S, Pons S, Fontaine JF, Tsetlin V, Maskos U, Ibanez-Tallon I. Aversion to nicotine is regulated by the balanced activity of beta4 and alpha5 nicotinic receptor subunits in the medial habenula. *Neuron.* 2011; 70:522–535. [PubMed: 21555077]
- Friedman A, Lax E, Dikshtein Y, Abraham L, Flaumenhaft Y, Sudai E, Ben-Tzion M, Ami-Ad L, Yaka R, Yadid G. Electrical stimulation of the lateral habenula produces enduring inhibitory effect on cocaine seeking behavior. *Neuropharmacology.* 2010; 59:452–459. [PubMed: 20600170]
- Friedman A, Lax E, Dikshtein Y, Abraham L, Flaumenhaft Y, Sudai E, Ben-Tzion M, Yadid G. Electrical stimulation of the lateral habenula produces an inhibitory effect on sucrose self-administration. *Neuropharmacology.* 2011; 60:381–387. [PubMed: 20955718]
- Gao DM, Hoffman D, Benabid AL. Simultaneous recording of spontaneous activities and nociceptive responses from neurons in the pars compacta of substantia nigra and in the lateral habenula. *Eur. J. Neurosci.* 1996; 8:1474–1478. [PubMed: 8758954]
- Glick SD, Ramirez RL, Livi JM, Maisonneuve IM. 18-Methoxycoronaridine acts in the medial habenula and/or interpeduncular nucleus to decrease morphine self-administration in rats. *Eur. J. Pharmacol.* 2006; 537:94–98. [PubMed: 16626688]
- Goncalves L, Sego C, Metzger M. Differential projections from the lateral habenula to the rostromedial tegmental nucleus and ventral tegmental area in the rat. *J. Comp. Neurol.* 2011 doi: 10.1002/cne.22787 [Epub ahead of print].
- Goto M, Canteras NS, Burns G, Swanson LW. Projections from the subfornical region of the lateral hypothalamic area. *J. Comp. Neurol.* 2005; 493:412–438. [PubMed: 16261534]
- Gottesfeld Z. Central and peripheral contributions of deafferentation-induced norepinephrine increase in the habenula. *Brain Res.* 1983; 268:359–361. [PubMed: 6871689]

- Hashimoto K, Amano T, Sakai N, Suzuki T, Narita M. Cell-dependent physiological synaptic action of morphine in the rat habenular nucleus: morphine both inhibits and facilitates excitatory synaptic transmission. *Neurosci. Lett.* 2009; 451:270–273. [PubMed: 19159664]
- Hentall ID, Budhrani VM. The interpeduncular nucleus excites the on-cells and inhibits the off-cells of the nucleus raphe magnus. *Brain Res.* 1990; 522:322–324. [PubMed: 2224530]
- Herkenham M, Nauta WJ. Afferent connections of the habenular nuclei in the rat. A horseradish peroxidase study, with a note on the fiber-of-passage problem. *J. Comp. Neurol.* 1977; 173:123–146. [PubMed: 845280]
- Herkenham M, Nauta WJ. Efferent connections of the habenular nuclei in the rat. *J. Comp. Neurol.* 1979; 187:19–47. [PubMed: 226566]
- Hikosaka O. GABAergic output of the basal ganglia. *Prog. Brain Res.* 2007; 160:209–226. [PubMed: 17499116]
- Hikosaka O. The habenula: from stress evasion to value-based decisionmaking. *Nat. Rev. Neurosci.* 2010; 11:503–513. [PubMed: 20559337]
- Hong S, Hikosaka O. The globus pallidus sends reward-related signals to the lateral habenula. *Neuron.* 2008; 60:720–729. [PubMed: 19038227]
- Hua Z, Shucheng H, Jiazhi F, Shao W. Responses of habenular neurons to both baroreceptive and noxious stimulations. *Chin. J. Neuroanat.* 1998; 14:179.
- Ide JS, Li CS. Error-related functional connectivity of the habenula in humans. *Front. Hum. Neurosci.* 2011; 5:25. [PubMed: 21441989]
- Ikemoto S. Brain reward circuitry beyond the mesolimbic dopamine system: a neurobiological theory. *Neurosci. Biobehav. Rev.* 2010; 35:129–150. [PubMed: 20149820]
- Jesuthasan S. Fear, anxiety and control in the zebrafish. *Dev. Neurobiol.* Jan 10.2011 Epub ahead of print.
- Ji H, Shepard PD. Lateral habenula stimulation inhibits rat midbrain dopamine neurons through a GABA(A) receptor-mediated mechanism. *J. Neurosci.* 2007; 27:6923–6930. [PubMed: 17596440]
- Kalen P, Strecker RE, Rosengren E, Bjorklund A. Regulation of striatal serotonin release by the lateral habenula-dorsal raphe pathway in the rat as demonstrated by in vivo microdialysis: role of excitatory amino acids and GABA. *Brain Res.* 1989; 492:187–202. [PubMed: 2473826]
- Kazi JA, Mori S, Kuchiiwa S, Nakagawa S. Prolonged expression of c-Fos protein in the lateral habenular nucleus of the Japanese monkey (*Macaca fuscata*) after eye enucleation. *Neurosignals.* 2004; 13:130–133. [PubMed: 15067200]
- Kim U. Topographic commissural and descending projections of the habenula in the rat. *J. Comp. Neurol.* 2009; 513:173–187. [PubMed: 19123238]
- Kim U, Chang SY. Dendritic morphology, local circuitry, and intrinsic electrophysiology of neurons in the rat medial and lateral habenular nuclei of the epithalamus. *J. Comp. Neurol.* 2005; 483:236–250. [PubMed: 15678472]
- King T, Vera-Portocarrero L, Gutierrez T, Vanderah TW, Dussor G, Lai J, Fields HL, Porreca F. Unmasking the tonic-aversive state in neuropathic pain. *Nat. Neurosci.* 2009; 12:1364–1366. [PubMed: 19783992]
- Kinsey AM, Wainwright A, Heavens R, Sirinathsinghji DJ, Oliver KR. Distribution of 5-HT(5A), 5-HT(5B), 5-HT(6) and 5-HT(7) receptor mRNAs in the rat brain. *Brain Res. Mol. Brain Res.* 2001; 88:194–198. [PubMed: 11295248]
- Klemm WR. Habenular and interpeduncularis nuclei: shared components in multiple-function networks. *Med. Sci. Monit.* 2004; 10:RA261–RA273. [PubMed: 15507867]
- Kowski AB, Veh RW, Weiss T. Dopaminergic activation excites rat lateral habenular neurons in vivo. *Neuroscience.* 2009; 161:1154–1165. [PubMed: 19374940]
- LaGraze SC, Borzan J, Peng YB, Fuchs PN. Selective regulation of pain affect following activation of the opioid anterior cingulate cortex system. *Exp. Neurol.* 2006; 197:22–30. [PubMed: 15996657]
- Lapicot O, Melin C, Modolo A, Nicolas C, Messaoudi Y, Monconduit L, Artola A, Luccarini P, Dalle R. Tonic and phasic descending dopaminergic controls of nociceptive transmission in the medullary dorsal horn. *Pain.* 2011; 152:1821–1831. [PubMed: 21514054]

- Lecourtier L, DeFrancesco A, Moghaddam B. Differential tonic influence of lateral habenula on prefrontal cortex and nucleus accumbens dopamine release. *Eur. J. Neurosci.* 2008; 27:1755–1762. [PubMed: 18380670]
- Lecourtier L, Neijt HC, Kelly PH. Habenula lesions cause impaired cognitive performance in rats: implications for schizophrenia. *Eur. J. Neurosci.* 2004; 19:2551–2560. [PubMed: 15128408]
- Lehner M, Taracha E, Skorzewska A, Wislowska A, Zienowicz M, Maciejak P, Szyndler J, Bidzinski A, Plaznik A. Sensitivity to pain and c-Fos expression in brain structures in rats. *Neurosci. Lett.* 2004; 370:74–79. [PubMed: 15489021]
- Li B, Piriz J, Mirrione M, Chung C, Proulx CD, Schulz D, Henn F, Malinow R. Synaptic potentiation onto habenula neurons in the learned helplessness model of depression. *Nature.* 2011; 470:535–539. [PubMed: 21350486]
- Li CS, Yan P, Chao HH, Sinha R, Paliwal P, Constable RT, Zhang S, Lee TW. Error-specific medial cortical and subcortical activity during the stop signal task: a functional magnetic resonance imaging study. *Neuroscience.* 2008; 155:1142–1151. [PubMed: 18674592]
- Luigjes J, van den Brink W, Feenstra M, van den Munckhof P, Schuurman PR, Schippers R, Mazaheri A, De Vries TJ, Denys D. Deep brain stimulation in addiction: a review of potential brain targets. *Mol. Psychiatry.* 2011
- Ma QP, Han JS. Neurochemical studies on the mesolimbic circuitry of antinociception. *Brain Res.* 1991; 566:95–102. [PubMed: 1814560]
- Ma QP, Shi YS, Han JS. Further studies on interactions between periaqueductal gray, nucleus accumbens and habenula in antinociception. *Brain Res.* 1992; 583:292–295. [PubMed: 1504835]
- Mahieux G, Benabid AL. Naloxone-reversible analgesia induced by electrical stimulation of the habenula in the rat. *Brain Res.* 1987; 406:118–129. [PubMed: 3032353]
- Mansour A, Khachaturian H, Lewis ME, Akil H, Watson SJ. Autoradiographic differentiation of mu, delta, and kappa opioid receptors in the rat forebrain and midbrain. *J. Neurosci.* 1987; 7:2445–2464. [PubMed: 3039080]
- Martin TJ, Miller M Jr, Dworkin SI, Smith JE, Porrino LJ. Alteration of local cerebral glucose utilization following intravenous administration of heroin in Fischer 344 rats. *Brain Res.* 1997; 755:313–318. [PubMed: 9175898]
- Matsumoto M. Role of the lateral habenula and dopamine neurons in reward processing. *Brain Nerve.* 2009; 61:389–396. [PubMed: 19378808]
- Matsumoto M, Hikosaka O. Lateral habenula as a source of negative reward signals in dopamine neurons. *Nature.* 2007; 447:1111–1115. [PubMed: 17522629]
- McQueen JK, Martin MJ, Harmar AJ. Local changes in cerebral 2-deoxyglucose uptake during alphaxalone anaesthesia with special reference to the habenulo-interpeduncular system. *Brain Res.* 1984; 300:19–26. [PubMed: 6203604]
- Meng H, Wang Y, Huang M, Lin W, Wang S, Zhang B. Chronic deep brain stimulation of the lateral habenula nucleus in a rat model of depression. *Brain Res.* 2011; 1422:32–38. [PubMed: 21978548]
- Meszaros J, Gajewska S, Tarchalska-Krynska B. Habenulo-interpeduncular lesions: the effects on pain sensitivity, morphine analgesia and open-field behavior in rats. *Pol. J. Pharmacol. Pharm.* 1985; 37:469–477. [PubMed: 3909126]
- Morilak DA, Barrera G, Echevarria DJ, Garcia AS, Hernandez A, Ma S, Petre CO. Role of brain norepinephrine in the behavioral response to stress. *Prog. Neuropsychopharmacol. Biol. Psychiatry.* 2005; 29:1214–1224. [PubMed: 16226365]
- Morris JS, Smith KA, Cowen PJ, Friston KJ, Dolan RJ. Covariation of activity in habenula and dorsal raphe nuclei following tryptophan depletion. *Neuroimage.* 1999; 10:163–172. [PubMed: 10417248]
- Murphy CA, DiCamillo AM, Haun F, Murray M. Lesion of the habenular efferent pathway produces anxiety and locomotor hyperactivity in rats: a comparison of the effects of neonatal and adult lesions. *Behav. Brain Res.* 1996; 81:43–52. [PubMed: 8950000]
- Neal CR Jr, Mansour A, Reinscheid R, Nothacker HP, Civelli O, Watson SJ Jr. Localization of orphanin FQ (nociceptin) peptide and messenger RNA in the central nervous system of the rat. *J. Comp. Neurol.* 1999; 406:503–547. [PubMed: 10205026]

- Neckers LM, Schwartz JP, Wyatt RJ, Speciale SG. Substance P afferents from the habenula innervate the dorsal raphe nucleus. *Exp. Brain Res.* 1979; 37:619–623. [PubMed: 520447]
- Nishikawa T, Fage D, Scatton B. Evidence for, and nature of, the tonic inhibitory influence of habenulointerpeduncular pathways upon cerebral dopaminergic transmission in the rat. *Brain Res.* 1986; 373:324–336. [PubMed: 2424555]
- Norris SK, Boden PR, Woodruff GN. Agonists selective for tachykinin NK1 and NK3 receptors excite subpopulations of neurons in the rat medial habenula nucleus in vitro. *Eur. J. Pharmacol.* 1993; 234:223–228. [PubMed: 7683268]
- Okamoto H, Agetsuma M, Aizawa H. Genetic dissection of the zebrafish habenula, a possible switching board for selection of behavioral strategy to cope with fear and anxiety. *Dev. Neurobiol.* 2011 doi:10.1002/dneu.20913 [Epub ahead of print].
- Omelchenko N, Bell R, Sesack SR. Lateral habenula projections to dopamine and GABA neurons in the rat ventral tegmental area. *Eur. J. Neurosci.* 2009; 30:1239–1250. [PubMed: 19788571]
- Parent A, Gravel S, Boucher R. The origin of forebrain afferents to the habenula in rat, cat and monkey. *Brain Res. Bull.* 1981; 6:23–38. [PubMed: 7470948]
- Paul MJ, Indic P, Schwartz WJ. A role for the habenula in the regulation of locomotor activity cycles. *Eur. J. Neurosci.* 2011; 34:478–488. [PubMed: 21777302]
- Paulson PE, Gorman AL, Yeziarski RP, Casey KL, Morrow TJ. Differences in forebrain activation in two strains of rat at rest and after spinal cord injury. *Exp. Neurol.* 2005; 196:413–421. [PubMed: 16182286]
- Paulson PE, Wiley JW, Morrow TJ. Concurrent activation of the somatosensory forebrain and deactivation of periaqueductal gray associated with diabetes-induced neuropathic pain. *Exp. Neurol.* 2007; 208:305–313. [PubMed: 17936273]
- Plassat JL, Boschert U, Amlaiky N, Hen R. The mouse 5HT5 receptor reveals a remarkable heterogeneity within the 5HT1D receptor family. *EMBO J.* 1992; 11:4779–4786. [PubMed: 1464308]
- Ploghaus A, Becerra L, Borras C, Borsook D. Neural circuitry underlying pain modulation: expectation, hypnosis, placebo. *Trends Cogn. Sci.* 2003; 7:197–200. [PubMed: 12757820]
- Ploghaus A, Tracey I, Clare S, Gati JS, Rawlins JN, Matthews PM. Learning about pain: the neural substrate of the prediction error for aversive events. *Proc. Natl. Acad. Sci. U.S.A.* 2000; 97:9281–9286. [PubMed: 10908676]
- Ploghaus A, Tracey I, Gati JS, Clare S, Menon RS, Matthews PM, Rawlins JN. Dissociating pain from its anticipation in the human brain. *Science.* 1999; 284:1979–1981. [PubMed: 10373114]
- Pobbe RL, Zangrossi H Jr. Involvement of the lateral habenula in the regulation of generalized anxiety- and panic-related defensive responses in rats. *Life Sci.* 2008; 82:1256–1261. [PubMed: 18513751]
- Pobbe RL, Zangrossi H Jr. The lateral habenula regulates defensive behaviors through changes in 5-HT-mediated neurotransmission in the dorsal periaqueductal gray matter. *Neurosci. Lett.* 2010; 479:87–91. [PubMed: 20478359]
- Pompeiano M, Palacios JM, Mengod G. Distribution of the serotonin 5-HT2 receptor family mRNAs: comparison between 5-HT2A and 5-HT2C receptors. *Brain Res. Mol. Brain Res.* 1994; 23:163–178. [PubMed: 8028479]
- Rainville J, Smeets RJ, Bendix T, Tveito TH, Poiraudéau S, Indahl AJ. Fear-avoidance beliefs and pain avoidance in low back pain-translating research into clinical practice. *Spine J.* 2011; 11:895–903. [PubMed: 21907633]
- Ray WJ, Slobounov S. Modulation and experience of external stimuli: toward a science of experience and interoception. *Acta Biol. Hung.* 2002; 53:551–558. [PubMed: 12501938]
- Ren K, Dubner R. Descending modulation in persistent pain: an update. *Pain.* 2002; 100:1–6. [PubMed: 12435453]
- Ren ZY, Shi J, Epstein DH, Wang J, Lu L. Abnormal pain response in pain-sensitive opiate addicts after prolonged abstinence predicts increased drug craving. *Psychopharmacology (Berl.).* 2009; 204:423–429. [PubMed: 19172249]

- Roiser JP, Levy J, Fromm SJ, Nugent AC, Talagala SL, Hasler G, Henn FA, Sahakian BJ, Drevets WC. The effects of tryptophan depletion on neural responses to emotional words in remitted depression. *Biol. Psychiatry*. 2009; 66:441–450. [PubMed: 19539268]
- Salas R, Baldwin P, de Biasi M, Montague PR. BOLD responses to negative reward prediction errors in human habenula. *Front. Hum. Neurosci*. 2010; 4:36. [PubMed: 20485575]
- Sarkis R, Saade N, Atweh S, Jabbur S, Al-Amin H. Chronic dizocilpine or apomorphine and development of neuropathy in two rat models. I: Behavioral effects and role of nucleus accumbens. *Exp. Neurol*. 2011; 228:19–29. [PubMed: 21146525]
- Sartorius A, Kiening KL, Kirsch P, von Gall CC, Haberkorn U, Unterberg AW, Henn FA, Meyer-Lindenberg A. Remission of major depression under deep brain stimulation of the lateral habenula in a therapy-refractory patient. *Biol. Psychiatry*. 2010; 67:e9–e11. [PubMed: 19846068]
- Sartorius N, Baghai TC, Baldwin DS, Barrett B, Brand U, Fleischhacker W, Goodwin G, Grunze H, Knapp M, Leonard BE, Lieberman J, Nakane Y, Pinder RM, Schatzberg AF, Svestka J, Baumann P, Ghalib K, Markowitz JC, Padberg F, Fink M, Furukawa T, Fountoulakis KN, Jensen P, Kanba S, Riecher-Rossler A. Antidepressant medications and other treatments of depressive disorders: a CINP Task Force report based on a review of evidence. *Int. J. Neuropsychopharmacol*. 2007; 10(Suppl. 1):S1–S207. [PubMed: 18096106]
- Savitz JB, Nugent AC, Bogers W, Roiser JP, Bain EE, Neumeister A, Zarate CA Jr. Manji HK, Cannon DM, Marrett S, Henn F, Charney DS, Drevets WC. Habenula volume in bipolar disorder and major depressive disorder: a high-resolution magnetic resonance imaging study. *Biol. Psychiatry*. 2011; 69:336–343. [PubMed: 21094939]
- Schultz W. Multiple dopamine functions at different time courses. *Annu. Rev. Neurosci*. 2007; 30:259–288. [PubMed: 17600522]
- Schultz W. Dopamine signals for reward value and risk: basic and recent data. *Behav. Brain Funct*. 2010; 6:24. [PubMed: 20416052]
- Schulz S, Schreff M, Nuss D, Gramsch C, Holtt V. Nociceptin/orphanin FQ and opioid peptides show overlapping distribution but not co-localization in pain-modulatory brain regions. *Neuroreport*. 1996; 7:3021–3025. [PubMed: 9116232]
- Schweinhardt P, Seminowicz DA, Jaeger E, Duncan GH, Bushnell MC. The anatomy of the mesolimbic reward system: a link between personality and the placebo analgesic response. *J. Neurosci*. 2009; 29:4882–4887. [PubMed: 19369556]
- Shepard PD, Holcomb HH, Gold JM. Schizophrenia in translation: the presence of absence: habenular regulation of dopamine neurons and the encoding of negative outcomes. *Schizophr. Bull*. 2006; 32:417–421. [PubMed: 16717257]
- Shumake J, Colorado RA, Barrett DW, Gonzalez-Lima F. Metabolic mapping of the effects of the antidepressant fluoxetine on the brains of congenitally helpless rats. *Brain Res*. 2010; 1343:218–225. [PubMed: 20470763]
- Skofitsch G, Jacobowitz DM. Autoradiographic distribution of ¹²⁵I calcitonin gene-related peptide binding sites in the rat central nervous system. *Peptides*. 1985; 6:975–986. [PubMed: 3001670]
- Smith CD, Lonstein JS. Contact with infants modulates anxiety-generated c-fos activity in the brains of postpartum rats. *Behav. Brain Res*. 2008; 190:193–200. [PubMed: 18374995]
- Smith WJ, Stewart J, Pfaus JG. Tail pinch induces fos immunoreactivity within several regions of the male rat brain: effects of age. *Physiol. Behav*. 1997; 61:717–723. [PubMed: 9145942]
- Sun M, Shong X, Goa J. Chagnes of behavior and CGRP, c-fos in habenula after chorinic compression of DRG in rats. *Heilongjiang Med. J*. 2002 doi:cnki: ISSN:1004-5775.0.2002-04-002.
- Sutherland RJ. The dorsal diencephalic conduction system: a review of the anatomy and functions of the habenular complex. *Neurosci. Biobehav. Rev*. 1982; 6:1–13. [PubMed: 7041014]
- Taraschenko OD, Shulan JM, Maisonneuve IM, Glick SD. 18-MC acts in the medial habenula and interpeduncular nucleus to attenuate dopamine sensitization to morphine in the nucleus accumbens. *Synapse*. 2007; 61:547–560. [PubMed: 17447255]
- Thornton EW, Bradbury GE. Effort and stress influence the effect of lesion of the habenula complex in one-way active avoidance learning. *Physiol. Behav*. 1989; 45:929–935. [PubMed: 2780877]

- Thornton EW, Evans JA, Harris C. Attenuated response to nomifensine in rats during a swim test following lesion of the habenula complex. *Psychopharmacology (Berl.)*. 1985; 87:81–85. [PubMed: 3933041]
- Ullsperger M, von Cramon DY. Error monitoring using external feedback: specific roles of the habenular complex, the reward system, and the cingulate motor area revealed by functional magnetic resonance imaging. *J. Neurosci.* 2003; 23:4308–4314. [PubMed: 12764119]
- Waeber C, Grailhe R, Yu XJ, Hen R, Moskowitz MA. Putative 5-HT receptors: localization in the mouse CNS and lack of effect in the inhibition of dural protein extravasation. *Ann. N.Y. Acad. Sci.* 1998; 861:85–90. [PubMed: 9928243]
- Wang QP, Nakai Y. The dorsal raphe: an important nucleus in pain modulation. *Brain Res. Bull.* 1994; 34:575–585. [PubMed: 7922601]
- Weissman AD, Marquis KL, Moreton JE, London ED. Effects of self-administered phencyclidine on regional uptake of 2-deoxy-D-[1-¹⁴C]glucose in brain. *Neuropharmacology*. 1989; 28:575–583. [PubMed: 2755563]
- Wise RA, Rompre PP. Brain dopamine and reward. *Annu. Rev. Psychol.* 1989; 40:191–225. [PubMed: 2648975]
- Wood PB. Role of central dopamine in pain and analgesia. *Expert Rev. Neurother.* 2008; 8:781–797. [PubMed: 18457535]
- Wright A, Luedtke KE, Vandenberg C. Duloxetine in the treatment of chronic pain due to fibromyalgia and diabetic neuropathy. *J. Pain Res.* 2011; 4:1–10. [PubMed: 21386950]
- Wu SS, Huang M, Cao XJ, Zhang CX, Wang S. The responses of pain-related neurons in habenula to nociceptive stimuli and morphine. *Zhongguo Ying Yong Sheng Li Xue Za Zhi.* 2005; 21:252–255. [PubMed: 21162187]
- Yu LC, Han JS. Habenula as a relay in the descending pathway from nucleus accumbens to periaqueductal grey subserving antinociception. *Int. J. Neurosci.* 1990; 54:245–251. [PubMed: 2265972]
- Zastawny RL, George SR, Nguyen T, Cheng R, Tsatsos J, Briones-Urbina R, O'Dowd BF. Cloning, characterization, and distribution of a mu-opioid receptor in rat brain. *J. Neurochem.* 1994; 62:2099–2105. [PubMed: 8189219]
- Zhang F, Zhou W, Liu H, Zhu H, Tang S, Lai M, Yang G. Increased c-Fos expression in the medial part of the lateral habenula during cue-evoked heroin seeking in rats. *Neurosci. Lett.* 2005; 386:133–137. [PubMed: 16002220]
- Zhou ZF, Du MY, Wu WY, Jiang Y, Han JS. Effect of intracerebral microinjection of naloxone on acupuncture- and morphine-analgesia in the rabbit. *Sci. Sin.* 1981; 24:1166–1178. [PubMed: 6272389]

Fig. 1.

Habenula neuroanatomy. (A) Rat habenula. A Nissl stain coronal segment of the rat brain, clearly demonstrates the relatively large (when compared with whole brain) rat habenula. This image is from an online atlas (<http://brainmaps.org/>, accessed May 2011). (B) Monkey habenula. A Nissl stain habenula in the monkey (Hikosaka, 2010 with permission). (C) Human habenula. (C1) An anatomical MRI image of the coronal, sagittal and horizontal view of the human habenula (from fsl). Numbers indicate the slice coordinates. (C2) Localization of the habenula in a coronal magnetic resonance imaging section (Savitz et al., 2011 with permission); (C3) Nissl and fiber staining of a coronal axial slice of the human brain. This image is from ZoomableHuman Brain Atlas (<http://zoomablebrain.bio.uci.edu/>, accessed May 2011).

Fig. 2.

Habenula afferent and efferent connections. *Key:* FrCtx = frontal cortex, NAc = nucleus accumbens, Lateral Hypo = lateral hypothalamus, EP = entopeduncular nucleus, CPu = caudate/putamen, Hippo = hippocampus, M = medial habenula, L = lateral habenula, P = pineal, IPN = interpeduncular nucleus, PAG = periaqueductal gray, Raphe = raphe nuclei, VTA = ventral tegmental area, and SNc = substantia nigra pars compacta. Adapted from Bianco and Wilson (2009) with permission.

Fig. 3.

Conceptual model of habenula circuits in chronic pain. (A) Normal condition. The figure shows habenula circuit-loops that confer potential functions of the structure. The formulation of the loops is based on known afferent outputs to structures that may have connections to other brain regions that send efferents to the habenula. Thus, four basic loops are shown: (1) *analgesia/hyperalgesic loop* – reflects the integration of outputs from the lateral habenula to regions of the brain that have known pain inhibitory or pain facilitatory action; (2) *reward-aversion loop* – outputs from the lateral habenula to the VTA through dopaminergic process and connections to the frontal cortex; (3) *stress loop* – medial habenula projections to the IPN send afferents to the hippocampus a region involved in a number of processes including stress response (but also memory); and the (4) *sensory-motor loop* – from the lateral habenula connections from the SNc project to basal ganglia regions; the complexity of function may include sensory, emotional and cognitive features and not simply sensory-motor integration. (B) Acute pain. Under acute pain conditions the activity of the habenula is shown to increase all circuits. This is a homeostatic response to a reversible physiological stimulus. (C) Chronic pain. Under chronic pain conditions loop homeostasis is disrupted. For example, the chronic pain state is an aversive state that is a hedonic deficit state where reward systems become dysfunctional; outputs from the pain modulator regions (normally inhibit pain) may now facilitate pain and thus signals from the periphery are enhanced. Stress loop may show enhanced drive with consequent alterations in the hippocampus that may include altered memory in chronic pain. Shown in the bottom right of each figure is the expected activation profile that is low in the normal/no pain condition, increases in response to evoked pain under acute pain conditions, and displays maximal drive in chronic pain conditions.

Fig. 4. Human habenula stimulation. The transversal and coronal view from fluorodeoxyglucose positron emission tomography (FDG-PET) imaging of the metabolic effects from LHb deep brain stimulation of a treatment resistant depression patient is used with permission (Sartorius et al., 2010). Note the regional correspondence to the anatomical image in Fig. 1.

Fig. 5.

Activation in habenula to pain. (A) Fos activation. Fos increase with pain representative digitized image of the LHb, is used with permission (Smith et al., 1997). *Key:* 3V = third ventricle, PVp = paraventricular thalamus, and fr = fasciculus retroflexus. (B) 2-DG measures of chronic pain activation. Spinal cord injury-induced activation in the rat habenula, is used with permission (Paulson et al., 2005); in the upper right corner contrast with a control rat to the left, and in diabetic neuropathic pain, is used with permission (Paulson et al., 2007); pictured on the bottom right versus a control rat habenula pictured to the immediate left. (C) fMRI activation to pain in humans. Coghill et al. (2003) with permission.