Published in final edited form as:

J Am Chem Soc. 2012 October 10; 134(40): 16845–16855. doi:10.1021/ja307796e.

Dynamics of Catalytic Resolution of 2-Lithio-*N*-Boc-piperidine by Ligand Exchange

Timothy K. Beng^a, William S. Tyree^a, Trent Parker^a, Chicheung Su^b, Paul G. Williard^b, and Robert E. Gawley^a

Robert E. Gawley: bgawley@uark.edu

^aDepartment of Chemistry and Biochemistry, University of Arkansas, Fayetteville, AR 72701

^bDepartment of Chemistry, Brown University, Providence, RI 02912

Abstract

The dynamics of the racemization, catalytic and stoichiometric dynamic resolution of 2-lithio-*N*-Boc-piperidine, **7**, have been investigated. The kinetic order in TMEDA, for both racemization and resolution of the title compound, and the kinetic order in resolving ligands, have been determined. The catalytic dynamic resolution is 0.5-order in chiral ligand **8**, 0.265 order in chiral ligand **10**, and second order in TMEDA. The X-ray crystal structure of ligand **10** shows it to be an octamer. Dynamic NMR studies of the resolution process were obtained. Some of the requirements for a successful catalytic dynamic resolution by ligand exchange have been identified.

Introduction

Organolithium compounds are widespread in organic synthesis.¹ The use of stoichiometric chiral ligands such as (–)-sparteine complexed to *sec*-BuLi, to effect enantioselective deprotonation represents one of the methods to generate scalemic organolithium compounds for use in asymmetric synthesis.² Unfortunately, effecting an asymmetric deprotonation on *N*-Boc-piperidines through the use of the chiral ligand (–)-sparteine is not as successful as with pyrrolidines,³ but partial success (up to 88:12 er in reasonable yields) has been achieved with O'Brien's diamine.⁴

In an important development, several chiral organolithium compounds have been discovered that are amenable to dynamic thermodynamic resolution (DTR – Scheme 1).⁵ In a DTR, a racemic organolithium is generated, and a chiral ligand is added which resolves the organolithium dynamically (an asymmetric transformation of the first kind).

Dynamic resolutions hold significant potential, since they obviate the need for an enantioselective deprotonation or an asymmetric synthesis of an enantioenriched precursor stannane. A disadvantage of a DTR can arise if the electrophile used to react with the organolithium is also reactive toward, and consumes, the chiral ligand, L*. Thus, a variation of a dynamic resolution using a catalytic amount of L* is desirable. Toru reported a catalytic dynamic resolution (CDR, Scheme 2), which is driven by the well-known phenomenon of crystallization-induced asymmetric transformation.⁶ In the Toru example, the resolution is accomplished by dynamic resolution of planar-chiral organolithium enantiomers by the

Correspondence to: Robert E. Gawley, bgawley@uark.edu.

Supporting Information Available. Full experimental details including kinetic plots, data analysis, and the NMR experiments. This information is available free of charge via the Internet at <http://pubs.acs.org>.

bisoxazoline ligand, followed by precipitation of homochiral aggregates (an asymmetric transformation of the second kind). Addition of aldehyde caused dissolution of the precipitate, and after quenching with Me₃SiCl and acidic workup, the product was isolated in good yield, with excellent diastereoselectivity and enantioselectivity.

In 2009, we reported the transmetalation of *N*-trimethylallylpyrrolidine, **1**, in the presence of *N,N*-diisopropylbispidine **3**,⁷ to afford racemic organolithium **2** (Scheme 3). Addition of 15 mol % of ligand **4**, and stirring at ~0 °C effected a catalytic resolution. Cooling to –78 °C and quenching with Me₃SiCl gave pyrrolidine **5** with an er of 93:7 (*R:S*).^{8,9}

In this catalytic resolution, the organolithium **2** is in excess of the chiral resolving ligand **4**. The remainder of the organolithium is presumably complexed to the achiral ligand **3**.¹⁰ In order to achieve a catalytic resolution, it is, at a minimum, necessary to ensure that the racemization of the organolithium is significantly slower than the dynamic resolution. Therefore, we studied the enantiomerization of **2** in the presence of the achiral bispidine ligand **3**,⁹ and the thermodynamic parameters were determined ($\Delta H^\ddagger = 25.1 \pm 1.4$ kcal/mol; $\Delta S^\ddagger = 8.1 \pm 5.0$ cal/mol·K). Similarly, the thermodynamic parameters for the resolution of **2** by stoichiometric quantities of **4** in the presence of **3** were measured ($\Delta H^\ddagger = 25.1 \pm 1.7$ kcal/mol; $\Delta S^\ddagger = 14.2 \pm 6.1$ cal/mol·K).¹¹ From these data, plots of $\Delta G^\ddagger$ vs. temperature¹² revealed that the resolution of **2** by **4** has a lower barrier than enantiomerization at all temperatures between –80 °C and +20 °C.⁹

Note that the catalytic resolution in Scheme 3 should not be called a dynamic *thermodynamic* resolution, because it does not employ stoichiometric amounts of the chiral ligand.^{5g,13} Neither can it be called a dynamic *kinetic* resolution, because this term applies to systems where the organolithium inverts rapidly, and resolution occurs by asymmetric substitution under Curtin-Hammett conditions. Thus, we refer to the process simply as a catalytic dynamic resolution, or CDR.

We wanted to know if a CDR was possible in the absence of the achiral ligand, but since conversion of stannane **1** to organolithium **2** by tin-lithium exchange is not possible in the absence of the achiral ligand, we tested the possibility by extrapolation using varying amounts of bispidine **3**. Figure 1 shows the fraction of the *R* enantiomer of **5** as a function of the number of equivalents of bispidine **3**, after treatment of *rac*-**2** with 15 mol% of **4** for 1 h at 0 °C.⁹ The y-intercept of 0.55 implies that the er in the absence of **3** would be 55:45. The catalytic resolution of **2** by **4** is accelerated by **3**, but it is also apparent that **3** is necessary to achieve a catalytic dynamic resolution. If 15% of **2** is resolved to 93:7 er, and the remaining 85% is racemic, the er of the solution would be 56:44, the same as the value of the y-intercept within experimental error. Therefore, a second requirement is that the chiral and achiral ligands must both be involved in the catalytic resolution. This could involve a total ligand exchange or a reorganization of the coordination of the organolithium to more than one ligand. Aggregation/deaggregation of the organolithium could be involved in the catalytic resolution by ligand exchange. The aggregation state of **2** has not been determined, but the aggregation state of the *N*-methyl analogue is a homochiral dimer and the *N*-ethyl analogue is a mixture of aggregates.¹⁴

Mechanistic details of a catalytic resolution by ligand exchange are of interest. Unfortunately, detailed mechanistic investigation of lithiopyrrolidine **2** is hampered by the fragility of the trimethylallyl group to the conditions of the reaction (tendency of the trimethylallyl group to rearrange, undergo proton transfer, and be lost during workup). Also, the presence of the achiral bispidine ligand, **3**,¹⁵ complicates product purification and recovery of the conjugate acid of catalyst **4**.

We wondered whether a catalytic resolution involving ligand exchange is unique to the *N*-trimethylallylpyrrolidine system in Scheme 3, or whether other systems could be identified which could be resolved catalytically. As noted above, we believe that it is necessary that the free energy barrier to resolution be lower than the free energy barrier to racemization. We therefore sought another system for investigation, and chose 2-lithio-*N*-Boc-piperidine, **7**, shown to undergo stoichiometric resolution by the Coldham group in the presence of TMEDA (Scheme 4),^{5f} and whose inversion dynamics were investigated jointly by our two groups.¹⁶

In lithiopiperidine **7**, we have an organolithium that is synthetically important and readily available. Moreover, the achiral ligand, TMEDA, is inexpensive. As it turned out (see Figure 14, below), the moderate final er of the Coldham resolution (77:23) permitted a crucial experiment that was not possible in investigating the more highly enantioselective catalytic resolution of **2**. An interesting fact uncovered previously was that both the stoichiometric resolution and the racemization of lithiopiperidine **7** are catalytic and first order in the *achiral* ligand TMEDA (from 0 to 2 equivalents of TMEDA for the DTR and 0.1 to 1.0 equivalents for racemization). Also, several other diamine ligands were tested, but failed to fill this catalytic role in the resolution – even TEEDA (tetraethyl ethylenediamine)!¹⁶

Herein we report the results of our detailed investigation into the dynamics of the catalytic resolution of **7** by **8**. During the course of this mechanistic investigation, a new pair of ligands, **10** and **11** were discovered, which effect resolution of **7** and which afford much higher enantioselectivity than **8** (Scheme 5).¹⁷ While many of these mechanistic studies were conducted with ligand **8**, some experiments are reported with ligand **10** which provide additional insight into the processes involved.

Results and Discussion

We have previously reported thermodynamic parameters for racemization and resolution of **7**.¹⁶ The stoichiometric DTR of *rac*-**7** by **8** in the presence of 2 equivalents of TMEDA has a large negative entropy of activation and a small enthalpy of activation ($\Delta H^\ddagger = 4.3 \pm 0.5$ kcal/mol; $\Delta S^\ddagger = -55.9 \pm 1.8$ cal/mol·K). The enantiomerization of **7** in the presence of 1 equivalent of TMEDA has $\Delta H^\ddagger = 18.1 \pm 0.7$ kcal/mol and $\Delta S^\ddagger = 5.0 \pm 3.2$ cal/mol·K. At -46°C , the free energy barrier to enantiomerization (in the presence of 2 equivalents of TMEDA) and the DTR (in the presence of 1 equivalent of TMEDA) are equal. Below this temperature, the free energy barrier for DTR is lower. Given this information, we tested the possibility of a CDR of **7** at -55°C , in the presence of TMEDA. The time evolution for catalytic dynamic resolution of *rac*-**7** was evaluated as illustrated in Figure 2. In oven-dried vials, stannane *rac*-**12** was treated with BuLi and 1.2 equiv of TMEDA at -78°C for 1 h to effect tin-lithium exchange, affording *rac*-**7**·TMEDA. The chiral ligand **8** (15 mol%) was then added and the reaction vessel was transferred to a thermostatted bath at -55°C (internal temperature). At various time intervals over 24 h, a vial was cooled to -78°C , quenched with Me₃SiCl and the er measured by CSP-GC. The final er of the product (77:23) is the same as that obtained by Coldham, et al, using ligand **8** in a stoichiometric DTR;^{5f} this experiment was duplicated in our hands in a control experiment. During the course of this experiment, no precipitate was observed in the reaction flask (compare Scheme 2).

These data are noteworthy for two reasons. First, a CDR is operative, and second, the *S* organolithium is configurationally stable for at least 12 hours under the reaction conditions, implying achievement of an equilibrium state. During the course of this mechanistic investigation, the CDR using ligands **10** and **11** was found to be very highly enantioselective, and synthetic applications have been reported.¹⁷

Further investigation of the structure, dynamics of enantiomerization, DTR and CDR of **7** by **8** and **10** were undertaken at carefully controlled temperatures, and revealed the following:

- Conclusive evidence of the solution structure of **7** was not obtained because the spectra are consistent with several possibilities, but evidence of dynamic exchange phenomena was found.
- There is variable configurational stability of **7** in the presence of different ligands at low temperature, when generated by tin-lithium exchange.
- Although TMEDA catalyzes racemization and DTR of lithiopiperidine **7** up to one equivalent,^{16,17} TMEDA retards racemization when in excess; thus, excess TMEDA increases the configurational stability of **7**.
- There is a second order dependence of the CDR on [TMEDA].
- We find a fractional order dependence for the CDR in the concentration of the chiral ligands [**8**] and [**10**].

These studies are described in detail below.

NMR studies

Before describing the dynamics of lithiopiperidine **7**, we begin with a study of its solution structure. It is known from *in situ* IR studies that the lithium ion of **7** is coordinated to the carbonyl oxygen, and the monomer structure in Figure 3 was proposed to be the structure in solution.^{4a} However, there are dimeric and polymeric structures that are consistent with the IR data. We used ⁶Li and ¹³C NMR to investigate the possible structures illustrated in Figure 3.

Transmetalation of stannanes *rac*-**12** and *S*-**12** with *n*-Bu⁶Li in ether/TMEDA provided {⁶Li}-**7**. The ⁶Li and ¹³C spectra were recorded at temperatures ranging from –80 to –30 °C. The ⁶Li spectra taken at several temperatures are shown in Figure 4. The racemate shows a single peak at 2.80 ppm assigned to the Li⁺ on **7** as well as small amounts of solvated lithium ion¹⁸ between 0.5 and 1.0 ppm at all temperatures between –30 and –80 °C. A single lithium signal is consistent with structures of the monomer, dimer-1, dimer-2, and the polymer shown in Figure 3. In contrast, the enantioenriched compound shows an approximately 1:1 ratio of **7** and solvated Li⁺ at –80 °C, which changes as the temperature is warmed. The two lithium signals observed at –80 °C are consistent with the structure of a triple ion and dimer-3. The growth of a shoulder on the upfield signal as the temperature is raised indicates a second type of solvated lithium ion, and the changes in line shape suggest a dynamic exchange process among solution species.

Partial ¹³C spectra of *rac*-**7** and *S*-**7** (93:7 er) at –81 °C (Figure 5, left and middle) show a peak at 56.2 ppm that is split into a 1:1:1 triplet (*J* = 11.9 Hz) due to ¹J coupling between ⁶Li and C-2 of a contact ion pair; at –30 °C, this coupling is no longer evident, consistent with a solvent-separated ion pair. Being indicative of a single lithium in contact with the carbanionic carbon, this triplet rules out dimer-1 and dimer-3. The resonance of the carbonyl carbon appears at 160.1 ppm in both *rac*-**7** and *S*-**7**. Thus, no change in the chemical shift of this peak could be detected between racemic or enriched **7** in TMEDA/Et₂O, even though we took care to reference the spectra to the chemical shift of *d*₄ methanol in a concentric capillary. These similarities suggest that the organolithium species being observed is the same, whether racemic or enantioenriched. Note however, that these observations do not preclude the presence of other organolithium species in the solution.

To summarize: for *rac*-**7**, the ^6Li and ^{13}C are consistent with the monomer, dimer-2, or polymer structures, but not the triple ion, dimer-1 or dimer-3. For *S*-**7**, the ^6Li spectrum is consistent with the triple ion or dimer-3 at $-80\text{ }^\circ\text{C}$, with other structures evident at temperatures $-60\text{ }^\circ\text{C}$. The coupling seen in the partial ^{13}C spectra are consistent with the monomer, triple ion, dimer-2, or polymer structures.

Attempts were made to monitor changes in solution structure during a catalytic resolution. To that end, a mixture of racemic stannane **12**, 4 equivalents of TMEDA, and 10 mol% of the conjugate acid of ligand **10** were dissolved in an NMR tube and cooled to $-78\text{ }^\circ\text{C}$. A solution of *n*-Bu ^6Li was added and the sample was transferred to an NMR probe cooled to $-80\text{ }^\circ\text{C}$. The temperature was maintained at $-80\text{ }^\circ\text{C}$ for 2 h to complete the tin-lithium exchange, then raised to $-45\text{ }^\circ\text{C}$ for 3 h to effect dynamic resolution, then cooled back to $-80\text{ }^\circ\text{C}$. We have previously determined that enantioenriched **7** is configurationally stable at $-80\text{ }^\circ\text{C}$.¹⁶ Four spectra were obtained, and are shown in Figure 6 (~1 h accumulation time in each case). Most noticeably, the 1:1:1 triplet indicating a contact ion pair comprising one carbon and one lithium atom is evident in all four spectra. There are also changes seen in the region between 50 and 53 ppm, where a series of unassigned peaks are barely evident in the top spectrum, disappear in the spectra at $-45\text{ }^\circ\text{C}$, and become much more prominent at the end of the resolution. Quenching the solution in the NMR tube with phenyl isocyanate and workup afforded an 89:11 ratio of pipecolic acid anilide enantiomers (see SI for details).^{17a}

X-ray crystal structure of ligand **10**

The dilithio ligand **10** formed beautiful crystals when an ether solution of the ligand was cooled in the absence of TMEDA. The ligand crystallized as an octamer that is formed by two cyclic tetramers stacking on top of each other. Each tetramer features a 16-membered ring consisting of four $(-\text{NMe}-\text{Li}-\text{O}-\text{Li}-)_4$ units. Figure 7a shows a ball and stick structure, including two ethers of crystallization. Figure 7b is a ChemDraw rendering of one of the 16-membered tetramer rings, with a monomer highlighted in blue. Note that for each monomer, one of the lithium atoms is chelated by two nitrogens and the oxygen, while the second lithium bridges the methylated nitrogen to the oxygen of the next monomer. Figure 7c shows the octamer, with the second tetramer rendered in red. Note the two types of bridging between the two 16-membered rings: the methylated nitrogens coordinate to the bridging lithium of second tetramer, while the chelated lithium is coordinated to the oxygen of the second tetramer. This results in eight 4-coordinate lithium atoms and eight 3-coordinate lithium atoms in the crystal. It is noteworthy that the eight N-methyl substituents of these chiral ligands completely fill the interior cavity of this octomer.

Configurational stability of **7**

In these studies, we chose to generate organolithium **7** by tin-lithium exchange from stannane **12** because we wanted to avoid the possibility of incomplete deprotonation of **6**, which could complicate the analysis of the product mixtures, and because we wished to generate **7** efficiently, while also having evidence of its formation.

Scheme 6 summarizes experiments used to evaluate configurational stability of **7** (formed by tin-lithium exchange) in the absence and presence of (–)-sparteine, bispidine **3**, and TMEDA. Previously, we found that tin-lithium exchange of *S*-**12** in the absence of any ligand at temperatures $-60\text{ }^\circ\text{C}$ resulted in racemic product.¹⁶ Below $-60\text{ }^\circ\text{C}$, the lithiation is too slow to be synthetically useful, but it was attempted at $-78\text{ }^\circ\text{C}$ to investigate configurational stability in the absence of ligands. At this temperature, the half-life for tin-lithium exchange is $>5\text{ h}$ (see SI for kinetic details). After 5 h of treatment of **12** with *n*-BuLi at $-78\text{ }^\circ\text{C}$, quenching with Me_3SiCl produced a mixture of **12** and **9**, but the latter was racemic. In the presence of either (–)-sparteine (**13**) or bispidine **3**, TLC monitoring showed

that transmetalation of *S*-**12** (73:27 er) to **7** was complete after 5 h; however, quenching with Me₃SiCl afforded racemic **9**. This result is in contrast to an 87:13 er of **7** that was obtained by deprotonation using *sec*-BuLi in the presence of (–)-sparteine,^{3a} perhaps reflecting different mechanisms for the formation of **7** involving different types of lithium ligation in different solvents or in the presence of different amines. In THF in the absence of diamines for example, tin-lithium exchange occurs with complete retention of configuration.^{4a} As noted previously,¹⁶ TMEDA plays a unique role in the racemization and DTR of lithiopiperidine **7**: both processes are first order in [TMEDA] between 0 and 1 molar equivalent of TMEDA. After treatment of stannane **12** of 73:27 er with *n*-BuLi for only 30 min at –78 °C in the presence of 1 equivalent of TMEDA, quenching with Me₃SiCl revealed complete transmetalation, and **9** was obtained in 73:27 er. Consistent with the uniqueness noted previously, of the ligands tested, only TMEDA is suitable for the CDR of **7** following tin-lithium exchange.¹⁶

DTR kinetics at –55 °C

Previously, the barriers to inversion in the DTR of *rac*-**7** by stoichiometric quantities of **8** were determined at four temperatures, ranging from –10 to +20 °C.¹⁶ Because we planned to investigate catalytic resolution at considerably lower temperatures, we decided to measure the kinetics of stoichiometric DTR at –55 °C. The time evolution of the stoichiometric DTR of *rac*-**7** by **8** (plus 1.2 equiv TMEDA) was measured, as illustrated in Figure 8. The evolution of the major enantiomer (in this case *S*) is given by the equation:

$$[S]_t = [S]_{eq} + ([S]_0 - [S]_{eq})e^{-(k_{obs})t},$$

where $[S]_t$ is the fraction *S* at time *t*, $[S]_0$ and $[S]_{eq}$ are the fractions of the *S* enantiomer at *t*=0 and *t*=∞ at this temperature, respectively. In this case, $[S]_0 = 0.5$ and $[S]_{eq} = 0.77$. The data reveal a half-life of ~3 hours at this temperature.

The effect of [8] on dynamic resolution

To evaluate the effect of varying amounts of chiral ligand **8** on dynamic resolutions, experiments were conducted using 15, 20, 50, 80 and 100 mol% catalyst. After 5 h at –55 °C, enantiomer ratios for **9** of 63:37, 66:34, 71:29, 75:25, and 78:22 were observed (Figure 9, diamonds). The data in Figure 9 clearly show that catalysis is occurring, since the observed er is always higher than that predicted if the system is at the equilibrium (Figure 9, circles), realized at higher temperatures (see Figure 15, below). This is all the more surprising, since the values should be *lower* than the predicted value if equilibrium was not yet reached. The supporting information has a similar plot for ligand **10**. Thus, quantitative evaluation of the rate of resolution with differing amounts of **8** and **10** was undertaken.

The dependence of the rate of resolution of lithiopiperidine **7** on chiral ligands **8** and **10** was investigated at –20 °C in the presence of TMEDA (1.0 equiv) in Et₂O (see SI for details). Figure 10 shows observed first order rate constants plotted against [8] and [10] from zero to one equivalent. A nonlinear fit of the data to the equation $k_{obs} = k[L^*]^n$ afforded *n* = 0.5, and $k = 1.59 \pm 0.06 \times 10^{-3} \text{ M}^{0.5} \text{ s}^{-1}$ for **8**. A similar fit of the observed rate constants plotted against [10] revealed *n* = 0.265 ± 0.022 and $k = 9.49 \pm 0.12 \times 10^{-4} \text{ M}^{-0.265} \text{ s}^{-1}$ (see SI for details). Thus, the reaction is half-order in [8] and ~1/4-order in [10]. A fractional order is indicative of a mechanism involving deaggregation of the chiral ligand.

The effect of [TMEDA] on CDR of **7**

As noted, TMEDA plays a unique role in the inversion dynamics of lithiopiperidine **7** (both enantiomerization and DTR).¹⁶ It was therefore of interest to explore the kinetic role of TMEDA in the CDR of **7**. The time evolution of *S*-**7** under CDR conditions in the presence of varying amounts of TMEDA (zero to four equivalents) was followed and the rate constants were measured (Figure 7). Upward curvature is emblematic of higher order dependence of rate on [TMEDA]. A nonlinear parametric fit of the equation $k_{obs} = k_d[\text{TMEDA}]^n$ reveals a second order dependence on [TMEDA] ($n = 1.99$; $k_c = 4.02 \pm 0.11 \times 10^{-3} \text{ M}^{-2} \text{ s}^{-1}$). This indicates that the transition structure of the rate limiting step in the CDR contains two TMEDA molecules.

Racemization of **7** with excess TMEDA

Since the dynamics of racemization of *S*-**7** were previously determined in the presence of 1 equiv TMEDA, and since we observed a second order dependence of the CDR on [TMEDA], we reexamined the racemization kinetics, this time in the presence of an excess of TMEDA. The enantiomerization was followed by generating the organolithium **7** (er 85:15 or 96:4 *S*:*R*) using tin–lithium exchange in Et₂O at –78 °C with BuLi and two or four equivalents of TMEDA, followed by warming to the desired temperature for different time periods, then cooling to –78 °C and quenching with excess Me₃SiCl as previously described (see SI for details).¹⁶ The rate constants were determined by nonlinear fits to the zero-order plots. Eyring analysis of the rate constants at their respective temperatures in the presence of varying amounts of TMEDA provided the thermodynamic parameters listed in Table 1. Plots of Gibbs free energy as a function of temperature¹¹ with these varying amounts of TMEDA are shown in Figure 12a, and reveal that racemization has a significantly higher energy barrier with four equivalents of TMEDA than with one or two equivalents, at all temperatures between –80 and 0 °C. Figure 12b shows that, in the presence of 2 equivalents of TMEDA, racemization and resolution of **7** by stoichiometric amounts of **8** have equal free energy barriers at –33 °C.¹¹ Below that temperature, resolution has a lower barrier.

In a preliminary communication, we showed that the rate of enantiomerization of **7** is first order in TMEDA up to one equivalent.¹⁶ However, the above experiments reveal that the barrier to enantiomerization is raised significantly in the presence of excess TMEDA. Because our CDR studies are carried out using excess TMEDA, we decided to study the rate of enantiomerization with excess TMEDA. The details are in the SI, but the results are plotted in Figure 13. There is an inverse dependence of racemization rate on [TMEDA] when TMEDA is in excess over **7**. Thus, for the purposes of developing the CDR, 2–4 equiv of TMEDA beneficially retards racemization. In the preparative experiments described later, and also those using ligands **10** and **11**,¹⁷ we employ four equivalents of TMEDA, which serves to accelerate resolution and retard racemization.

Approach to the –55 °C equilibrium from either direction

The zero order plots of the CDR, starting from *rac*-**7**, approach but never exceed the er value that is observed in the stoichiometric DTR, 77:23. In our study of the CDR of trimethylallyl pyrrolidine **2**, we showed that *R*-**2** (93:7 er) maintained configurational stability under the catalytic conditions for 90 minutes.⁹ For lithiopiperidine **7**, we have stannylpiperidine *S*-**12** of 96:4 er,^{17a} and it was of interest to see whether this er is maintained under the catalytic conditions, or whether the er would drop to the 77:23 er that is observed when starting with racemate. We subjected both enantiomers to catalytic conditions (15 mol % **8**, 2 equiv TMEDA, –55 °C), and recorded the evolution of er, as shown in Figure 14. In both experiments, the final er matched the er in a stoichiometric DTR. The rate constant, k_{SR} , for equilibration of *S*-**7** (96:4 er) with 2 equiv TMEDA and 15 mol% **8** is $1.93 \pm 0.11 \times 10^{-5} \text{ s}^{-1}$.

Similarly, the rate constant, k_{RS} , for CDR of *R*-**7** (72:28 er) is $6.24 \pm 0.45 \times 10^{-5} \text{ s}^{-1}$. The ratio k_{RS}/k_{SR} is 3.23, which is similar to the ratio of *S*-**7** to *R*-**7** in a DTR (3.35 for er 77:23).

Effect of temperature on the position of the equilibrium

The above discussion emphasizes the importance of staying below the crossover temperature, such that the free energy barrier for resolution is below the barrier for racemization. Two control experiments demonstrate the importance of temperature control. Racemic lithiopiperidine **7** was subjected to catalytic conditions: 4 equivalents of TMEDA and 15 mol % **8** or **10**, at -55°C . The er was measured at several time intervals until the er reached 76:24 for **8** and 95:5 for **10**. The temperature was then raised to 0°C and the er monitored again. The dramatic results are illustrated in Figure 10. At 0°C , the er drops dramatically to about 55:45. Clearly, careful control of temperature is necessary to achieve a CDR.

Optimized conditions for CDR

Given the above findings, we decided on the optimum conditions of 5–15 mol% **10** and 4 equivalents of TMEDA, and have used them in several synthetic applications. Both direct electrophilic substitutions and transmetalations with ZnCl_2 followed by Cu(I) or Pd(0) cross-couplings were demonstrated with numerous examples with a high degree of enantioselectivity, usually 95:5 er.¹⁷ Examples using trimethylsilyl chloride and phenyl isocyanate as electrophiles are given in Scheme 7 for comparison of ligands **8** and **10**.

Summary and conclusions

There are several important details that emerge from these studies:

1. *NMR data are inconclusive as to the solution structure of **7**, although differences in the ^6Li spectra of rac-**7** and *S*-**7** are evident.*
2. *The ^6Li spectra of *S*-**7** exhibit dynamic phenomena as the temperature is raised to -60°C and above.*
3. *Splitting of the ^{13}C NMR signal for the carbanionic carbon persists throughout the CDR. Other signals appear in the same region, suggesting the presence of other species.*
4. *Consistent with faster organolithium dynamics at higher temperatures, rapid racemization is observed when the reaction mixture containing resolved **7** is raised to 0°C .*
5. *Chiral ligands **8** and **10** increase the rate of CDR. The kinetic order for CDR of lithiopiperidine **7** is 0.5-order in [**8**] and 0.265-order in [**10**]. Fractional orders implicate deaggregation of the chiral catalysts. Ligand **10** is an octamer in the solid state.*
6. *Excess TMEDA accelerates the CDR. Although the stoichiometric DTR is first order in [TMEDA],¹⁶ the CDR (0.15 equiv **8**) is 2nd order in [TMEDA].*
7. *Excess TMEDA retards racemization. The rate constant for enantiomerization of lithiopiperidine **7** is at a maximum when the ratio **7**:TMEDA = 1. Racemization is first order in [TMEDA] up to 1 equivalent of TMEDA (however racemization is very fast in the absence of TMEDA), and inverse order when excess TMEDA is present, up to 4 equivalents.*

8. The final er of the stoichiometric DTR and the catalytic dynamic resolution are the same.
9. Temperature control is critical to the success of the catalytic dynamic resolution.
10. The rate law for the CDR using L^* as catalyst, and which converts *R*-**7** to *S*-**7**, is:

$$\frac{d[S - 7]}{dt} = k[R - 7][\text{TMEDA}]^2[L^*]^n, \text{ where } n = 0.5 \text{ for } \mathbf{8} \text{ and } 0.265 \text{ for } \mathbf{10}.$$

We have observed CDR for five combinations of organolithium, catalyst and achiral ligand (Figure 16): 2-lithio-*N*-trimethylallylpyrrolidine, **2**, using either diamino lithium alkoxide **4** or (–)-sparteine as catalyst with *N,N*-diisopropylbispidine **3** as the achiral ligand,⁹ and 2-lithio-*N*-Boc-piperidine, **7**, with either diamino lithium alkoxide **8**, or dilithio ligands **10** or **11** as catalysts and TMEDA as the achiral ligand.¹⁷ What is clear from these studies is that ligand exchange is an important process in dynamic resolutions of lithiopiperidine **7** and trimethylallyl pyrrolidine **2**, under both stoichiometric and catalytic conditions. The effect of a ligand on the rate of carbanion inversion of a chiral organolithium species can change with the molar ratio of ligand to the organolithium. It is also clear that an achiral ligand and careful control of temperature are necessary to achieve a catalytic resolution. As a guide to choosing a temperature, we have used the “crossover temperature”, where the free energy barrier for racemization and dynamic resolution are equal: below this temperature, catalytic resolution can occur, but above this temperature, CDR cannot occur. Further refinement of temperature is based on the free energy barrier for the CDR, such that a reasonable half-life can be achieved.

If the organolithium species and the chiral ligand are monomeric, catalytic resolution is not possible, because the process would violate the principle of microscopic reversibility. However as Toru has shown,⁶ irreversible population of a homochiral aggregate from a racemic monomer or a heterochiral aggregate by catalytic dynamic resolution is possible. In the case of the piperidines under study in this work, there is some evidence of differences in solution structure between racemic and enantioenriched **7**. In this system, we find no visible evidence of aggregate precipitation, but we do see evidence of variable solution structure that is dependent on temperature. Furthermore, we also observe a change in final enantiomer ratio that is dependent on temperature. In addition, the fractional order in L^* of the CDR provides clear evidence that de-aggregation of the resolving ligand is operative in the dynamic resolution of **7**. We have also demonstrated that the catalytic resolution requires exchange with an achiral ligand, in this case TMEDA.¹⁹ These observations may be due to a change in solution structure between racemic and enantioenriched **7**, such as the formation of a homochiral aggregate at low temperature. At a higher temperature such as 0 °C, new pathways for racemization become operative, such that the organolithium that is presumably complexed to TMEDA racemizes rapidly, while the remainder, complexed to L^* , remains enantioenriched. Additionally, the fractional kinetic order in ligands **8** and **10** implicate changes in aggregation state of the resolving ligand in the rate determining step. Our working hypothesis of this phenomenon is that a successful CDR depends on ligand exchange with an achiral ligand *and* on aggregation-deaggregation phenomena of the species in solution, including the chiral organolithium and/or the resolving ligand.

Experimental Section

For general experimental conditions, see the Supporting Information.

Effect of varying [8] on DR of *N*-Boc-2-lithiopiperidine 5 in the presence of TMEDA

General Procedure—In oven-dried tubes equipped with a stir bar and capped with rubber septa, *rac*-**12** (0.06 M in ether) was treated with BuLi (1.2 equiv) and TMEDA (1.2 equiv) at $-78\text{ }^{\circ}\text{C}$ for 1 hour under argon to effect tin-lithium exchange, affording *rac*-**7**·TMEDA. The concentration of **8** was varied (0.0 to 100 mol %) and the tubes were quickly transferred to the $-55\text{ }^{\circ}\text{C}$ bath. After 5 h at this temperature, the tubes were transferred to the $-78\text{ }^{\circ}\text{C}$ bath and rapidly quenched with excess Me_3SiCl . The silanes were subsequently analyzed by CSP-SFC monitoring at 210 nm. See Supporting Information for details.

Activation Parameters for Racemization of **7** with 2.0 equiv TMEDA

Typical kinetic run—In oven-dried septum capped tubes equipped with a stir bar, *S*-**12** (85:15 er, 0.06 M in ether) and TMEDA (2.0 equiv) were treated with BuLi (1.2 equiv.) at $-78\text{ }^{\circ}\text{C}$ for 30 min under nitrogen to effect tin-lithium exchange. The tubes were quickly transferred to a second thermostatted bath and warmed to the desired temperature. At various time intervals a tube was cooled to $-78\text{ }^{\circ}\text{C}$ and rapidly quenched with excess Me_3SiCl . After 4 h, MeOH (2 mL) was added and the mixture was extracted into Et_2O . The silanes were subsequently analyzed by CSP-SFC monitoring at 210 nm. The rate constants were determined by nonlinear fits to the zero-order plots. See Supporting Information for details.

Details of X-ray crystal structure determination of ligand **10**

A clear colorless chunk-like specimen of $\text{C}_{96}\text{H}_{192}\text{Li}_{16}\text{N}_{16}\text{O}_8\cdot(\text{C}_4\text{H}_{10}\text{O})_2$ approximate dimensions $0.280\text{ mm} \times 0.300\text{ mm} \times 0.340\text{ mm}$, was used for the X-ray crystallographic analysis. The X-ray intensity data were measured on a Bruker Apex 1K diffractometer. Integration of the data using a triclinic unit cell yielded a total of 21236 reflections to a maximum θ angle of 26.48° ($0.80\text{ \AA}$ resolution), of which 18549 were independent (average redundancy 1.145, completeness = 83.5%, $R_{\text{int}} = 3.81\%$, $R_{\text{sig}} = 7.20\%$) and 14814 (79.86%) were greater than $2\sigma(F^2)$. The final cell constants of $a = 11.668(4)\text{ \AA}$, $b = 17.093(6)\text{ \AA}$, $c = 18.174(6)\text{ \AA}$, $\alpha = 64.950(5)^{\circ}$, $\beta = 76.278(5)^{\circ}$, $\gamma = 88.581(6)^{\circ}$, volume = $3178.2(19)\text{ \AA}^3$, are based upon the refinement of the XYZ-centroids of reflections above $20\sigma(I)$. Data were corrected for absorption effects using the multi-scan method (SADABS). The ratio of minimum to maximum apparent transmission was 0.534 – 0.711.

The structure was solved and refined using the Bruker SHELXTL Software Package, using the space group P 1, with $Z = 1$ for the formula unit, $\text{C}_{104}\text{H}_{212}\text{Li}_{16}\text{N}_{16}\text{O}_{10}$. The final anisotropic full-matrix least-squares refinement on F^2 with 1343 variables converged at $R1 = 7.53\%$, for the observed data and $wR2 = 22.77\%$ for all data. The goodness-of-fit was 1.036. The largest peak in the final difference electron density synthesis was $0.869\text{ e}^{-}/\text{\AA}^3$ and the largest hole was $-0.564\text{ e}^{-}/\text{\AA}^3$ with an RMS deviation of $0.076\text{ e}^{-}/\text{\AA}^3$. On the basis of the final model, the calculated density was 1.023 g/cm^3 and $F(000)$, 1076 e^{-} .

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank the National Science Foundation (CHE 0616352 and 1011788) and the Arkansas Biosciences Institute for direct support of this work. TP thanks the NSF-REU for a summer fellowship (CHE 0851505). Core facilities were funded by the Arkansas Biosciences Institute and the National Institutes of Health (P30 RR031154 and P30 GM103450). We also thank Professors Donna Blackmond, Iain Coldham, David Collum, and Jonathan Clayden for helpful discussions.

References and Notes

- (a) Clayden, J. *Organolithiums: Selectivity for Synthesis*. Pergamon; 2002. (b) Hodgson, DM., editor. *Organolithiums in Enantioselective Synthesis*. Vol. Vol. 5. Berlin Heidelberg: Springer-Verlag; 2003. (c) Rappoport, Z.; Marek, I., editors. *The Chemistry of Organolithium Compounds*. New York: Wiley; 2004. (d) Majewski, M.; Snieckus, V., editors. *Organometallics: Compounds of Group 1 (Li ... Cs)*. Vol. Vol. 8a. Stuttgart: Thieme; 2006. (e) Gawley, RE., editor. *Stereochemical Aspects of Organolithium Compounds*. Vol. Vol. 26. Weinheim: Verlag Helvetica Chimica Acta; 2010.
- (a) Hoppe D, Hense T. *Angew. Chem. Int. Ed.* 1997; 36:2283–2316. (b) Hoppe, D.; Christoph, G. *The chemistry of organolithium compounds*. Rappoport, Z.; Marek, I., editors. Oxford: John Wiley and Sons, Ltd.; 2004. p. 1055-1164. (c) Beak P, Johnson TA, Kim DD, Lim SH. *Top. Organomet. Chem.* 2003; 5:139–176. (d) Kizirian J-C. *Top. Stereochem.* 2010; 26:189–251. (e) Mitchell EA, Peschiulli A, Lefevre N, Meerpoel L, Maes BUW. *Chem.-Eur. J.* 2012; 18:10092–10142. [PubMed: 22829434]
- (a) Bailey WF, Beak P, Kerrick ST, Ma S, Wiberg KB. *J. Am. Chem. Soc.* 2002; 124:1889–1896. [PubMed: 11866600] (b) Coldham I, O'Brien P, Patel JJ, Raimbault S, Sanderson AJ, Stead D, Whittaker DTE. *Tetrahedron: Asymmetry.* 2007; 18:2113–2119. (c) Metallinos C, Dudding T, Zaifman J, Chaytor JL, Taylor NJ. *J. Org. Chem.* 2007; 72:957–963. [PubMed: 17253816]
- (a) Stead D, Carbone G, O'Brien P, Campos KR, Coldham I, Sanderson A. *J. Am. Chem. Soc.* 2010; 132:7260–7261. [PubMed: 20462193] (b) Dixon AJ, McGrath MJ, O'Brien P. *Org. Synth.* 2009; Coll. Vol. 11:25–37.
- (a) Beak P, Anderson DR, Curtis MD, Laumer JM, Pippel DJ, Weisenburger GA. *Acc. Chem. Res.* 2000; 33:715–727. [PubMed: 11041836] (b) Coldham I, Dufour S, Haxell TFN, Howard S, Vennall GP. *Angew. Chemie Int. Ed.* 2002; 41:3887–3889. (c) Coldham I, Dufour S, Haxell TFN, Vennall GP. *Tetrahedron.* 2005; 61:3205–3220. (d) Whisler MC, MacNeil S, Snieckus V, Beak P. *Angew. Chemie Int. Ed.* 2004; 43:2206–2225. (e) Coldham I, Dufour S, Haxell TFN, Patel JJ, Sanchez-Jimenez G. *J. Am. Chem. Soc.* 2006; 128:10943–10951. [PubMed: 16910691] (f) Coldham I, Raimbault S, Chovatia PT, Patel JJ, Leonori D, Sheikh NS, Whittaker DTE. *Chem. Commun.* 2008:4174–4176. (g) Coldham I, Sheikh NS. *Top. Stereochem.* 2010; 26:253–293. (h) Coldham I, Raimbault S, Whittaker DTE, Chovatia PT, Leonori D, Patel JJ, Sheikh NS. *Chem. Eur. J.* 2010; 16:4082–4090. [PubMed: 20175161]
- (a) Nakamura S, Hirata N, Kita T, Yamada R, Nakane D, Shibata N, Toru T. *Angew. Chem. Int. Ed.* 2007; 46:7648–7650. (b) Nakamura S, Hirata N, Yamada R, Kita T, Shibata N, Toru T. *Chem. Eur. J.* 2008; 14:5519–5527. [PubMed: 18461587]
- Bispidine **3** was first used as an exchangeable achiral ligand in catalytic deprotonations: McGrath MJ, O'Brien P. *J. Am. Chem. Soc.* 2005; 127:16378–16379. [PubMed: 16305208]
- Assuming retentive substitution, this implies accumulation of the *R* organolithium.
- Beng TK, Yousaf TI, Coldham I, Gawley RE. *J. Am. Chem. Soc.* 2009; 131:6908–6909. [PubMed: 19405472]
- Tin-lithium exchange was used to generate **2** from **1**, a process which fails in the absence of a diamine ligand.
- In all cases, the best fits of the kinetic data were to reversible, 1st order plots.
- Plots of $\Delta G^\ddagger$ vs. Temperature over a large temperature range assume that $\Delta H^\ddagger$ and $\Delta S^\ddagger$ are constant over the entire range. Since our Eyring plots were obtained over a small temperature range of 20–30 degrees, the values of $\Delta H^\ddagger$ and $\Delta S^\ddagger$ are valid in the temperature range where the measurements were made, and we assume that they offer a first approximation over the larger range.

13. Lee WK, Park YS, Beak P. *Acc. Chem. Res.* 2009; 42:224–234. [PubMed: 19152329]
14. (a) Low E, Gawley RE. *J. Am. Chem. Soc.* 2000; 122:9562–9563. (b) Gawley RE, Klein R, Ashweek NJ, Coldham I. *Tetrahedron.* 2005; 61:3271–3280.
15. Barker G, O'Brien P, Campos KR. *Arkivoc.* 2011:217–229.
16. Coldham I, Leonori D, Beng TK, Gawley RE. *Chem. Commun.* 2009:5239–5240. corrigendum with correct enthalpy and entropy values: **2010**, 9267–9268.
17. (a) Beng TK, Gawley RE. *J. Am. Chem. Soc.* 2010; 132:12216–12217. [PubMed: 20806976] (b) Beng TK, Gawley RE. *Org. Lett.* 2011; 13:394–397. [PubMed: 21174392] (c) Beng TK, Gawley RE. *Heterocycles.* 2012; 84:697–718. [PubMed: 22408286]
18. Reich HJ, Sikorski WH, Gudmundsson BÖ, Dykstra RR. *J. Am. Chem. Soc.* 1998; 120:4035–4036.
19. In the catalytic resolution of **2** by **4**, the achiral ligand required is bispidine **3**. See ref. 9.

Figure 1.
Resolution of **2** by **4**, with varying amounts of bispidine **3** (0 °C, 1 h).⁹

Figure 2.
Catalytic dynamic resolution of *rac*-7 by **8**.

Figure 3. Possible solution structures for **7** (neglecting the absolute and relative configuration of the metal-bearing carbons).

Figure 4. ^6Li NMR spectra of **7** obtained by transmetalation of **12** in Et_2O in the presence of 4 equivalents of TMEDA; the peaks at 0 ppm are external 1.0 M $^6\text{LiCl}$ in CD_3OD in a concentric capillary. *Left:* $\{^6\text{Li}\}$ - rac-7 . *Right:* enantioenriched $\{^6\text{Li}\}$ - S-7 .

Figure 5. Partial ^{13}C spectra of racemic (left), enantioenriched (middle, 93:7 er) $\{\text{}^6\text{Li}\}$ -**7** in TMEDA/ Et_2O at -80°C , showing the metal-bearing carbon at 56.2 ppm, referenced to external CD_3OD at 49.0 ppm.

Figure 6.
Partial ^{13}C spectra of a CDR conducted in an NMR tube.

Figure 7.

X-ray crystal structure of dilithio ligand **10**. (a) Ball and stick rendering of the octameric crystal structure comprised of stacked 16-membered rings and two molecules of diethyl ether; (b) one 16-membered ring tetramer with a monomer colored blue; (c) stacked tetramers, with the rear tetramer colored red.

Figure 8.

Evolution of the fraction of *S* enantiomer in the stoichiometric DTR of **7** by **8** (1.0 equiv) in the presence of TMEDA (1.2 equiv.) at $-55\text{ }^{\circ}\text{C}$. A nonlinear least-squares fit of the data, using $[S]_{eq} = 0.77$ and $[S]_0 = 0.5$, yielded a rate constant, $k_{obs} = 7.48 \pm 0.63 \times 10^{-5} \text{ s}^{-1}$.

Figure 9.

The effect of varying **[8]** on the dynamic resolution of *rac-7* with TMEDA (1.2 equiv.) at -55°C in ether for 5 h, after quenching with Me_3SiCl . The circles and the straight line indicate a calculated fraction if the system contains *rac-7*·TMEDA and **7·8** having 77:23 er. The diamonds indicate the experimental values observed after 5 h.

Figure 10.

Observed rate constants vs $[\mathbf{8}]$ (top) and $[\mathbf{10}]$ (bottom) for the dynamic resolution of *rac-7* (0.06 M) in the presence of TMEDA (1.0 equiv) at $-20\text{ }^{\circ}\text{C}$ in Et_2O (the points at the origin are not experimental).

Figure 11.

Effect of varying [TMEDA] on CDR of **7** (0.06 M) in the presence of **8** (0.15 equiv) in ether at -55°C for 4 h. The curve describes the equation $k_{obs} = k_c[\text{TMEDA}]^n$, where $k_c = 4.02 \pm 0.11 \times 10^{-3} \text{ M}^{-2}\text{s}^{-1}$ and $n = 1.99$.

Figure 12.

(a) Plots of $\Delta G^\ddagger$ vs. temperature for the enantiomerization free energy barrier of **7**, with varying amounts of TMEDA. (b) Plots of $\Delta G^\ddagger$ vs. temperature for racemization and stoichiometric resolution of **7** in the presence of 2 equivalents of TMEDA.¹¹

Figure 13.
Effect of 1–4 equiv TMEDA on rates of racemization of **7** (0.25 M) at $-40\text{ }^{\circ}\text{C}$.

Figure 14.
Evolution of CDR starting from enantioenriched *R* or *S* **12**.

Figure 15.

Proof of a low-temperature process that effects a catalytic resolution at -55°C , but which fails at 0°C . The CDR of **7** in the presence of TMEDA (4.0 equiv) at -55°C , followed by warming to 0°C for 30 min results in rapid racemization. (a) ligand **8**; (b) ligand **10**.

Figure 16.
Organolithium and ligand combinations amenable to catalytic dynamic resolution.

Scheme 1.A dynamic thermodynamic resolution, or DTR, using stoichiometric chiral ligand, L^* .

Scheme 2.
Toru's catalytic dynamic resolution.

Scheme 3.
CDR of 2-lithio-*N*-trimethylallylpyrrolidine.⁹

Scheme 4.
Dynamic thermodynamic resolution of **7**.^{5f}

Scheme 5.
Dynamic thermodynamic resolution of **7** with ligands **10** and **11**.^{17a}

S-12
73:27 er

13, (–)-sparteine

ligand	time	er 9
none	5h	50:50*
13	5h	50:50
3	5h	50:50
TMEDA	30 min	73:27

* plus recovered **S-12**

Scheme 6.

Transmetalation and configurational stability studies.

Scheme 7.
Optimized CDR conditions.

Table 1Thermodynamic parameters for racemization of **7**.

Entry	RLi-L	Description	$\Delta H^\ddagger$ (kcal/mol)	$\Delta S^\ddagger$ (cal/mol·K)
1 ^a	7 -TMEDA	Racemization of 7 with 1.0 equiv TMEDA	18.1 ± 0.7	5.0 ± 3.2
2	7 -TMEDA	Racemization of 7 with 2.0 equiv TMEDA	20.1 ± 1.8	9.9 ± 7.6
3	7 -TMEDA	Racemization of 7 with 4.0 equiv TMEDA	22.5 ± 2.0	16.7 ± 8.8

^aData taken from reference 14.