Published in final edited form as:

Immunol Rev. 2012 November ; 250(1): 303–316. doi:10.1111/j.1600-065X.2012.01160.x.

Structural Insights into the Common γ -Chain Family of Cytokines and Receptors from the Interleukin-7 Pathway

Scott T. R. Walsh

Institute for Bioscience and Biotechnology Research, W. M. Keck Laboratory for Structural Biology, Department of Cell Biology and Molecular Genetics, Rockville, MD, USA

SUMMARY

Over the past 13 years, numerous crystal structures of complexes of the common γ -chain (γ_c) cytokine receptors and their cytokines have been solved. Even with the remarkable progress in the structural biology of γ_c receptors and their cytokines or interleukins, there are valuable lessons to be learned from the structural and biophysical studies of interleukin-7 (IL-7) and its α -receptor (IL-7R α) and comparisons to other γ_c family members. The structure of the IL-7/IL-7R α complex teaches that interfaces between the γ_c interleukins and their receptors can vary in size, polarity, and specificity, and that significant conformational changes might be necessary for complexes of interleukins and their receptors to bind the shared, activating γ_c receptor. Binding, kinetic, and thermodynamic studies of IL-7 and IL-7R α show that glycosylation and electrostatics can be important to interactions between ILs and their receptor, even where the glycans and charged residues are distant from the interface. The structure of the IL-7R α homodimer is a reminder that often-ignored non-activating complexes likely perform roles just as important to signaling as activating complexes. And last but not least, the structural and biophysical studies help explain and potentially treat the diseases caused by aberrant IL-7 signaling.

Keywords

interleukin-7; interleukin-7 receptor; common γ -chain family; structures; cancer

Introduction

Cells communicate with one another through extracellular signaling proteins known as cytokines. Each cytokine binds to the extracellular domain(s) of either one or two matching cell-surface receptors denoted as α - and β -chain receptors (Fig. 1). Certain cytokines and their matching receptors also bind to another membrane-bound receptor, the common γ -chain (γ_c) (CD132). The γ_c family of cytokines or interleukins includes interleukin-2 (IL-2), IL-4, IL-7, IL-9, IL-15, and IL-21 (reviewed in 1, 2). The interaction of the cytokine, its matching receptor(s), and γ_c enables Janus kinase (JAK) and signal transducer and activator of transcription (STAT) proteins to interact with the intracellular domains of the cytokine receptors and trigger the classical JAK/STAT signaling pathway, which ultimately activates transcription of target genes (2). Careful regulation of the signaling cascade initiated by the interactions among γ_c interleukins and their receptors is fundamental to development, proliferation, and homeostasis of B, T, and natural killer (NK) cells of the immune system (2).

Correspondence to: Scott T.R. Walsh, Institute for Bioscience and Biotechnology Research, W. M. Keck Laboratory for Structural Biology, Department of Cell Biology and Molecular Genetics, 9600 Gudelsky Drive, Rockville, MD, 20850 USA, Tel.: +1 240 314 6478, Fax: +1 240 314 6225, swalsh12@umd.edu.

The author has no conflicts of interest to declare.

IL-7 is in some ways a model example of a γ_c interleukin that induces signaling through receptor heterodimerization. IL-7 binds to its α -receptor, IL-7R α (CD127), and γ_c through their extracellular domains (ECDs) to form a ternary complex that activates the JAK/STAT, phosphoinositol 3-kinase (PI3K)/Akt, or SRC pathways (reviewed in 1). IL-7 and IL-7R α exhibit structural features similar to the other γ_c interleukins and their receptors, and IL-7 interacts with IL-7R α using the same secondary structures used by other γ_c interleukins and their receptors. Both IL-7 and IL-7R α , like the other γ_c interleukins and their receptors, are glycoproteins comprised of numerous asparagines that can be attached to N-linked glycans, or serines/threonines attached to O-linked glycans, or the first tryptophan of the WSXWS sequence motif attached to a C-mannose (3). Also similar to other γ_c IL-specific receptors (4–7), IL-7R α self-associates to form homodimers incapable of signaling (8). Lastly, mutations in IL-7R α , as seen for other mutated γ_c IL-specific receptors, can result in disease, such as autoimmune conditions (e.g. multiple sclerosis, type 1 diabetes, and colitis) (reviewed in 9), severe combined immunodeficiency (SCID) (reviewed in 10), and cancers (e.g. breast, leukemias, and lymphomas) (11–13).

IL-7 is in other ways unique and potentially creates a new paradigm for cytokine-induced receptor heterodimerization signaling. The interface between IL-7 and IL-7R α is relatively small, more apolar, less charged, and less specific than the interfaces between other γ_c interleukins and their receptors, which may be important for IL-7R α 's ability to bind partners besides IL-7. The α -helices of IL-7 and the angular geometries of IL-7R α differ so much from the other γ_c interleukins and their receptors that either IL-7 and IL-7R α undergo conformational changes to bind γ_c , or γ_c binds in a different conformation (14, 15). Glycosylation, although generally thought to be unimportant for γ_c interleukin/receptor interactions, significantly influences the binding affinity of IL-7R α for IL-7 (14). Receptor-receptor association, also generally underappreciated among γ_c IL-specific receptors, likely regulates IL-7 signaling by sequestering the IL-7 binding surface and requiring dissociation and re-orientation of IL-7R α and γ_c to bind IL-7. Lastly, the IL-7R α homodimer and IL-7/IL-7R α structures suggest rationales for the mutations causing B-cell acute lymphoblastic leukemia (B-ALL) and T-cell acute lymphoblastic leukemia (T-ALL) and SCID and potential diagnostic and therapeutic strategies to treat these diseases (15).

A comparison of the IL-7/IL-7R α structure to other γ_c family members

Complex structures of the γ_c interleukins and their receptors have been solved: IL-2 bound to its α -receptor, IL-2R α , its β -receptor, IL-2R β , and γ_c (16, 17); IL-4 bound to its α -receptor (IL-4R α) (18); IL-4 bound to IL-4R α and γ_c (19); IL-7 bound its α -receptor, IL-7R α (14); IL-15 bound to its α -receptor, IL-15R α (20, 21); and IL-21 bound to its α -receptor, IL-21R α (3) (Fig. 2). IL-2R α and IL-15R α adopt sushi domain structures and are not homologous to any component of the IL-7 signaling pathway and therefore are not covered in this review. These complex structures comprise the receptors' ECDs. Furthermore, there are several unbound structures of IL-2 (22, 23), IL-4 (24–27), and IL-21 (28).

The structure of the IL-7/IL-7R α complex resembles the structures of IL-2/IL-2R β as seen in the quaternary complex, IL-4/IL-4R α as seen in the binary and ternary complexes, and IL-21/IL-21R α (Fig. 2). Similar to other γ_c interleukins and their receptors, IL-7 adopts an up-up-down-down four-helix bundle topology with two cross-over loops, and IL-7R α forms an L-shaped architecture with each arm of the 'L' comprised of a fibronectin type III (FNIII) domain, denoted as D1 and D2, connected by a 3_{10} -helical linker. The D1 domain of IL-7R α contains conserved cysteine residues across the receptors forming disulfide bonds and the D2 domain of IL-7R α contains the highly conserved WSXWS sequence motif as well. Also similar to other γ_c interleukin/receptor complexes, IL-7 sits at the elbow region

of the D1 and D2 domains of IL-7R α and interacts with these domains at *site 1* through residues on helices A and C. Yet, the global similarities of these complexes give way to meaningful differences at the interface between IL-7 and IL-7R α and the potential interfaces between IL-7/IL-7R α and γ_c .

The IL-7/IL-7R α interface

The IL-7/IL-7R α interface is more apolar and less specific than the interfaces of the other γ_c interleukin/receptor interfaces (Table 1). The IL-7/IL-7R α interface comprises on average 47% apolar residues, 34% polar residues, and 5 hydrogen bonds. In contrast, the IL-2/IL-2R β interface comprises 24% apolar residues, 39% polar residues, and 8 hydrogen bonds. The IL-4/IL-4R α interface comprises on average 27% apolar residues, 43% polar residues, and 15 hydrogen bonds, and the IL-21/IL-21R α interface comprises 32% apolar residues, 30% polar residues, and 17 hydrogen bonds. The IL-7/IL-7R α interface predominantly involves non-specific hydrophobic and van der Waals contacts with a shape complementarity (S_c) value on average of 0.67 (S_c value gives how well packed an interface is with values ranging from 0, poorly packed, to 1, perfectly packed)(29). Unlike the IL-7/IL-7R α interface, the IL-2/IL-2R β interface involves a mixture of hydrophobic and hydrophilic residues with a S_c of 0.74. The IL-4/IL-4R α interface is dominated by side chain-specific hydrophilic contacts with a S_c of on average 0.72, and the IL-21/IL-21R α interface includes both hydrophobic and hydrophilic residues with a S_c of 0.70.

Of all the observed characteristics of the γ_c interleukin/receptor interfaces, the size of the interface correlates best with the binding affinities of these complexes (Table 2). The obvious trend is that the larger the buried surface area (BSA), the stronger the equilibrium binding dissociation constant (K_d): IL-21/IL-21R α (BSA = 1000 Å², K_d = 70 pM) > IL-4/IL-4R α (BSA = 807 Å², K_d = 150 pM) > IL-7/IL-7R α (BSA = 717 Å², K_d = 59 nM) > IL-2/IL-2R β (BSA = 670 Å², K_d = 530 nM). There do not appear to be any trends correlating the strength of the binding affinities with the other measured parameters, such as the composition of residues at the interfaces.

The IL-7/IL-7R α interface consists of the fewest charged residues of the γ_c interleukin/receptor complexes, but electrostatics still significantly influence association (30). The IL-7/IL-7R α interface comprises only 20% charged residues, whereas the IL-4/IL-4R α , IL-2/IL-2R β , and IL-21/IL-21R α interfaces comprise 30%, 37%, and 38% charged residues, respectively. Yet, the binding affinity of IL-7 and IL-7R α decreased significantly as a function of increasing sodium chloride concentrations, as measured by surface plasmon resonance spectroscopy (30). Fitting these data using an ion linkage mechanism (31), the number of ions consumed or released upon binding (2.0–2.8 ions) was much greater than that measured for the electrostatically driven IL-4/IL-4R α interaction (0.8 ions) (32) and was within the range of ions typically associated with highly charged nucleic acid-protein interactions (31, 33). Given the lack of complementary charged residues in the IL-7/IL-7R α interface, global long-range electrostatic charges are likely responsible for these impressive electrostatic parameters. By primary sequence analysis, IL-7 is a basic protein with an isoelectric point (pI) of 8.5, and the IL-7R α ECD is an acidic protein with a pI of 5.4. The negative charge and field potential of IL-7R α is distributed throughout the molecule, but mainly concentrated on the backside of the receptor away from the binding face of IL-7 (30). The positive charge and field potential of IL-7 is localized to a few residues at the amino-terminus and to a patch on the top part of the molecule (30).

The atypical interface between IL-7 and IL-7R α demonstrates that the interactions between γ_c interleukins and their receptors can vary widely in size, polarity, charge, and specificity. Diverse characteristics at the γ_c IL/receptor interface is desirable, because they allow these molecules to bind to multiple partners, not unlike γ_c . For instance, both IL-4R α and IL-7R α

participate in type I and type II interactions. In type I interactions, the α -receptor is the interleukin-specific receptor; in type II interactions, it is the activating receptor, playing the role that γ_c performs in the type I interactions. Thus, IL-4R α binds not only IL-4 in a type I interaction but also IL-13/IL-13R α in a type II interaction (19), and IL-7R α binds not only IL-7 in a type I interaction, but also thymic stromal lymphopoietin (TSLP) and thymic stromal lymphopoietin receptor (TSLPR) in a type II interaction (34). The adaptive nature of the interactions between γ_c interleukins and receptors that are involved in type I and type II complexes is exemplified by the increase in size, polarity, and specificity of the IL-4/IL-4R α interface as compared to the IL-13/IL-4R α interface in the IL-13 ternary complex (19). It remains to be seen how the IL-7/IL-7R α interface will compare to the TSLP/IL-7R α interface. Nevertheless, for the γ_c interleukins and interleukin-specific receptors to bind different partners, they must be flexible and able to interact through interfaces with distinguishing attributes.

The potential IL-7/IL-7R α interface with γ_c

The structures of the IL-2 quaternary complex and IL-4 ternary complex demonstrate that γ_c interacts with both the interleukins and interleukin-specific receptors (Fig. 2). The contacts between IL-2 and IL-4 with γ_c at *site 2a* are rather small, polar, and specific (Table 1). The IL-2/ γ_c interface buries 497 Å² of surface area, comprises 58% polar and 29% apolar residues, and exhibits a S_c of 0.84. The IL-4/ γ_c interface buries 536 Å² of surface area, comprises 59% polar and 18% apolar residues, and exhibits a S_c of 0.82. The *site 2a* interfaces involve residues from helices A and D in IL-2 and IL-4 and various elbow loop residues of γ_c . The surfaces of IL-2 and IL-4 that bind to γ_c , are characterized as apolar ‘canyons’ surrounded by specific peripheral polar interactions, and the surface of γ_c that binds to the interleukins is described as ‘rigid’ and ‘flat’ (19).

The contacts between IL-2R β and IL-4R α with γ_c at *site 2b* are more extensive than those between the interleukins and γ_c at *site 2a*. The BSAs of the IL-2R β / γ_c and IL-4R α / γ_c interfaces are 874 and 675 Å², respectively. Both interfaces are more polar than apolar: the IL-2R β / γ_c interface comprises 41% polar and 33% apolar residues, and the IL-4R α / γ_c interface comprises 57% polar and 24% apolar residues. Yet, the IL-2R β / γ_c interface exhibits much higher shape complementarity (S_c of 0.64) than the IL-4R α / γ_c interface (S_c of 0.49), but the IL-4R α / γ_c interface displays more hydrogen bonds than the IL-2R β / γ_c interface (21 *versus* 14), which suggests that receptor/receptor contacts can enhance the specificity of complex formation. The two receptors interact through their D2 domains and are related by almost a twofold symmetry (19).

The structure of the IL-7/IL-7R α complex shows that IL-7 is positioned to bind to γ_c via helices A and D, but not in the same orientation as IL-2 and IL-4. Backbone C α superimpositions of the four-helix bundles of IL-2, IL-4, and IL-21 onto IL-7 result in root mean squared deviations (rmsds) of 2.5 Å, on average 3.2 Å, and 2.4 Å, respectively (Fig. 3). Helix D of IL-7 is approximately 7°–12° displaced from helix D of IL-2 and IL-21 and 14°–26° displaced from helix D of IL-4. The overall poor superimposition of the four-helix bundle and angular displacements of helix D in particular suggest that IL-7 presents a different orientation of helices to the flat, rigid surface of γ_c (Fig. 4A).

In addition to using a different orientation to bind γ_c , IL-7 likely uses residues at the *site 2a* interface that were not predicted to interact with γ_c . The residues on IL-7 that were predicted to constitute a γ_c recognition motif include M17, I19, and L23 on helix A and W142, among others, on helix D (19, 35, 36). However, a turn of a π -helix in helix A that is unique to IL-7 changes the alignment of residues, such that S13, M17, and D23 of IL-7 align with the residues in IL-2 and IL-4 that comprise the predicted γ_c recognition motif (Fig. 4B). W142 of helix D also was reported to be critical to the interaction between IL-7 and γ_c ,

but W142 is buried into the hydrophobic core of the four-helix bundle and not interacting with γ_c . Accordingly, mutation of W142 likely causes a folding defect in helix D and/or in IL-7 that, in turn, causes the reported decreases in signaling (35, 36).

The differences in the structures of IL-7R α , IL-2R β , IL-4R α , and IL-21R α indicate that conformational changes must occur before the IL-7/IL-7R α complex interacts with γ_c . Backbone superimpositions of the D1, D2, or both domains of IL-7R α onto IL-2R β result in rmsds from as low as 1.8 Å to as high as 2.7 Å (Fig. 4A,C,D). None of the receptors possesses angular geometries between the D1 and D2 domains similar to IL-7R α (Table 3). Although IL-7R α , IL-2R β , and IL-21R α all share the same elbow (ϵ) angle between the D1 and D2 domains ($\Delta\epsilon = 75^\circ$), the twist (τ) and swivel (σ) angles show that the D1 and D2 domains of IL-7R α are rotated away from IL-2R β ($\Delta\tau = 13^\circ$ and $\Delta\sigma = 24^\circ$) and IL-21R α ($\Delta\tau = 17^\circ$ and $\Delta\sigma = 23^\circ$). The D1 and D2 domains of IL-7R α and IL-4R α have very similar twist angles ($\Delta\tau \sim 1^\circ$) but differ in their elbow angles ($\Delta\epsilon = 12^\circ$) and dramatically in their swivel angles ($\Delta\sigma = 33^\circ$). The poor superimposition and considerably different angular geometries of the IL-7R α FNIII domains prevent the binding of γ_c onto the IL-7/IL-7R α structure. Superimposing the D1, D2, or both D1/D2 domains of the IL-7/IL-7R α structure onto the corresponding domains of the IL-2 quaternary or IL-4 ternary structures yields steric clashes between the D2 domains of IL-7R α and γ_c (Fig. 4C), steric clashes between IL-7 and γ_c , (Fig. 4D), or IL-7 being too distant to contact γ_c (Fig. 4A, C).

The structure of the IL-21/IL-21R α complex is also not compatible with the γ_c conformation from the IL-2 and IL-4 structures but for different reasons. The angular geometries of IL-21R α are nearly identical to those of IL-2R β ($\Delta\epsilon = 0^\circ$, $\Delta\tau = 4^\circ$, and $\Delta\sigma = 1^\circ$) (Table 3), and a superimposition of the IL-21/IL-21R α structure onto the D2 domain of IL-2R β in the quaternary structure results in an rmsd of 0.6 Å (Fig. 4C). IL-21, however, is nowhere near γ_c in the superimposition, most probably due to the extensive interface between IL-21 and IL-21R α (BSA = 1000 Å²) drawing helix D away from γ_c . Therefore, either the IL-7/IL-7R α and IL-21/IL-21R α complexes undergo structural rearrangements or γ_c uses a different conformation than the one observed in the IL-2 and IL-4 structures to form the IL-7 and IL-21 ternary complexes.

The influence of glycosylation on the interactions of γ_c family members

The γ_c interleukins and their receptors are all glycoproteins. Despite its universal presence in the γ_c family, glycosylation appears to play varying roles in the interactions between the interleukins and their receptors. At one end of the spectrum is IL-4 and its α -receptor. Glycosylation of either the IL-4 or IL-4R α does not influence the binding affinity or their ability to signal (18, 32, 37, 38). Also at this end of the spectrum is IL-2. Glycosylation of the IL-2 does not influence the binding affinity to its receptors or its ability to signal (39, 40). There are no studies reporting the effects of glycosylation of the IL-2 receptors.

Towards the other end of the spectrum is IL-21 and its receptor. The structure of the IL-21/IL-21R α reveals that the glycan chain of N-glycosylated N54 in the D1 domain of IL-21R α forms a bridge to the mannose glycan attached to the sidechain of W195 (C-mannosylation), the first tryptophan in the highly conserved WSXWS motif in the D2 domain of IL-21R α (3). Although there are no studies reporting the impact of glycosylation on binding or signaling, the glycan chain attached to N54 of IL-21R α is required for proper production and secretion from HEK293 cells, and the structure indicates that the N- and C-linked glycan bridge may stabilize the two fibronectin domains (3).

Also at the other end of the spectrum is IL-7 and its α -receptor. Glycosylation of IL-7R α clearly does influence its binding to IL-7. The binding affinity of IL-7 to unglycosylated IL-7R α is 18 μ M, whereas the binding affinity of IL-7 to glycosylated IL-7R α is on average

59 nM (Table 2). This approximately 300-fold enhancement in binding affinity results from an at least 5200-fold increase in the association rate. An increase in the on-rate, especially of this magnitude, is highly unusual not just for glycosylated cytokine-receptor interactions but for glycosylated protein-protein interactions as a whole (reviewed in 33, 41). Moreover, the faster association rate is wholly accounted for by the proximal N-acetyl glucosamine groups attached to the asparagine residues of IL-7R α , and thus is indifferent to the type of glycans and the extent of branching from expression from insect or mouse cell lines.

The structures of IL-7/IL-7R α do not provide a clear structural basis for the impact of glycosylation on the binding kinetics and affinity (Fig. 5A). The global structural differences between the complexes of glycosylated and unglycosylated IL-7R α with IL-7 are fairly small and subtle. These structures superimpose with rmsds of on average 0.59 Å. Locally, none of the six potential N-linked glycosylation sites (N29, N45, N131, N162, N212, and N213) on IL-7R α are in the interface with IL-7. The glycans extend away from the structure and generally do not make contacts with residues besides their attached asparagines.

One possible way in which glycosylation of IL-7R α could contribute to its interaction with IL-7 is through the overall electrostatic potential of IL-7R α . A long-range electrostatic attraction drives the interaction between IL-7 (pI 8.4) and IL-7R α (pI 5.3) through complementary overall charges of the molecules (30). Even though there are very few charged residues at the IL-7/IL-7R α interface, the IL-7R α residues interspersed throughout the protein but away from the interface are mainly acidic, and the IL-7 residues above and away from the interface are mainly basic and predicted to interact with negatively charged glycosaminoglycans, such as heparin (42). Glycosylation, in theory, could modulate those charges through the negatively charged sialic acid groups at the ends of complex N-glycans produced by vertebrates. This mechanism of action seems unlikely, however, because the degree of branching and the types of glycans typically determine to what extent glycosylation affects electrostatic potential and the IL-7/IL-7R α interaction is insensitive to variations in branching and glycans (14).

A more likely way in which glycosylation of IL-7R α may contribute to its interaction with IL-7 is by shifting the equilibrium of conformations sampled by unbound IL-7R α . Unbound IL-7R α is likely undergoing conformational exchange between at least two states: IL-7R α unable and able to bind IL-7. The existence of these two states is consistent with the fact that the binding kinetics of IL-7 and IL-7R α best fit to a two-state exchange mechanism and that the binding kinetics of IL-4 and IL-4R α , which are unaffected by glycosylation, fit to a single-step reaction (also the other IL/receptor binding kinetics) (32, 40, 43, 44). Unpublished results from my laboratory demonstrate an increased thermodynamic stability of glycosylated IL-7R α relative to unglycosylated IL-7R α , which also supports glycosylation's effect on the IL-7R α states. We know through thermodynamic analysis the role of glycosylation of IL-7R α is affecting the unbound states of IL-7R α versus the bound state with IL-7 (unpublished results). By shifting the equilibrium toward the IL-7R α state that is able to bind IL-7, glycosylation very well could increase the binding k_1 on-rate and affinity measured for the IL-7/IL-7R α interaction. Further unpublished results by my laboratory reveal extensive interplay among the N-glycans acting synergistically (positive binding cooperativity) to enhance its binding affinity to IL-7. The more difficult question of 'how' are the N-glycans on IL-7R α enhancing its binding affinity to IL-7 are being tackled now.

The crystal structure of the unbound state of IL-7R α ECD identified the first O-linked glycosylation site of this receptor (15). The unbound IL-7R α structure comprised two receptor molecules in the asymmetric unit. Chain B of the IL-7R α displayed clear difference electron density around the sidechain of S133 that fit well with a N-acetyl galactosamine

(GalNAc) glycan. Mass spectrometry confirmed that S133 was glycosylated with a GalNAc. Mass spectrometry further identified that T132 also is O-glycosylated with a GalNAc. Chain A of IL-7R α showed no signs of these O-glycans on these residues. The N-acetyl galactosamine of S133 forms numerous hydrogen bonds with receptor residues and water-mediated interactions. Backbone superimposition of the two receptor chains shows that the O-glycan changes the conformation from residues 133–143 to adopt two turns of an α -helix relative to chain A. The importance of O-linked glycosylation of the IL-7R α is an area of active investigation.

The role of unliganded receptor/receptor complexes in γ_c signaling

Research on the γ_c family has focused primarily on the formation of the complexes involved in the stepwise cytokine-induced heterodimerization mechanism (Fig. 1). According to this mechanism, the complexes of interest for IL-7 signaling are IL-7/IL-7R α and IL-7/IL-7R α / γ_c . Yet, lurking in the research background have been observations of γ_c complexes not contemplated by this textbook mechanism. For instance, there have been reports of homodimers of IL-7R α , IL-2R β , IL-4R α , and IL-9R α , and heterodimers of these receptors with γ_c (4–8, 45). None of these complexes has been reported to induce signaling, but their inability to signal should not lead to a conclusion that they are irrelevant or undeserving of attention. To the contrary, these unliganded receptor/receptor complexes likely perform important roles in γ_c family signaling, such as protecting the γ_c interleukin machinery from degradation, localizing the right γ_c family members to specific cellular niches, and modulating the timing and degree of signaling. They also may explain and serve as new targets for treating diseases implicating IL-7 signaling.

The only structure of a γ_c unliganded receptor/receptor complex available is the IL-7R α homodimer, although there is a non- γ_c unliganded homodimer structure of erythropoietin receptor not discussed here (15, 46). The individual IL-7R α molecules in the homodimer are very similar to the IL-7R α structure seen in the IL-7/IL-7R α complex (Fig. 6A). Backbone superimpositions give an average rmsd of less than 1 Å. IL-7R α self-associates with a binding affinity of 610 μ M, which is weaker than the IL-2R β homodimer ($K_d = 3 \mu$ M) and the IL-7R α / γ_c heterodimer ($K_d = 3 \mu$ M) (5, 15). Consistent with these *in vitro* measurements, full-length IL-7R α and IL-2R β bind more tightly to γ_c than to themselves *in vivo* as well (8, 45).

IL-7 binding is likely inhibited by the unliganded homodimer because all of the residues participating in the homodimer interface are also involved in IL-7 binding, including certain elbow loop residues. Because only a subset of residues are involved, the IL-7R α homodimer interface buries less surface area, is more apolar, and less specific than the IL-7/IL-7R α interface (Table 1). The homodimer buries on average 584 Å²; comprises 56% apolar residues, 17% polar residues, 27% charged residues, and 1 hydrogen bond; and exhibits a S_c of 0.60; whereas the IL-7/IL-7R α complex buries on average 717 Å²; comprises on average 47% apolar residues, 34% polar residues, 20% charged residues, and 5 hydrogen bonds; and exhibits S_c of on average 0.67. The IL-7R α homodimer interface is also smaller and includes many more apolar residues and much fewer hydrogen bonds than the IL-2R β / γ_c and IL-4R α / γ_c interfaces. The IL-2R β / γ_c interface buries 874 Å² of surface area and comprises 33% apolar residues and 14 hydrogen bonds, whereas the IL-4R α / γ_c interface buries 675 Å² of surface area and comprises 24% apolar residues and 21 hydrogen bonds.

The IL-7R α homodimer structure as a whole is oriented entirely differently with respect to the membrane than any of the other γ_c complex structures (compare Figs. 2 and 6B). The N- and C- termini of the receptors are located on opposite ends such that the receptors form an 'X' when looking down onto the cell surface. The average distance between the C-terminal

domains is 110 Å (Fig. 6C). In stark contrast, the average distance between the C-terminal domains of IL-2R β and IL-4R α to γ_c is 27 Å. This 83 Å distance between the C-terminal domains of the non-activating IL-7R α homodimer and activating IL-2R β and IL-4R α complexes is likely the reason why IL-7R α association does not activate the JAK/STAT pathway. The JAK1 molecules bound to the intracellular domains of IL-7R α are physically kept apart in the IL-7R α homodimer and can only be activated when brought together.

The binding constants of the IL-7R α homodimer and IL-7R α / γ_c heterodimer, the residues involved in their interfaces, and their orientation on the membrane lead to a proposed mechanism for IL-7 activation, and possibly γ_c family signaling generally, not envisioned by the stepwise, cytokine-induced heterodimerization mechanism (Fig. 6C). During inactivation, both the IL-7R α homodimer and IL-7R α / γ_c heterodimer are preformed at the membrane with their C-termini 110 Å apart, and the JAK1 and JAK3 molecules attached to the their respective intracellular domains are separated. IL-7 cannot bind to IL-7R α in either the homodimer or heterodimer because the IL-7R α loop residues that bind IL-7 also bind to IL-7R α in the homodimer and heterodimer (author's unpublished results). Accordingly, these complexes must dissociate to activate the pathway. Because IL-7R α binds γ_c more tightly than itself and the elbow loop residues not involved in the IL-7R α interface are involved in the IL-7R α / γ_c interface, the IL-7R α homodimer dissociation will require less energy than the IL-7R α / γ_c dissociation. Upon dissociation, IL-7R α and γ_c are able to bind IL-7 and rotate 90° away from the cell surface. This rotation brings the C-termini of IL-7R α and γ_c within less than 30 Å from each other so that JAK1 and JAK3 attached to the intracellular domains are in close proximity to each other and may activate signal transduction.

Diseases associated with IL-7 signaling

Gain-of-function mutations of the IL-7R α

Gain-of-function mutations in the IL-7R α have been identified from cancer patients with B-ALL and T-ALL by two independent research groups (12, 13). Given the importance of the IL-7 pathway in the survival of memory B and T cells (reviewed in 1, 47), it is not too surprising to finally isolate IL-7R α sequences in leukemia patients. In one study, Barata, Durum, and coworkers (13) determined 14 T-ALL mutations and tested a couple of them for their *in vitro* and *in vivo* effects relative to wildtype IL-7R α . These T-ALL mutations consist of insertions and/or deletions of residues at the N-terminal region of the IL-7R α TMD (Fig. 7A). In another study, Israeli and coworkers (12) determined BALL mutations that localize to either the ECD (S165-to-C165) or the TMD (multiple insertions and/or deletions) of the IL-7R α . These B-ALL mutations were tested in *in vitro* experiments relative to wildtype IL-7R α . These researchers also genotyped 30 different T-ALL mutations in the IL-7R α TMD but did not test their properties (12). These B- and T-ALL mutations were different from the first study by Zenatti *et al.* (13). A common feature of the B-ALL and T-ALL mutations between the two studies involves the incorporation of an additional cysteine residue either in the IL-7R α ECD or in the N-terminal region of the IL-7R α TMD. Not all the T-ALL mutants contain an extra cysteine residue. Experimentally, a subset of the cysteine containing B- and T-ALL mutations displayed activation of the IL-7 pathway independent of IL-7 or γ_c (12, 13). Zenatti *et al.* (13) further demonstrated T-cell leukemogenesis in mouse models of a subset of the cysteine containing T-ALL mutations. The mechanism described above for the IL-7 signaling pathway can be applied to provide a structural rationale for the ALL mutations.

I posit that the ALL mutations position either the ECDs or the TMDs of two or more copies of the IL-7R α to a distance less than 30 Å between their C-termini and in a structural geometry allowing for the activation of the JAK1 molecules on the IL-7R α ICDs in an IL-7-

and γ_c -independent manner. Structural models have been constructed of the S165-to-C165 mutation in the IL-7R α ECD in B-ALL and one of the cysteine containing insertion sequences near the IL-7R α TMD in T-ALL (T2 mutation and also called JPSP7) to support this hypothesis. Fig. 7B displays a disulfide linked S165-to-C165 IL-7R α ECDs. S165 is located on a solvent exposed loop formed by β -strands C'2 and E2 of the D2 domain and can easily be accessible to form a disulfide bond between two IL-7R α molecules when mutated to a cysteine. The C-termini of the two IL-7R α molecules in the structural model are separated by a distance of 29 Å, well within the predicted range to activate the signaling cascade independent of the α -receptors' ligands.

Before I discuss a structural model of the T2 T-ALL mutation, further background information is presented to provide credence of our structural model and mechanism. First, biochemical treatment of the T1 and T2 sequences with reducing agents (dithiothreitol or β -mercaptoethanol) on the cell surface reduced the aggregation states of these sequences and the level of STAT5 phosphorylation, strongly indicating disulfide-bond formation between two IL-7R α molecules (13). Second, mutation of the cysteine residues in the T1 and T2 sequences to either an alanine or serine abrogated STAT5 phosphorylation in cells (13). Third, unpublished results indicate mutation of the C-terminally positioned proline (P226) relative to the cysteine residue (C225) in T2 to either a serine or glycine abrogates cell survival (Dr. Scott Durum personal communication). These results suggest that it is more than just dimerization of this region, but there is also an important angular geometry at play across the bilayer, leading to activation of the pathway. Fourth, the crystal structures we have determined of the IL-7R α ECD consist of residues 1-219 of the full-length receptor. We have only been able to observe electron density to build residues in the range of 209–212. The residues between 210 (or 213) to 219, termed the juxtamembrane region, are thus highly flexible and allow mobility of the α -receptor on the cell surface. Fifth, the IL-7R α TMD is predicted to adopt a membrane-spanning α -helix. The T-ALL mutations and the wildtype IL-7R α TMD were fed into a computational design algorithm developed by DeGrado and coworkers (48) to optimize packing geometries of α -helices in a lipid bilayer, based on known membrane crystal structures. It was clear from this analysis that the majority of T-ALL sequences (and the B-ALL sequences) cannot fit entirely within the lipid bilayer and the N-termini of these sequences will be solvent exposed on the extracellular side. With this current knowledge, a plausible structural model of the T2 T-ALL mutation was constructed.

Fig. 7C shows a structural model of the T2 T-ALL mutation. Residues of PILTTCPT of the T2 mutation were solvent exposed and I228 being the first residue in the lipid bilayer. Thus, it is reasonable for a disulfide bond to be formed between C225 of chains A and B. Disulfide bond formation of cysteine residues within the lipid bilayer does occur readily (e.g. the wildtype IL-7R α TMD contains a cysteine at the C-terminal end of the sequence) (49). The juxtamembrane and transmembrane regions were built with a disulfide bond between C225 of chains A and B. The distance between the C α atoms of the W247 residues in the TMD is 11 Å, well within the range to self-activate the JAK1 kinases independent of IL-7 and γ_c . It should be noted that not all the ALL mutations contained an unpaired cysteine residue at the N-terminal region. We are currently pursuing the crystal structures of several of the ALL mutations, the wildtype IL-7R α TMD, and understanding their binding energetics with membrane environments. Therapeutically, it may be conceivable to isolate conformationally specific antibodies that can recognize a disulfide linked IL-7R α T-ALL mutant over wildtype IL-7R α on the cell surface. These experiments are underway between the Walsh and Durum laboratories.

SCID mutations

Mutations in the IL-7R α ECD have been identified in patients suffering from SCID (Fig. 8A). The human phenotype of IL-7R α SCID is T $^+$ B $^+$ NK $^+$ (10, 50). No IL-7R α SCID mutations have been identified in the TMD or ICD. Each of the IL-7R α ECD mutations map onto residues outside the binding epitope with IL-7 and the predicted binding epitope with γ_c . Instead, these mutations map onto residues in the hydrophobic cores of the D1 and D2 domains, cysteines of disulfide bonds, or the highly conserved WSXWS sequence motif. Five of the SCID mutations (G8R, S24R, L35R, C54Y, and C98Y) are located in the D1 domain. The G8R mutation could not be mapped onto the IL-7R α structure, because there was no electron density for this residue. The S24R mutation eliminates a hydrogen bond between S24 and L130 and likely requires movement to accommodate the bulky, polar arginine side chain, both of which may destabilize the linker connecting the D1 and D2 domains. The L35R mutation forces a bulky, polar side chain into the hydrophobic core of the D1 domain and presumably unfolds it. The two cysteine mutations, C54Y and C98Y, eliminate a disulfide bond and, in turn, disrupt the folding and/or stability of the D1 domain.

The remaining SCID mutations (P112H, P112S, L115R, R186stop, W197stop, W200C, S198N, W200C, S201I) are located in the D2 domain. The P112H, P112S, and L115R likely destabilize the hydrophobic core of the D2 domain. The R186 and W197 mutations convert these residues to stop codons, leading to premature termination of the mRNA. The S198N, W200C, and S201I mutations are in or near the WSXWS motif and potentially disrupt the extensive π -cation sidechain stacking interactions of this motif and its interactions with other with sidechains. Thus, all of the SCID mutations in IL-7R α probably result in folding defects or destabilize the α -receptor, limiting its ability to interact with its ligands and signal.

To date, 344 mutations have been identified in the γ_c receptor in patients suffering from X-linked SCID (<http://research.nhgri.nih.gov/scid/>). The human phenotype of γ_c SCID is T $^-$ B $^+$ NK $^-$ (reviewed in 10). Unlike the IL-7R α SCID mutations, the γ_c SCID mutations span the entire length of the receptor including the extracellular, transmembrane, and intracellular domains. Fig. 8B pinpoints the γ_c SCID mutations on the ECD. Similar to the IL-7R α SCID mutations, the majority of the SCID mutations in the ECD localize to residues involved in hydrophobic cores of the domains, cysteines in disulfide bonds, and areas at or near the WSXWS sequence motif. These γ_c SCID mutations likely cause protein-folding defects, resulting in loss of binding activity and signal transduction. Unlike the IL-7R α SCID mutations, a series of γ_c SCID mutations map to the elbow loop residues, Y103, C160, L208, C209, and G210, which interact directly with the interleukins at the *site 2a* interface. Mutagenesis studies have shown these residues to be important for γ_c interactions with all the interleukins (51, 52). For both IL-7R α and γ_c SCIDs, the current treatment strategy is bone marrow transplantation.

Acknowledgments

I thank Dr. Julie Dohm, Esq. for critical reading and comments on the manuscript. The IL-7 research project is supported by NIH grant AI72142.

References

1. Mazzucchelli R, Durum SK. Interleukin-7 receptor expression: intelligent design. *Nat Rev Immunol.* 2007; 7:144–154. [PubMed: 17259970]
2. Rochman Y, Spolski R, Leonard WJ. New insights into the regulation of T cells by gamma(c) family cytokines. *Nat Rev Immunol.* 2009; 9:480–490. [PubMed: 19543225]

3. Hamming OJ, et al. Crystal structure of interleukin-21 receptor (IL-21R) bound to IL-21 reveals that sugar chain interacting with WSXWS motif is integral part of IL-21R. *J Biol Chem.* 2012; 287:9454–9460. [PubMed: 22235133]
4. Kammer W, et al. Homodimerization of interleukin-4 receptor alpha chain can induce intracellular signaling. *J Biol Chem.* 1996; 271:23634–23637. [PubMed: 8798580]
5. Rose T, Moreau JL, Eckenberg R, Theze J. Structural analysis and modeling of a synthetic interleukin-2 mimetic and its interleukin-2Rbeta2 receptor. *J Biol Chem.* 2003; 278:22868–22876. [PubMed: 12676936]
6. Malka Y, et al. Ligand-independent homomeric and heteromeric complexes between interleukin-2 or -9 receptor subunits and the gamma chain. *J Biol Chem.* 2008; 283:33569–33577. [PubMed: 18829468]
7. Pillet AH, et al. Human IL-Rbeta chains form IL-2 binding homodimers. *Eur Cytokine Netw.* 2008; 19:49–59. [PubMed: 18299274]
8. Rose T, et al. Interleukin-7 compartmentalizes its receptor signaling complex to initiate CD4 T lymphocyte response. *J Biol Chem.* 2010; 285:14898–14908. [PubMed: 20167604]
9. Mazzucchelli RI, Riva A, Durum SK. The human IL-7 receptor gene: Deletions, polymorphisms and mutations. *Semin Immunol.* 2012; 24:225–230. [PubMed: 22425228]
10. Kovanen PE, Leonard WJ. Cytokines and immunodeficiency diseases: critical roles of the gamma(c)-dependent cytokines interleukins 2, 4, 7, 9, 15, and 21, and their signaling pathways. *Immunol Rev.* 2004; 202:67–83. [PubMed: 15546386]
11. Barata JT, Cardoso AA, Nadler LM, Boussiotis VA. Interleukin-7 promotes survival and cell cycle progression of T-cell acute lymphoblastic leukemia cells by down-regulating the cyclin-dependent kinase inhibitor p27(kip1). *Blood.* 2001; 98:1524–1531. [PubMed: 11520803]
12. Shochat C, et al. Gain-of-function mutations in interleukin-7 receptor-alpha (IL7R) in childhood acute lymphoblastic leukemias. *J Exp Med.* 2011; 208:901–908. [PubMed: 21536738]
13. Zenatti PP, et al. Oncogenic IL7R gain-of-function mutations in childhood T-cell acute lymphoblastic leukemia. *Nat Genet.* 2011; 43:932–939. [PubMed: 21892159]
14. McElroy CA, Dohm JA, Walsh ST. Structural and biophysical studies of the human IL-7/IL-7Ralpha complex. *Structure.* 2009; 17:54–65. [PubMed: 19141282]
15. McElroy CA, et al. Structural reorganization of the interleukin-7 signaling complex. *Proc Natl Acad Sci USA.* 2012; 109:2503–2508. [PubMed: 22308406]
16. Wang X, Rickert M, Garcia KC. Structure of the quaternary complex of interleukin-2 with its alpha, beta, and gammac receptors. *Science.* 2005; 310:1159–1163. [PubMed: 16293754]
17. Stauber DJ, Debler EW, Horton PA, Smith KA, Wilson IA. Crystal structure of the IL-2 signaling complex: paradigm for a heterotrimeric cytokine receptor. *Proc Natl Acad Sci USA.* 2006; 103:2788–2793.
18. Hage T, Sebald W, Reinemer P. Crystal structure of the interleukin-4/receptor alpha chain complex reveals a mosaic binding interface. *Cell.* 1999; 97:271–281. [PubMed: 10219247]
19. LaPorte SL, et al. Molecular and structural basis of cytokine receptor pleiotropy in the interleukin-4/13 system. *Cell.* 2008; 132:259–272. [PubMed: 18243101]
20. Olsen SK, et al. Crystal Structure of the interleukin-15. interleukin-15 receptor alpha complex: insights into trans and cis presentation. *J Biol Chem.* 2007; 282:37191–37204. [PubMed: 17947230]
21. Chirifu M, et al. Crystal structure of the IL-15-IL-15Ralpha complex, a cytokine-receptor unit presented in trans. *Nat Immunol.* 2007; 8:1001–1007. [PubMed: 17643103]
22. Brandhuber BJ, Boone T, Kenney WC, McKay DB. Three-dimensional structure of interleukin-2. *Science.* 1987; 238:1707–1709. [PubMed: 3500515]
23. Arkin MR, et al. Binding of small molecules to an adaptive protein-protein interface. *Proc Natl Acad Sci USA.* 2003; 100:1603–1608. [PubMed: 12582206]
24. Walter MR, et al. Crystal structure of recombinant human interleukin-4. *J Biol Chem.* 1992; 267:20371–20376. [PubMed: 1400355]

25. Powers R, Garrett DS, March CJ, Frieden EA, Gronenborn AM, Clore GM. Three-dimensional solution structure of human interleukin-4 by multidimensional heteronuclear magnetic resonance spectroscopy. *Science*. 1992; 256:1673–1677. [PubMed: 1609277]
26. Smith LJ, et al. Human interleukin 4. The solution structure of a four-helix bundle protein. *J Mol Biol*. 1992; 224:899–904. [PubMed: 1569578]
27. Wlodawer A, Pavlovsky A, Gustchina A. Crystal structure of human recombinant interleukin-4 at 2. 25 Å resolution. *FEBS Lett*. 1992; 309:59–64. [PubMed: 1511746]
28. Bondensgaard K, et al. The existence of multiple conformers of interleukin-21 directs engineering of a superpotent analogue. *J Biol Chem*. 2007; 282:23326–23336. [PubMed: 17565991]
29. Lawrence MC, Colman PM. Shape complementarity at protein/protein interfaces. *J Mol Biol*. 1993; 234:946–950. [PubMed: 8263940]
30. Walsh ST. A biosensor study indicating that entropy, electrostatics, and receptor glycosylation drive the binding interaction between interleukin-7 and its receptor. *Biochemistry*. 2010; 49:8766–8778. [PubMed: 20815339]
31. Record MT Jr, Anderson CF, Lohman TM. Thermodynamic analysis of ion effects on the binding and conformational equilibria of proteins and nucleic acids: the roles of ion association or release, screening, and ion effects on water activity. *Q Rev Biophys*. 1978; 11:103–178. [PubMed: 353875]
32. Shen BJ, Hage T, Sebald W. Global and local determinants for the kinetics of interleukin-4/interleukin-4 receptor alpha chain interaction. A biosensor study employing recombinant interleukin-4-binding protein. *Eur J Biochem*. 1996; 240:252–261. [PubMed: 8797861]
33. Schreiber G, Haran G, Zhou HX. Fundamental Aspects of Protein-Protein Association Kinetics. *Chem Rev*. 2009; 109:839–860. [PubMed: 19196002]
34. Leonard WJ. TSLP: finally in the limelight. *Nat Immunol*. 2002; 3:605–607. [PubMed: 12087416]
35. Cosenza L, Rosenbach A, White JV, Murphy JR, Smith T. Comparative model building of interleukin-7 using interleukin-4 as a template: a structural hypothesis that displays atypical surface chemistry in helix D important for receptor activation. *Protein Sci*. 2000; 9:916–926. [PubMed: 10850801]
36. vanderSpek JC, Sutherland JA, Gill BM, Gorgun G, Foss FM, Murphy JR. Structure function analysis of interleukin 7: requirement for an aromatic ring at position 143 of helix D. *Cytokine*. 2002; 17:227–233. [PubMed: 12027403]
37. Wang Y, Shen BJ, Sebald W. A mixed-charge pair in human interleukin 4 dominates high-affinity interaction with the receptor alpha chain. *Proc Natl Acad Sci USA*. 1997; 94:1657–1662. [PubMed: 9050834]
38. Hage T, Reinemer P, Sebald W. Crystals of a 1:1 complex between human interleukin-4 and the extracellular domain of its receptor alpha chain. *Eur J Biochem*. 1998; 258:831–836. [PubMed: 9874253]
39. Landgraf BE, et al. Recombinant interleukin-2 analogs. Dynamic probes for receptor structure. *J Biol Chem*. 1992; 267:18511–18519. [PubMed: 1526987]
40. Myszka DG, Arulanantham PR, Sana T, Wu Z, Morton TA, Ciardelli TL. Kinetic analysis of ligand binding to interleukin-2 receptor complexes created on an optical biosensor surface. *Protein Sci*. 1996; 5:2468–2478. [PubMed: 8976555]
41. Schreiber G, Walter MR. Cytokine-receptor interactions as drug targets. *Curr Opin Chem Biol*. 2010; 14:511–519. [PubMed: 20619718]
42. Zhang F, et al. Biophysical characterization of glycosaminoglycan-IL-7 interactions using SPR. *Biochimie*. 2012; 94:242–249. [PubMed: 22085638]
43. Liparoto SF, Myszka DG, Wu Z, Goldstein B, Laue TM, Ciardelli TL. Analysis of the role of the interleukin-2 receptor gamma chain in ligand binding. *Biochemistry*. 2002; 41:2543–2551. [PubMed: 11851401]
44. Zhang JL, Foster D, Sebald W. Human IL-21 and IL-4 bind to partially overlapping epitopes of common gamma-chain. *Biochem Biophys Res Commun*. 2003; 300:291–296. [PubMed: 12504082]

45. Pillet AH, Laverigne V, Pasquier V, Gesbert F, Theze J, Rose T. IL-2 induces conformational changes in its preassembled receptor core, which then migrates in lipid raft and binds to the cytoskeleton meshwork. *J Mol Biol.* 2010; 403:671–692. [PubMed: 20816854]
46. Livnah O, Stura EA, Middleton SA, Johnson DL, Jolliffe LK, Wilson IA. Crystallographic evidence for preformed dimers of erythropoietin receptor before ligand activation. *Science.* 1999; 283:987–990. [PubMed: 9974392]
47. Khaled AR, Durum SK. Death and Baxes: mechanisms of lymphotropic cytokines. *Immunol Rev.* 2003; 193:48–57. [PubMed: 12752670]
48. Senes A, Chadi DC, Law PB, Walters RF, Nanda V, Degrado WF. E(z), a depth-dependent potential for assessing the energies of insertion of amino acid side-chains into membranes: derivation and applications to determining the orientation of transmembrane and interfacial helices. *J Mol Biol.* 2007; 366:436–448. [PubMed: 17174324]
49. White SH, Wimley WC. Membrane protein folding and stability: physical principles. *Annu Rev Biophys Biomol Struct.* 1999; 28:319–365. [PubMed: 10410805]
50. Puel A, Ziegler SF, Buckley RH, Leonard WJ. Defective IL7R expression in T(-)B(+)NK(+) severe combined immunodeficiency. *Nat Genet.* 1998; 20:394–397. [PubMed: 9843216]
51. Olosz F, Malek TR. Three loops of the common gamma chain ectodomain required for the binding of interleukin-2 and interleukin-7. *J Biol Chem.* 2000; 275:30100–30105. [PubMed: 10887198]
52. Olosz F, Malek TR. Structural basis for binding multiple ligands by the common cytokine receptor gamma-chain. *J Biol Chem.* 2002; 277:12047–12052. [PubMed: 11815609]
53. Kossiakoff AA, deVos AM. Structural basis for cytokine hormone-receptor recognition and receptor activation. *Adv Protein Chem.* 1998; 52:67–108. [PubMed: 9917918]
54. Wang X, Lupardus P, Laporte SL, Garcia KC. Structural biology of shared cytokine receptors. *Annu Rev Immunol.* 2009; 27:29–60. [PubMed: 18817510]
55. DeLano, W. The PyMOL Molecular Graphics System. 2002.
56. Jones S, Thornton JM. Protein-protein interactions: a review of protein dimer structures. *Prog Biophys Mol Biol.* 1995; 63:31–65. [PubMed: 7746868]
57. Letzelter F, Wang Y, Sebald W. The interleukin-4 site-2 epitope determining binding of the common receptor gamma chain. *Eur J Biochem.* 1998; 257:11–20. [PubMed: 9799097]
58. Deivanayagam CC, et al. Novel fold and assembly of the repetitive B region of the *Staphylococcus aureus* collagen-binding surface protein. *Structure.* 2000; 8:67–78. [PubMed: 10673425]

Fig. 1. Schematic representation of cytokine-induced receptor heterodimerization signaling mechanism

The cytokine receptor is a transmembrane protein comprising an extracellular ligand binding domain (ECD), a single pass transmembrane domain (TMD), and an intracellular domain (ICD). The ECD folds into two fibronectin type III (FNIII) domains, D1 and D2. The γ_c receptors are members of the cytokine receptor homology class I (CRH) family that include receptors for growth hormone, erythropoietin, prolactin, and various interleukins (53, 54). The defining features of the CRH I family include conserved cysteine residues (depicted by yellow bars) involved in disulfide bonds in the D1 domain and a WSXWS primary sequence motif (depicted by red bars) in the D2 domain. During the initiation step, the cytokine or interleukin (IL) in this case interacts with the ECD of its α/β -chain receptor forming the *site 1* interface. This leads to the intermediate step where the 1:1 complex can associate with the shared common gamma-chain (γ_c) receptor. The binding of the γ_c receptor involves an interface between the interleukin and γ_c called *site 2a* and an interface between D2 regions of the α/β -chain and γ_c receptors called *site 2b*. The bringing together of the α/β -chain and γ_c receptors by the interleukin activates the classical janus kinase and signal transducers and activators of transcription (JAK/STAT) pathway and other signaling pathways.

Fig. 2. Side views relative to the cell membrane of γ_c complex structures

Ribbon diagrams of the (A) IL-7/IL-7R α complex (3di3.pdb) (14), (B) IL-2/IL-2R α /IL-2R β / γ_c complex (2bfi.pdb) (16), (C) IL-4/IL-4R α / γ_c (3bpl.pdb) (19), and (D) IL-21/IL-21R α (3tgx.pdb) (3). There is also a crystal structure of the IL-4/IL-4R α binary complex (1iar.pdb) (18), but it is highly similar to the IL-4 ternary complex above. The different chains of each structure are colored with different shades of green (IL-7 proteins), blue (IL-2 proteins), yellow (IL-4 proteins), and magenta (IL-21 proteins). Red sticks in the cytokine receptors are the disulfide bonds. The six loop regions, *L1-L6*, which constitutes the cytokine binding epitopes on the receptors are labeled for the IL-7R α structure. The interleukins adopt a 4-helix bundle topology and are labeled *HA-HD* for IL-7. MH denotes a turn of an α -helix in the first cross-over loop of IL-7. The *site 1* interaction involves HA and HC of the interleukins to their α - β -receptors. The *site 2a* interaction involves HA and HD of the interleukins to γ_c and the *site 2b* interaction involves the D2 domains of the α - β -receptors and γ_c . PyMOL was used to generate the structural pictures (55).

Fig. 3. Structural superimpositions of γ_c interleukins to IL-7

Three different views of backbone superimpositions of the 4-helix bundle of each known interleukin structure onto IL-7. The colors for each interleukin are labeled along with the corresponding root mean squared deviations (rmsds) to IL-7 performed with PyMOL. The left figure highlights the *site 1* interface (HA and HC) of the interleukins to their high affinity cytokine receptors (IL-7R α , IL-2R β , IL-4R α , and IL-21R α) and the *site 2a* interface with γ_c (HA). The middle view is looking down the 4-helix bundles. The right view highlights the *site 2a* binding interface of HD of the interleukins to γ_c .

Fig. 4. Structural superimpositions of the γ_c receptors

(A) Backbone superimpositions of the IL-7Ra, IL-4Ra, and IL-21Ra onto IL-2R β using both the D1 and D2 domains. (B) Primary sequence alignments based on the structure of IL-7 to IL-2 and IL-4. Red boxes indicate contact residues observed in the IL-2/IL-4/ γ_c structures and predicted for IL-7 and IL-21. The shaded box highlights residues buried in the hydrophobic cores of the 4-helix bundles of the interleukins. (C, D) Further backbone superimpositions of the IL-7Ra, IL-4Ra, and IL-21Ra onto IL-2R β using either D1 domains (C) or D2 domains (D). The circles indicate steric clashes or residues being too distant in the potential interactions of IL-7 or IL-7Ra to γ_c for sites 2a or 2b.

Fig. 5. Structural views of glycosylation of the IL-7Ra

(A) Structural superimposition of the IL-7 complex with an unglycosylated IL-7Ra, onto the IL-7 complex with a glycosylated IL-7Ra, indicating that N-linked glycosylation does not participate directly in the *site 1* interface with IL-7 and does not induce large conformational changes between the two structures. The six asparagines of the IL-7Ra that can be N-linked glycosylated (N29, N45, N131, N162, N212, and N213) are labeled and two N-acetyl glucosamines attached to N29, N45, and N131 were experimentally observed. The secondary elements of the IL-7Ra molecules superimpose with an rmsd of 0.6 Å. (B) Experimental observation of an O-linked glycosylation of an N-acetyl galactosamine to S133 and the conformational changes induced by the glycan from residues 133-143 of the two IL-7Ra structures (15).

Fig. 6. Proposed signaling mechanism(s) of the IL-7 pathway and the γ_c family

(A) Ribbon diagram of an IL-7Rα homodimer structure viewed as looking down onto the cell membrane (3up1.pdb) (15). (B) Side view of the IL-7Rα homodimer structure modeled on a lipid bilayer showing that there is plenty of room for the C-termini of both chains to reach the top of the cell membrane. (C) Proposed signaling mechanism(s) of the IL-7 and γ_c pathways of preassembly of the receptors before interleukin binding. The IL-7Rα/ γ_c is a model generated by superimposing the γ_c structure onto chain B of the IL-7Rα homodimer structure. Mutagenesis data indicate that the elbow loop residues of IL-7Rα are important to its interaction with γ_c independent of IL-7 (unpublished results from my laboratory). The IL-7 ternary complex is also a structural model made by docking the γ_c structure close to the IL-7/IL-7Rα binary complex (14).

Fig. 7. Mutations identified in IL-7R α from cancer patients

(A) Sequence alignments of the mutations identified from patients with T-cell acute lymphocytic leukemia (T-ALL) (designated with a 'T') and B-ALL (designated with a 'B') in the N-terminal region of the juxtamembrane/transmembrane region of IL-7R α (12, 13). (B) A model of the S165C mutation identified from patients with B-ALL. The S165C mutation induced homodimerization of the IL-7R α ECDs and activates the IL-7 pathway independent of IL-7 and γ_c . The C-termini of the two IL-7R α chains are less than 30 Å apart. (C) A model generated of the homodimerized IL-7R α with the 'T2' mutation that was demonstrated to activate the IL-7 pathway independent of IL-7 and γ_c and induced oncogenesis and death in mouse models. The C-termini of the two transmembrane domains are less than 30 Å apart.

Fig. 8. IL-7R α and γ_c severe combined immunodeficiency (SCID) mutations from patients
 (A) IL-7R α SCID mutations are depicted as red spheres and labeled. (B) γ_c X-linked SCID mutations are shown as red spheres or pink sticks. The γ_c SCID mutations colored pink are involved in the *site 2a* interactions with the interleukins. For both figures, the dashed boxes represent the residues involved in the *site 2b* interaction between the D2 domains of γ_c and α -/ β -receptors. The WSXWS sequence motifs of both structures that form extensive π -cation stacking interactions are displayed as sticks.

Table 1

Summary of the binding interfaces of Type 1 γ_c complexes^a

Interface	<i>b</i> BSA (Å ²)	H-bonds	^c S _c	% Polar	% Apolar	% Charged
IL-7/unglyco IL-7Rα ^d	728	5	0.69	32	50	18
IL7/glyco IL-7Rα	705	5	0.65	36	43	21
IL-7Rα/IL-7Rα	584	1	0.60	17	56	27
IL-2/IL-2Rβ	670	8	0.74	39	24	37
IL-2/γ _c	497	5	0.84	58	29	13
IL-2Rβ/γ _c	874	14	0.64	41	33	26
IL-4/IL-4Rα binary	835	15	0.74	40	31	29
IL-4/IL-4Rα ternary	778	14	0.70	46	22	32
IL-4/γ _c	536	10	0.82	59	18	23
IL-4Rα/γ _c	675	21	0.49	57	24	19
IL-21/IL-21Rα	1,000	17	0.70	30	32	38

^aValues calculated from PISA, CCP4, and the protein-protein interaction server at <http://www.bioinformatics.sussex.ac.uk/protorp/index.html>. The % type of residues is defined according to Thornton and coworkers (56). Polar = N, C, Q, H, S, T, W, Y. Apolar = A, G, I, L, M, F, P, V. Charged = D, R, K, E.

^bAveraged buried surface area between the cytokine/receptor and receptor at the interface.

^cShape complementarity of the interface.

^dAveraged numbers for the two IL-7/unglycosylated IL-7Rα complexes in the asymmetric unit.

Table 2Binding affinities of γ_c family members

<i>interaction</i>	<i>K_d</i>
IL-7/unglyco IL-7R α	18 μ M ^a
IL-7/glyco IL-7R α	59 nM ^a
IL-7R α /IL-7R α	0.6 mM ^b
IL-7R α / γ_c	3 μ M ^b
IL-2/IL-2R β	530 nM ^c
IL-2R β /IL-2R β	3 μ M ^c
IL-2R β / γ_c	NB ^d
IL-4/IL-4R α	150 pM ^e
IL-4R α /IL-4R α	ND ^f
IL-4R α / γ_c	NB ^g
IL-21/IL-21R α	70 pM ^h
IL-21R α /IL-21R α	ND
IL-21R α / γ_c	NB ^h

^aSPR measurement at 25 °C (14).^bSPR measurement at 25 °C (15).^cSPR measurement at 25 °C (43).^dSedimentation equilibrium measurement at 25 °C (5).^eSPR measurement at 25 °C (32).^fND = not determined or measured.^gNB = no detectable binding by SPR (57).^hSPR measurement at 25 °C (44).

Table 3Receptor domain orientations^a

<i>receptor</i>	elbow angle (ϵ , °)	twist angle (τ , °)	swivel angle (σ , °)
unglyco IL-7R α	75	159	112
glyco IL-7R α	74	158	113
IL-7R α dimer A	77	162	108
IL-7R α dimer B	79	164	107
IL-2R β	75	172	88
γ_c (IL-2)	89	158	112
IL-4R α (binary)	87	155	78
IL-4R α (ternary)	84	160	79
γ_c (IL-4)	94	167	103
IL-21R α	75	176	89

^aThe angles relate the D2 domains relative to the D1 domains. The angular geometries of the FNIII domains were determined using methods described previously (58). The elbow angle defines the angle between the two domains forming the “L” shape architecture. The twist or roll angle defines the angle between the x axes of the D1 and D2 domains. The swivel or spin of the D2 domain in the x-z plane defines the swivel angle.