Published in final edited form as:

Mol Reprod Dev. 2012 October ; 79(10): 680–688. doi:10.1002/mrd.22073.

The forkhead transcription factor FoxY regulates Nanos

Jia L. Song^{1,2} and Gary M. Wessel^{1,*}

¹Department of Molecular Biology, Cellular Biology and Biochemistry, Brown University, Providence, RI 02912 USA

Abstract

FoxY is a member of the forkhead transcription factor family that appeared enriched in the presumptive germ line of sea urchins (Ransick et al., 2002, *Dev Biol* 246:132). Here we test the hypothesis that FoxY is involved in germ line determination in this animal. We found two splice forms of FoxY that share the same DNA-binding domain but vary in the carboxy-terminal trans-activation/repression domain. Both forms of the FoxY protein are present in the ovary and in the early embryo, and their mRNAs accumulate to their highest levels in the small micromeres and adjacent non-skeletogenic mesoderm. Knockdown of FoxY resulted in a dramatic decrease in the Nanos mRNA and protein levels as well as a loss of coelomic pouches in the 2-week-old larvae. Our results indicate that FoxY positively regulates Nanos at the transcriptional level and is essential for reproductive potential in this organism.

Keywords

winged-helix proteins; germ line; multipotent cells; sea urchin

INTRODUCTION

Forkhead proteins are transcription factors characterized by their highly conserved forkhead (fh) DNA-binding domains (Kaufmann and Knochel 1996). They have diverse functions in a wide variety of animals and fungi, including cell fate specification and differentiation, regulation of the cell cycle, and metabolism (reviewed in Carlsson and Mahlapuu 2002). Forkhead proteins are thought to bind to DNA as monomers, with binding sites consisting of 15–17 bp (reviewed in Carlsson and Mahlapuu 2002). The number of forkhead-box genes, or Fox genes, correlates with the anatomical complexity of the organism. For example, 4 forkhead transcription factors are found in the yeast *Saccharomyces*, 22 in the purple sea urchin *Strongylocentrotus purpuratus* (Tu et al. 2006), and 39 in humans (reviewed in Carlsson and Mahlapuu 2002).

The *Strongylocentrotus purpuratus* FoxY (FoxC-like) gene was first identified by Ransick et al. (2002) in a differential screen for endomesoderm effectors. Ransick et al. compared the mRNA of embryos treated with LiCl, known to “vegetalize” the embryo, to mRNA from “animalized” embryos over-expressing a dominant-negative cadherin that blocks the nuclearization of β -catenin (Miller and McClay 1997). This approach was used to search for transcription factors involved in endomesodermal specification. The FoxY mRNA localized prominently with small micromeres of the early embryo and in the coelomic pouch of the larvae, a pattern of special interest to us because it overlaps with the localization of both

*Corresponding Author: 185 Meeting Street, Box G, Brown University, Providence, RI 02912, USA, rhet@brown.edu, T: 401.863.1051.

²Current address: Department of Biological Sciences, University of Delaware, Newark, 19716, USA

nanos and *vasa* mRNAs and proteins (Juliano et al. 2006; Juliano et al. 2010b; Voronina et al. 2008). *Nanos* and *Vasa* are genes involved in germ line function and in multipotency (Juliano et al. 2010a).

Nanos contains two CCHC zinc fingers and acts with Pumilio as a translational repressor by binding the Nanos response element (NRE) located in the 3'-untranslated regions (3'-UTRs) of Nanos-regulated mRNAs. The role of Nanos was first described in *Drosophila*, where Nanos and Pumilio are required together to translationally repress *hunchback* mRNA to promote posterior patterning (Murata and Wharton 1995; Wharton and Struhl 1991). In the sea urchin, Nanos is required to maintain the small micromeres and adult rudiment formation (Juliano et al. 2010b). While many studies focus on the function of Nanos as a translational repressor, its transcriptional regulation is not well understood. We tested FoxY as a potential regulator of *nanos* based on the previously reported FoxY in situ hybridization pattern (Ransick et al. 2002) and found that FoxY positively regulates *nanos* transcription.

RESULTS

Two splice forms of FoxY

We found two splice forms of *foxy* mRNA (Fig. 1, S1) that share the same DNA-binding region in the amino-terminus. Within the sixth exon, the short form has a segment of 76 nucleotides not present in the long form, resulting in a frame shift of the carboxyl terminus of the protein. As a consequence, the isoelectric point and molecular weight of FoxY-S (short splice form) and FoxY-L (long splice form) differ significantly: 6.56 / 62.7 kDa and 8.3 / 71 kDa, respectively.

Both splice forms are most abundant in early blastula and mesenchyme blastula and decrease in gastrula (Fig. 1), which is similar to a previous study detailing a high resolution time course for all forms of *foxy* mRNA (Materna et al. 2010). Using probes against both forms of *foxy*, we observed *foxy* mRNAs accumulation in cells of the small micromere lineage, as indicated by Vasa protein immunolabeling, and in the adjacent non-skeletogenic mesoderm (Fig. 2). By the larval stage, *foxy* mRNA decreases in abundance by real-time, quantitative PCR (QPCR), and is not detectable any longer by RNA in situ hybridization (Fig. 1 and data not shown).

FoxY protein is present maternally

To identify FoxY protein, we generated two different peptide antibodies against the FoxY splice forms and tested them by both immunoblotting and in situ immunolabeling. By immunoblot analysis, we detected both forms of the FoxY proteins of the expected sizes present in the ovary, blastula, and gastrula stages (Fig. 3). Unfortunately, these antibodies do not label embryos in situ, so we do not know the spatial distribution of the FoxY protein. At this point we cannot decipher if FoxY-L and FoxY-S have different mRNA and/or protein localizations, since the in situ RNA probes of whole mount embryos do not distinguish against these splice forms and the antibodies do not work in whole mount immunolabeling.

The peptide sequences used for antibody generation were designed outside of the conserved Forkhead domain. We blasted these peptide sequences in the annotated genome databases (SpBase.org) and do not find similarities to any other Fox protein (data not shown), suggesting that it's unlikely that these antibodies recognize other Fox protein family members. By immunoblot, we observed a 48% decrease in the level of FoxY protein in *foxy* morpholino antisense oligonucleotide (MASO)-knockdown embryos, indicating that the antibodies are largely selective, that the knockdown is not complete, and/or that the FoxY protein accumulates stably in the embryo (Fig. 4A). We also tested the efficacy of a *foxy* MASO with reporter constructs containing a complementary sequence to the *foxy* MASO

(Fig. 4B,C). Results indicate that only the mCherry reporter construct with the antisense *foxy* MASO sequence was repressed by the *foxy* MASO, whereas a control eGFP reporter construct without the target *foxy* sequence co-injected into the embryo was not affected by the presence of the MASO (Fig 4B,C). This result suggests that the injected *foxy* MASO is specifically binding to the target sequence and is functional in the embryo.

Knockdown of FoxY leads to a decrease in nanos mRNA and protein

To test the function of FoxY, we knocked down FoxY in embryos. We first assayed the level of *foxy* mRNAs in these embryos at 11, 17, and 24 hours post-fertilization (hpf) by QPCR, and observed that FoxY positively regulates its own mRNA (Fig. 5).

Next, we examined the effect of knockdown on Nanos2, which is present exclusively in the small micromere lineage. *nanos* mRNA was not significantly altered with FoxY knockdown at 11 and 17 hpf; by 24 hpf, however, *nanos* mRNA was significantly decreased by QPCR (Fig. 5A). As a fluorescent dye was coinjected with the *foxy* MASO as a marker for injection and also as a proxy of the amount of *foxy* MASO injected into the newly fertilized embryo, we were also able to determine by in situ hybridization that *nanos* mRNA quantities remaining in an embryo depended on the amount of *foxy* MASO introduced (Fig. 5B). *foxy* knockdown embryos also showed a significant decrease in Nanos protein by 24 hpf (Fig. 6).

FoxY knockdown leads to a loss of coelomic pouches

FoxY depletion in the embryo resulted in no obvious developmental defects up to the pluteus stage. After 2 weeks of culturing, however, the overall size of the larvae was significantly smaller than the mock-injected embryos of the same age (Fig. 7). In addition, coelomic pouches that were present in the pluteus stage regressed and eventually were lost. This *foxy*-knockdown phenotype is similar to the *nanos*-knockdown phenotype (Juliano et al. 2010b), suggesting that FoxY is epistatic to Nanos.

DISCUSSION

FoxY splice variants may have different downstream regulatory effects

Results indicated that the two splice forms of *foxy* are differentially expressed. In the ovary, it appears that the FoxY-S form is more abundant than FoxY-L, whereas in later embryonic stages, the FoxY-L is more abundant (Fig. 3). These two splice variants have different carboxy-terminal amino acid composition, thus each variant has its own *trans*-activation/repression domains and may differentially regulate its target genes. The functional difference between these two FoxY isoforms is currently not known, although previous studies have shown that the function of forkhead proteins may depend on their phosphorylation sites (Chen et al. 2009; Singh et al. ; Tan et al. 1998). As FoxY-L has 7 additional predicted serine phosphorylation sites and 1 additional threonine and tyrosine phosphorylation sites, we speculate that the FoxY transcription factor isoforms may recruit different cofactors or have its activity modified depending on the status and location of its phosphorylation sites.

FoxY knockdown results in a decrease of *nanos2* mRNA and protein

Both forms of *foxy* mRNA peak around the blastula stage (Fig. 1), and FoxY protein is present in great abundance in the ovaries (Fig. 3). From previous studies, *nanos* mRNA and protein are also abundant in the oocytes and later restricted to the small micromere descendants in the embryo (Juliano et al. 2006; Juliano et al. 2010b). Although several studies have focused on the post-translational regulation of Nanos, the *cis*-regulatory transcriptional control of the *nanos* gene is mostly unknown. Recent study on the *Drosophila nanos* identified an enhancer element located between -108 and +97 of its transcription start

site that is sufficient to drive Nanos-GFP expression in the germline stem cells (Ali et al. 2009). Maternal transcription factors Ovo and initiation complexes bip2 and Trf2 are required for germline-specific gene expression of *nanos* and *vasa* in the germ plasm in *Drosophila* (Yatsu et al. 2008). Ovo is a transcriptional activator involved in female germline maintenance in *Drosophila* (Mevel-Ninio et al. 1991; Oliver et al. 1987; Staab and Steinmann-Zwicky 1996). The sea urchin *ovo* mRNA is localized to the vegetal plate of the mesenchyme blastula, but is undetectable later (Juliano et al. 2006). Since *nanos* mRNA is upregulated in the early blastula stage and maintained at the same levels until the pluteus stage, other transcription factors are likely involved in the transcriptional activation of *nanos*. The majority of forkhead proteins bind to a seven-nucleotide core consensus sequence (RYMAAYA [(R=A or G; Y = C or T; M = A or C)]) that is found several times in the *nanos* promoter, suggesting that forkhead transcription factors can directly regulate *nanos* (Kaufmann et al. 1995; Overdier et al. 1994; Pierrou et al. 1994). *nanos* is also subjected to post-transcriptional control, such as translational regulation by RNA localization, as observed in *Drosophila* (Gavis and Lehmann 1994) and by Deadend RNA binding protein in zebrafish (Kedde et al. 2007); microRNA-mediated deadenylation (Mishima et al. 2006); and translational repression by a RNA secondary structure element as in *Xenopus* (Luo et al. 2011).

Functions of FoxY in adulthood and embryogenesis

The small micromere descendants that become part of the coelomic pouches contribute to the adult rudiment. In adult tissues, *foxy* mRNA is present in the gonads and its protein is present in the ovary (data not shown and Fig. 3). FoxY may function during oogenesis (in adult tissues) and in early development by regulating *nanos*. It may also regulate endomesodermal genes. Previous studies indicated that *foxy* mRNA is found only in the small micromeres (Materna and Davidson 2012; Ransick et al. 2002); however, using Vasa protein as a reference for small micromeres (Voronina et al., 2008), we find that *foxy* mRNA has a broader distribution within the small micromeres and in the adjacent non-skeletogenic mesoderm in the blastula and gastrula stages (Fig. 2). This suggests that FoxY may potentially regulate other genes within other lineages, although one dominant function appears to be the transcriptional regulation of *nanos*.

MATERIALS AND METHODS

Animals

Adult *Strongylocentrotus purpuratus* were obtained from Patrick Leahy (Point Loma Invertebrate Lab, Lakeside, CA) and kept in 15°C artificial seawater until needed. Artificial seawater was generated from Instant Ocean (Aquarium Systems, Mentor, OH). Eggs and sperm were collected by intracoelomic injection of 0.5M KCl. Eggs were collected in filtered seawater and sperm was collected dry.

Cloning of *foxy* spliced forms—Nested *foxy* primers were used in the cloning of full-length *foxy* in a PCR reaction using gastrula cDNA. Primers used in the first and second rounds of PCR were: FoxY5'-103For (5'-CTGCACTGACTCTGCCTACA) with FoxY-3'+2038Rev (5'-TGTCTGTTGGATCCAGCAGT) and FoxY5'-77For (5'-GCTTCACAAATCTCGCCTCA) with FoxY-TGAend (5'-TCACATACTGTGTATTCGTGT). Splice forms were detected using nested primers FoxY-F1 (5'-CATCCTAACTTGCCATGCAC) with 3'FoxY-3'end (5'-TCACATACTGTGGTATTCGTGT) and FoxYF2 (5'-TTTGATCAGTGCGATCGACTC) with FoxY R1 (5'-CTTGAGATCGTCCTGATGTC) using cDNAs from the ovary, egg, 4-cell, 32-cell, early blastula, mesenchyme blastula, gastrula, and pluteus. PCR products were run on 1.5% agarose gel. Bands were cut out, gel purified using QiaQuick columns (Qiagen,

Valencia, CA), and cloned into pGEM T-easy (Promega). Recombinant clones were sequenced.

Whole mount in situ hybridization—*foxy* antisense probes were labeled with digoxigenin (DIG RNA Labeling Kit (SP6/T7); Roche Diagnostics Corporation, Indianapolis, IN, USA). PCR primers used to clone the RNA probe were Fox-1F (5'-ATGGAGGACCAGGAGGATGATG) and Fox3' Rev (5'-GAATTCACCTTTGATGATGTACGCTCAGGTC). The plasmid was linearized with BamHI and in vitro-transcribed with Sp6 RNA polymerase. This probe was used to detect native transcript in eggs and embryos at a concentration of 0.1 µg probe/mL, according to previously published protocols (Minokawa et al. 2004).

Real-time, quantitative PCR (QPCR)—A total of 500 uninjected or 100 injected embryos were collected at various time points, and total RNA were extracted from them using the Qiagen RNeasy microkit according to manufacturer's instructions (Qiagen Inc., Valencia, CA). cDNA was amplified using the TaqMan Reverse Transcription Reagents kit (Applied Biosystems, Foster City). QPCR was performed using the 7300 Real-Time PCR system (Applied Biosystems, Foster City). QPCR primers were designed using the Primer3 program (Rozen and Skaletsky 2000). Ten uninjected or two injected embryo-equivalents were used for each QPCR reaction with the SYBER Green PCR Master Mix (Invitrogen, Carlsbad, CA). The estimated numbers of transcripts are calculated based on the level of ubiquitin in various developmental stages, as described previously (Materna and Davidson 2012). For injected embryos, results were first normalized to ubiquitin levels and expressed as fold-difference compared to the uninjected eggs. Primers QPCR-L-For (5'-CTTCTGATCGAATCCATGCC) and QPCR-exonRev (5'-TTGTCTGGATGTGTGTCTGGA) were used to detect FoxY-L and primers QPCR-S-F (5'-GATGATGAGAGACAACCAAC) and QPCR-exonRev were used to detect FoxY-S.

Antibody generation—Antiserum was raised in rabbits against polypeptides indicated in Fig. S1. N-term (DAIRATDVTAESRHC) and C-term (CPPSPFTTVSPPDTH) peptide polyclonal antibodies were generated by Sigma Genosys (Woodlands, Texas). For affinity purifications, peptides were immobilized using the Pierce AminoLink Plus Immobilization Kit (ThermoFisher Scientific; Rockford, IL) according to the manufacturer's instructions. Heat-inactivated serum was passed over the antigen-immobilized column, and bound antibodies were eluted with 1 mL 100 mM glycine (pH2.5) into 50 µl of 1 M Tris (pH 9.5).

Fluorescence quantification—Confocal Z-stack images were collected on an LSM 510 laser scanning confocal microscope (Carl Zeiss, Inc.; Thornwood, NY). Projection images of the Z-stacks were generated for quantitative analysis. Quantitative pixel-intensity analysis of whole embryos was conducted using the Metamorph program (Molecular Devices, Downingtown, PA) to determine the levels of Nanos2 immunolabeling in untreated, mock, and *foxy* MASO-injected embryos.

Western blotting—Embryo extracts were prepared in 2X sample buffer with 1 mM dithiothreitol and boiled for 5 minutes. Samples were separated by one-dimensional polyacrylamide gel electrophoresis (PAGE) in pre-cast 4–20% polyacrylamide Tris-Glycine gels (Nu-Sep Incorporated, Lawrenceville, GA, USA). Total protein was transferred onto nitrocellulose membrane (Fisher Scientific, Pittsburgh, PA, USA) and blocked with Blotto (3% non-fat dry milk (w/v), 170 mM NaCl, 50 mM Tris, 0.05% Tween20 (v/v) pH 8). Ten micrograms of each affinity purified N-term and C-term primary antibody were diluted separately in ___? mL Blotto, followed by either alkaline phosphatase- or horseradish peroxidase-conjugated secondary antibodies diluted in Blotto (1:5000 dilution (v/v); Jackson

Immunoresearch Laboratories Incorporated, West Grove, PA, USA). Blots were washed 2 times with Blotto for 15 min, followed by one wash in 1X phosphate-buffered saline (PBS)-0.5% Tween20. Immunoreactivity was detected in developmental series by the alkaline phosphatase colorimetric reaction using 5-bromo-4-chloro-3-indolyl phosphate/nitro blue tetrazolium (BCIP/NBT). For the *foxy* MASO Western blots, proteins were detected by horseradish peroxidase-dependent chemiluminescence using luminol / hydrogen peroxide. The FoxY protein was normalized to the yolk protein YP30 and quantitated by densitometry using Image J (Abramoff et al. 2004).

Microinjection—Morpholino antisense oligonucleotides (MASO) against *foxy*, (5'-CATGGCTCCAAGTGCAGAACTAC) was ordered from GeneTools (Philomath, OR). Microinjections were performed as previously described (Cheers and Ettensohn 2004), with modifications. MASO oligonucleotides were resuspended in sterile water and heated to 60°C for 10 minutes prior to use. Injection solutions contain 20% sterile glycerol, 2 mg/ml 10,000MW Texas Red lysine charged dextran (Molecular Probes, Carlsbad, CA), and varying concentrations of MASO. Eggs were dejellied in acidic seawater (pH 5.2) for 10 minutes on ice, followed by seawater washes. Dejellied eggs were rowed onto protamine sulfate-coated (4% w/v) 60×15 mm petri dishes. Eggs were fertilized with sperm in the presence of 1 mM 3-amino-triazol (Sigma, St. Louis, MO). Injections were performed using a Femto Jet® injection system (Eppendorf; Hamburg, Germany). Injection needles 1×90 mm glass capillaries with filaments (Narashige; Tokyo, Japan) were pulled on a vertical needle puller (Narishige; Tokyo, Japan). The injected *foxy* MASO corresponds to approximately 12 nM.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We gratefully acknowledge support from the National Institutes of Health (HD028152) and the National Science Foundation (IOS-1120972).

We thank present and past members of PRIMO for helpful discussions. This work is supported by NRSA awarded to JLS and NIH grant awarded to GMW.

Abbreviations

hpf	hours post-fertilization
MASO	morpholino antisense oligonucleotide
QPCR	real-time, quantitative PCR
UTR	untranslated region

References

- Abramoff MD, Magalhães PJ, Ram SJ. Image Processing with ImageJ. Biophotonics International. 2004; 11(7):36–42.
- Ali I, Ur Rehman M, Rashid F, Khan S, Iqbal A, Laixin X, Ahmed NU, Swati AZ. Cis-regulatory elements affecting the Nanos gene promoter in the germline stem cells. J Biotechnol. 2009
- Carlsson P, Mahlapuu M. Forkhead transcription factors: key players in development and metabolism. Dev Biol. 2002; 250(1):1–23. [PubMed: 12297093]

- Cheers MS, Etensohn CA. Rapid microinjection of fertilized eggs. *Methods Cell Biol.* 2004; 74:287–310. [PubMed: 15575612]
- Chen YJ, Dominguez-Brauer C, Wang Z, Asara JM, Costa RH, Tyner AL, Lau LF, Raychaudhuri P. A conserved phosphorylation site within the forkhead domain of FoxM1B is required for its activation by cyclin-CDK1. *J Biol Chem.* 2009; 284(44):30695–30707. [PubMed: 19737929]
- Gavis ER, Lehmann R. Translational regulation of nanos by RNA localization. *Nature.* 1994; 369(6478):315–318. [PubMed: 7514276]
- Juliano CE, Swartz SZ, Wessel GM. A conserved germline multipotency program. *Development.* 2010a; 137(24):4113–4126. [PubMed: 21098563]
- Juliano CE, Voronina E, Stack C, Aldrich M, Cameron AR, Wessel GM. Germ line determinants are not localized early in sea urchin development, but do accumulate in the small micromere lineage. *Dev Biol.* 2006; 300(1):406–415. [PubMed: 16970939]
- Juliano CE, Yajima M, Wessel GM. Nanos functions to maintain the fate of the small micromere lineage in the sea urchin embryo. *Dev Biol.* 2010b; 337(2):220–232. [PubMed: 19878662]
- Kaufmann E, Knochel W. Five years on the wings of fork head. *Mech Dev.* 1996; 57(1):3–20. [PubMed: 8817449]
- Kaufmann E, Muller D, Knochel W. DNA recognition site analysis of *Xenopus* winged helix proteins. *J Mol Biol.* 1995; 248(2):239–254. [PubMed: 7739038]
- Kedde M, Strasser MJ, Boldajipour B, Oude Vrielink JA, Slanchev K, le Sage C, Nagel R, Voorhoeve PM, van Duijse J, Orom UA, Lund AH, Perrakis A, Raz E, Agami R. RNA-binding protein Dnd1 inhibits microRNA access to target mRNA. *Cell.* 2007; 131(7):1273–1286. [PubMed: 18155131]
- Luo X, Nerlick S, An W, King ML. *Xenopus* germline nanos1 is translationally repressed by a novel structure-based mechanism. *Development.* 2011; 138(3):589–598. [PubMed: 21205802]
- Materna SC, Davidson EH. A comprehensive analysis of Delta signaling in pre-gastrular sea urchin embryos. *Dev Biol.* 2012; 364(1):77–87. [PubMed: 22306924]
- Materna SC, Nam J, Davidson EH. High accuracy, high-resolution prevalence measurement for the majority of locally expressed regulatory genes in early sea urchin development. *Gene Expr Patterns.* 2010; 10(4–5):177–184. [PubMed: 20398801]
- Mevel-Ninio M, Terracol R, Kafatos FC. The ovo gene of *Drosophila* encodes a zinc finger protein required for female germ line development. *Embo J.* 1991; 10(8):2259–2266. [PubMed: 1712294]
- Miller JR, McClay DR. Characterization of the role of cadherin in regulating cell adhesion during sea urchin development. *Dev Biol.* 1997; 192(2):323–339. [PubMed: 9441671]
- Minokawa T, Rast JP, Arenas-Mena C, Franco CB, Davidson EH. Expression patterns of four different regulatory genes that function during sea urchin development. *Gene Expr Patterns.* 2004; 4(4):449–456. [PubMed: 15183312]
- Mishima Y, Giraldez AJ, Takeda Y, Fujiwara T, Sakamoto H, Schier AF, Inoue K. Differential regulation of germline mRNAs in soma and germ cells by zebrafish miR-430. *Curr Biol.* 2006; 16(21):2135–2142. [PubMed: 17084698]
- Murata Y, Wharton RP. Binding of pumilio to maternal hunchback mRNA is required for posterior patterning in *Drosophila* embryos. *Cell.* 1995; 80(5):747–756. [PubMed: 7889568]
- Oliver B, Perrimon N, Mahowald AP. The ovo locus is required for sex-specific germ line maintenance in *Drosophila*. *Genes Dev.* 1987; 1(9):913–923. [PubMed: 3428601]
- Overdier DG, Porcella A, Costa RH. The DNA-binding specificity of the hepatocyte nuclear factor 3/ forkhead domain is influenced by amino-acid residues adjacent to the recognition helix. *Mol Cell Biol.* 1994; 14(4):2755–2766. [PubMed: 8139574]
- Pierrou S, Hellqvist M, Samuelsson L, Enerback S, Carlsson P. Cloning and characterization of seven human forkhead proteins: binding site specificity and DNA bending. *Embo J.* 1994; 13(20):5002–5012. [PubMed: 7957066]
- Ransick A, Rast JP, Minokawa T, Calestani C, Davidson EH. New early zygotic regulators expressed in endomesoderm of sea urchin embryos discovered by differential array hybridization. *Dev Biol.* 2002; 246(1):132–147. [PubMed: 12027439]
- Rozen S, Skaletsky H. Primer3 on the WWW for general users and for biologist programmers. *Methods Mol Biol.* 2000; 132:365–386. [PubMed: 10547847]

- Singh A, Ye M, Bucur O, Zhu S, Santos MT, Rabinovitz I, Wei W, Gao D, Hahn WC, Khosravi-Far R. Protein Phosphatase 2A Reactivates FOXO3a through a Dynamic Interplay with 14-3-3 and AKT. *Mol Biol Cell*.
- Staab S, Steinmann-Zwicky M. Female germ cells of *Drosophila* require zygotic ovo and otu product for survival in larvae and pupae respectively. *Mech Dev*. 1996; 54(2):205–210. [PubMed: 8652413]
- Tan PB, Lackner MR, Kim SK. MAP kinase signaling specificity mediated by the LIN-1 Ets/LIN-31 WH transcription factor complex during *C. elegans* vulval induction. *Cell*. 1998; 93(4):569–580. [PubMed: 9604932]
- Tu Q, Brown CT, Davidson EH, Oliveri P. Sea urchin Forkhead gene family: phylogeny and embryonic expression. *Dev Biol*. 2006; 300(1):49–62. [PubMed: 17081512]
- Voronina E, Lopez M, Juliano CE, Gustafson E, Song JL, Extavour C, George S, Oliveri P, McClay D, Wessel G. Vasa protein expression is restricted to the small micromeres of the sea urchin, but is inducible in other lineages early in development. *Dev Biol*. 2008; 314(2):276–286. [PubMed: 18191830]
- Wharton RP, Struhl G. RNA regulatory elements mediate control of *Drosophila* body pattern by the posterior morphogen nanos. *Cell*. 1991; 67(5):955–967. [PubMed: 1720354]
- Yatsu J, Hayashi M, Mukai M, Arita K, Shigenobu S, Kobayashi S. Maternal RNAs encoding transcription factors for germline-specific gene expression in *Drosophila* embryos. *Int J Dev Biol*. 2008; 52(7):913–923. [PubMed: 18956321]

Fig. 1.

Two splice forms of *foxy*. (A) *foxy* splice forms both share the same conserved forkhead DNA-binding domain, but differ in the sixth exon near the C-terminus domain of the protein. (B) QPCR quantitation of FoxY-L and FoxY-S transcripts. Calculation of the estimated number of transcripts is based on ubiquitin levels measured during corresponding developmental stages (Materna et al. 2010).

Fig. 2. *foxy* mRNA has broader accumulation than the Vasa protein. Embryos were first labeled with *foxy* mRNA in situ probe, followed by immunostaining for Vasa protein (red). *foxy* mRNAs has broader expression in the non-skeletogenic Vegetal-1 (Veg1) and Vegetal-2 (Veg2) regions.

Fig. 3. FoxY protein expression. FoxY protein isoforms are detected in the ovary, blastula, and gastrula embryonic extracts. The N-term antibody detects both forms of FoxY whereas the C-term antibody specifically detects FoxY-S.

Fig. 4. *foxy* MASO control experiment. (A) FoxY protein levels decreased by approximately 50%, as measured by densitometry using imageJ. The FoxY protein is normalized to YP30 yolk protein. (B) mCherry reporter transcript containing complimentary sequence to the *foxy* MASO was repressed in the presence of *foxy* MASO whereas the control eGFP reporter, whose coding sequence was flanked by β -globin UTRs, expressed in the presence or absence of *foxy* MASO. (C) The fluorescence level of mCherry was normalized to eGFP signals to adjust for the amount of injected mRNAs. The difference in mCherry/eGFP fluorescence measurements observed in control vs. *foxy* MASO-treated sets was statistically significant ($p=1.89658 \times 10^{-24}$, Student's *t*-test).

Fig. 5. *foxy* knockdown results in a decrease of *nanos* mRNA at 24 hpf. (A) QPCR was used to measure the level of transcript levels in *foxy* MASO-injected embryos. Transcript levels of *nanos*, *foxy-L*, and *foxy-S* were assayed at 11, 17, and 24 hpf. Results are normalized to mock-injected (Texas Red) controls. (B) Whole mount RNA in situ hybridization was used to detect the level of *nanos* mRNAs in control and *foxy* MASO-injected embryos at 24 hpf. The depletion in *nanos* mRNA is dependent on the amount of *foxy* MASO injected (indicated by the dashed circle), as reflected by the intensity of co-injected Texas Red dextran.

A.

B.

Fig. 6.

foxy-knockdown embryos have decreased Nanos protein. Embryos injected with *foxy* MASO were immunolabeled with the Nanos antibody at the blastula stage. (A) Fluorescent signals using the Nanos antibody (green) was quantitated by pixel intensity in uninjected and *foxy* MASO-treated blastulae. The Hoescht dye indicates nuclear staining. (B) Fluorescent signals were quantified with Metamorph as described in the Methods and Materials. The horizontal bar indicates averaged fluorescent signals. ($P=0.06$ by the Student's *t*-test).

Fig. 7.

foxy MASO-injected embryos are smaller and lose their coelomic pouches in the late larval stage (2 weeks after fertilization). (A) *foxy* MASO-injected embryos were smaller in size and have a loss of coelomic pouches. (B) Three independent sets of embryos were injected with Texas Red control or *foxy* MASO and cultured for 2 weeks. The *foxy* MASO-injected embryos have significantly fewer normal embryos. Normal embryos are defined as having 6 arms, skeletal structures, a functional gut, and coelomic pouches. Student's *t*-test was used to determine the significance of the knockdown compared to the mock control ($P=0.038$).