Published in final edited form as:

Chem Sci. 2012 January 1; 3(1): 72–76. doi:10.1039/C1SC00515D.

C(sp³)–F reductive elimination from alkylgold(III) fluoride complexes†

Neal P. Mankad and F. Dean Toste*

Department of Chemistry, University of California-Berkeley, Berkeley, CA, 94720, USA

Abstract

Rare examples of C(sp³)–F reductive elimination were observed from several *cis*-F₂Au(R)(IPr) intermediates generated by oxidation of (IPr)AuR complexes with XeF₂. For R groups bearing β-hydrogens, β-hydride elimination was competitive with C(sp³)–F reductive elimination. For strained cyclic R groups and most acyclic R groups lacking β-hydrogens, carbocation-like rearrangements occurred prior to C(sp³)–F reductive elimination. Kinetics of the decay of one *cis*-F₂Au(R)(IPr) species, stereochemical analysis of reductive elimination with a chiral R group, and DFT analysis collectively suggest C(sp³)–F reductive elimination proceeding through transient cationic [(IPr)Au(F)(R)]⁺ intermediates with significant ionization of the Au–alkyl bonds.

Introduction

Installation of C–F bonds by transition metal catalysis is particularly fascinating both because of significant interest in fluorocarbon products for important applications,¹ but also because of the fundamental challenges posed by designing C–F bond-forming catalytic cycles.² One main difficulty involves mediating C–F reductive elimination, which is likely to be the key turnover step in many such catalytic cycles. C–F reductive elimination remains rare compared to other types of C–X reductive elimination (X = *e.g.* C, N, O, Cl, Br, I) despite the large thermodynamic driving force provided by formation of strong C–F bonds (H₃C–F BDE = 110 kcal mol^{–1}).³ Key advances in stoichiometric C(sp²)–F reductive elimination⁴ as well as fluorination of transition metal π-complexes⁵ have led directly to recent discoveries of corresponding C(sp²)–F and C(allyl)–F bond-forming catalysts.^{2,4d,5,6} On the other hand, stoichiometric C(sp³)–F reductive elimination from discrete σ-organometallic intermediates has not been extensively studied⁷ despite a number of reports of stable, high-valent alkyl–metal fluoride complexes.⁸ Moreover, the reverse process, C(sp³)–F oxidative addition, has only been observed directly in a single recent report.⁹ Identification of systems that allow for C(sp³)–F reductive elimination and careful study of the principles that govern this process promise to guide the design of future C(sp³)–F bond-forming catalysts.¹⁰

Results and discussion

Fluorination of Au–C(sp³) bonds

Alkylgold(III) intermediates are well known to undergo facile C–X reductive elimination (X = O, Cl, Br, or I). Recently a mechanistic study was conducted on carbon–halogen reductive

†Electronic supplementary information (ESI) available: Synthetic, spectroscopic, kinetic, crystallographic, and computational details. CCDC reference numbers 837541. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/c1sc00515d

elimination from $X_2Au(Me)(IPr)$ complexes ($IPr = 1,3\text{-bis}(2,6\text{-diisopropylphenyl})\text{imidazol-2-ylidene}$; $X = Cl, Br, \text{ or } I$), which produce CH_3X even at low temperature.¹¹ In contrast, we recently reported the oxidation of $(IPr)AuMe$ (**1**) with XeF_2 to yield the difluoride analogue $cis\text{-}F_2Au(Me)(IPr)$ (**1-F₂**), which is remarkably robust towards thermal decomposition by CH_3F elimination.¹²

We reasoned that ground-state destabilization of $F_2Au(R)(IPr)$ complexes might lead to successful $C(sp^3)\text{-}F$ bond formation. It is well known that steric pressure facilitates reductive elimination processes in this way.¹³ Indeed, during the course of our ongoing studies on the role of gold(III) fluorides in catalysis,^{12,14} we sought to oxidize the more hindered complexes $(IPr)Au(tBu)$ (**2**) and $(IPr)AuCH_2CH_2tBu$ (**3**) with XeF_2 in order to study the reactivity of the intermediates $cis\text{-}F_2Au(tBu)(IPr)$ (**2-F₂**) and $cis\text{-}F_2Au(CH_2CH_2tBu)(IPr)$ (**3-F₂**) with arylboronic acid coupling partners. During those studies we found that, if starved of coupling partners, these intermediates decomposed to yield product mixtures containing the gold(I) fluoride complex $(IPr)AuF$, as well as alkenes resulting from formal $\beta\text{-H}$ elimination and fluoroalkanes presumably derived from $C(sp^3)\text{-}F$ reductive elimination (Table 1).¹⁵ Reaction of $(IPr)AuCH_2CD_2tBu$ (**4**), in which the β position is perdeuterated, with XeF_2 led to a product mixture only slightly perturbed from that of **3** (Table 1) and in which $\beta\text{-H(D)}$ elimination still outcompeted $C\text{-}F$ elimination.

We next examined complexes bearing cyclic alkyl groups in order to disfavor $\beta\text{-H}$ elimination (Table 2). For cyclohexyl derivative **5**, $C\text{-}F$ reductive elimination indeed outcompeted $\beta\text{-H}$ elimination by a ratio of $\sim 5 : 2$. The bulkier (–)-menthyl derivative **6** produced (–)-menthyl fluoride as the exclusive product with perfect stereoretention and with no evidence of alkene products.¹⁶ Surprisingly, $\beta\text{-H}$ elimination outcompeted $C\text{-}F$ reductive elimination for cyclopentyl derivative **7**. For the more strained cycloalkyl derivatives **8** and **9**, $\beta\text{-H}$ elimination pathways were marginalized but carbocation-like rearrangement processes were observed as minor pathways relative to direct $C\text{-}F$ reductive elimination. Specifically, cyclopropylmethyl fluoride was observed as a minor product from fluorination of cyclobutyl derivative **8**, and allyl fluoride was observed as a minor product from fluorination of cyclopropyl derivative **9** (Table 2).

We next studied complexes lacking $\beta\text{-hydrogens}$ (Table 3). Reaction of the *neo*-pentyl complex **10** with XeF_2 also led to carbocation-like behavior, with the rearranged tertiary alkyl fluoride product (*tert*-pentyl fluoride) dominating over the expected primary alkyl fluoride product (*neo*-pentyl fluoride). This apparent methyl migration was also observed for the trimethylsilylmethyl derivative **11**, with the rearrangement pathway outcompeting the direct $C\text{-}F$ reductive elimination pathway to yield ethylfluorodimethylsilane as the major product (Table 3). Alkyl migration even occurred for the 1-adamantylmethyl complex **12** upon reaction with XeF_2 , further highlighting the preference for Au^{III} -mediated $C(sp^3)\text{-}F$ reductive elimination to produce tertiary rather than primary fluoroalkanes when possible. By contrast, reaction of $(IPr)AuCH_2Ph$ (**13**) with XeF_2 yielded FCH_2Ph in good yield (Table 3).

The $(IPr)AuF$ byproduct of these reductive elimination processes was accessed on preparative scale by reaction of **13** with XeF_2 and studied by X-ray crystallography. The solid-state structure of $(IPr)AuF \cdot 2CH_2Cl_2$ revealed an extended hydrogen-bonding network, with the CH_2Cl_2 solvent molecules acting as hydrogen bond donors towards the terminal $Au\text{-}F$ moiety (Fig. 1). Similar hydrogen bonding interactions exist for $(SIPr)AuF \cdot 2CH_2Cl_2$ ($SIPr = 1,3\text{-bis}(2,6\text{-diisopropylphenyl})\text{imidazolin-2-ylidene}$).¹⁷ The $Au\text{-}F$ distance in $(IPr)AuF$ (2.071 (2) Å) is slightly shorter than that in $(SIPr)AuF$ (2.082(2) Å).

Kinetics of the decay of an alkylgold(III) difluoride intermediate

For several of the fluorination reactions the *cis*-F₂Au(R)(IPr) intermediates were easily identified by their characteristic ¹⁹F NMR properties (Table S1†),¹² and it was thus established that (IPr)AuF and the organic products depicted in Tables 1–3 appeared over time as these *cis*-F₂Au(R)(IPr) complexes disappeared. One such intermediate, **11-F₂**, was chosen for kinetics studies. The disappearance of intermediate **11-F₂** obeyed first-order kinetics (Figure S1†). An Eyring plot (Figure S2†) revealed activation parameters of $\Delta H^\ddagger = 12.7 \text{ kcal mol}^{-1}$ and $\Delta S^\ddagger = -30 \text{ e.u.}$, and thus an activation free energy of $\Delta G^\ddagger = 21.7 \text{ kcal mol}^{-1}$ at 298 K. When samples were spiked with various concentrations of added tetrabutylammonium fluoride (TBAF), the decay of **11-F₂** exhibited an approximately inverse first-order relationship with fluoride concentration (Fig. 2 and S3†).

A proposed mechanistic scheme accounting for the various observed reaction pathways is shown in Scheme 1. Consistent with the inverse rate dependence on fluoride concentration, we propose initial fluoride dissociation from an alkylgold(III) difluoride complex to reveal an activated, 3-coordinate complex of the type [(IPr)Au(F)(R)]⁺. Such a complex with R = Me has been characterized crystallographically in dimeric form.¹² This key intermediate can then mediate R–F reductive elimination *via* a classical 3-centered transition state, consistent with the observed retention of stereochemistry observed for **6**, followed by trapping of the [(IPr)Au]⁺ fragment with the dissociated fluoride. The coordinatively unsaturated nature of a [(IPr)Au(F)(R)]⁺ species would also allow for β -H elimination either by a traditional Au-mediated mechanism¹⁸ or by deprotonation of the β position by F[–]. We further propose that such a 3-coordinate, cationic species places a significant degree of positive charge character in the Au^{III}–R bond, resulting in the alkyl migrations and other carbocation-like rearrangements described herein *in lieu* of direct C–F reductive elimination. The large and negative $\Delta S^\ddagger$ for the decay of **11-F₂** is unusual for a dissociative process and could be indicative of a highly ordered transition state necessary for tandem methyl migration/reductive elimination. It is also possible that solvent reorganization is necessary for solvation of F[–] and contributes to this entropic term.¹⁹

Iodination of Au–C(sp³) bonds

It has been established previously¹¹ that CH₃I reductive elimination from *trans*-I₂Au(Me)(IPr) similarly proceeds through transient [(IPr)Au(Me)(I)]⁺. We thus chose to examine the degree of carbocation-like character possessed by such cationic alkylgold(III) iodide intermediates relative to their alkylgold(III) fluoride analogues.

Product mixtures derived from halogenation of selected (IPr)AuR complexes with I₂ are shown in Table 4. First it is worth reiterating that though C–F reductive elimination was not observed even at elevated temperatures for R = Me, clean C–I reductive elimination was observed at room temperature.^{11,12} While β -H elimination and cyclobutyl ring contraction/opening did occur to some extent for the bulkier iodide complexes, certain carbocation-like rearrangement such as methyl migration and cyclopropyl ring-opening which occurred readily for the fluoro complexes did not occur to any detectable extent for the iodo complexes. For example, the *neo*-pentyl and trimethylsilylmethyl derivatives produced the primary rather than tertiary iodoalkane products exclusively (Table 4). It follows that the carbocationic character of alkylgold(III) complexes is accentuated by fluoride ligands. Lastly, we note that, like C–F reductive elimination, C–I reductive elimination from gold(III) complexes has been shown previously to proceed with stereoretention.¹⁴ For no case other than (IPr)AuMe was the alkylgold(III) diiodide intermediate observable by NMR spectroscopy prior to iodoalkane extrusion.¹¹

DFT analysis of relevant alkylgold(III) intermediates

In order to examine the nature of the cationic [(IPr)Au(F)(R)]⁺ intermediates, we initiated a preliminary DFT study. The optimized structure of *cis*-F₂Au(CH₂*t*Bu)(IPr) (**10-F**₂) exhibited the expected square planar geometry with a Au–CH₂ distance of 2.110 Å and Au–F distances of 2.014 and 2.112 Å (*cis* and *trans* to the alkyl group, respectively).

Minimization of [(IPr)Au(CH₂*t*Bu)(F)]⁺ (**10-F**⁺) from various starting geometries converged to an optimized T-shaped geometry, with the vacant coordination site occurring *trans* to the alkyl ligand. The Au–CH₂ distance in **10-F**⁺ was elongated to 2.180 Å despite dissociation of the *trans* ligand and onset of positive molecular charge. The calculated LUMO (LUMO = lowest unoccupied molecular orbital) of **10-F**⁺, shown in Fig. 3, was σ* (Au–alkyl) in character and exhibited a through-space antibonding overlap between the σ* (Au–alkyl) contribution and that from an adjacent σ(C–CH₃) bond, suggesting an electronic mechanism for methyl migration. Mulliken population analysis indicated that the LUMO of **10-F**⁺ had larger %C contribution (43%) and smaller %Au contribution (20%) than the LUMO of **10-F**₂ (26% C, 31% Au).

In order to understand the special carbocation-like behavior of [(IPr)Au(F)(R)]⁺ intermediates, we analyzed the entire halide series [(IPr)Au(X)(CH₂*t*Bu)]⁺ (**10-X**⁺, X = F, Cl, Br, and I) by analogous DFT methods. The entire series of complexes was predicted to have roughly similar Au–C_{alkyl} distances, electrostatic charges on CH₂, and LUMO compositions (Table S4[†]). A striking relationship was observed, however, between the identity of the halide ligand X and the natural charge on Au (Table 5). According to natural population analysis, the electrostatic charge on Au ranged from 0.48e in **10-I**⁺ to 0.86e in **10-F**⁺. Because the natural charge on the alkyl CH₂ unit remained roughly constant (0.14e for **10-I**⁺ to 0.11e for **10-F**⁺) across the series, this translated to a particularly polarized Au–C_{alkyl} bond for **10-F**⁺, as measured by the difference in charge between Au and CH₂ (Table 5). In other words, the fluoro ligand imparts a build-up of positive charge on Au, thereby partially ionizing the Au–C_{alkyl} bond and making (IPr)AuF a particularly effective leaving group. These factors are likely contributors to the unusual carbocation-like behaviour observed in Au-mediated C(sp³)–F reductive elimination reactions.

Conclusions

Alkylgold(III) fluoride complexes are proposed as the active electrophiles in many Au(I)/Au(III) catalytic cycles involving nucleophilic coupling partners.²⁰ For such reactions it has been proposed^{12,14,21} that:

1. C–C coupling is enabled by the relatively long lifetime of alkylgold(III) fluoride intermediates towards C–F reductive elimination
2. C–C coupling occurs by direct attack of nucleophiles on the alkyl group rather than by transmetalation to Au followed by reductive elimination
3. the presence of fluoride ligands enhances the reactivity of alkylgold(III) intermediates towards cross-coupling.

The results communicated herein are fully consistent with these assertions. Gold(III)-mediated C(sp³)–F reductive elimination clearly is less facile than, for example, gold(III)-mediated C(sp³)–I reductive elimination, based both on the differing reactivity of F₂Au(Me)(IPr) and I₂Au(Me)(IPr) as well as the longer lifetimes of F₂Au(R)(IPr) intermediates relative to the corresponding I₂Au(R)(IPr) intermediates. The predominantly C_{alkyl}-centric nature of the LUMO in complexes like **10-F**⁺ provides an electronic basis for direct attack of nucleophiles onto C_{alkyl} rather than Au. Apparently one role of fluoride in such cross-coupling reactions, in addition to forming strong B–F or Si–F bonds, is to accentuate

carbocation-like character by having a highly ionic interaction with Au^{III} relative to other, more covalent Au^{III}–X bonds (such as X = I). This discrepancy is evident both in the differing electronic structures of **10-F**⁺ and **10-I**⁺ as well as in their distinct reductive elimination reactivity patterns (Tables 1–4). We therefore suspect that Au^{III}–F bonds will continue to play an important role in Au(I)/Au(III) catalysis development, and attempts to incorporate C–F reductive elimination into such cycles are currently underway.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by NIHGM (RO1 GM073932). N.P.M. was supported by a NIH Kirchstein-NRSA postdoctoral fellowship. The UC Berkeley Molecular Graphics and Computational Facility is supported by the NSF (CHE-0840505). Prof. Robert Bergman provided helpful suggestions.

Notes and references

1. Müller K, Faeh C, Diederich F. *Science*. 2007; 317:1881. [PubMed: 17901324]
2. Furuya T, Hamlet AS, Ritter T. *Nature*. 2011; 473:470. [PubMed: 21614074]
3. McMillen DF, Golden DM. *Annu. Rev. Phys. Chem.* 1982; 33:493.
4. Selected examples: Grushin VV. *Acc. Chem. Res.* 2010; 43:160. [PubMed: 19788304] Ball ND, Sanford MS. *J. Am. Chem. Soc.* 2009; 131:3796. [PubMed: 19249867] Furuya T, Benitez D, Tkatchouk E, Strom A, Tang P, Goddard WA III, Ritter T. *J. Am. Chem. Soc.* 2010; 132:3793. [PubMed: 20196595] Watson DA, Su M, Teverovskiy G, Zhang Y, Garcia-Fortanet J, Kinzel T, Buchwald SL. *Science*. 2009; 25:1661. [PubMed: 19679769]
5. Selected examples: Akana JA, Bhattacharyya KX, Müller P, Sadighi JP. *J. Am. Chem. Soc.* 2007; 129:7736. [PubMed: 17547409] Katcher MH, Doyle AG. *J. Am. Chem. Soc.* 2010; 132:17402. Hollingworth C, Hazari A, Hopkinson MN, Tredwell M, Benedetto E, Huiban M, Gee AD, Brown JM, Gouverneur V. *Angew. Chem., Int. Ed.* 2011; 50:2613.
6. For example: Furuya T, Ritter T. *Org. Lett.* 2009; 11:2860. [PubMed: 19507870] Hull KL, Anani WQ, Sanford MS. *J. Am. Chem. Soc.* 2006; 128:7134. [PubMed: 16734446] Wang X, Mei T-S, Yu J-Q. *J. Am. Chem. Soc.* 2009; 131:7520. [PubMed: 19435367]
7. (a) Zhao S-B, Becker JJ, Gagné MR. *Organometallics*. 2011; 30:3926. [PubMed: 21869853] (b) Kaspi AW, Goldberg I, Vigalok A. *J. Am. Chem. Soc.* 2010; 132:10626. [PubMed: 20681679]
8. Selected examples: Anderson CM, Crespo M, Ferguson G, Lough AJ, Puddephatt RJ. *Organometallics*. 1992; 11:1177. Hughes RP, Laritchev RB, Zakharov LN, Rheingold AL. *Organometallics*. 2005; 24:4845.
9. Choi J, Wang DY, Kundu S, Choliy Y, Emge TJ, Krogh-Jespersen K, Goldman AS. *Science*. 2011; 332:1545. [PubMed: 21700870] See also: Terao J, Todo H, Watabe H, Ikumi A, Shinohara Y, Kambe N. *Pure Appl. Chem.* 2008; 80:941. and references therein.
10. C(sp³)–F reductive elimination can be hypothesized to occur for certain existing catalytic cycles. For example: Wu T, Yin G, Liu G. *J. Am. Chem. Soc.* 2009; 131:16354. [PubMed: 19856929]
11. Scott VJ, Labinger JA, Bercaw JE. *Organometallics*. 2010; 29:4090.
12. Mankad NP, Toste FD. *J. Am. Chem. Soc.* 2010; 132:12859. [PubMed: 20726525]
13. Hartwig JF. *Inorg. Chem.* 2007; 46:1936. [PubMed: 17348724]
14. Tkatchouk E, Mankad NP, Benitez D, Goddard WA III, Toste FD. *J. Am. Chem. Soc.* 2011; 133:14293. [PubMed: 21861448]
15. C–C homocoupling rather than C–F coupling occurs when similar reactions are carried out with R₃PAuR⁺ complexes: Hashmi ASK, Ramamurthi TD, Rominger F. *J. Organomet. Chem.* 2009; 694:592. (b) ref. 12.
16. Zuzek AA, Reynolds SC, Glueck DS, Golen JA, Rheingold AL. *Organometallics*. 2011; 30:1812.
17. Laitar DS, Müller P, Gray TG, Sadighi JP. *Organometallics*. 2005; 24:4503.

18. Tamaki A, Magennis SA, Kochi JK. J. Am. Chem. Soc. 1974; 96:6140.
19. Selected examples of reductive elimination with large and negative $\Delta S^\ddagger$: Packett DL, Trogler WC. Inorg. Chem. 1988; 27:1768. Brown MP, Puddephatt RJ, Upton CEE. J. Chem. Soc., Dalton Trans. 1974:2457.
20. Recent reviews: Engle KM, Mei T-S, Wang X, Yu J-Q. Angew. Chem., Int. Ed. 2011; 50:1478. Hopkinson MN, Gee AD, Gouverneur V. Chem.-Eur. J. 2011; 17:8248. [PubMed: 21678513]
21. (a) Brenzovich WE Jr, Benitez D, Lackner AD, Shunatona HP, Tkatchouk E, Goddard WA III, Toste FD. Angew. Chem., Int. Ed. 2010; 49:5519. (b) Brenzovich WE Jr, Brazeau J-F, Toste FD. Org. Lett. 2010; 12:4728. [PubMed: 21028911] (c) Melhado AD, Brenzovich WE Jr, Lackner AD, Toste FD. J. Am. Chem. Soc. 2010; 132:8885. [PubMed: 20557048]

Fig. 1. Extended solid-state structure of (IPr)AuF·2CH₂Cl₂, as 50% ellipsoids. Only solvent hydrogen atoms (in calculated positions) are shown. Selected bond lengths (Å) and angles (°): Au–F, 2.071(2); Au–C, 1.954(5); C–Au–F, 177.65(15).

Fig. 2.

Dependence of the rate of decay of *cis*-F₂Au(CH₂SiMe₃)(IPr) (**11-F₂**) on F⁻ concentration, as determined by the method of initial rates. $k_{\text{obs}} = (\Delta[\mathbf{11-F_2}]/\Delta t)/[\mathbf{11-F_2}]_0$, $[\mathbf{11-F_2}]_0 = 32.4$ mM, TBAF = tetrabutylammonium fluoride. The red line is a smoothed-curve guide.

Fig. 3. Calculated LUMO (BPV86/LANL2DZ/6-311G++**, implicit CH₂Cl₂ solvation, 0.04 isocontour) of [(IPr)Au(CH₂tBu)(F)]⁺ (**10-F⁺**).

Scheme 1.
Proposed mechanistic scenario.

Table 1Fluorination of selected alkylgold(i) complexes^a

$\text{IPr} \cdot \text{Au} \cdot \text{R} \xrightarrow[\text{- Xe}]{\text{XeF}_2} \text{IPr} \cdot \text{Au} \cdot \begin{array}{c} \text{F} \\ \\ \text{R} \end{array} \xrightarrow{\text{- IPr} \cdot \text{Au} \cdot \text{F}} \text{products}$ $\text{1 - 4} \qquad \qquad \qquad \text{1-F}_2 \text{ - 4-F}_2$		
Number	R	Product mixture ^b
1 ^c		None
2		+ 17% 66%
3 ^c		+ 11% 56%
4 ^c		+ 15% 33%

^a 30 mM (IPr)AuR, 1.1 equiv XeF₂, CDCl₃ solvent, room temp.^b Determined by ¹H and ¹⁹F NMR. Several NMR yields should be considered lower limits due to product volatility.^c *cis*-F₂Au(R)(IPr) intermediate observed by ¹⁹F NMR.

Fluorination of cycloalkylgold(I) complexes^a^c *cis*-F₂Au(R)(IPr) intermediate observed by ¹⁹F NMR.

Table 3Fluorination of alkylgold(i) complexes lacking β -hydrogens^a

$\text{IPr-Au-R} \xrightarrow[\text{- Xe}]{\text{XeF}_2} \text{IPr-Au-F} \xrightarrow[\text{- IPr-Au-F}]{\text{10-F}_2 \text{ - 13-F}_2} \text{products}$		
Number	R	Product mixture ^b
10 ^c		 12% 63% 16%
11 ^c		 11% 75%
12		 11% 73%
13		 85%

^a 30 mM (IPr)AuR, 1.1 equiv XeF₂, CDCl₃ solvent, room temp.^b Determined by ¹H and ¹⁹F NMR. Several NMR yields should be considered lower limits due to product volatility.^c *cis*-F₂Au(R)(IPr) intermediate observed by ¹⁹F NMR.

Table 4

Iodination of selected alkylgold(I) complexes^a

$\text{IPr-Au-R} \xrightarrow{\text{I}_2} \text{IPr-Au(R)I} \xrightarrow{-\text{IPr-Au-I}} \text{products}$	
R	Product mixture ^b
I-CH_3 quant. ^c
+ 55% 18%
+ + 60% 31% 5%
93%
84%
95%

^a30 mM (IPr)AuR, 1.1 equiv I₂, CDCl₃ solvent.^bDetermined by ¹H NMR. Several NMR yields should be considered lower limits due to product volatility.^cTaken from ref. 11.

Table 5Charge distribution of **10-X⁺** as a function of X^a

X	$q(\text{Au})^b$	$q(\text{X})^b$	$q(\text{CH}_2)^b$	$\Delta q(\text{Au}-\text{CH}_2)^c$
F	0.86	-0.62	0.11	0.75
Cl	0.69	-0.53	0.14	0.55
Br	0.61	-0.42	0.14	0.47
I	0.48	-0.29	0.14	0.34

^a Natural charge based on NBO analysis (BPV86/LANL2DZ/6-311G++**, implicit CH₂Cl₂ solvation).^b q = charge.^c Δq = difference in charge between Au and CH₂.