Published in final edited form as:

Headache. 2012 October ; 52(Suppl 2): 102–106. doi:10.1111/j.1526-4610.2012.02241.x.

Migraine: Maladaptive Brain Responses to Stress

Nasim Maleki, Lino Becerra, and David Borsook

Center for Pain and the Brain, Children's Hospital Boston, Harvard Medical School

Abstract

Migraine offers a unique model to understand the consequences of repeated stressors on the brain. Repeated stressors can alter the normal response of physiological systems and this concept has been termed 'allostatic load'. In the case of the brain, the effects of repeated stress may lead to alteration in brain networks both functionally and structurally. As a result the brain responds abnormally to environmental conditions (psychological or physiological). Here we present an alternative perspective on migraine disease and propose that changes in brain states may occur as a result of repeated migraine attacks through maladaptive coping mechanisms. The cascade of these effects can lead to further deterioration of adaptation and thus lead to transformation or chronification of the disease.

Introduction

Migraine is a disabling primary headache disorder affecting millions of individuals, by some estimates 45 million Americans¹. As a condition it is complex and may be considered in a continuum not only in terms of the peri-headache changes in episodic migraine but also the progression to high frequency and chronic daily headache. Thus it is a major perturber of the brain systems in a number of ways. These include behavioral and/or physiologic responses such as pain², cardiovascular changes³, immunological changes⁴.

The responses to a migraine attack (e.g, pain, stress hormones, affective changes, nausea/vomiting etc) are mediated through increased activity of mediators that are in excess of normal adaptive processes^{5,6}. Over time these stressors could lead to an altered brain state characterized by increased cortical excitability⁷, changes in brain morphology and changes in behavior^{8–11}. Such stressors are typically repeated periodically in both episodic migraine and chronic daily headache. As a result the brain will respond abnormally to environmental conditions (psychological or physiological). If viewed in this context, pathophysiological changes in brain structure and function in migraine may contribute to an improved understanding of the condition that frequently affects individuals for a significant portion (>15 years¹²) of their lives.

Stress and Migraine

Stress as a trigger for migraine attacks is present in nearly 70% of individuals¹³. High levels of stress are reported in migraine patients, particularly in those suffering from chronic daily migraine¹⁴. Both endogenous (e.g., hormone) and exogenous (e.g., physical stressors (e.g., light) or psychological stressors) stressors add to the burden of the disease. Emotional or physical trauma (e.g., abuse, particularly in childhood), socioeconomic or social stress¹⁵ are examples of psychological stressors. Stressors could also be of physiological origin such as menstrual period in women. Some women experience menstrual migraines or their

migraines around menstrual period could be more intense or even last longer. In addition, migraine is comorbid with numerous brain diseases that are associated with stress and a potential contributing factor (e.g., anxiety, depression¹⁶). Even in the pediatric population, stress plays an important role. Children with migraine have increased pulse rate, higher diastolic blood pressure and higher low-frequency/high-frequency ratio after a 5-min recovery from an emotional stressor¹⁷. When the frequency or severity of these stressors escalate, the protective adaptive responses (allostatic responses) that maintain a stable state for the brain become overused and dysregulated. As a result there is wear and tear on the brain (“allostatic load”) which may alter brain networks both functionally and structurally. As a result the brain responds abnormally to stressors and reaches an altered state, in which both behavior and systemic physiology are altered in ways that can lead to further allostatic load, Figure 1.

Effects of Repeated Migraines on the Brain

Are there any biological consequences in the brain as a result of migraine attacks? While there is no clear answer to this question yet, there are multiple lines of evidence that suggest that the migraine brain is different compared to normal brain^{18–21}. The migraine brain is hyperexcitable during the migraine-free periods between the attacks (interictal phase)^{22–24,25}. During the attacks the brain’s response to some stimuli is abnormal. For example, in many patients light, smell or noise could exacerbate the migraine pain when they are experiencing an attack. Hyperexcitability of the brain⁷ could be due to a decrease in inhibitory neurotransmitters (e.g., GABA), or an increase in excitatory neurotransmitters (e.g., glutamate) in the cortical regions. Impaired or diminished habituation to stimuli in the interictal period in migraineurs^{26, 27} is another characteristic of migraine, which also supports maladaptive brain changes. Increasingly, evidence suggests abnormal inhibitory control of pain with migraine that may result from deficiencies in inhibition or increased facilitation²⁸. A variety of stressful stimuli have been shown to affect pain sensitivity suggesting altered higher centers have a significant influence on pain modulation. Neural substrates mediating active versus passive emotional coping have been identified within distinct, longitudinal neuronal columns of the midbrain periaqueductal gray (PAG) region²⁹. In addition to pain modulation³⁰, the PAG has been implicated in responses to defensive and autonomic regulation³¹.

The basal ganglia have also been implicated in migraine. In general, the basal ganglia play an important role in pain processing (reviewed in³²). Migraineurs with more advanced forms of the disease, reduced response to noxious thermal stimulation in the caudate, putamen and pallidum associated with larger volume in caudate⁸. These may be explained by ongoing inflammation, enhanced iron accumulation or increased dendritic complexity. Hypometabolism in the caudate has been reported for familial hemiplegic migraine³³.

Decreased grey matter in anterior cingulate cortex^{9, 10, 34} and insula^{9, 10} in migraine patients has been reported. These changes correlate with the frequency of migraine attacks or duration of the disease^{9, 10}. Decreased cortical gray matter volume in the cingulate cortex is reported in patients with depression³⁵ too. Interestingly, migraine is also associated with an increased rate of affective illness including depression³⁶ with patients who have higher frequency headaches as in daily chronic headache³⁷ at an increasing higher risk of depression.

Impairment of stress response adaptation has been associated with inactivation of the medial prefrontal cortex activity in rats³⁸. Evidence of altered function in this area has been supported by fMRI studies of migraine patients stimulated with a painful heat stimulus showing increased activation in the perigenual cortex vs. healthy controls³⁹. A connection

between stress and medial prefrontal cortical regions has also been reported in resting state functional connectivity studies of adolescents undergoing social stress: cortisol responsivity was shown to be correlated with “salience” network that includes the anterior insula and medial prefrontal cortex ⁴⁰. The region is involved in processing negative emotion including anxiety.

Conclusion

Migraine is considered a brain disease with widespread alterations in a number of distributed functions that include both cortical and subcortical systems. Multiple brain changes are reflective of alterations in the migraine brain that may be a result of the repetitive nature of the disease state, a predisposition in these patients, genetic factors or a combination of all of these. There is evidence for co-localized morphometric and functional cortical differences as a function of the number of to repeated migraine attacks ¹¹ specifically in the response patterns in the sensory vs. affective processing regions in the brain. While response to acute stressors is mostly protective but chronic stressors do produce damage ⁴¹ and thus a model disease of allostatic load ⁴².

Acknowledgments

The work was supported in by grants from NIH (K24 NS064050 (NINDS), R01-NS056195 (NINDS) and R01-NS073997 to DB.

References

1. Stewart WF, Shechter A, Rasmussen BK. Migraine prevalence. A review of population-based studies. *Neurology*. 1994; 44:S17–23. [PubMed: 8008222]
2. Kelman L. Pain characteristics of the acute migraine attack. *Headache*. 2006; 46:942–953. [PubMed: 16732840]
3. Melek IM, Seyfeli E, Duru M, Duman T, Akgul F, Yalcin F. Autonomic dysfunction and cardiac repolarization abnormalities in patients with migraine attacks. *Med Sci Monit*. 2007; 13:RA47–49. [PubMed: 17325646]
4. Pradalier A, Launay JM. Immunological aspects of migraine. *Biomed Pharmacother*. 1996; 50:64–70. [PubMed: 8761711]
5. McEwen BS. Protective and damaging effects of stress mediators: central role of the brain. *Dialogues Clin Neurosci*. 2006; 8:367–381. [PubMed: 17290796]
6. McEwen BS. Physiology and neurobiology of stress and adaptation: central role of the brain. *Physiol Rev*. 2007; 87:873–904. [PubMed: 17615391]
7. Chen WT, Wang SJ, Fuh JL, Lin CP, Ko YC, Lin YY. Persistent ictal-like visual cortical excitability in chronic migraine. *Pain*. 2011; 152:254–258. [PubMed: 21145169]
8. Maleki, NBL.; Nutile, L.; Burstein, R.; Borsook, D. Structural differences in migraine brain are related to the attack frequency. Society for Neuroscience; San Diego: 2010a.
9. Kim JH, Suh SI, Seol HY, et al. Regional grey matter changes in patients with migraine: a voxel-based morphometry study. *Cephalalgia*. 2008; 28:598–604. [PubMed: 18422725]
10. Valfre W, Rainero I, Bergui M, Pinessi L. Voxel-based morphometry reveals gray matter abnormalities in migraine. *Headache*. 2008; 48:109–117. [PubMed: 18184293]
11. Maleki N, Becerra L, Brawn J, Bigal M, Burstein R, Borsook D. Concurrent Functional and Structural Cortical Alterations in Migraine. *Cephalalgia: an international journal of headache*. 2012 In Press.
12. Kelman L. Migraine changes with age: IMPACT on migraine classification. *Headache*. 2006; 46:1161–1171. [PubMed: 16866720]
13. Theeler BJ, Kenney K, Prokhorenko OA, Fideli US, Campbell W, Erickson JC. Headache triggers in the US military. *Headache*. 2009; 50:790–794. [PubMed: 19925623]

14. Dodick DW. Review of comorbidities and risk factors for the development of migraine complications (infarct and chronic migraine). *Cephalalgia*. 2009; 29 (Suppl 3):7–14. [PubMed: 20017749]
15. Chyu L, Upchurch DM. Racial and ethnic patterns of allostatic load among adult women in the United States: findings from the National Health and Nutrition Examination Survey 1999–2004. *J Womens Health (Larchmt)*. 2011; 20:575–583. [PubMed: 21428732]
16. Amouroux R, Rousseau-Salvador C. Anxiety and depression in children and adolescents with migraine: a review of the literature. *Encephale*. 2008; 34:504–510. [PubMed: 19068340]
17. Huss D, Derefinko K, Milich R, Farzam F, Baumann R. Examining the stress response and recovery among children with migraine. *J Pediatr Psychol*. 2009; 34:707–715. [PubMed: 18922826]
18. Kruit MC, van Buchem MA, Hofman PA, et al. Migraine as a risk factor for subclinical brain lesions. *JAMA*. 2004; 291:427–434. [PubMed: 14747499]
19. Kruit MC, van Buchem MA, Launer LJ, Terwindt GM, Ferrari MD. Migraine is associated with an increased risk of deep white matter lesions, subclinical posterior circulation infarcts and brain iron accumulation: the population-based MRI CAMERA study. *Cephalalgia*. 2010; 30:129–136. [PubMed: 19515125]
20. Welch KM, Nagesh V, Aurora SK, Gelman N. Periaqueductal gray matter dysfunction in migraine: cause or the burden of illness? *Headache*. 2001; 41:629–637. [PubMed: 11554950]
21. Schmitz N, Admiraal-Behloul F, Arkink EB, et al. Attack frequency and disease duration as indicators for brain damage in migraine. *Headache*. 2008; 48:1044–1055. [PubMed: 18479421]
22. Ozkul Y, Uckardes A. Median nerve somatosensory evoked potentials in migraine. *Eur J Neurol*. 2002; 9:227–232. [PubMed: 11985630]
23. Schoenen J. Neurophysiological features of the migrainous brain. *Neurol Sci*. 2006; 27 (Suppl 2):S77–81. [PubMed: 16688634]
24. Afra J. Intensity dependence of auditory evoked cortical potentials in migraine. Changes in the peri-ictal period. *Funct Neurol*. 2005; 20:199–200. [PubMed: 16483461]
25. Bussone G. Pathophysiology of migraine. *Neurol Sci*. 2004; 25 (Suppl 3):S239–241. [PubMed: 15549546]
26. Coppola G, Pierelli F, Schoenen J. Habituation and migraine. *Neurobiol Learn Mem*. 2009; 92:249–259. [PubMed: 18675928]
27. Perrotta A, Serrao M, Sandrini G, et al. Reduced habituation of trigeminal reflexes in patients with episodic cluster headache during cluster period. *Cephalalgia*. 2008; 28:950–959. [PubMed: 18624808]
28. Vanegas H, Schaible HG. Descending control of persistent pain: inhibitory or facilitatory? *Brain Res Brain Res Rev*. 2004; 46:295–309. [PubMed: 15571771]
29. Keay KA, Bandler R. Parallel circuits mediating distinct emotional coping reactions to different types of stress. *Neurosci Biobehav Rev*. 2001; 25:669–678. [PubMed: 11801292]
30. Heinricher MM, Tavares I, Leith JL, Lumb BM. Descending control of nociception: Specificity, recruitment and plasticity. *Brain Res Rev*. 2009; 60:214–225. [PubMed: 19146877]
31. Lovick TA. The periaqueductal gray-rostral medulla connection in the defence reaction: efferent pathways and descending control mechanisms. *Behav Brain Res*. 1993; 58:19–25. [PubMed: 8136045]
32. Borsook D, Upadhyay J, Chudler EH, Becerra L. A key role of the basal ganglia in pain and analgesia--insights gained through human functional imaging. *Mol Pain*. 6:27. [PubMed: 20465845]
33. Gutschalk A, Kollmar R, Mohr A, et al. Multimodal functional imaging of prolonged neurological deficits in a patient suffering from familial hemiplegic migraine. *Neurosci Lett*. 2002; 332:115–118. [PubMed: 12384224]
34. Schmidt-Wilcke T, Ganssbauer S, Neuner T, Bogdahn U, May A. Subtle grey matter changes between migraine patients and healthy controls. *Cephalalgia*. 2008; 28:1–4. [PubMed: 17986275]
35. Salvatore G, Nugent AC, Lemaitre H, et al. Prefrontal cortical abnormalities in currently depressed versus currently remitted patients with major depressive disorder. *Neuroimage*. 2011; 54:2643–2651. [PubMed: 21073959]

36. Frediani F, Villani V. Migraine and depression. *Neurol Sci.* 2007; 28 (Suppl 2):S161–165. [PubMed: 17508165]
37. Antonaci F, Nappi G, Galli F, Manzoni GC, Calabresi P, Costa A. Migraine and psychiatric comorbidity: a review of clinical findings. *J Headache Pain.* 2011; 12:115–125. [PubMed: 21210177]
38. Weinberg MS, Johnson DC, Bhatt AP, Spencer RL. Medial prefrontal cortex activity can disrupt the expression of stress response habituation. *Neuroscience.* 2010; 168:744–756. [PubMed: 20394807]
39. Tessitore A, Russo A, Esposito F, et al. Interictal cortical reorganization in episodic migraine without aura: an event-related fMRI study during parametric trigeminal nociceptive stimulation. *Neurol Sci.* 2011; 32 (Suppl 1):S165–167. [PubMed: 21533737]
40. Thomason ME, Hamilton JP, Gotlib IH. Stress-induced activation of the HPA axis predicts connectivity between subgenual cingulate and salience network during rest in adolescents. *J Child Psychol Psychiatry.* 2011
41. McEwen BS, Seeman T. Protective and damaging effects of mediators of stress. Elaborating and testing the concepts of allostasis and allostatic load. *Ann N Y Acad Sci.* 1999; 896:30–47. [PubMed: 10681886]
42. Borsook D, Maleki N, Becerra L, McEwen B. Understanding migraine through the lens of maladaptive stress responses: a model disease of allostatic load. *Neuron.* 2012; 73:219–234. [PubMed: 22284178]

Figure 1.

Brain in the normal state responds adaptively (adaptive allostasis) to stressors ranging from physiologic to genetic in order to maintain the same brain state. In migraine the effect of stressors may be additive or cumulative and the brain's response to the same stressors becomes maladaptive and thus changes to the brain may happen changing state of the brain to a new state.