Published in final edited form as:

Neuroscience. 2012 October 25; 223: 447–456. doi:10.1016/j.neuroscience.2012.07.040.

Sex Differences in U50,488H-induced Phosphorylation of p44/42 MAP kinase in the Guinea Pig Brain

Khampaseuth Rasakham², Katherine L. McGillivray, and Lee-Yuan Liu-Chen¹

Department of Pharmacology and Center for Substance Abuse Research, Temple University School of Medicine, Philadelphia, PA

Abstract

Recently there has been a widespread interest in the development of kappa opioid receptor (KOPR) ligands for treatment of pain, depression and anxiety, and prevention of stress-induced drug relapse. However, most of these preclinical studies have been conducted using male experimental animals. In the present study we examined if sex differences exist in neural activity induced by the KOPR agonist U50,488H. Here, we used immunohistochemistry to detect activation (phosphorylation) of p44/42 MAPK as an indicator of neural activity. Following habituation to injection for 3 days, adult guinea pigs received a single injection of U50,488H (5 mg/kg, s.c.) and perfused thirty to forty-five minutes later. U50,488H-induced an increase in the number of cells immuno-positive for phosphorylated p44/42 MAPK in subregions of the amygdala, thalamus, paraventricular nucleus of the hypothalamus, periaqueductal grey, and dorsal raphe nuclei. In contrast, U50,488H-induced a decrease in immuno-positive cells in the ventrolateral and lateral orbital cortex. Pretreatment with the KOPR antagonist norBNI (10 mg/kg, i.p.) 18 hrs prior to U50,488H significantly reversed the effects of U50,488H in most regions. In addition, we observed a notable sex difference in the basolateral amygdala; in males, U50,488H-induced an increase in immuno-positive cell numbers but a decrease in females. However, across other brain regions males were generally more sensitive to U50,488H-induced alterations than females. These results suggest the need to include female subjects in studies examining emotional responses to KOPR ligands.

Keywords

Kappa opioid receptor; sex difference; guinea pig; p44/42 MAP kinase; neural activity

The kappa opioid receptor is one of three major types of opioid receptors, which include μ , δ and κ or MOPR, DORP, and KOPR, respectively. Activation of the KOPR produces many effects including analgesia, dysphoria, water diuresis, antipruritis and attenuates cocaine craving in addicts [reviewed in (Liu-Chen, 2004)]. KOPRs are localized throughout the rostral-caudal axis of the mammalian brain. Radioligand binding studies in rodents reveal that KOPR is enriched in the claustrum, hypothalamus, endopiriform nucleus, caudate

© 2012 IBRO. Published by Elsevier Ltd. All rights reserved.

¹Corresponding Author: Lee-Yuan Liu-Chen, Ph.D., Professor, Department of Pharmacology, Center for Substance Abuse Research, Temple University School of Medicine, 3420 N. Broad Street, Philadelphia, PA 19140, phone: (215) 707-4188; fax: (215) 707-7068.

²Present address: Department of Radiology, Athinoula A. Martinos Center for Biomedical Imaging, Massachusetts General Hospital 149 13th St, Charlestown, MA 02129

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

putamen, nucleus accumbens and neocortex (Quirion et al., 1987; Mansour et al., 1988; Quirion and Pilapil, 1991; Wang et al., 2011). Of interest, studies indicate that KOPR distribution in the human brain is more similar to the guinea pig brain than to the rat or mouse brain. For example, in guinea pig and human brains, but not the rat or mouse brain, KOPR is abundant in the cerebellum, in deep layers (layer V and VI) of the cortex and is found in striosomes (having patchy distribution) within the striatum (Quirion et al., 1987; Mansour et al., 1988; Quirion and Pilapil, 1991; Wang et al., 2011). Therefore, we consider the guinea pig to be a more appropriate small animal model for KOPR studies.

Early studies by Levine and colleagues (Gear et al., 1996b; Gear et al., 1996a) found that in patients undergoing molar teeth extraction partial KOPR agonists, such as pentazocine, butorphanol and nalbuphine, produced greater analgesic efficacy in women than men. In contrast, studies in rodents and non-human primates have revealed that selective full KOPR agonists produced greater antinociceptive effects in males than females [(Rasakham and Liu-Chen, 2011; Craft, 2003) and references therein].

Recently, we demonstrated endpoint-dependent sex differences in behavioral pharmacology of KOPR in guinea pigs (Wang et al., 2011). Males showed greater analgesia/antinociception and abnormal postural changes in response to the KOPR agonist U50,488H than females. In contrast, U50,488H inhibited cocaine-induced hyperactivity to greater extents in females than in males. Moreover, anatomical KOPR distribution and receptor-G protein coupling supported the behavioral differences observed between the sexes (Wang et al., 2011). [³H]U69,593, binding studies showed that males had higher levels of KOPR in the periaqueductal grey (PAG), a critical brain region involved in pain. In addition, males showed higher KOPR levels and U50,488H-stimulated [³⁵S]GTPγS binding in several areas related to pain, sedation, and motor control.

Although KOPR distribution and *in vitro* receptor functionality provided support for an anatomical basis for the observed behavioral differences, the downstream effect of KOPR agonist on neural activity has not been examined.

In the present study we aimed to investigate potential sex differences in neural activity induced by the KOPR agonist U50,488H. We use phosphorylation of p44/42 MAPK (also known as extracellular regulated kinases 1 and 2, ERK1/2) as an indicator of neural activity *in vivo*. Activation of p44/42 MAP kinase via phosphorylation (P-p44/42 MAPK) has been used as a marker of altered neural activity since the administration of many drugs, including morphine, psychostimulants, and antipsychotic drugs, produce distinct and specific patterns of changes in P-p44/42 MAPK in select brain regions [for example, see (Eitan et al., 2003; Valjent et al., 2004)]. In addition, p44/42 MAPK activity has been shown to underlie many forms of synaptic plasticity, such as learning and memory (Roberson et al., 1999), drug addiction (Valjent et al., 2004) and mu opioid analgesic tolerance (Eitan et al., 2003). It is important to note that enhancement in P-p44/42 MAPK in brain regions after U50,488H is distinctly different from increased P-p44/42 MAPK in cells expressing the KOPR *in vitro*, which typically has an initial peak at 5–10 min (Li et al., 1999; Bohn et al., 2000).

In the present study we use P-p44/42 MAPK as an indicator of neuronal activity, similar to the use of c-Fos expression (Chang et al., 1988; Liu et al., 1994; Bot and Chahl, 1996). Importantly, P-p44/42 MAPK is an ideal target for our studies since, unlike c-Fos, P-p44/42 MAPK activity occurs rapidly and is upstream to the induction of c-Fos expression. Induction of c-Fos expression commonly occurs 2–3 h after drug administration and is more related to long-term adaptive changes.

Experimental Procedures

Animals

Age-matched male and female Dunkin-Hartley guinea pigs (450–600 g) were used for these studies. Animals were housed at the Temple University School of Medicine animal facility and used in compliance with IACUC regulations. Guinea pigs were housed in same sex groups of 3 per cage (54 × 38 × 28 cm). Animals were allowed to acclimate to animal facilities for at least 1 week prior to any handling and experiments. Animals were placed into a standard rat cage (40 × 20 × 20 cm), 2 (same sex) per cage, for transport to and from the animal facility on a covered animal transport cart. Animals were allowed to acclimate to the laboratory for at least 90 min prior to handling and remained in the laboratory for at least 1 hour prior to returning to the central animal facilities.

Chemicals

The KORP agonist U50,488H and KOPR antagonist norbinaltorphimine (norBNI) was kindly provided by the National Institute on Drug Abuse (NIDA) (Bethesda, MD). Sodium pentobarbital was purchased from Ovation Pharmaceuticals, Inc (Deerfield, IL) through the Central Animal Facilities at Temple University. Rabbit phosphorylation specific p44/42 (Thr202/Tyr204) MAP kinase antibody was from Cell Signaling (Beverly, MA). Phosphorylation specific p44/42 MAP kinase recognizes only the phosphorylated (activated) form of p44/42 MAPK. Biotinylated goat anti-rabbit IgG, and avidin biotin-peroxidase complex (ABC) were from Vector Labs (Burlingham, CA). 3,3'-Diaminobenzidine (DAB) was from Sigma Co. (St. Louis, MO). All other reagent-grade chemicals were obtained from Sigma Chemical Co (St Louis, MO) or Fisher Scientific (Pittsburgh, PA).

Drug treatment and perfusion

Guinea pigs were habituated to daily subcutaneous (s.c.) injections of saline (1 ml/kg) for 3 days (2x per day, between 9:00–12:00 and 16:00–18:00). On the day of perfusion, animals were injected with U50,488H at 5 mg/kg in a volume of 1 ml/kg or 0.9 % saline (1 ml/kg). We examined the effect at the dose of 5 mg/kg as this dose produced strong behavioral effects in guinea pigs (Wang et al., 2011). A separate group of guinea pigs were treated with norBNI approximately 18 hrs prior to U50,488H or saline treatment. Twenty seven minutes following drug administration, guinea pigs were anesthetized with sodium pentobarbital [150 mg/kg, intraperitoneally, (i.p.)] then transcardially perfused approximately 30–45 minutes after U50,488H injection with 800 ml of 4% paraformaldehyde (PFA) in 0.1M phosphate buffer (PB) (pH 7.4). We chose this time point given that peak pharmacological effects at this dose were observed previously in the guinea pig (Wang et al., 2011). Brains were removed and post-fixed in 4% PFA in 0.1M PB overnight at 4°C and then incubated at 4°C in 30% sucrose for 2 days for cryoprotection.

Immunohistochemistry

Coronal sections of the tissue were cut at 50 µm. Free-floating sections were processed by immunohistochemistry immediately after sectioning. Sections were washed 3 × 5 minutes in 0.01M PBS. Sections were then treated for 30 minutes with a solution of 10% methanol and 0.3% H₂O₂ in 0.01M PBS to inactivate endogenous peroxidases followed by washing in 0.01M PBS 3 × 5 minutes and blocking with 5% normal goat serum (NGS)/1% bovine serum albumin (BSA)/0.2% Triton X-100 solution for 1 hour at room temperature. Sections were then incubated with rabbit phospho-specific 44/42 primary antibody (1:2000 in 1% NGS/1%BSA/0.2% Triton X-100) overnight on a shaker at 4°C. The sections were washed 3 × 5 minutes followed by incubation in secondary antibody, biotinylated goat anti-rabbit IgG (1:100) in 1%NGS/1%BSA/0.2% Triton X-100 for 90 minutes. Sections were washed and

then incubated in ABC (1:100) in 0.01M PBS for 1 hour. Sections were then washed and stained using DAB (0.02%) and 0.0072% H₂O₂ prepared in distilled water. Following staining, tissue sections were rinsed once in distilled water for 5 minutes and then washed 3 × 5 minutes in 0.01M phosphate buffer (PB). The sections were mounted on subbed slides with 0.01M PB and air-dried completely. The sections were dehydrated 1 minute each through 70%, 80%, 95%, and 100% ethanol, followed by de-fatting in 2 changes of xylene 5 minutes each. Slides were immediately cover-slipped using DPX mounting media (Sigma, St. Louis, MO).

Image and data analysis

All slides were examined with a Nikon TE-300 inverted light microscope at the same light intensity. Photos of each section were taken with a Nikon Optronics camera using the software MagnaFire with a consistent exposure time for concurrently processed sections. The images were analyzed with NIH ImageJ 1.42. Specifically, a threshold level was chosen from a 256-level gray scale. The threshold was selected to allow detection of nuclei, cells and fibers stained with moderate to high intensity, while suppressing lightly stained nuclei, cells and fibers. A region of interest box (ROI) was defined for each select brain region and used to sample from each region. Immuno-positive phospho-p44/42 MAP kinase stained cells were manually counted from several sections ranging from 1–4 sections, depending on the brain region. For each animal a single average value was calculated from bilateral readings from the sections. More specifically, the number of sections quantified for each region from each animal ranged as follows: cingulate cortex, 1–2 sections; infralimbic cortex, 2–4; dorsal pendicular cortex, 2–4; ventrolateral orbital cortex, 2–4; lateral orbital cortex, 2–4; claustrum, 2–4; nucleus accumbens, 3–4; lateral septum, 3–4; paraventricular nucleus, 2–4; ventromedial hypothalamus, 3–4; central amygdala, 1–3; basolateral amygdala, 1–3; medial amygdala, 2–4; thalamus, 2–4; periaqueductal grey, 3–4; and dorsal raphe nuclei, 2–4. Comparison of drug treatment effects and sex group effects were examined by two-way analysis of variance (ANOVA), where $p < 0.05$ was deemed significant. Significant effects were further analyzed by Tukey *post hoc* tests. Blockade of U50,488H by norBNI pre-treatment and nor-BNI treatment alone effects were analyzed using Student's *t* test ($p < 0.05$).

Results

Analysis of P-p44/42 MAPK immuno-positive cells and fibers were assessed in select ROIs throughout the rostral to caudal parts of the guinea pig brain as shown in Figure 1. We present the results in three anatomical subdivisions: telencephalon (cortex, claustrum and striatum), diencephalon (thalamus, hypothalamus, amygdala structures) and midbrain (periaqueductal grey region & dorsal raphe nuclei). Table 1 depicts the quantitative analysis.

Telencephalon

Representative images of telencephalon regions (lateral orbital cortex, claustrum, and nucleus accumbens) are shown in figure 2.

Males vs Females—In the infralimbic cortex, ventrolateral orbital cortex (VLO) and lateral orbital cortex (LO) U50,488H treatment significantly reduced the number of immuno-positive cells ($p < 0.05$). However, there were no main effects of sex in these areas ($p = 0.81, 0.92$ and 0.20 , respectively). Pretreatment with nor-BNI did not reverse the effects of U50,488H in any of these regions, as nBNI+U50,488H was not significantly different from U50,488H only ($p > 0.05$) (Table 1). Notably, pretreatment with nor-BNI prior to saline (nBNI+Saline) was not significantly different than saline treatment ($p = 0.20, 0.31$ and 0.53 in infralimbic cortex, VLO and LO, respectively).

In the cingulate cortex, dorsal pedicular cortex, nucleus accumbens, claustrum, dorsal lateral septum, and ventral lateral septum no significant differences between sex or drug treatment were found. However, we did observe a trend in drug treatment effects in a few of these areas, such as the cingulate cortex and nucleus accumbens ($p = 0.08$ and $p = 0.07$, respectively). Interestingly, compared to U50,488H alone, norBNI pretreatment reduced the number of U50,488H-induced immuno-positive cells in the dorsal pedicular cortex. However, this was not statistically significant ($p = 0.06$). Treatment with norBNI prior to saline was not statistically different from saline in any of these regions.

Diencephalon

Representative images of diencephalon regions, the paraventricular nucleus (PVN), central amygdala (CeA), medial amygdala (MeA), basolateral amygdala (BLA) and thalamus), are shown in figure 3.

Males vs Females—U50,488H induced a significant increase in immuno-positive cells in the CeA, MeA, and THA ($p < 0.05$). There was no significant main effect of sex in any of these regions ($p = 0.21, 0.63$, and 0.67 , respectively). Notably, nor-BNI pretreatment blocked the effects of U50,488H in each of these regions ($p < 0.05$).

In the BLA U50,488H increased the number of immuno-positive cells in males while in females U50,488H caused a reduction. Moreover, females had a significantly greater number of immuno-positive cells than males in the BLA. Pretreatment with norBNI produced a trend of reduction from the effects of U50,488H alone ($p = 0.06$, Table 1). However, norBNI treatment prior to saline induced a significant increase in immuno-positive cells compared to saline ($p = 0.02$).

In the PVN U50,488H increased the number of immuno-positive cells in males but not females ($p < 0.05$) and this effect was blocked by nor-BNI pretreatment. Saline-treated females showed a greater number of immuno-positive cells than saline-treated males, but it did not reach statistical significance. There were no significant effects in the ventromedial hypothalamus (VMH).

Midbrain

We analyzed immuno-labeling in the periaqueductal grey (PAG) region and dorsal raphe nuclei (DRN) as shown in figure 4. We took cell counts of the total periaqueductal grey region as well as the lateral subdivision.

Males vs Females—U50,488H increased the number of immuno-positive cells in the PAG and DRN. A main effect of sex was found in the DRN with females having more immuno-positive cells than males but no sex difference was found in the PAG. However, while nor-BNI treatment reduced the effects of U50,488H it did not reach statistical significance in the PAG, lateral PAG and DRN ($p = 0.35, 0.14$ and 0.06 , respectively; see Fig 4 and Table 1).

Discussion

In the present study we found that U50,488H induced an increase in phosphorylation of p44/42 MAPK in brain areas that regulate pain, diuresis, mood, stress and substance addiction. In contrast, in regions involved in cognition we observed that U50,488H-induced a decrease in p44/42 MAPK activity. U50,488H generally induces a similar pattern of P-p44/42 MAPK activity in both sexes; however, in some regions males show greater responses than females. Moreover, in most brain regions the effects of U50,488H were

reversed by norBNI in male guinea pigs. It is likely that norBNI antagonism of U50,488H will be similar in both sexes as reported previously (Craft et al., 2000; Craft and McNiel, 2003). Altogether, these results correlate with the sex difference in level and G-protein coupling of KOPR we found previously and provide a neuroanatomical mechanism to explain our previous behavioral observations.

P-p44/42 as a measure of neuronal activity

P-p44/42 was used as a measure of neuronal activity since its activation is known to regulate a host of cellular processes (Pearson et al., 2001). The basal tone of P-p44/42 may be higher than the more commonly used marker of neural activity, cFos, due to its activation by endogenous neurotrophic factors, hormones and neurotransmitters; therefore, we can assess both reductions and enhancement of activity induced by KOPR activation. Since p44/42 MAPK activation occurs rapidly and is upstream of long-term adaptive processes, it allows us to detect immediate changes of neural (cellular) activation at a time point when peak behavioral effects of a KOPR agonist occur. Our results on U50,488H-caused elevations in P-p44/42 MAPK in male guinea pigs are generally consistent with previous findings on U50,488H-induced increase in cFos expression in male guinea pigs (Bot and Chahl, 1996), indicating that P-p44/42 MAPK is a valid marker for neural activity.

While activation of KOPR modulates P-p44/42 MAPK, the changes may not occur in cells expressing the KOPR. Rather, P-p44/42 MAPK may be modulated in neurons innervated by those expressing KOPR. For example, as KOPR activation results in inhibition of neuronal activity its activation on a GABAergic cell would lead to a disinhibition of downstream target cells (Li et al., 2012). Indeed, previous studies have found that alterations in P-p44/42 MAPK induced by morphine occurred primarily in cells that did not express the MOPR (Eitan et al., 2003). In the present study we did not conduct co-localization experiments because of the limited availability of antibodies that can specifically detect the guinea pig KOPR. In our previous study we mapped KOPR localization and receptor-G-protein coupling in the guinea pig using [³H]U69,593 and U50,488H-induced GTP γ S binding (Wang et al., 2011). At the level of brain regions, in some areas P-p44/42 MAPK matches KOPR localization and receptor-G-protein coupling, e.g. thalamus and hypothalamus, while in some regions with high levels of KOPR and receptor-G-protein coupling we did not detect any changes in U50,488H-induced P-p44/42 MAPK, e.g. septal and deep cortical layers of cortex. Alternative explanations for the lack of correlation between P-p44/42 MAPK and receptor localization and G-protein coupling (Wang et al., 2011) include competition between several intracellular signaling pathways, G protein coupling may not signal to p44/42 MAPK in these regions, or different temporal activity in these regions.

KOPR and Antinociception

Activation of KOPR in PAG and DRN contributes to antinociception (Chen et al., 2007; Kanarek et al., 2001; Land et al., 2009). The U50,488H induced increase in neural activity in the PAG and DRN we observed is consistent with previous reports measuring cFos induction by U50,488H in male guinea pigs (Bot and Chahl, 1996) and metabolic imaging studies in rats after treatment with the KOPR agonist salvinorin A (Hooker et al., 2009). In the present study females generally had higher levels of P-p44/42 MAPK than males in the DRN, while in the lateral and entire PAG this difference did not reach statistical significance ($p=0.31$ and 0.17 , respectively). This may mask the ability to detect a drug treatment effect. Normalizing against the number of activated cells found in the saline groups, we found U50,488H-induced an average 7.6- and 7.3-fold increase in the lateral PAG and DRN, respectively of males, while in females a 4.3- and 1.9-fold increase was observed; this suggests lower effects of U50,488H in females. These results may in part explain the greater antinociceptive effects of U50,488H in male than female guinea pigs (Wang et al., 2011).

KOPR, emotional states and drug addiction

KOPR agonists acutely oppose reinforcing, locomotor stimulating and locomotor sensitizing effects of drugs of abuse in males [for a review see (Shippenberg et al., 2007)]. While U50,488H-induced a trend in increase in neural activity ($p=0.07$) in the nucleus accumbens, a key region involved in psychostimulant-induced hyperactivity (Koob, 1992; Sharp et al., 1987), we did not find a sex difference in this increase ($p=0.11$ for sex x treatment interaction). We also did not detect any difference in KOPR levels or KOPR-mediated G protein activation (Wang et al., 2011). Together, this suggests that sex differences in other regions, e.g. ventral tegmentum area, that project into the nucleus accumbens or its efferent projections, e.g. caudate putamen, may be involved in the sex difference in U50,488H-induced inhibition of cocaine hyperactivity (Wang et al., 2011).

Activation of KOPR produces dysphoria in humans and aversion, stress- and depressive-like behaviors in animals (Pfeiffer et al., 1986; McLaughlin et al., 2003; McLaughlin et al., 2006), while blockade of KOPR or deletion of KOPR produces anxiolytic- and antidepressant-like effects (Knoll et al., 2007; Carr and Lucki, 2010; McLaughlin et al., 2003; McLaughlin et al., 2006). KOPR activation in the nucleus accumbens, amygdala, and DRN are implicated in depressive- and anxiety-like behaviors (Carlezon, Jr. et al., 2006; Knoll et al., 2011; Bruchas et al., 2009). However, studies to date have been conducted only in male animals.

The U50,488H-induced neural activation in the central and medial amygdala subregions in both sexes is consistent with our findings on KOPR levels and KOPR-mediated G protein activation in the whole amygdala (Wang et al., 2011). However, in the BLA females had greater basal activity than males and U50,488H induced an increase in neural activity in males, but a reduction in females. This sexual dimorphism may be attributed to differences in organization of other neurotransmitter systems modulating p44/42 MAPK activity in the BLA. Alternatively, there may be sexual dimorphism in KOPR-G protein coupling in cells in the BLA or in cells that project to the BLA. Recently, Bangasser et al. (2010) found that females had higher G_s protein coupling to corticotrophin-releasing factor receptor in the locus coeruleus than males. Since KOPR activity in the BLA is required for fear and anxiety-like behaviors (Knoll et al., 2011) and for stress-induced anxiety (Bruchas et al., 2009) our results may explain in part the findings that stress produces sex-dependent effects on anxiety-like behavior (Beck and Luine, 2002; Mitsushima et al., 2006; Waddell et al., 2008). Future studies are needed to clarify the observed functional sex difference of KOPR activity in the BLA.

KOPR antagonists have been found to block stress-induced reinstatement of cocaine seeking (McLaughlin et al., 2006). Antagonism of KOPR in DRN blocked stress-induced reinstatement of cocaine seeking in males (Land et al., 2009). As females had greater neural activity in the DRN but a lower net change in activity by U50,488H than males, KOPR antagonists may have different effects in females.

KOPR, diuresis and neuroendocrine functions

In the present study we found that males showed greater U50,488H-induced p44/42 activation in the paraventricular nucleus (PVN), while no sex differences were observed in the ventromedial hypothalamus (VMH). Our present results are consistent with previous reports that U50,488 induced an increase in c-Fos expression in the PVN of male guinea pigs (Bot and Chahl, 1996) and rats (Laorden et al., 2000; Laorden et al., 2003). In contrast, we previously found a trend in higher U50,488H-stimulated [35 S] GTP γ S binding in the hypothalamus of female compared to male guinea pigs (Wang et al., 2011). As females had

slightly higher receptor-G protein coupling in hypothalamus than males this suggests that KOPR-G-protein signaling in the PVN is not associated with p44/42 MAPK activity.

The PVN is critically involved in diuresis and in the control of neuroendocrine activity (release of stress hormones) (Kiss et al., 1986; Mezey and Kiss, 1985). KOPR agonists induce water diuresis (von Voigtlander et al., 1983). KOPR and dynorphin are highly co-localized in vasopressin-containing cells in the PVN of guinea pigs (Shuster et al., 2000). Selective KOPR agonists have been found to induce greater diuresis in males than females (Elliott et al., 2006; Craft et al., 1998). The greater U50,488H-induced activation of cells in the PVN in males than females may underlie the observed sex difference in diuresis.

KOPR activity modulates release of CRF, adrenocorticotrophic hormone (ACTH), and cortisol (Taylor et al., 1996; Calogero et al., 1996). In non-human primates no sex differences were observed in U50,488-induced release of ACTH and cortisol (Pascoe et al., 2008). The present findings suggest that KOPR modulation of P-p44/42 MAPK activity in the PVN does not correlate with the sex effects of U50,488H-induced ACTH and cortisol release.

KOPR agonists induce greater prolactin release in females than males (Manzanares et al., 1993; Butelman et al., 2007; Kreek et al., 1999). Prolactin release is controlled primarily by the tuberoinfundibular dopamine pathway projecting from the arcuate nucleus to median eminence (Arita and Porter, 1984). However, release of thyrotrophin-releasing hormone from hypophysiotropic neurons in the PVN (Taylor et al., 1990) can also modulate prolactin secretion from the pituitary (Gershengorn, 1986). Our findings suggest that KOPR-induced p44/42 MAPK activation in the PVN does not underlie sex differences in prolactin release.

KOPR and cognition

KOPR agonists have been found to produce deficits in attention (Shannon et al., 2007; Paine et al., 2007; Nemeth et al., 2010). The infralimbic, ventrolateral and lateral orbital cortices are thought to be involved in the control of cognitive processes such as attention, impulsivity and reasoning (Hoover and Vertes, 2011). Reduced neural activity in these regions may underlie the observed deficits in attention induced by KOPR agonists. However, since norBNI pretreatment did not reverse the effects of U50,488H and norBNI itself had no effects compared to saline these results suggest a non-KOPR mediated effect on p44/42 MAPK activity by U50,488H in these regions, such as blockade of potassium channels (Pugsley et al., 1994). It must be noted that our findings are in contrast to previous studies that have found that KOPR agonists induced an increase in neural activity in the orbital cortex (Bot and Chahl, 1996; Hooker et al., 2009). The reason for the discrepancy is unclear but could be due to differences in basal activity.

Role of estrous cycle

The guinea pig's estrous cycle is much longer than the rat or mouse; each cycle lasting 10–17 days: proestrus, 2–4 days; estrus, 9–11 hours; metestrus, 2–4 days; diestrus 8–10 days (Lilley et al., 1997). We did not examine estrous cycle effects since in our preliminary experiments we did not find that guinea pigs in proestrus/estrus differed from those in diestrus in their behavioral responses to U50,488H (Wang et al., 2011). Our goal here was to identify if organizational differences between the sexes could explain our behavioral findings. Thus, whether sex hormones have activational effect on U50,488H-induced neural activity remains to be investigated. However, our studies (Wang et al., 2011) and others suggest that sex hormones do not consistently or significantly play a role in KOPR agonist effects (Craft et al., 1998; Craft and Bernal, 2001; Gear et al., 1996a; Gioiosa et al., 2008).

Contributions of drug distribution and pharmacokinetics

The effects of U50,488H on p44/42 MAPK activity is not likely to be due to sex differences in drug distribution and pharmacokinetics. First, results of our previous study found that the direction of sexual dimorphism in the behaviors tested were not consistent across the endpoints (Wang et al., 2011). Second, although males generally showed greater neural activity changes induced by U50,488H, in some regions, e.g. thalamus, the effects were equivalent between the sexes, consistent with our previous study showing no difference in KOPR levels and KOPR-G protein coupling (Wang et al., 2011). Lastly, it has been demonstrated that brain distribution after systemic administration of [³H]U69,593 did not differ between male and female rats (Craft et al., 1998).

Conclusions

In summary, the present study provides a neuroanatomic mapping of circuitry activated by a KOPR agonist in guinea pigs. Together with our previous KOPR localization and KOPR-G protein coupling studies (Wang et al., 2011), this study implicates a further need to explore sex differences in KOPR-mediated behaviors, including stress, anxiety and drug addiction. Pharmacological and biochemical evidence from male subjects provide strong support for the therapeutic potential of KOPR ligands. However, since sex differences in KOPR distribution, activation and functional circuitry exist, further studies to determine the extent and mechanisms of sexual dimorphism are warranted.

Acknowledgments

This research was supported by NIH/NIDA grant DA17302 (LYLC) and T32 DA 07237-20 (KR). We thank Mr. Chongguang Chen for technical assistance.

Abbreviations

ANOVA	analysis of variance
BLA	basolateral amygdala
CeA	central amygdala
CLA	claustrum
DRN	dorsal raphe nuclei
HPA	hypothalamic-pituitary-adrenal
KOPR	kappa opioid receptor
i.p	intraperitoneally
LO	lateral orbital cortex
MAPK	mitogen activated protein kinase
MeA	medial amygdala
MOPR	mu opioid receptor
NAc	nucleus accumbens
norBNI	norbinaltorphimine
PAG	periaqueductal grey
PVN	paraventricular nucleus of the hypothalamus
ROI	region of interest

s.c	subcutaneously
THA	thalamus
U50, 488H (U50)	trans-(±)-3,4-dichloro-N-methyl-N-(2-[1-pyrrolidinyl]-cyclohexyl) benzeneacetamide methanesulfonate
U69, 593	(5a, 7a, 8b)-(-)- <i>N</i> -methyl- <i>N</i> -(7-(1-pyrrolidinyl)1-oxaspiro (4,5) dec-8-yl) benzeacetamide
VLO	ventrolateral orbital cortex
VMH	ventromedial hypothalamus

References

- Arita J, Porter JC. Relationship between dopamine release into hypophysial portal blood and prolactin release after morphine treatment in rats. *Neuroendocrinology*. 1984; 38:62–67. [PubMed: 6320031]
- Bangasser DA, Curtis A, Reyes BA, Bethea TT, Parastatidis I, Ischiropoulos H, Van Bockstaele EJ, Valentino RJ. Sex differences in corticotropin-releasing factor receptor signaling and trafficking: potential role in female vulnerability to stress-related psychopathology. *Mol Psychiatry*. 2010; 15:877, 896–877, 904. [PubMed: 20548297]
- Beck KD, Luine VN. Sex differences in behavioral and neurochemical profiles after chronic stress: role of housing conditions. *Physiol Behav*. 2002; 75:661–673. [PubMed: 12020731]
- Bohn LM, Belcheva MM, Coscia CJ. Mitogenic signaling via endogenous kappa-opioid receptors in C6 glioma cells: Evidence for the involvement of protein kinase C and the mitogen-activated protein kinase signaling cascade. *J Neurochem*. 2000; 74:564–573. [PubMed: 10646507]
- Bot G, Chahl LA. Induction of Fos-like immunoreactivity by opioids in guinea-pig brain. *Brain Res*. 1996; 731:45–56. [PubMed: 8883853]
- Bruchas MR, Land BB, Lemos JC, Chavkin C. CRF1-R activation of the dynorphin/kappa opioid system in the mouse basolateral amygdala mediates anxiety-like behavior. *PLoS One*. 2009; 4:e8528. [PubMed: 20052275]
- Butelman ER, Mandau M, Tidgewell K, Prisinzano TE, Yuferov V, Kreek MJ. Effects of salvinorin A, a kappa-opioid hallucinogen, on a neuroendocrine biomarker assay in nonhuman primates with high kappa-receptor homology to humans. *J Pharmacol Exp Ther*. 2007; 320:300–306. [PubMed: 17060493]
- Calogero AE, Scaccianoce S, Burrello N, Nicolai R, Muscolo LA, Kling MA, Angelucci L, D'Agata R. The kappa-opioid receptor agonist MR-2034 stimulates the rat hypothalamic-pituitary-adrenal axis: studies in vivo and in vitro. *J Neuroendocrinol*. 1996; 8:579–585. [PubMed: 8866244]
- Carlezon WA Jr, Beguin C, Dinieri JA, Baumann MH, Richards MR, Todtenkopf MS, Rothman RB, Ma Z, Lee DY, Cohen BM. Depressive-Like Effects of the {kappa}-Opioid Receptor Agonist Salvinorin A on Behavior and Neurochemistry in Rats. *J Pharmacol Exp Ther*. 2006; 316:440–447. [PubMed: 16223871]
- Carr GV, Lucki I. Comparison of the kappa-opioid receptor antagonist DIPPA in tests of anxiety-like behavior between Wistar Kyoto and Sprague Dawley rats. *Psychopharmacology (Berl)*. 2010; 210:295–302. [PubMed: 20369354]
- Chang SL, Squinto SP, Harlan RE. Morphine activation of c-fos expression in rat brain. *Biochem Biophys Res Comm*. 1988; 157:698–704. [PubMed: 3144275]
- Chen X, Geller EB, Rogers TJ, Adler MW. The chemokine CX3CL1/fractalkine interferes with the antinociceptive effect induced by opioid agonists in the periaqueductal grey of rats. *Brain Res*. 2007; 1153:52–57. [PubMed: 17459345]
- Craft RM. Sex differences in opioid analgesia: “from mouse to man”. *Clin J Pain*. 2003; 19:175–186. [PubMed: 12792556]
- Craft RM, Bernal SA. Sex differences in opioid antinociception: kappa and ‘mixed action’ agonists. *Drug Alcohol Depend*. 2001; 63:215–228. [PubMed: 11418226]

- Craft RM, Kruzich PJ, Boyer JS, Harding JW, Hanesworth JM. Sex differences in discriminative stimulus and diuretic effects of the kappa opioid agonist U69,593 in the rat. *Pharmacol Biochem Behav.* 1998; 61:395–403. [PubMed: 9802834]
- Craft RM, McNeil DM. Agonist/antagonist properties of nalbuphine, butorphanol and (–)-pentazocine in male vs. female rats. *Pharmacol Biochem Behav.* 2003; 75:235–245. [PubMed: 12759132]
- Craft RM, Ulibarri CM, Raub DJ. Kappa opioid-induced diuresis in female vs. male rats. *Pharmacol Biochem Behav.* 2000; 65:53–59. [PubMed: 10638636]
- Eitan S, Bryant CD, Saliminejad N, Yang YC, Vojdani E, Keith D Jr, Polakiewicz R, Evans CJ. Brain region-specific mechanisms for acute morphine-induced mitogen-activated protein kinase modulation and distinct patterns of activation during analgesic tolerance and locomotor sensitization. *J Neurosci.* 2003; 23:8360–8369. [PubMed: 12967998]
- Elliott JC, Picker MJ, Sparrow AJ, Lysle DT. Dissociation between sex differences in the immunological, behavioral, and physiological effects of kappa- and delta-opioids in Fischer rats. *Psychopharmacology (Berl).* 2006; 185:66–75. [PubMed: 16397747]
- Gear RW, Gordon NC, Heller PH, Paul S, Miaskowski C, Levine JD. Gender difference in analgesic response to the kappa-opioid pentazocine. *Neurosci Lett.* 1996a; 205:207–209. [PubMed: 8852594]
- Gear RW, Miaskowski C, Gordon NC, Paul SM, Heller PH, Levine JD. Kappa-opioids produce significantly greater analgesia in women than in men. *Nat Med.* 1996b; 2:1248–1250. [PubMed: 8898754]
- Gershengorn MC. Mechanism of thyrotropin releasing hormone stimulation of pituitary hormone secretion. *Annu Rev Physiol.* 1986; 48:515–526. [PubMed: 3010825]
- Gioiosa L, Chen X, Watkins R, Umeda EA, Arnold AP. Sex chromosome complement affects nociception and analgesia in newborn mice. *J Pain.* 2008; 9:962–969. [PubMed: 18635401]
- Hooker JM, Patel V, Kothari S, Schiffer WK. Metabolic changes in the rodent brain after acute administration of salvinorin A. *Mol Imaging Biol.* 2009; 11:137–143. [PubMed: 19132449]
- Hoover WB, Vertes RP. Projections of the medial orbital and ventral orbital cortex in the rat. *J Comp Neurol.* 2011; 519:3766–3801. [PubMed: 21800317]
- Kanarek RB, Mandillo S, Wiatr C. Chronic sucrose intake augments antinociception induced by injections of mu but not kappa opioid receptor agonists into the periaqueductal gray matter in male and female rats. *Brain Res.* 2001; 920:97–105. [PubMed: 11716815]
- Kiss JZ, Kanyicska B, Nagy GY. The hypothalamic paraventricular nucleus has a pivotal role in regulation of prolactin release in lactating rats. *Endocrinol.* 1986; 119:870–873.
- Knoll AT, Meloni EG, Thomas JB, Carroll FI, Carlezon WA Jr. Anxiolytic-like effects of kappa-opioid receptor antagonists in models of unlearned and learned fear in rats. *J Pharmacol Exp Ther.* 2007; 323:838–845. [PubMed: 17823306]
- Knoll AT, Muschamp JW, Sullivan SE, Ferguson D, Dietz DM, Meloni EG, Carroll FI, Nestler EJ, Konradi C, Carlezon WA Jr. Kappa opioid receptor signaling in the basolateral amygdala regulates conditioned fear and anxiety in rats. *Biol Psychiatry.* 2011; 70:425–433. [PubMed: 21531393]
- Koob GF. Drugs of abuse: anatomy, pharmacology and function of reward pathways. *Trends Pharmacol Sci.* 1992; 13:177–184. [PubMed: 1604710]
- Kreek MJ, Schluger J, Borg L, Gunduz M, Ho A. Dynorphin A1–13 causes elevation of serum levels of prolactin through an opioid receptor mechanism in humans: gender differences and implications for modulation of dopaminergic tone in the treatment of addictions. *J Pharmacol Exp Ther.* 1999; 288:260–269. [PubMed: 9862779]
- Land BB, Bruchas MR, Schattauer S, Giardino WJ, Aita M, Messinger D, Hnasko TS, Palmiter RD, Chavkin C. Activation of the kappa opioid receptor in the dorsal raphe nucleus mediates the aversive effects of stress and reinstates drug seeking. *Proc Natl Acad Sci USA.* 2009; 106:19168–19173. [PubMed: 19864633]
- Laorden ML, Castells MT, Martinez MD, Martinez PJ, Milanes MV. Activation of c-fos expression in hypothalamic nuclei by mu- and kappa-receptor agonists: correlation with catecholaminergic activity in the hypothalamic paraventricular nucleus. *Endocrinol.* 2000; 141:1366–1376.

- Laorden ML, Castells MT, Milanes MV. Effects of U-50488H and U-50488H withdrawal on c-fos expression in the rat paraventricular nucleus. Correlation with c-fos in brainstem catecholaminergic neurons. *Br J Pharmacol*. 2003; 138:1544–1552. [PubMed: 12721110]
- Li C, Pleil KE, Stamatakis AM, Busan S, Vong L, Lowell BB, Stuber GD, Kash TL. Presynaptic inhibition of gamma-aminobutyric Acid release in the bed nucleus of the stria terminalis by kappa opioid receptor signaling. *Biol Psychiatry*. 2012; 71:725–732. [PubMed: 22225848]
- Li J-G, Luo LY, Krupnick JG, Benovic JL, Liu-Chen L-Y. U50,488H-induced internalization of the human kappa opioid receptor involves a beta-arrestin- and dynamin-dependent mechanism. Kappa receptor internalization is not required for mitogen-activated protein kinase activation. *J Biol Chem*. 1999; 274:12087–12094. [PubMed: 10207034]
- Lilley KG, Epping RJ, Hafner LM. The guinea pig estrous cycle: correlation of vaginal impedance measurements with vaginal cytologic findings. *Lab Anim Sci*. 1997; 47:632–637. [PubMed: 9433700]
- Liu J, Nickolenko J, Sharp FR. Morphine induces c-fos and junB in striatum and nucleus accumbens via D1 and N-methyl-D-aspartate receptors. *Proc Natl Acad Sci USA*. 1994; 91:8537–8541. [PubMed: 8078918]
- Liu-Chen L-Y. Agonist-induced regulation and trafficking of kappa opioid receptors. *Life Sci*. 2004; 75:511–536. [PubMed: 15158363]
- Mansour A, Khachaturian H, Lewis ME, Akil H, Watson SJ. Anatomy of CNS opioid receptors. *Trends Neurosci*. 1988; 11:308–314. [PubMed: 2465635]
- Manzanares J, Wagner EJ, Moore KE, Lookingland KJ. Kappa opioid receptor-mediated regulation of prolactin and alpha-melanocyte-stimulating hormone secretion in male and female rats. *Life Sci*. 1993; 53:795–801. [PubMed: 8394964]
- McLaughlin JP, Land BB, Li S, Pinar JE, Chavkin C. Prior activation of kappa opioid receptors by U50,488 mimics repeated forced swim stress to potentiate cocaine place preference conditioning. *Neuropsychopharmacology*. 2006; 31:787–794. [PubMed: 16123754]
- McLaughlin JP, Marton-Popovici M, Chavkin C. Kappa opioid receptor antagonism and prodynorphin gene disruption block stress-induced behavioral responses. *J Neurosci*. 2003; 23:5674–5683. [PubMed: 12843270]
- Mezey E, Kiss JZ. Vasoactive intestinal peptide-containing neurons in the paraventricular nucleus may participate in regulating prolactin secretion. *Proc Natl Acad Sci USA*. 1985; 82:245–247. [PubMed: 3881755]
- Mitsushima D, Yamada K, Takase K, Funabashi T, Kimura F. Sex differences in the basolateral amygdala: the extracellular levels of serotonin and dopamine, and their responses to restraint stress in rats. *Eur J Neurosci*. 2006; 24:3245–3254. [PubMed: 17156385]
- Nemeth CL, Paine TA, Rittiner JE, Beguin C, Carroll FI, Roth BL, Cohen BM, Carlezon WA Jr. Role of kappa-opioid receptors in the effects of salvinorin A and ketamine on attention in rats. *Psychopharmacology (Berl)*. 2010; 210:263–274. [PubMed: 20358363]
- Paine TA, Tomasiewicz HC, Zhang K, Carlezon WA Jr. Sensitivity of the five-choice serial reaction time task to the effects of various psychotropic drugs in Sprague-Dawley rats. *Biol Psychiatry*. 2007; 62:687–693. [PubMed: 17343834]
- Pascoe JE, Williams KL, Mukhopadhyay P, Rice KC, Woods JH, Ko MC. Effects of mu, kappa, and delta opioid receptor agonists on the function of hypothalamic-pituitary-adrenal axis in monkeys. *Psychoneuroendocrinology*. 2008; 33:478–486. [PubMed: 18325678]
- Paxinos, G.; Watson, C. *The Rat Brain Stereotaxic Atlas*. Academic Press; 1998.
- Pearson G, Robinson F, Beers GT, Xu BE, Karandikar M, Berman K, Cobb MH. Mitogen-activated protein (MAP) kinase pathways: regulation and physiological functions. *Endocr Rev*. 2001; 22:153–183. [PubMed: 11294822]
- Pfeiffer A, Brantl V, Herz A, Emrich HM. Psychotomimesis mediated by kappa opiate receptors. *Science*. 1986; 233:774–776. [PubMed: 3016896]
- Pugsley MK, Saint DA, Walker MJ. An electrophysiological basis for the antiarrhythmic actions of the kappa-opioid receptor agonist U-50,488H. *Eur J Pharmacol*. 1994; 261:303–309. [PubMed: 7813552]

- Quirion, R.; Pilapil, C. Distribution of multiple opioid receptors in the human brain. In: Mendelsohn, FAO.; Paxinos, G., editors. *Receptors in the human nervous system*. Academic Press, Inc; San Diego, CA: 1991. p. 103-121.
- Quirion R, Pilapil C, Magnan J. Localization of kappa opioid receptor binding sites in human forebrain using [3H]U69,593: comparison with [3H]bremazocine. *Cell Mol Neurobiol*. 1987; 7:303–307. [PubMed: 2830975]
- Rasakham K, Liu-Chen LY. Sex differences in kappa opioid pharmacology. *Life Sci*. 2011; 88:2–16. [PubMed: 20951148]
- Roberson ED, English JD, Adams JP, Selcher JC, Kondratieck C, Sweatt JD. The mitogen-activated protein kinase cascade couples PKA and PKC to cAMP response element binding protein phosphorylation in area CA1 of hippocampus. *J Neurosci*. 1999; 19:4337–4348. [PubMed: 10341237]
- Shannon HE, Eberle EL, Mitch CH, McKinzie DL, Statnick MA. Effects of kappa opioid receptor agonists on attention as assessed by a 5-choice serial reaction time task in rats. *Neuropharmacol*. 2007; 53:930–941.
- Sharp T, Zetterstrom T, Ljungberg T, Ungerstedt U. A direct comparison of amphetamine-induced behaviours and regional brain dopamine release in the rat using intracerebral dialysis. *Brain Res*. 1987; 401:322–330. [PubMed: 3815099]
- Shippenberg TS, Zapata A, Chefer VI. Dynorphin and the pathophysiology of drug addiction. *Pharmacol Ther*. 2007; 116:306–321. [PubMed: 17868902]
- Shuster SJ, Riedl M, Li X, Vulchanova L, Elde R. The kappa opioid receptor and dynorphin co-localize in vasopressin magnocellular neurosecretory neurons in guinea-pig hypothalamus. *Neuroscience*. 2000; 96:373–383. [PubMed: 10683577]
- Taylor CC, Wu D, Soong Y, Yee JS, Szeto HH. kappa-Opioid agonist, U50,488H, stimulates ovine fetal pituitary-adrenal function via hypothalamic arginine-vasopressin and corticotrophin-releasing factor. *J Pharmacol Exp Ther*. 1996; 277:877–884. [PubMed: 8627570]
- Taylor T, Wondisford FE, Blaine T, Weintraub BD. The paraventricular nucleus of the hypothalamus has a major role in thyroid hormone feedback regulation of thyrotropin synthesis and secretion. *Endocrinol*. 1990; 126:317–324.
- Tindal JS. The forebrain of the guinea pig in stereotaxic coordinates. *J Comp Neurol*. 1965; 124:259–266. [PubMed: 14330744]
- Valjent E, Pages C, Herve D, Girault JA, Caboche J. Addictive and non-addictive drugs induce distinct and specific patterns of ERK activation in mouse brain. *Eur J Neurosci*. 2004; 19:1826–1836. [PubMed: 15078556]
- von Voigtlander PF, Lahti RA, Ludens JH. U-50,488: a selective and structurally novel non-Mu (kappa) opioid agonist. *J Pharmacol Exp Ther*. 1983; 224:7–12. [PubMed: 6129321]
- Waddell J, Bangasser DA, Shors TJ. The basolateral nucleus of the amygdala is necessary to induce the opposing effects of stressful experience on learning in males and females. *J Neurosci*. 2008; 28:5290–5294. [PubMed: 18480285]
- Wang YJ, Rasakham K, Huang P, Chudnovskaya D, Cowan A, Liu-Chen LY. Sex Difference in {kappa}-Opioid Receptor (KOPR)-Mediated Behaviors, Brain Region KOPR Level and KOPR-Mediated Guanosine 5'-O-(3-[35S]Thiotriphosphate) Binding in the Guinea Pig. *J Pharmacol Exp Ther*. 2011; 339:438–450. [PubMed: 21841040]

Highlights

- Kappa opioid agonist U50,488H alters p44/42 MAPK activity in the guinea pig.
- Males generally show greater changes induced by U50,488H than females.
- Pretreatment with norBNI antagonized the effects of U50,488H in most brain regions.
- In basolateral amygdala, U50 increased P-MAPK in males but reduced it in females.
- These results highlight the need to include females in kappa opioid studies.

Figure 1.

Brain regions of interest (ROI). Areas of interest are those that displayed high levels of P-p44/42 immunoreactivity and were determined based on (A) the rat brain atlas of Paxinos and Watson (Paxinos and Watson, 1998) for rostral regions as there is no guinea pig atlas for these areas and (B) the guinea pig atlas of Tindal (Tindal, 1965) for caudal regions. ROIs sampled for each brain region are highlighted in grey. Abbreviations: BLA basolateral amygdala; CeA, central amygdala; Cgl, cingulate cortex; CLA, claustrum; DP, dorsal peduncular cortex; DRN, dorsal raphe nuclei; IL, infralimbic cortex; LO, lateral orbital cortex; LSd, lateral septum dorsal; LSv, lateral septum ventral; VO, ventrolateral orbital cortex; MeA, medial amygdala; NAc, nucleus accumbens; PAG, periaqueductal grey region; PVN, paraventricular nucleus; THA, thalamus; VMH, ventromedial hypothalamus

Figure 2.

Representative images of phospho-p44/42 MAP kinase immunolabeling in the telencephalon. Images shown are female (A) and male (B) guinea pigs treated with (left to right) saline, U50,488H, norBNI + saline (in males only), norBNI + U50,488H (in males only). From top to bottom, lateral orbital cortex, claustrum, and nucleus accumbens. Scale bar = 100 μ m.

Figure 3.

Representative images of phospho-p44/42 MAP kinase immunolabeling in the diencephalon. Images shown are female (A) and male (B) guinea pigs treated with (left to right) saline, U50,488H, norBNI + saline (in males only), norBNI + U50,488H (in males only). From top to bottom, paraventricular nucleus, central amygdala, medial amygdala, and thalamus. Scale bar = 100 μ m.

Figure 4.

Representative images of phospho-p44/42 MAP kinase immunolabeling in the midbrain. Images shown are female (A) and male (B) guinea pigs treated with (left to right) saline, U50,488H, norBNI + saline (in males only), norBNI + U50,488H (in males only). From top to bottom, lateral periaqueductal gray region and dorsal raphe nuclei. Scale bar = 100 μ m.

Table 1
Quantitation of Phospho-p44/42 MAPK immunopositive labeling

Bright-field images of coronal sections were analyzed. Quantitation was performed using NIH ImageJ 1.42 software. Prior to quantification, a threshold level was chosen from a 256-level gray scale. The threshold was selected to allow detection of nuclei, cells and fibers stained with moderate to high intensity, while suppressing lightly stained nuclei, cells and fibers. An ROI was defined for each select brain region and used to sample from each region. Bilateral counts of phospho-p44/42 MAPK immuno-positive cells and averages from 1–4 sections, depending on the brain region, were calculated for each animal with each animal representing a single observation. Comparisons between sex and treatment groups were made. Data represents mean ($\pm$ sem). Male-saline (n = 3), Male-U50,488 (n = 3–4), Male-nBNI+U50,488H (n = 3–4), Male-nBNI+U50,488H (n = 3) Female-saline (n = 4–5), and Female-U50,488 (n = 5), where each n is an independent animal. Scale bar = 100 μ m

Brain Region	Female Saline	U50,488	Male Saline	U50,488	nBNI+Saline	nBNI+U50,488
Cingulate cortex	9.4 $\pm$ 4.6	1.8 $\pm$ 1.4	7.0 $\pm$ 2.9	3.0 $\pm$ 1.2	3.4 $\pm$ 1.3	1.1 $\pm$ 0.9
Infralimbic cortex	12.0 $\pm$ 3.0	5.4 $\pm$ 1.9 *	14 $\pm$ 6.6	5.0 $\pm$ 1.5 *	3.8 $\pm$ 1.1	2.5 $\pm$ 1.8
Dorsal peduncular cortex	22.4 $\pm$ 6.2	13.0 $\pm$ 3.4	18.0 $\pm$ 5.1	12.7 $\pm$ 4.3	10.7 $\pm$ 0.9	1.3 $\pm$ 0.8§
Ventrolateral orbital cortex	7.0 $\pm$ 2.1	1.7 $\pm$ 0.9 **	7.7 $\pm$ 2.2	0.7 $\pm$ 0.3 **	4.8 $\pm$ 1.1	0.1 $\pm$ 0.1
Lateral orbital cortex	8.6 $\pm$ 1.8	4.2 $\pm$ 0.9 **	6.7 $\pm$ 1.8	1.0 $\pm$ 0.6 **	5.0 $\pm$ 1.6	0.5 $\pm$ 0.5
Clastrum	12.6 $\pm$ 7.3	15.8 $\pm$ 3.9	5.3 $\pm$ 1.5	21.7 $\pm$ 4.9	5.9 $\pm$ 1.1	2.0 $\pm$ 0.9§
Nucleus accumbens	11.6 $\pm$ 4.2	16.2 $\pm$ 8.1	1.0 $\pm$ 0.0	25.3 $\pm$ 3.7	2.9 $\pm$ 1.2	1.3 $\pm$ 0.8§
Lateral septum (dorsal)	26.8 $\pm$ 5.8	23.0 $\pm$ 2.1	26.3 $\pm$ 5.8	25.3 $\pm$ 8.7	18.1 $\pm$ 2.7	6.8 $\pm$ 2.7
Lateral septum (ventral)	25.6 $\pm$ 4.6	26.2 $\pm$ 3.9	28.3 $\pm$ 10.8	18.3 $\pm$ 5.2	22.3 $\pm$ 3.3	8.5 $\pm$ 1.9
Paraventricular nucleus	80.6 $\pm$ 19.5	62.8 $\pm$ 9.8	32.6 $\pm$ 9.7	81.9 $\pm$ 9.5 ^A	29.9 $\pm$ 6.4	44.3 $\pm$ 4.1§
Ventromedial hypothalamus	35.4 $\pm$ 6.5	34.2 $\pm$ 2.0	29.9 $\pm$ 5.6	36.8 $\pm$ 5.0	15.2 $\pm$ 5.2	24.4 $\pm$ 7.3
Central amygdala	38.2 $\pm$ 8.6	63.4 $\pm$ 16.5 *	36.7 $\pm$ 3.0	93.5 $\pm$ 15.9 *	25.8 $\pm$ 3.8	39.5 $\pm$ 4.8§
Medial amygdala	31.4 $\pm$ 6.6	45.2 $\pm$ 7.6 **	25.0 $\pm$ 6.1	54.0 $\pm$ 2.6 **	20.0 $\pm$ 3.8	22.4 $\pm$ 5.0§
Basolateral amygdala	31.0 $\pm$ 5.2 ⁺	16.4 $\pm$ 3.7 ⁺	11.0 $\pm$ 0.6	17.3 $\pm$ 1.4 *	22.5 $\pm$ 2.8 §	10.0 $\pm$ 2.7
Thalamus	31.6 $\pm$ 3.4	79.4 $\pm$ 14.7 **	25.3 $\pm$ 7.3	86.0 $\pm$ 8.2 **	24.4 $\pm$ 8.7	35.8 $\pm$ 5.9§
Periaqueductal grey region	38.3 $\pm$ 21.4	108.0 $\pm$ 24.7 *	17.7 $\pm$ 7.2	60.3 $\pm$ 26.8 *	12.4 $\pm$ 4.9	30.6 $\pm$ 16.5
Lateral PAG	5.6 $\pm$ 2.8	24.2 $\pm$ 4.5 *	2.3 $\pm$ 0.3	17.8 $\pm$ 6.5 *	1.9 $\pm$ 0.5	6.2 $\pm$ 2.8
Dorsal raphe nuclei	50.8 $\pm$ 11.0 ⁺⁺	90.6 $\pm$ 12.3 ^{***++}	7.0 $\pm$ 2.3	50.8 $\pm$ 10.7 **	7.9 $\pm$ 4.1	21.3 $\pm$ 8.3

* $p < 0.05$,

** $p < 0.01$ vs saline, main effect of treatment (by 2×2 ANOVA);

- ⁺ $p < 0.05$,
⁺⁺ $p < 0.01$ vs male, main effect of sex (by 2×2 ANOVA);
[^] $p < 0.05$ vs same-sex, significant interaction (by 2×2 ANOVA);
[§] $p < 0.05$ vs male-U50,488H same region (Student's t-test);
[§] $p < 0.05$, vs male-saline same region (Student's t-test).