T.O. Cheng, M.D.

THE DEVELOPMENT OF ECHOCARDIOGRAPHY IN CHINA: THE PIONEERING ROLE OF XIN-FANG WANG

Tsung O. Cheng, M.D.

The George Washington University Medical Center, Washington, D.C.

Introduction

I cannot discuss the history of the development of echocardiography in China without describing the pioneering role of Xin-fang Wang (王新房) from Wuhan, China, who is the “father of modern echocardiography.”¹ Although Inge Edler from Sweden²⁻⁵ and Harvey Feigenbaum from the United States⁶ were also oftentimes referred to as the fathers of echocardiography, both Edler⁷ and Feigenbaum⁸ recognized that the Chinese used cardiac ultrasonography in the very early years (Figure 1).

Development of Echocardiography in China

In the late 1950s, investigators in both Shanghai and Wuhan were already making and using ultrasonic devices to examine the hearts of their patients.⁹ In 1958, research on ultrasound was very popular in all industries throughout China. The pioneer of using ultrasound technology to perform medical diagnosis was Shi An (安适)(Figure 2A), who was not a physician but a customs officer and later a librarian in Shanghai.¹⁰ Working with human subjects, An successfully used a Type 1 converted ultrasound metal flaw detector of Shanghai Jiangnan Shipyard with a frequency of 2.5 megahertz (Figure 2B).⁹

It might be of some interest to point out here how Edler and Hertz developed echocardiography. The son of Gustav Hertz, the famous 1925 Nobel Prize Laureate in Physics,^{4,7} C. Hellmuth Hertz was not a physician but had a deep interest in ultrasonography. He was aware of an ultrasonic reflectoscope at a shipyard in Malmö and borrowed the device during a weekend in May 1953.^{3,7} When Edler and Hertz placed the transducer of the reflectoscope over the chest, they could detect moving echoes, obviously originating from the movements of the heart. Whereas Hertz knew there was no possibility of obtaining research grants for buying an ultrasonic reflectoscope, he discovered that the Siemens Corporation in Germany was building ultrasonic reflectoscopes for nondestructive material testing and flaw detection at the shipyard.⁷ Since Hertz’s father had been the director of Siemens & Halske AG’s Research Laboratory before the end of World War II, he contacted director

Wolfgang Gellinek of the Siemens medical branch in Erlangen, Germany. Hertz demonstrated the discovery using a Siemens reflectoscope in the factory laboratory. Immediately recognizing its potential for application in cardiology, Gellinek ordered one of their reflectoscopes to be sent to Lund as a loan for a year, despite the fact that some of their regular customers had to wait.⁷ Due to later development, this loan was never returned. It is, indeed, fascinating to note how both Hertz’s and An’s discoveries about echocardiography originated from a shipyard!

In 1958, An founded the Shanghai Ultrasound Medical Research Group at the Sixth People’s Hospital of Shanghai, and his team included Tao-Hsin Wang, a classmate of mine who later became a well-known neurosurgeon, and Zhi Zhang Xu, a.k.a. Chih Chang Hsu. This special research team made further studies of patients and published a preliminary report in Chinese in 1960.¹¹ In 1962, his team reported the results of its application in 4,618 patients, studied from December 1958 to October 1961, in the English edition of *Chinese Medical Journal*.¹² The organs they examined included the liver, stomach, uterus, breast, brain, lungs, and heart. This was a seminal publication as far as the hydatidiform mole was concerned because it had not been previously reported in the medical literature.

Role of Xin-fang Wang

Xin-fang Wang was born in Henan, China in 1934. Xin-fang in Chinese means “new house” (新房); Wang’s father gave him this

Figure 1. China’s early contribution to echocardiography from Shanghai and Wuhan (where Wang is) acknowledged by both (A) Edler⁷ and (B) Feigenbaum.⁸

Figure 2. (A) Shih An (courtesy XF Wang) and (B) China’s earliest ultrasonic diagnostic device, the A-mode scanner, made in Jiangnan Shipyard in Shanghai and used in 1958 by An for diagnosing various diseases in patients at the Shanghai Sixth People’s Hospital (courtesy B Hu and XF Wang).

name because they had just moved into a new house. However, I tend to think that his father had the foresight of recognizing his son's ultimate career in echocardiography, which in essence deals with new knowledge about the heart including, among other things, the "atrial chamber" (心房), which is homophonous with "new house" (新房). Wang received his medical degree in 1958 from Tongji Medical University in Wuhan. He became associate professor of medicine and director of the echocardiographic department at Union Hospital of the Tongji Medical University in 1980. In 1986, he was promoted to professor of medicine and deputy director of the Institute of Cardiovascular Disease at the Tongji Medical University, now called the Cardiovascular Institute of Huazhong University of Science and Technology.

Wang began to study medical ultrasound technology in 1961. He found ultrasound to be very effective in examining pathological alterations of accumulated body fluid, which appears as a fluid flat segment in A-mode ultrasound image and a fluid dark area in B-mode ultrasound image. Based on these characteristics, he then classified human tissues into four categories: (1) tissues with non-ultrasound reflection, such as urine, ascites, pleural effusion, and pericardial effusion; (2) tissues with little ultrasound reflection, such as liver and spleen; (3) tissue with strong ultrasound reflection, such as the mammary glands; and (4) tissue interface with almost complete ultrasound reflection, such as the interface between soft tissue like chest and abdominal wall, and the air-filled space like lungs. Using this classification of human tissues by ultrasound reflections, he then introduced a new ultrasonic imaging method to diagnose liver abscess, pleural effusion, and pericardial effusion, each of which appears as a dark region because there is no reflection of ultrasound in liquids. Since 1963, Wang concentrated on the application of ultrasound to study the heart, realizing that the heart chambers were full of blood with little ultrasound reflection and thus depicted as dark regions, while the heart walls were solid tissues with strong ultrasound reflection resulting in a clear delineation between them on the ultrasound image.

Wang's Pioneering Work

Fetal Echocardiography

Wang in the early 1960s raised this question: If ultrasound could be used to detect fetal movement, why could it not monitor fetal heart beat in accordance with the differences of ultrasound reflection of the amniotic fluid and the fetus?¹ In 1963 at a national ultrasound meeting in Wuhan, he reported echocardiographic findings of normal fetuses in pregnant women.¹³ His 1964 report of the M-mode echocardiographic study of the fetal heart rate and size in 140 pregnant women was the earliest such report in the world literature (Figure 3).¹⁴ This seminal publication appeared 8 years before such application was reported in the English literature by Winsberg in 1972.¹⁵

Mitral Stenosis

In the early 1960s, Wang pioneered the prototype echocardiographic equipment synchronized to electrocardiographic and phonocardiographic signals.¹ In addition to duplicating the

Figure 3. World's first fetal echocardiogram (courtesy XF Wang).

Figure 4. (A) World's first demonstration of coincidence of the E-peak (double-headed arrow) on echocardiogram with the opening snap (arrow) followed by the flat EA (now called EF) slope corresponding to a long diastolic murmur (DM) in a simultaneously recorded phonocardiogram in a patient with mitral stenosis (courtesy XF Wang). (B) Echocardiogram of a normal-sized left atrium in a normal subject. The distance from the closed mitral valve (C) to the posterior wall of the left atrium (N) is the size of the left atrium, which is 8 mm here (courtesy XF Wang). (C) The world's first echocardiographic demonstration of enlarged left atrium in mitral stenosis. The distance from C to N is widened to 22 mm in this patient. Note also the flat EA curve that is typical for mitral stenosis (courtesy XF Wang). O: transmission pulse; MV: mitral valve curve; N: posterior wall of left atrium near atrioventricular ring.

findings of Edler with regard to mitral stenosis,^{16,17} he clarified the mechanisms of various points, peaks, and segments of the echocardiograms in mitral stenosis. He was the first in the world to show that the opening snap in mitral stenosis coincides with the E-peak of the echocardiogram (Figure 4A).¹⁷

Wang was also the first in the world in 1965 to measure the size of the left atrium by echocardiography.¹⁷ He showed that, in contrast to a normal subject (Figure 4B), patients with mitral stenosis showed an increased left atrial size (Figure 4C). In recent years, assessment of left atrial size has gained increasing importance in clinical practice, because left atrial enlargement is an important marker of cardiovascular disease and a predictor of such adverse events as atrial fibrillation, stroke, congestive heart failure, and even mortality in a wide range of cardiovascular conditions including obesity.¹⁸ Although volume measurement of the left atrium should be made nowadays from two-dimensional (2D) or three-dimensional (3D) echocardiograms because of asymmetric left atrial enlargement due to constraint by surrounding structure, the pioneering work of Wang in using M-mode to assess the left atrial size should not be forgotten.

Intracardiovascular Smoke-Like Echo

Intravascular smoke-like echo always appears in regions of stasis, but the exact mechanism of its production remained unclear until 1992, when Wang reported a close correlation between intravascular smoke-like echo in the left atrium and erythrocyte rouleaux formation (Figure 5).¹⁹ This new finding has considerable significance in the interpretation of echocardiographic findings in terms of hemodynamic and cellular mechanisms, especially in patients with mitral stenosis.

Red blood cells aggregate to form rouleaux in patients with severe mitral stenosis because of blood flow stasis and decreased shear force; therefore, smoke-like echo is observed in the left atrium.²⁰⁻²² When blood flow passes through the stenotic mitral orifice into the left ventricle, flow velocity and shear force are increased, rouleaux disperse, and smoke-like echo disappears.¹⁹ Smoke-like echo in the left atrium has also abated

Figure 5. From a preagonal dog, (A) smoke-like echoes in both left ventricle (LV) and right ventricle (RV) as seen on the ultrasonogram, and (B) red blood cells as a dense network of rouleaux seen under a microscope (courtesy XF Wang).

Contrast Echocardiography

Wang's publications in 1978 and 1979 anteceded the 1983 article of Gaffney et al. by 5 years.²⁴⁻²⁸ Xu²⁹ from Shanghai in 1981 and Hua³⁰ from Wuxi in 1982 later each reported the use of carbon dioxide as an echocardiographic contrast agent. Roelandt's group from the Netherlands published their work in 1981 as well.³¹

Pericardial Effusion

Wang was also the first in the world to use echocardiography in guiding pericardiocentesis in August of 1963 (Figure 6).³⁷ The patient was a 21-year-old male. His echocardiogram revealed pericardial fluid located 1.5–6.0 cm beneath the skin in the left sixth intercostal space at the midclavicular line with a wave pulsing with the heart beat at 6.0 cm. Wang diagnosed a large amount of pericardial effusion and recommended that pericardiocentesis be performed, with the aspirating needle inserted to a depth of 3.0–5.0 cm beneath the skin (Figure 6A). On August 21, 1963, pericardiocentesis was performed using sterile

technique and under local anesthesia, with the aspirating needle inserted to a depth of 3.0 cm beneath the skin through the sixth left intercostal space midway between the midclavicular line and the anterior axillary line. A total amount of 370 ml of grass-green fluid was withdrawn. The procedure was successful and uneventful (Figure 6B).

Diagnosis of constrictive pericarditis is clinically important, therapeutically challenging, and oftentimes quite difficult. The diagnosis of constrictive pericarditis is usually based on the measurement of pericardial thickness and hemodynamic alterations. However, it is difficult to measure the pericardial thickness by echocardiography, and a few patients with constrictive pericarditis had normal pericardial thickness. Furthermore, similar symptoms and hemodynamic changes may exist in other disease states such as restrictive cardiomyopathy.

Other Major Contributions

There are more than 50 textbooks on echocardiography published in China. Wang published China's first textbook on

Figure 6. Original handwritten notes of world's first pericardiocentesis under echocardiographic guidance performed in Wuhan, China in August 1963. (A) Echocardiographic findings on the day before pericardiocentesis; (B) Procedure notes on the following day during and following pericardiocentesis (courtesy XF Wang). See text for detailed discussion.

Figure 7. Quantitative tissue Doppler imaging in (A) a normal subject, (B) a patient with constrictive pericarditis, and (C) a patient with restrictive cardiomyopathy. Yellow line: parietal pericardium; blue line: outer-layer myocardium; red line: inner-layer myocardium (courtesy XF Wang). See text for discussion.

echocardiography in 1975, although his name never appeared anywhere in the book; during the period of China's Cultural Revolution, only the name of the institution, and not the author, was allowed. The first edition of his book was formally published in 1981. Its second, third, and fourth editions were published in 1985, 1999, and 2009, respectively. Wang also published China's first color Doppler textbook in 1991. His classic publication in 1995 on dynamic 3D echocardiography, also known as 4D echocardiography, was the first in China if not the first in the world.⁵²

All of Wang's outstanding work and publications, which numbered over 200, received high praises both at home and abroad. Consequently, he was awarded many national and international honors, such as the "History of Medical Ultrasound Pioneer Award" by the World Federation of Ultrasound in Medicine and Biology and the American Institute of Ultrasound in Medicine (1988) — along with Yong Chiang Chou, Wang Xue Guo, Zhi Zhang Xu, and Ruo Feng — and other prestigious awards by the State Council of China (1979, 1998, 2005), the State Education Commission (1992), and the Ministry of Health (1992). His most recent honor is an Honorary Fellowship in the American Society of Echocardiography awarded in July of this year. This honor has been awarded previously to only 11 recipients from around the world, namely, Jos Roelandt from the Netherlands, Liv Hatle from Norway, Junichi Yoshikawa from Japan, Fausto Pinto from Portugal, Namsik Chung from Korea, Alessandro Distanto from Italy, Shintaro Beppu from Japan, Jorge A. Lowenstein from Argentina, Chuwa Tei from Japan, Jean G. Dumesnil from Canada, and Raimund Erbel from Germany.

In short, echocardiography is continuing to advance under the very able and resourceful leadership of Wang, who certainly deserves the title of "Father of Modern Echocardiography." He may even be worthy of the title "Grandfather of Modern Echocardiography" since he has trained many well-known echocardiographers in China who in turn train younger generations of echocardiographers. But despite his "grandfather"

status, Wang is not prepared to retire anytime soon: he continues to respond to my letters and e-mails within 24 hours, goes to his university library and echocardiographic laboratory every day, attends important ultrasonic meetings, and continues to publish.⁴⁹⁻⁵¹

Conclusion

Echocardiography has revolutionized the practice of cardiology in the 21st century. It is the most popular and widely used cardiac imaging technique in the world and is the second most-frequently ordered cardiologic test after ECG. It is indispensable in the practice of medicine, and continues to benefit countless patients around the world. The extraordinary progress made in echocardiography in China under very limited resources is indeed a magnificent tribute to the dedication of Wang and his colleagues. All of us owe a great deal to Wang for his many pioneering contributions to echocardiography. As the Chinese saying goes, "When drinking the water, think of the man who dug the well."¹ So whenever we perform or order an echocardiogram for diagnostic, prognostic, or therapeutic purposes, always think of, and also be thankful to, Xin-fang Wang in Wuhan, China.

Epilogue

Both Edler and Wang are pioneers in echocardiography, and I had the good fortune of knowing them both. I met Edler in Atlanta during the 40th annual convocation of the American College of Cardiology in 1991, when he was awarded an Honorary Fellowship along with two other awardees — Bakken, the founder of Medtronic, and Tung, the "Paul D. White of China" (Figure 8A).⁹ William Winters, the editor-in-chief of this journal and then-president of the American College of Cardiology, served as Edler's marshal while I served as Tung's marshal (Figure 8B). I sat with Edler and other dignitaries on the dais (Figure 8C), and we had a long conversation afterwards. I found Edler not only extremely knowledgeable but also very friendly and unassuming. According to Winters in his presentation speech, Edler was responsible for his own interest in echocardiography dating back to the early 1960s.

Figure 8. Fortieth annual convocation of the American College of Cardiology, during which three honorary fellowships were awarded. (A) Names of recipients and their marshals. (B) Dr. C.L. Tung, Shanghai, China (left), William Winters, Houston, Texas (center) and this author (right). (C) Seated on the dais, Edler (short arrow) and this author (long arrow) next to Tung and other dignitaries.

Figure 9. Meetings of this author with Wang in China on various occasions: (A) in 1979 with Wang (right) and his mentor Y.S. Li (left) in Li's home; (B) in 1992 in Tiananmen Square in Beijing; (C) in 1994 discussing an interesting echocardiogram in Wang's laboratory. Note the contrast of Wang's attire between A and B; Wang wore a Mao jacket in 1979 but a western suit in 1992 in front of the portrait of Chairman Mao in Tiananmen Square.

Edler, likewise, paid tribute to Winters for his contributions in the echocardiographic evaluation of mitral stenosis, post-mitral and aortic valve replacement, and pericardial effusion and atrial septal defect.⁷

I first met Wang in Wuhan, China in 1979 during the 4-week visit of the first teaching delegation of the American College of Physicians to China, which I led (Figure 9A).⁹ I was instantly impressed by his intelligence, brilliance, innovativeness, diligence, earnestness, and humility. Because we share many common interests, we become very close friends and have collaborated on many publications and projects. I visited him in China often and always learned a great deal from him at each visit (Figures 9B-9C).

Wang is also adept in photography and Chinese poetry. He frequently combined both to compose beautiful and touching stanzas that he would send to me periodically. I am particularly moved by the following that he sent to me in 2010, the Year of the Tiger (Figure 10). I was born in the Year of the Tiger. It reads:

*"Dedicated to Professor Tsung O. Cheng
(English translation courtesy Lü Fei, Second Secretary, Chinese Embassy,
Washington, D.C.)*

*Like the tigers crouching on the summit,
Marvelous is your achievement in the study of heart.
With students everywhere in the world,
Your accomplishment will run through the vast universe.
Congratulations with respect
From your student Wang Xin-fang"*

I consider myself extremely fortunate to have made my acquaintances with Wang — my best friend, my most respected confidant, and my most inspiring colleague.

Figure 10. Pictorial stanza this author received from Wang in 2010, the Year of the Tiger, which was this author's year of birth. In the Chinese calendar, every lunar year is represented by an animal, and the cycle repeats every 12 years. See text for English translation.

Conflict of Interest Disclosure: The author has completed and submitted the *Methodist DeBakey Cardiovascular Journal* Conflict of Interest Statement and none were reported.

Funding/Support: The author has no funding disclosures.

Keywords: echocardiography, Xin-fang Wang, cardiac ultrasonography

References

- Cheng TO. Professor Xin-fang Wang from China is the father of modern echocardiography. *Int J Cardiol.* 2012 Jan 26;154: 225-31.
- Holmer NG, Lindström K, Lundström N-R, Marsal K, Olsson B, Pettersson H, et al. In memoriam: Inge Edler—the father of echocardiography. *Eur J Ultrasound.* 2001 Mar;13(1):179-82.
- In memoriam. Inge Edler: the father of echocardiography. *Eur J Echocardiogr.* 2001 Mar;2(1):1-2.
- Acierno LJ, Worrell LT. Inge Edler: father of echocardiography. *Clin Cardiol.* 2002 Apr;25(4):197-9.
- Singh S, Goyal A. The origin of echocardiography. A tribute to Inge Edler. *Tex Heart Inst J.* 2007;34(4):431-8.
- American Heart Association [Internet]. Dallas: American Heart Association; c2005-2012. Previous Distinguished Scientists; 2005[cited 2012 Jul 16]. Available from: http://my.americanheart.org/professional/Councils/DistinguishedScientists/Previous-Years-Distinguished-Scientists_UCM_321415_SubHomePage.jsp
- Edler I, Lindström K. The history of echocardiography. *Ultrasound Med Biol.* 2004 Dec; 30(12): 1565-644.
- Feigenbaum H, Armstrong WF, Ryan T. Feigenbaum's Echocardiography. 6th edition. Baltimore: Lippincott Williams & Wilkins; 2004. p. 1-10.
- Cheng TO. The evolution of cardiology in China. In: Wan S, Yim APC, editors. *Cardiothoracic surgery in China: past, present and future.* Hong Kong: The Chinese University Press; 2007. p. 204-305.
- Wang, Xin-fang. (deputy director of the Cardiovascular Institute of Huazhong University of Science and Technology, Wuhan, China). Personal communication with Tsung O. Cheng, M.D. (George Washington University Medical Center, Washington, D.C.). 2005 Feb 23.
- An S, Wang TH, An SY, et al. Application of ultrasound in clinical diagnosis: a preliminary report. *Chin Med J (Chinese edition).* 1960;46:48-55.
- An S, Wang TH, An SY, et al. The use of pulsed ultrasound in clinical diagnosis. *Chin Med J.* 1962 May;81:315-25.
- Wang, Xin-fang. (deputy director of the Cardiovascular Institute of Huazhong University of Science and Technology, Wuhan, China). Personal communication with Tsung O. Cheng, M.D. (George Washington University Medical Center, Washington, D.C.). 2011 Jul 10.

14. Wang XF, Xiao JP. The use of fetal echocardiography for pregnancy diagnosis. *Chin J Obstet Gynecol.* 1964;10(4):267-9.
15. Winsberg F. Echocardiography of the fetal and newborn heart. *Invest Radiol.* 1972 May-Jun;7(3):152-8.
16. Wang XF, Sheng RT. Clinical application of ultrasonic cardiography: recording methodology. In: Wuhan Branch of Chinese Medical Association, editors. *Corpus of ultrasonic thesis*; 1963. p. 21-25.
17. Gao Y, Wang XF, Gao RY, et al. Characteristics of normal ultrasonic cardiogram and its changes in patients with mitral stenosis. *Chin J Intern Med.* 1965;13:710-2, 763.
18. Roelandt J. The increasing importance of left atrial size assessment. *J Cardiovasc Med (Hagerstown).* 2011 Feb;12(2):147.
19. Wang XF, Liu L, Cheng TO, Li ZA, Deng YB, Wang JE. The relationship between intracardiovascular smoke-like echo and erythrocyte rouleaux formation. *Am Heart J.* 1992 Oct;124(4):961-5.
20. Daniel WG, Nellessen U, Schröder E, Nonnast-Daniel B, Bednarski P, Nikutta P, et al. Left atrial spontaneous echo contrast in mitral valve disease: an indicator for an increased thromboembolic risk. *J Am Coll Cardiol.* 1988 Jun;11(6):1204-11.
21. Iliceto S, Antonelli G, Sorino M, Biasco G, Rizzon P. Dynamic intracavitary left atrial echoes in mitral stenosis. *Am J Cardiol.* 1985 Feb;55(5):603-6.
22. Beppu S, Nimura Y, Sakakibara H, Nagata S, Park YD, Izumi S. Smoke-like echo in the left atrial cavity in mitral valve disease: its features and significance. *J Am Coll Cardiol.* 1985 Oct;6(4):744-9.
23. Chow WH, Chow LTC, Cheung KL. Resolution of spontaneous echocardiographic contrast after percutaneous balloon mitral commissurotomy. *Int J Cardiol.* 1992 Feb;34(2):209-10.
24. Wang XF, Wang J, Huang Y, Cai C. Contrast echocardiography with hydrogen peroxide: I. Experimental study. *Chin Med J (Engl).* 1979 Sep;92(9):595-9.
25. Wang XF, Wang JN, Chen HR, Lu CF. Contrast echocardiography with hydrogen peroxide. II. Clinical application. *Chin Med J (Engl).* 1979 Oct;92(10):693-702.
26. Wang XF, Wang JE, Lu CF, Chen HR, Huang YZ. Contrast echocardiography with hydrogen peroxide in identifying cardiac anatomic structures. *Acta Academiae Medicinae Wuhan.* 1978;6:36-9.
27. Wang XF, Wang JE, Huang YZ, Chen HR, Lu CF, Cao C. Experimental study of hydrogen peroxide echocardiography. *Acta Academiae Medicinae Wuhan.* 1978;6:31-35.
28. Gaffney FA, Lin JC, Peshock RM, Bush L, Buja LM. Hydrogen peroxide contrast echocardiography. *Am J Cardiol.* 1983 Sep 1;52(5):607-9.
29. Xu ZZ, Shen XD, Jiang L, Su BJ, Xu ZX. Study of contrast echocardiography with CO₂ produced by vitamin C and sodium bicarbonate. *Chin J Physical Med.* 1981;3:193-9.
30. Hua ZQ, Du YZ, Qian YG, et al. Experimental study and clinical use of contrast echocardiography with acetic acid and sodium bicarbonate. *Chin J Physical Med.* 1982;4:1-4.
31. Meltzer RS, Serruys PW, Hugenholtz PG, Roelandt J. Intravenous carbon dioxide as an echocardiographic contrast agent. *J Clin Ultrasound.* 1981 Mar;9(3):127-31.
32. Feigenbaum H, Waldhausen JA, Hyde LP. Ultrasound diagnosis of pericardial effusion. *J Am Med Assoc.* 1965 Mar 1;191:711-4.
33. Cheng TO, Xu ZZ, Wang XF. The first use of echocardiograms in the diagnosis of pericardial effusion. *Int J Cardiol.* 2009 May 1;134(1):1-5.
34. Cheng TO. Echocardiography in the diagnosis of pericardial effusion: a historical perspective. *Methodist Debaque Cardiovasc J.* 2011 Oct-Dec;7(4):49-50.
35. Hsu CC. Ultrasonic diagnostics. Shanghai: Shanghai Sci Tech Press; 1961. p. 167.
36. Hsu CC. Preliminary studies on ultrasonics in cardiological diagnosis. II. The use of A-scope ultrasound apparatus in the diagnosis of heart disease. *Acta Acad Med Prim Shanghai.* 1964;2:335-43.
37. Wang XF, Wang JE, Ultrasound Department, First Affiliated Hospital of Wuhan Medical College. The use of ultrasound in diagnosis of pericardial effusion. *Chin Med J.* 1973;7(7):411-4.
38. Lu XF, Wang XF, Cheng TO, Xie MX, Lu Q. Diagnosis of constrictive pericarditis by quantitative tissue Doppler imaging. *Int J Cardiol.* 2009 Sep 11;137(1):22-8.
39. Cao QL, Jiang L. The standard views and clinical use of transesophageal echocardiography. *Chin J Med Imaging Technol.* 1989;5(2):8-11.
40. Wang XF, Li ZA, Cheng TO, Deng YB, Wang JE, Yang Y. Biplane transesophageal echocardiography: an anatomic-ultrasonic-clinical correlative study. *Am Heart J.* 1992 Apr; 123(4 Pt 1):1027-38.
41. Wang XF, Li ZA, Cheng TO, Deng YB, Zheng LH, Hu G, et al. Clinical application of three-dimensional transesophageal echocardiography. *Am Heart J.* 1994 Aug;128(2):380-8.
42. Cheng TO, Wang XF, Zheng LH, Li ZA, Lu P. Three-dimensional transesophageal echocardiography in the diagnosis of mitral valve prolapse. *Am Heart J.* 1994 Dec;128(6 Pt 1):1218-24.
43. Wang XF, Li ZA, Cheng TO, Xie MX, Hu G, Deng YB, et al. Four-dimensional echocardiography: methods and clinical application. *Am Heart J.* 1996 Sep;132(3):672-84.
44. Cheng TO, Xie MX, Wang XF, Li ZA, Hu G. Evaluation of mitral valve prolapse by four-dimensional echocardiography. *Am Heart J.* 1997 Jan;133(1):120-9.
45. Ji R, Wang X, Cheng TO, Liu W, Li Z, Liu L. Experimental study of assessment on ventricular activation origin and contraction sequence by Doppler tissue imaging. *J Huazhong Univ Sci Technolog Med Sci.* 2002;22(1):52-7.
46. Cheng TO, Xie MX, Wang XF, Wang Y, Lu Q. Real-time 3-dimensional echocardiography in assessing atrial and ventricular septal defects: an echocardiographic-surgical correlative study. *Am Heart J.* 2004 Dec;148(6):1091-5.
47. Xie MX, Wang XF, Cheng TO, Wang J, Lu Q. Comparison of accuracy of mitral valve area in mitral stenosis by real-time, three-dimensional echocardiography versus two dimensional echocardiography versus Doppler pressure half-time. *Am J Cardiol.* 2005 Jun 15;95(12):1496-9.
48. Xie MX, Wang XF, Cheng TO, Lu Q, Yuan L, Liu X. Real-time 3-dimensional echocardiography: a review of the development of the technology and its clinical application. *Progr Cardiovasc Dis.* 2005 Nov-Dec;48(3):209-25.
49. Cheng TO, Wang XF, Zhang J, Xie MX. Recent advances in the echocardiographic diagnosis of mitral valve prolapse. *Int J Cardiol.* 2010 Apr 1;140(1):1-11.
50. Yang YL, Wang XF, Cheng TO, Xie MX, Lü Q, He L, et al. Echocardiographic characteristics of the criss-cross heart. *Int J Cardiol.* 2010 Apr 15;140(2):133-7.
51. Yang YL, Xie MX, Cheng TO, Wang XF, Lü Q, Ghoorah D. Left coronary sinus of Valsalva aneurysm ruptured into the left ventricle: diagnosis by two-dimensional and real time three-dimensional echocardiography. *Int J Cardiol.* 2011 Sep 1;151(2):e35-6.
52. Wang XF. Four-dimensional echocardiography: imaging methods and clinical applications. *J Clin Cardiol.* 1995;11:383-4.