Published in final edited form as:

Tetrahedron Lett. 2012 November 28; 53(48): 6475–6478. doi:10.1016/j.tetlet.2012.09.026.

Acid and base catalyzed Davis-Beirut reaction: experimental and theoretical mechanistic studies and synthesis of novel 3-amino-2*H*-indazoles

Belem Avila^a, Mohammad H. El-Dakdouki^b, Musa Z. Nazer^c, Jason G. Harrison^a, Dean J. Tantillo^a, Makhluaf J. Haddadin^b, and Mark J. Kurth^a

^aDepartment of Chemistry, University of California, One Shields Avenue, Davis, California 95616

^bDepartment of Chemistry, American University of Beirut, Lebanon

^cDepartment of Chemistry, University of Jordan, Amman, Jordan

Abstract

The Davis-Beirut reaction, which provides an efficient synthesis of 2*H*-indazoles and, subsequently, indazolones, is shown to proceed rapidly from *o*-nitrosobenzaldehyde and primary amines under both acid or base catalysis. Experimental and theoretical evidence in support of a reaction mechanism is provided in which *o*-nitrosobenzylidene imine is a pivotal intermediate in this N,N-bond forming heterocyclization reaction. The Davis-Beirut reaction is also shown to effectively synthesize a number of novel 3-amino-2*H*-indazole derivatives.

Keywords

Davis-Beirut reaction; 2*H*-indazole; *o*-nitrosobenzaldehyde; mechanism; catalysis

Indazoles and indazolones are privileged heterocycles with many biological applications, which have prompted the development of numerous methods for the construction of these heterocyclic systems.¹ While the chemistries to and of 1*H*-indazoles are well developed,² 2*H*-indazoles are considerably less well studied.³ In recent years, we have reported the syntheses of a number of pyrazolo-based heterocycles (**1**), 2*H*-indazoles (**2**) and, subsequently, 1*H*-indazolones (**3a/b**) that were obtained via the Davis-Beirut reaction (Figure 1).⁴ We previously suggested that *o*-nitrosobenzylidene was an important intermediate in this chemistry. As part of our ongoing interest in the development of the Davis-Beirut reaction, we report here both acid and base-catalyzed examples that provide mechanistic insight into the reactions shown in Scheme 1. Computational studies are also reported, which provided for a detailed understanding of this intriguing reaction.

We have previously shown that 3-alkoxy-2*H*-indazoles (**6**) and 5*H*-indazolo[3,2-*b*]benzo[*d*]-1,3-oxazines (**2**) can be obtained from either 2-nitrobenzaldehyde (**4**) or 1-(bromomethyl)-2-nitrobenzene (**5**) by Davis-Beirut reactions mediated by KOH and water in

© 2012 Elsevier Ltd. All rights reserved.

Supplementary data

Supplementary data (additional computational details) associated with this article can be found, in the online version, at <http://###>.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

alcoholic solvent (Scheme 1).^{4b,c} As depicted in Scheme 2, we propose that *o*-nitrobenzylamine **7** delivers hemiaminal nitroso intermediate **8** under basic conditions and subsequent loss of water renders *o*-nitrosobenzylidene imine **9**. This *o*-nitrosobenzylidene imine is an important intermediate that engages two possible pathways to the 2*H*-indazole product. Path (a) proceeds by stepwise *N,N*-cyclization (**9** → **10**), alkoxyaminal formation (**10** → **11**), and 1,4-elimination of water (**11** → **12** → **2**). The path (b) variant has *N,N*-cyclization and alkoxyaminal formation occurring in a concerted manner (**9** → **11**) followed by 1,4-elimination of water. Given this framework, it was envisioned that *o*-nitrosobenzaldehyde (**13a**) and 2-aminobenzyl alcohol (**14**) could serve as precursors to intermediate **9** and, by a Davis-Beirut reaction, lead to 2*H*-indazole **2**.

Experimental studies to probe this possibility started with the photolysis of *o*-nitrobenzyl alcohol (**15**) to give *o*-nitrosobenzaldehyde (**13a**) in 48% yield (Scheme 3;^{5a} isolation and handling of **13a** requires attention as it decomposes readily – especially upon heating in solution or exposure to light). We also investigated the hydrolysis of *o*-nitrobenzyl tosylates and mesoylates as routes to **13a**,^{5b} but the yields were poor and variable in our hands. Subjecting **13a** and 2-aminobenzyl alcohol (**14**) to typical Davis-Beirut heterocyclization conditions (aqueous KOH in MeOH) failed to deliver indazolobenzoxazine **2** at room temperature; indeed, the starting materials were found to be still present (with stoichiometry-based evidence of *o*-nitrosobenzaldehyde decomposition). In contrast, catalytic acetic acid in MeOH at room temperature led to the rapid formation (~1 min.) of **2** in 85% yield (Scheme 3; Method A). This transformation constitutes the first example of an acid-catalyzed Davis-Beirut reaction.

With these acid catalysis conditions in hand, we next tested a one-pot sequential protocol where step one was photolysis of a methanolic solution of **15** (14 h) followed by addition of **14** in catalytic acetic acid (step two). This method, which avoids the isolation of **13a**, delivered **2**, but in 50% yield (Scheme 3, Method B). ¹H-NMR analysis of the crude reaction mixture established that starting material **15** remained (incomplete photolysis; prolonged photolysis reduced the amount of **15**, but did not improve the yield of **2** due to the instability of **13a**). Method B circumvents the isolation of the rather unstable *o*-nitrosobenzaldehyde and, additionally, proceeds in a slightly higher overall yield than the two independent steps of Method A. Finally, these results with **13a** + **14** lend credence to the formation of hemiaminal nitroso intermediate **8** and its conversion to *o*-nitrosobenzylidene imine **9** in the Davis-Beirut reaction.

We next investigated a one-pot–one-step photolysis/Davis-Beirut reaction (Method C, Scheme 3). For this method, a methanolic solution of **14** and **15** plus catalytic acetic acid was photolyzed (14 h), giving **2** but in only 10% yield. While low yielding, it is interesting to note that this one-pot–one-step reaction (**14** + **15** → **2**) is a clean process, giving only product and recovered starting materials.

In order to gain further insight into the reaction mechanism, density functional theory calculations⁷ were performed on the intermediates of the proposed reaction mechanism as shown in Figure 2. The results of these calculations indicate that all proposed intermediates are thermodynamically viable under the experimental conditions. Further, formation of final product **2** is extremely exergonic (nearly 60 kcal/mol lower in energy than the individual reactants).⁸ Also of interest was the nature of the key transition state(s) involved in the conversion of **9** to **11**. Our calculations suggest that a concerted attack [see Scheme 2, path (b) after proton exchange] is possible,⁹ with the alkoxide oxygen attacking the carbon of the imine, whose nitrogen lone-pair attacks the nitrogen of the protonated nitroso group. The transition state structure for this reaction (Figure 3a)¹⁰ is predicted to be associated with a barrier of only 3.7 kcal/mol. A stepwise process is also possible, but this has a higher barrier

(the transition state structure for the first step, associated with a predicted barrier of 16.8 kcal/mol, is shown in Figure 3b). Promoting the reaction by protonation or deprotonation was also examined. The former is predicted to lead to a stepwise process with a barrier of 9.6 kcal/mol (the transition state structure for the first step is shown in Figure 3c). The latter leads to another concerted process (transition state structure shown in Figure 3d; note that this process is less synchronous than that shown in Figure 3a⁹), also with a low predicted barrier, 6.5 kcal/mol. Overall, concerted double cyclization is predicted to predominate under neutral or basic conditions, while a stepwise process is predicted for acidic conditions.

Encouraged by these initial studies, we carried out the Davis-Beirut reactions outlined in Scheme 4. Treatment of the *o*-nitrobenzaldehydes **16a–c** with 6-methoxybenzo[d]thiazol-2-amine (**17**) followed by reduction of the resulting imines with NaBH₄ gave *o*-nitrobenzylamines **18a–c** (Scheme 4a; Method 1) in yields ranging from 86–93%. These *o*-nitrobenzylamines **18a–b** were then reacted with DBU in THF under anhydrous conditions at room temperature to affect Davis-Beirut heterocyclizations to 2-amino-2*H*-indazole **19a** and **19b** in yields of 41% and 50%, respectively. The presence of the 6-methoxy-2-aminobenzothiazole moieties at the 3-position of 2*H*-indazoles **19a/b** was, initially, surprising. However, mechanistic considerations lead us to speculate that 6-methoxy-2-aminobenzothiazole (**17**) is generated during the reaction of *o*-nitrobenzylamines **18a/b** with DBU. In Scheme 4b, we offer an explanation for how **17** is formed. In support of this mechanism, it was found that addition of excess **17** to the Method 1 reaction of **18a/b** (DBU/THF) increased the yields of **19a** and **19b** to 60–75%.

Finally, we extended the photolysis method for the preparation of *o*-nitrosobenzaldehyde (**13a**) to the preparation of *o*-nitroso-benzaldehyde analogs **13b/c**. Treating compounds **13a/b** with two equivalents of 6-methoxybenzothiazol-2-amine (**17**) under basic conditions delivered **19a/b** in yields of 41% and 50%, respectively (Scheme 4c; Method 2). It is interesting to note that **19c** could not be isolated from either procedure (Scheme 4a/c) – presumably because the *o*-nitrosbenzylidene intermediate (analog of **9**) was not formed under the reaction conditions.

In summary, the acid- and base-catalyzed Davis-Beirut reactions to 2*H*-indazoles were studied experimentally and theoretically (DFT calculations) and several unique 2-amino-2*H*-indazoles were synthesized. All evidence supports the reaction proceeding through an *o*-nitrosbenzylidene imine intermediate, which undergoes a concerted *N,N*-bond forming heterocyclization reaction. Indeed, the mechanistic insights reported here may provide the basis for developing other substrates for the Davis-Beirut reaction.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial support was provided by the National Science Foundation (CHE-0910870) and the National Institutes of Health (GM089153). We also acknowledge the National Science Foundation's Partnership for Advanced Computational Infrastructure (Pittsburgh Supercomputer Center). J.G.H. thanks the United States Department of Education for a GAANN Fellowship and B.A. thanks the Bristol-Myers Squibb Graduate Chemist Fellowship and Alfred P. Sloan Minority Ph.D. Program for fellowship support.

References and notes

1. (a) Haddadin MJ, Conrad WB, Kurth MJ. Mini-Rev Med Chem. In press. (b) Cankarova N, Hlavac J, Krchnak V. Org Prep Proc Int. 2010; 42:433–465. (c) Schmidt AB, Snovydyovych B. Euro J Org

- Chem. 2008;4073–4095.(d) Huang LJ, Shih ML, Chen HS, Pan SL, Teng CM, Lee FY, Kuo SC. *Bioorg Med Chem.* 2006; 14:528–536. [PubMed: 16202611] (d) De Angelis M, Stossi F, Carlson KA, Katzenellenbogen BS, Katzenellenbogen JA. *J Med Chem.* 2005; 48:1132–1144. [PubMed: 15715479]
2. Jung N, Brase S. *Science of Synthesis.* 2010; 41:613.
 3. (a) Stadlbauer, W.; Camp, N. *Science of Synthesis: Houben-Weyl, Methods of Molecular Transformations.* Bellus, D.; Ley, SV.; Noyori, R.; Regitz, M.; Schaumann, E.; Shinkai, E.; Thomas, EJ.; Trost, BM.; Reider, PJ., editors. Vol. 12. Thieme; Stuttgart: 2002. p. 227(b) Wu C, Fang Y, Larock RC, Shi F. *Org Lett.* 2010; 12:2234–2237. [PubMed: 20394430]
 4. (a) Kurth MJ, Olmstead MM, Haddadin MJ. *J Org Chem.* 2005; 70:1060–1062. [PubMed: 15675871] (b) Mills AD, Nazer MZ, Haddadin MJ, Kurth MJ. *J Org Chem.* 2006; 71:2687–2689. [PubMed: 16555821] (c) Mills AD, Maloney P, Hassanein E, Haddadin MJ, Kurth MJ. *J Comb Chem.* 2007; 9:171–177. [PubMed: 17206845] (d) Butler JD, Solano DM, Robins LI, Haddadin MJ, Kurth MJ. *J Org Chem.* 2008; 73:234–240. [PubMed: 18052193] (e) Solano DM, Butler JD, Haddadin MJ, Kurth MJ. *Org Syn.* 2010; 87:339–349.(f) Avila B, Solano DM, Haddadin MJ, Kurth M. *J Org Lett.* 2011; 13:1060–1063.(g) Oakdale JS, Solano DM, Fettinger JC, Haddadin MJ, Kurth MJ. *Org Lett.* 2009; 11:2760–2763. [PubMed: 19505119] (h) Donald MB, Conrad WE, Oakdale JS, Butler JD, Haddadin MJ, Kurth MJ. *Org Lett.* 2010; 12:2524–2527. [PubMed: 20438102] (i) Conrad WE, Fukazawa R, Haddadin MJ, Kurth MJ. *Org Lett.* 2011; 13:3138–2141. [PubMed: 21612219] (j) Conrad WE, Rodriguez KX, Nguyen HH, Fettinger JC, Haddadin MJ, Kurth MJ. *Org Lett.* 2012; 14:3870–3873. [PubMed: 22823414]
 5. (a) Bamberger E. *Chem Ber.* 1918; 51:606–612.(b) Chen L-J, Burka LT. *Tetrahedron Lett.* 1998; 39:5351–5354.
 6. (a) **Method A.** *o*-Nitrobenzyl alcohol (2.61 mmol) was dissolved in MeOH (400 mL) and was placed under a 500W halogen lamp for 14 h. The solution was concentrated by rotary evaporation at a temperature below 30 °C, and purified by flash chromatography (100% DCM) to afford *o*-nitrosobenzaldehyde **13** as a white solid (168 mg, 48% yield). Spectral data is in accord with literature values.⁹ *5H*-Benzo[4,5][1,3]oxazino[3,2-*b*]indazole (**2**). To a solution of **13** (0.44 mmol) and **14** (0.44 mmol) in MeOH (4.4 mL) in a round-bottom flask equipped with a stir bar was added acetic acid (26 µL). The reaction mixture was allowed to stir until TLC indicated the disappearance of *o*-nitrosobenzaldehyde. The reaction was complete after 1 min., after which saturated bicarbonate was added and extracted with ethyl acetate (3 × 10 mL). The combined organic extracts were washed with brine, dried over sodium sulfate, filtered, and concentrated. The crude material was purified by flash chromatography yielding **2**, a white solid (85 mg, 85% yield). Spectral data is in accord with literature values.^{3d,e} (b) **Method B.** Synthesis of *5H*-Benzo[4,5][1,3]oxazino[3,2-*b*]indazole (**3**). *o*-Nitrobenzyl alcohol (0.74 mmol) was dissolved in MeOH (100 mL) in a glass beaker and was placed under a 500W halogen lamp for 14 h. After photolysis, the solution was placed in round-bottom flask equipped with a stir bar, after which, **14** (0.74 mmol) and acetic acid (24 µL) was added to the solution. The reaction mixture was allowed to stir until TLC indicated the disappearance of *o*-nitrosobenzaldehyde. The resulting mixture was then diluted with saturated bicarbonate was added and extracted with ethyl acetate (3 × 5 mL). The combined organic extracts were washed with brine, dried over sodium sulfate, filtered, and concentrated. The crude material was purified by flash chromatography yielding **3**, a white solid (82 mg, 50% yield). Spectral data is in accord with literature values.^{3d,e} (c) **Method C.** Synthesis of *5H*-Benzo[4,5][1,3]oxazino[3,2-*b*]indazole (**3**). To a solution of *o*-nitrobenzyl alcohol (0.74 mmol) and **14** in MeOH (100 mL) in a glass beaker was added acetic acid (24 µL) and was placed under a 500W halogen lamp for 14 h. The resulting mixture was then diluted with saturated bicarbonate was added and extracted with ethyl acetate (3 × 10 mL). The combined organic extracts were washed with brine, dried over sodium sulfate, filtered, and concentrated. The crude material was purified by flash chromatography yielding **3**, a white solid (16 mg, 10% yield). Spectral data is in accord with literature values.^{3d,e}
 7. All calculations were performed using *Gaussian09*. Structures were fully optimized at the M06-2X/6-31+G(d,p) level with an implicit solvent continuum model (CPCM) in methanol. Full references and coordinates can be found in the Supporting Information.
 8. As the reaction proceeds, there are several equivalents of water lost. To compensate for this in the calculations, the energy of isolated water molecules were calculated and their energy added to those

of the various intermediates as indicated; this allows all structures to be compared on the same energy scale.

9. See supporting information for intrinsic reaction coordinate and brief discussion. Leading references on concerted but highly asynchronous reactions: Tantillo DJ. *J Phys Org Chem.* 2008; 21:561–570. Williams A. *Concerted Organic and Bio-Organic Mechanisms.* CRC Press 2000 Dewar MJS. *J Am Chem Soc.* 1984; 106:209–219.
10. See Supporting Information for further discussion.
11. (a) **Preparation of 18a–c.** Equimolar amounts of *o*-nitrobenzaldehyde derivatives **16a** [or **b** or **c**] and 2-amino-6-methoxybenzo-thiazole (**17**) were mixed thoroughly and heated at 110 °C. The Schiff base, initially a viscous red liquid, solidified upon cooling and was dissolved in the minimal volume of methanol for reduction with NaBH₃ at room temperature. The pure (TLC) *o*-nitrobenzylamine derivatives **18a** [or **b** or **c**] precipitated as yellow a solid, which was collected and washed with cold methanol. The structures of **18a–c** were supported by IR, ¹H-NMR, ¹³C-NMR and MS. (b) **Preparation of 19a/b.** (i) **Method 1.** From **18a/b**. Equimolar amounts of *o*-nitrobenzylamine derivative **18a** [or **b**] and 2-amino-6-methoxybenzothiazole (**17**) were dissolved in dry THF in a round bottom flask wrapped with aluminum foil. Five drops of DBU were added the reaction mixture was stirred for 4 h at room temperature. The solution was concentrated by rotary evaporation and purified by column chromatography using 9:1 DCM:n-hexane as eluent. The green fluorescing fraction was collected and the solvent was evaporated under vacuum at 35 °C to yield **19a** [or **b**] as a yellow solid in 41% yield [50% for **19b**]. (ii) **Method 2.** From *o*-nitrosobenzaldehyde **13a/b**. One equivalent of *o*-nitrosobenzaldehyde derivative **13a** [or **b**] and two equivalents of 2-amino-6-methoxybenzothiazole were dissolved in dry THF in a round bottom flask wrapped with aluminum foil. Five drops of DBU were added to the reaction mixture, which was then stirred for 3 h at room temperature (reaction monitored by TLC). When all of the *o*-nitrosobenzaldehyde had been consumed, THF was evaporated at room temperature. The resulting dark brown mixture was purified by column chromatography using 9:1 DCM:n-hexane as eluent. The green fluorescing fraction was collected by evaporation of the solvent using a rotary evaporator set at 35 °C. The yellow solid was collected and placed in a vacuum oven over night. The structures of **19a/b** prepared by **Methods 1** and **2** were verified by IR, ¹H-NMR, ¹³C-NMR and MS.

Figure 1.
2H-Indazole and indazolone examples.

Figure 2. Overall reaction free energetics in kcal/mol, calculated at the M06-2X/6-31+G(d,p) level in implicit solvent (CPCM, methanol).

Figure 3. Key transition state structures for cyclization processes (M06-2X/6-31+G(d,p) in implicit solvent (CPCM, methanol)). Key distances are shown in Angstroms. (a) Concerted, neutral. (b) Stepwise, neutral. (c) Stepwise, protonated. (d) Concerted, deprotonated.

Scheme 1.
The Davis-Beirut Reaction.

Scheme 2.

Proposed mechanism for the formation of 2H-indazole **2** by the Davis-Beirut reaction.

Scheme 3.
Acid-catalyzed Davis-Beirut reactions to **2**.⁶

a) Method 1 ($R^1/R^2 = H/H$; $R^1/R^2 = H/Cl$; $R^1/R^2 = OMe/OMe$):

b) Proposed mechanism to **17** and **19a**.

c) Method 2 ($R^1/R^2 = H/H$; $R^1/R^2 = H/Cl$; $R^1/R^2 = OMe/OMe$):

Scheme 4.
Base-catalyzed Davis-Beirut reactions.¹¹