Published in final edited form as:

J Addict Med. 2012 December ; 6(4): 280–286. doi:10.1097/ADM.0b013e318262a97a.

Financial Factors and the Implementation of Medications for Treating Opioid Use Disorders

Hannah K. Knudsen, Ph.D.^{a,b} and Paul M. Roman, Ph.D.^{c,d}

^a Department of Behavioral Science, University of Kentucky, Lexington, KY

^b Center on Drug and Alcohol Research, University of Kentucky, Lexington, KY

^c Center for Research on Behavioral Health and Human Services Delivery, Owens Institute for Behavioral Research, University of Georgia, Athens, GA

^d Department of Sociology, University of Georgia, Athens, GA

Abstract

Objectives—Despite the established effectiveness of pharmacotherapies for treating opioid use disorders, implementation of medications for addiction treatment (MAT) by specialty treatment programs is limited. This research examined relationships between organizational factors and the program-level implementation of MAT, with attention paid to specific sources of funding, organizational structure, and workforce resources.

Methods—Face-to-face structured interviews were conducted in 2008–2009 with administrators of 154 community-based treatment programs affiliated with the National Institute on Drug Abuse's Clinical Trials Network (CTN); none of these programs exclusively dispensed methadone without offering other levels of care. Implementation of MAT was measured by summing the percentages of opioid patients receiving buprenorphine maintenance, methadone maintenance, and tablet naltrexone. Financial factors included the percentages of revenues received from Medicaid, private insurance, criminal justice, the Federal block grant, state government, and county government. Organizational structure and workforce characteristics were also measured.

Results—Implementation of MAT for opioid use disorders was low. Greater reliance on Medicaid was positively associated with implementation after controlling for organizational structure and workforce measures, while the association for reliance on criminal justice revenues was negative.

Conclusions—The implementation of MAT for opioid use disorders by specialty addiction treatment programs may be facilitated by Medicaid, but may be impeded by reliance on funding from the criminal justice system. These findings point to the need for additional research that considers the impact of organizational dependence on different types of funding on patterns of addiction treatment practice.

Send correspondence and reprint requests to: Hannah K. Knudsen, Ph.D., Department of Behavioral Science, University of Kentucky, 109 Medical Behavioral Science Building, Lexington, KY 40536-0086; Tel: 859-323-3947; Fax: 859-323-5350; Hannah.Knudsen@uky.edu.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

medications; pharmacotherapy; implementation science; opioid use disorders; treatment programs; Medicaid

Opportunities to effectively treat opioid use disorders (OUDs) are enhanced by the availability of pharmacological treatments. Unlike substance use disorders (SUDs) associated with stimulants or cannabis where such treatments are not yet available, there are effective pharmacological treatments for OUDs, namely methadone, buprenorphine, tablet naltrexone, and most recently, a depot formulation of naltrexone. The diffusion of buprenorphine and naltrexone has been remarkably slow, in part because of the financial resources required to implement pharmacotherapies.

For the past 40 years, specialized opioid treatment programs (OTPs) offering methadone have been the primary setting for treating opioid-dependent patients. Historically, the nation's OTPs have been inadequate in number to meet the demand for treatment. In 2000 (prior to the Food and Drug Administration's (FDA) approval of buprenorphine, OTPs constituted just 9% of US treatment facilities for SUDs, and methadone was included in the treatment plans of just 40% of all opioid treatment admissions (SAMHSA, 2002). The combination of the FDA's approval of newer medications (Ling and Smith, 2002; Saxon and McCarty, 2005) and regulatory changes allowing for the prescription of buprenorphine in non-OTP settings (Jaffe and O'Keeffe, 2003) has increased the potential expanding medications for addiction treatment (MAT) throughout the treatment system (Roman et al., 2011).

While the number of medications for OUDs has expanded, studies on the "research-to-practice gap" (Lamb et al., 1998) have repeatedly shown that such medications have not been widely adopted by treatment programs outside of the OTP system (Knudsen et al., 2006; Koch et al., 2006; Ducharme et al., 2007; Knudsen et al., 2007; Heinrich and Hill, 2008; Thomas et al., 2008; Friedmann et al., 2010; Knudsen et al., 2010; Wallack et al., 2010; Knudsen et al., 2011; Savage et al., 2012). Little is known about the implementation, or routine use, of MAT. To date, health services research has largely focused on medication adoption, measured by a dichotomous measure of availability, rather than implementation (Garner, 2009). A single study of MAT implementation for OUDs revealed limited implementation within adopting programs, with just 34.4% of patients with OUDs actually receiving MAT (Knudsen et al., 2011). The study did not analyze variations in implementation across programs in the sample.

Financial factors are often described as barriers to the delivery of pharmacotherapies in addiction treatment (Rieckmann et al., 2010; Knudsen et al., 2011). Resource dependence theory points to the relevance of funding sources for MAT implementation. This theoretical perspective contends that organizational strategy represents efforts to manage relationships with external entities that provide important resources (Pfeffer, 1987; Fennell and Warnecke, 1988). Treatment organizations must manage multiple external relationships since they typically receive funding from a mixture of sources, including Federal block grants, Medicaid, state and local government allocations, contracts with criminal justice, and private insurance reimbursement (Cartwright and Solano, 2003; Mark et al., 2005; Kubiak et al., 2009; Rieckmann et al., 2009; Stewart and Horgan, 2011; Thomas et al., 2011).

This heterogeneity of funding, with each funder having its own priorities and goals, likely has implications for the implementation of MAT. Although states currently vary in the services covered by Medicaid (Garfield et al., 2010), previous research has demonstrated

that greater reliance on Medicaid is positively associated with the adoption of SUD medications (Knudsen et al., 2010). Other research revealed that programs receiving any government funds were less likely to adopt buprenorphine, but that having at least one managed care contract was positively associated with adoption (Ducharme and Abraham, 2008). Historically, the criminal justice system has not embraced the use of medications to treat addiction. Correctional institutions have been slow to adopt medications for treating OUDs (Smith-Rohrberg et al., 2004; Rich et al., 2005), and less than one-third of administrators of Departments of Corrections indicate that medications are an ancillary service provided to offenders receiving community-based services (Kubiak et al., 2009).

This research examines the implementation of MAT for OUDs in a sub-sample of treatment programs within the National Drug Abuse Treatment Clinical Trials Network (CTN). The CTN partners university-based researchers with specialty addiction treatment programs across the US to conduct multi-site clinical trials of treatment interventions in heterogeneous settings and to disseminate evidence-based treatment practices (Hanson et al., 2002). It includes the full range of drug treatment modalities: outpatient “drug free” programs, therapeutic communities, OTPs, inpatient hospital-based programs, and complex organizations delivering varying levels of care at multiple sites (Ducharme et al., 2007; McCarty et al., 2008).

Drawing on data from 152 treatment centers within the CTN, this research addresses two questions. First, is greater reliance on certain types of financial resources associated with MAT implementation? Second, are financial factors associated with the implementation of MAT after controlling for organizational structure and workforce characteristics?

METHODS

Sample and Data Collection

Data are drawn from the ongoing National Treatment Center Study (NTCS), which includes research on the community-based addiction treatment organizations affiliated with the CTN (Roman et al., 2010). To be eligible for this study, organizations were required to offer either methadone maintenance as an opioid treatment program (OTP) or a level of treatment at least equivalent to the American Society of Addiction Medicine (ASAM) definition of outpatient services (Mee-Lee et al., 1996). Some organizations were housed in a single facility, while others consisted of multiple treatment centers, defined as organizational units with autonomous administrators who held discretionary control over their unit’s budget. When multiple treatment centers were embedded within a single organization, data were collected from the administrator and/or clinical director at each center. In 2008-2009, we identified 238 eligible treatment centers. Administrators and/or clinical directors of 198 treatment centers participated in face-to-face interviews (response rate = 84.7%). The research design was approved by the Institutional Review Boards of the University of Georgia and University of Kentucky.

For this analysis, 42 OTPs that did not offer other ASAM-defined levels of care were excluded because their implementation of MAT is essentially 100% (i.e., universal implementation is accomplished because MAT is their *raison d’être*). Treatment centers offering methadone as well as other ASAM-defined levels of care ($n = 11$) were included because MAT was not the only treatment option for patients with OUDs, as evidenced by an average rate of MAT below 50% in these 11 centers. These criteria restricted the analytic sample to 156 treatment centers.

Measures

The measure of implementation of MAT for OUDs was constructed by summing the percentages of patients with OUDs receiving buprenorphine for maintenance, tablet naltrexone, and methadone maintenance. At the time of the study, the depot injectable formulation of naltrexone had not received FDA approval for OUD treatment. Implementation for each medication was first measured, with programs not offering a particular medication being coded as zero. Then, the three indicators were summed to yield an aggregate measure of MAT implementation.

For sources of revenues, administrators were asked what percentage of past-fiscal year revenues came from Medicaid, private insurance, the criminal justice system, the Federal block grant, state government (excluding block grant funding), and county government. To maximize accuracy, administrators were sent these questions prior to the interview so that they could consult financial records.

Organizational variables consisted of both structural and workforce measures. Three structural variables were measured: location within a hospital setting (1 = hospital-based, 0 = non-hospital), profit status (1 = for-profit, 0 = non-profit), and delivery of outpatient-only services (1 = outpatient-only, 0 = offers inpatient/residential services). Workforce characteristics included measures of both physicians and counselors. We constructed a measure of physicians on staff by summing the numbers of psychiatrists and other physicians who were employed by the center (i.e., on the payroll). Workforce professionalism was measured by the percentage of counselors who holding at least a Master's level degree and the percentage of counselors who were personally in recovery.

Statistical Analysis

To address missing data on the covariates, we used multiple imputation by chained equations. If we had used listwise deletion, 13.9% of treatment centers would have been excluded from the analysis. Consistent with the recommendation of Allison (2009), we excluded 4 centers with missing data for the dependent variable. Imputation was performed using the “ice” command in Stata 12.0 (Royston, 2005; Royston, 2005), which generated 20 imputed data sets.

Our models of MAT implementation relied upon negative binomial regression (Long and Freese, 2006). The decision to use negative binomial regression rather than ordinary least squares (OLS) regression was informed by three factors. First, respondents reported the percentage of patients receiving medications as a whole number; while OLS regression assumes truly continuous variables (i.e., not whole numbers), negative binomial regression is an appropriate method for dependent variables represented by counts. Second, negative values for MAT implementation were impossible, which violated another assumption of OLS regression. Finally, negative binomial regression can handle substantially skewed data where a sizeable proportion of cases are zero on the dependent variable (Beck and Tolnay, 1995; Long, 1997).

An additional analytic challenge was that the measures of financial factors were not truly independent from each other. For example, a higher percentage of revenues from Medicaid meant that the other sources of revenues could only be lower. Therefore, we estimated a set of bivariate negative binomial regression models for each of the six financial factors. Then, we estimated multivariate models for each financial factor that was significant at the bivariate-level ($p < .05$, two-tailed test) with the organizational and workforce variables. For all models, we used the “mi estimate” command, which pooled the estimates from the 20 imputed datasets into a single set of results (Barnard and Rubin, 1999; Royston, 2004). We

included the “cluster” option (Long and Freese, 2006), which yields robust standard errors to account for data nesting (i.e., multiple treatment centers nested within a single organization).

RESULTS

In these treatment centers, 17.0% had adopted buprenorphine, 9.3% had adopted naltrexone, and 7.2% offered methadone for the treatment of OUDs. Overall, the average level of implementation of medications for addiction treatment (MAT) was low, reaching only an average of 9.6% of patients with opioid use disorders (OUDs). This low level of implementation was driven by the sizeable percentage (76.3%; $n = 116$) of treatment centers that had not implemented MAT for any OUD patients. Implementation was considerably greater in the 36 programs that offered at least one of the medications for OUDs, with an average of 40.4% of OUD patients receiving MAT ($SD = 34.9$).

Other descriptive statistics appear in Table 1. In the average treatment center, the two largest sources of revenues were state governments (meaning, allocations apart from the Federal block grant) and Medicaid. All of these measures had large standard deviations, indicating the heterogeneity of treatment center funding. The vast majority were located outside of hospital settings and operated on a non-profit basis. About half of these centers only offered outpatient treatment. The average number of physicians was slightly less than one, reflecting the fact that about half (53.0%; $n = 79$) did not have any physicians on staff.

Table 2 presents bivariate negative binomial (NB) regression models for each of the six financial factors. Three of the financial factors were significantly associated with MAT implementation. First, there was a positive association between the percentage of past-fiscal year revenues from Medicaid and MAT implementation. Greater reliance on revenues from the criminal justice system was negatively associated with MAT implementation. There was a negative association between MAT implementation and reliance on funding from county governments.

Next, organizational and workforce variables were added to the models for these three significant financial factors (Table 3). Controlling for organizational structure and workforce characteristics, reliance on Medicaid continued to be positively associated with MAT implementation (Model 1). While the unstandardized coefficient may appear small, the magnitude of this association became clearer when the change in the expected count of MAT implementation was calculated for a standard deviation increase in reliance on Medicaid. The expected count of MAT implementation increased by about 96.1% for a standard deviation ($SD = 24.06$) change in the percentage of revenues from Medicaid [$96.1 = (100)(e^{.028 \times 24.06} - 1)$]. Consistent with the bivariate model, there was a negative association between reliance on funding from the criminal justice system and MAT implementation (Model 2). A standard deviation ($SD = 13.95$) increase in the percentage of revenues from the criminal justice system reduced the expected count of MAT implementation by nearly half [$-48.1 = (100)(e^{-.047 \times 13.95} - 1)$]. As seen in Model 3, reliance on revenues from county governments trended towards, but did not achieve, statistical significance once structural and workforce variables were added to the model.

While our main goal was to examine sources of revenues in relation to MAT implementation, we also examined the associations for organizational and workforce characteristics because of the paucity of prior MAT implementation studies. Across the three models, location in a hospital setting was associated with MAT implementation. Using Model 1 as an example, the difference in the expected count of MAT implementation between hospital-based and non-hospital programs was about 375.9% [$375.9 = (100)(e^{1.560} - 1)$]. For-profit programs reported significantly lower MAT implementation than non-profit

programs in two of the three models; in the model of criminal justice funding, the p -value for profit status was .051. Based on Model 1, the expected count for MAT implementation was about 90.2% lower in for-profit centers compared to non-profit centers [$90.2 = (100)(e^{-2.321} - 1)$]. The number of physicians was positively associated with MAT implementation in two of the three models; the association in Model 1 trended towards, but did not achieve, significance ($p = .087$). Similarly, workforce professionalism, as measured by the percentage of Master's-level counselors, was positively associated with implementation in two of the three models. In Model 3, workforce professionalism was not statistically significant, although the direction of the association was consistent with the other two models ($p = .091$).

DISCUSSION

This study pioneers the examination of the relationships between financial factors and the implementation of medications for addiction treatment (MAT) for opioid use disorders (OUDs). The scope of implementation varies by the perspective taken on the data. MAT implementation, defined as the percentage of OUD patients receiving pharmacotherapy, was particularly low when addiction treatment centers that had adopted no medications were included in the analysis. Restricting the dataset to only those centers that had adopted at least one OUD medication yielded a rate of implementation (40%) that was slightly higher than a prior study of privately funded addiction treatment programs (34%) (Knudsen et al., 2011).

Multivariate models indicated that greater reliance on Medicaid for funding was positively associated with MAT implementation. It is important to note that Medicaid itself has historically been implemented differently across states in terms of which treatment services are covered (Garfield et al., 2010; Terry-McElrath et al., 2011). There has been a tendency within the SUD treatment field to regard Medicaid policies as a hindrance to innovation adoption, a perspective not supported by these data. A vital direction for future research is to link state-level data regarding Medicaid policy with program-level data on MAT implementation. Collection of Medicaid policy data was beyond the scope of the current study, so we were unable to control for whether programs were located in states where Medicaid provided coverage for MAT and included these medications on its formulary. Resource dependence theory suggests that supportive state policies would likely result in greater implementation of MAT (Kubiak et al., 2009; Thomas et al., 2011), although the presence of a supportive state policy is no guarantee that programs will accept Medicaid payment (Terry-McElrath et al., 2011).

The positive association between Medicaid funding and MAT implementation may suggest that the impending roll-out of key provisions in the Patient Protection and Affordable Care Act of 2010 (ACA; i.e., health reform) could be beneficial for OUD patients. Under ACA, the number of people covered by Medicaid is poised to expand considerably (Holahan and Headen, 2010) and will likely include significant numbers of individuals with substance use disorders (Buck, 2011; Stewart and Horgan, 2011). Notably, the ACA requires parity, such that Medicaid will offer the essential benefits that are available via the state insurance exchanges. It is expected that these changes will increase funding for SUD treatment via Medicaid (Buck, 2011), which may increase access and implementation of MAT given our current findings regarding Medicaid. Future research should examine whether the ACA has these intended consequences, and whether there are unanticipated consequences for the implementation of MAT and other services.

In contrast to the positive association for Medicaid, greater reliance on criminal justice funding was negatively associated with MAT implementation. This finding was not

unexpected, given the work of Kubiak and colleagues (2009) who surveyed state Department of Corrections leaders and found that medications were generally not a highly prioritized treatment service for these purchasers of treatment services. Historically, the criminal justice system has been slow to view MAT as an acceptable part of the treatment process (Chandler et al., 2009; Marlowe, 2011), and few correctional institutions have actively attempted to refer opioid-dependent offenders to MAT in the community (Rich et al., 2005; Nunn et al., 2009). Despite the consistency between our findings and these prior studies, more data are needed to develop an in-depth understanding of the reasons behind the limited adoption and implementation of medications for individuals involved with the criminal justice system. Diversion programs, drug courts, and the delivery of treatment under correctional auspices involve an intermingling of medical and justice paradigms that may suggest conflicts between key assumptions of these two systems. Addressing the complexities of increasing implementation of MAT for individuals under community corrections supervision (i.e., probation or parole) is a focus of the Criminal Justice Drug Abuse Treatment Studies 2 (CJDATS 2) initiative, which may yield additional insights regarding how to improve access to MAT for drug-involved offenders (Friedmann et al., 2012).

The trend of county funding being negatively associated with MAT implementation appears as a somewhat unique finding. It may suggest a lower level of exposure to newer treatment modalities at this level of decision-making. It may also represent greater influence at the county level of local persons from the recovery community, perhaps with a stronger orientation towards traditional 12-step and drug-free treatment modalities.

Our analysis focused on treatment organizations, but factors beyond financing are likely to explain additional variance in MAT implementation. Studies are needed about how physicians make prescribing decisions. Data from pharmacies has shown increases in the numbers of MAT prescriptions over time (Mark et al., 2009), but little is known about the factors that physicians consider when deciding whether to prescribe MAT to their patients. For example, what is the impact of a patient's primary source of payment for treatment on prescribing decisions? Previous work examining barriers to prescribing naltrexone for patients with alcohol use disorders found that patients' inability to afford medication was a prevalent barrier (Mark et al., 2003). It is less clear whether physicians are more likely to prescribe MAT for insured patients, such as those with Medicaid.

In addition, research is needed regarding the preferences of patients. There is little reason to assume readiness for public acceptance of MAT, except for the possibility of positive attitudes generated from knowing individuals who have benefitted from methadone treatment. To date, diffusion of information about the technology of MAT into public media has been minimal in that it has not been the target of any organized educational campaign. Receiving treatment outside of the OTP system may be a signal that some patients with OUDs prefer to not include pharmacotherapy in their treatment plan. However, where patients receive treatment is constrained by availability of treatment in the local context. Furthermore, patients cannot exert their preferences when programs do not offer MAT. Nonetheless, the dearth of available data means that we cannot untangle the complexities of how much of the variance in MAT implementation can be traced back to prescribers' behaviors and patients' preferences for treatment.

Several limitations of the research design should be noted. First, these data are cross-sectional, which limits our ability to determine causal relationships between financial factors and the implementation of MAT for OUDs. We are currently collecting another round of data from CTPs in the CTN, so we plan to conduct longitudinal analyses once data collection is complete. While diverse in the types of treatment centers (e.g., outpatient,

hospital inpatient, residential, etc.) that are affiliated with it (McCarty et al., 2008), the CTN does not constitute a random sample, which may limit the extent to which these findings may generalize to the rest of the US addiction treatment system. An additional limitation is that our measures were self-reported by administrators, which may increase the risk of recall or social desirability bias. We attempted to mitigate recall bias for the financial measures by sending administrators a worksheet containing these questions prior to the interview. The relatively low rate of MAT implementation seems to suggest that administrators were not exaggerating their center's use of MAT.

CONCLUSIONS

The limited implementation of medications for the treatment of opioid use disorders (OUDs) in specialty programs points to the continued significance of the research-to-practice gap. This research demonstrated that financial factors, specifically reliance on Medicaid and criminal justice funding, are associated with medication implementation even after controlling for key resources, such as access to physicians. The impending changes in the US health care system under the ACA, particularly with regard to the anticipated expansion of Medicaid, may have a positive impact on implementation. However, additional longitudinal research is needed to empirically demonstrate whether Medicaid expansion actually has this effect. Furthermore, longitudinal data would allow for consideration of whether criminal justice funding remains a persistent barrier to the implementation of medications for OUDs. Given the evolving policy and funding environment, additional studies are needed about how financial factors are related to the use of pharmacotherapy in the treatment of patients with opioid use disorders.

Acknowledgments

This research was supported by funding from the National Institute on Drug Abuse, National Institutes of Health (R01DA14482).

REFERENCES

- Allison, PD. Missing data. In: Millsap, RE.; Maydeu-Olivares, A., editors. *The SAGE Handbook of Quantitative Methods in Psychology*. Sage; Thousand Oaks, CA: 2009. p. 72-89.
- Barnard J, Rubin DR. Small sample degrees of freedom with multiple imputation. *Biometrika*. 1999; 86:948–955.
- Beck EM, Tolnay SE. Analyzing historical count data: Poisson and negative binomial regression models. *Hist Methods*. 1995; 28:125–131.
- Buck JA. The looming expansion and transformation of public substance abuse treatment under the Affordable Care Act. *Health Aff (Millwood)*. 2011; 30:1402–1410. [PubMed: 21821557]
- Cartwright WS, Solano PL. The economics of public health: Financing drug abuse treatment services. *Health Policy*. 2003; 66:247–260. [PubMed: 14637010]
- Chandler RK, Fletcher BW, Volkow ND. Treating drug abuse and addiction in the criminal justice system: Improving public health and safety. *JAMA*. 2009; 301:183–190. [PubMed: 19141766]
- Ducharme LJ, Abraham AJ. State policy influence on the early diffusion of buprenorphine in community treatment programs. *Subst Abuse Treat Prev Policy*. 2008; 3:17. [PubMed: 18570665]
- Ducharme LJ, Knudsen HK, Roman PM, et al. Innovation adoption in substance abuse treatment: Exposure, trialability, and the Clinical Trials Network. *J Subst Abuse Treat*. 2007; 32:321–329. [PubMed: 17481455]
- Fennell, ML.; Warnecke, RB. *The Diffusion of Medical Innovations: An Applied Network Analysis*. Plenum Press; New York: 1988.
- Friedmann PD, Hoskinson R Jr, Gordon M, et al. Medication-assisted treatment in criminal justice agencies affiliated with the Criminal Justice-Drug Abuse Treatment Studies (CJ-DATS): Availability, barriers, and intentions. *Subst Abus*. 2012; 33:9–18. [PubMed: 22263709]

- Friedmann PD, Jiang L, Alexander JA. Top manager effects on buprenorphine adoption in outpatient substance abuse treatment programs. *J Behav Health Serv Res*. 2010; 37:322–337. [PubMed: 19296223]
- Garfield RL, Lave JR, Donohue JM. Health reform and the scope of benefits for mental health and substance use disorder services. *Psychiatr Serv*. 2010; 61:1081–1086. [PubMed: 21041345]
- Garner BR. Research on the diffusion of evidence-based treatments within substance abuse treatment: A systematic review. *J Subst Abuse Treat*. 2009; 36:376–399. [PubMed: 19008068]
- Hanson GR, Leshner AI, Tai B. Putting drug abuse research to use in real-life settings. *J Subst Abuse Treat*. 2002; 23:69–70. [PubMed: 12220602]
- Heinrich CJ, Hill CJ. Role of state policies in the adoption of naltrexone for substance abuse treatment. *Health Serv Res*. 2008; 43:951–970. [PubMed: 18454775]
- Holahan, J.; Headen, I. [Accessed April 23, 2012] Medicaid coverage and spending in health reform: National and state-by-state results for adults at or below 133% FPL. 2010. <http://www.kff.org/healthreform/upload/Medicaid-Coverage-and-Spending-in-Health-Reform-National-and-State-By-State-Results-for-Adults-at-or-Below-133-FPL.pdf>
- Jaffe JH, O'Keeffe C. From morphine clinics to buprenorphine: Regulating opioid agonist treatment of addiction in the United States. *Drug Alcohol Depend*. 2003; 70:S3–S11. [PubMed: 12738346]
- Knudsen HK, Abraham AJ, Oser CB. Barriers to the implementation of medication-assisted treatment for substance use disorders: The importance of funding policies and medical infrastructure. *Eval Program Plann*. 2011; 34:375–381. [PubMed: 21371752]
- Knudsen HK, Abraham AJ, Roman PM. Adoption and implementation of medications in addiction treatment programs. *J Addict Med*. 2011; 5:21–27. [PubMed: 21359109]
- Knudsen HK, Ducharme LJ, Roman PM. Early adoption of buprenorphine in substance abuse treatment centers: Data from the private and public sectors. *J Subst Abuse Treat*. 2006; 30:363–373. [PubMed: 16716852]
- Knudsen HK, Ducharme LJ, Roman PM. The adoption of medications in substance abuse treatment: Associations with organizational characteristics and technology clusters. *Drug Alcohol Depend*. 2007; 87:164–174. [PubMed: 16971059]
- Knudsen HK, Roman PM, Oser CB. Facilitating factors and barriers to the use of medications in publicly funded addiction treatment organizations. *J Addict Med*. 2010; 4:99–107. [PubMed: 20835350]
- Koch AL, Arfken CL, Schuster CR. Characteristics of US substance abuse treatment facilities adopting buprenorphine in its initial stage of availability. *Drug Alcohol Depend*. 2006; 83:274–278. [PubMed: 16413146]
- Kubiak SP, Arfken CL, Gibson ES. Departments of corrections as purchasers of community-based treatment: A national study. *J Subst Abuse Treat*. 2009; 36:420–427. [PubMed: 19004602]
- Lamb, S.; Greenlick, MR.; McCarty, D., editors. Bridging the Gap between Practice and Research: Forging Partnerships with Community-Based Drug and Alcohol Treatment. National Academy Press; Washington, D.C.: 1998.
- Ling W, Smith D. Buprenorphine: Blending practice and research. *J Subst Abuse Treat*. 2002; 23:87–92. [PubMed: 12220606]
- Long, JS. Regression Models for Categorical and Limited Dependent Variables. Sage; Thousand Oaks, CA: 1997.
- Long, JS.; Freese, J. Regression Models for Categorical Dependent Variables Using Stata. 2nd edition. StataCorp; College Station, TX: 2006.
- Mark, TL.; Coffey, RM.; McKusick, D., et al. National Estimates of Expenditures for Mental Health Services and Substance Abuse Treatment, 1991–2001. SAMHSA; Rockville, MD: 2005.
- Mark TL, Kassed CA, Vandivort-Warren R, et al. Alcohol and opioid dependence medications: Prescription trends, overall and by physician specialty. *Drug Alcohol Depend*. 2009; 99:345–349. [PubMed: 18819759]
- Mark TL, Kranzler HR, Song X. Understanding US addiction physicians' low rate of naltrexone prescription. *Drug Alcohol Depend*. 2003; 71:219–228. [PubMed: 12957340]
- Marlowe DB. Evidence-based policies and practices for drug-involved offenders. *Prison J*. 2011; 91:27S–47S.

- McCarty D, Fuller B, Kaskutas LA, et al. Treatment programs in the National Drug Abuse Treatment Clinical Trials Network. *Drug Alcohol Depend.* 2008; 92:200–207. [PubMed: 17875368]
- Mee-Lee, D.; Gartner, L.; Miller, MM., et al. *Patient Placement Criteria for the Treatment of Substance-Related Disorders*. 2nd ed. American Society of Addiction Medicine; Chevy Chase, MD: 1996.
- Nunn A, Zaller N, Dickman S, et al. Methadone and buprenorphine prescribing and referral practices in US prison systems: Results from a nationwide survey. *Drug Alcohol Depend.* 2009; 105:83–88. [PubMed: 19625142]
- Pfeffer, J. A resource dependence perspective on intercorporate relations. In: Mizruchi, MS.; Schwartz, M., editors. *Intercorporate Relations: The Structural Analysis of Business*. Cambridge University Press; Cambridge, England: 1987. p. 25-55.
- Rich JD, Boutwell AE, Shield DC, et al. Attitudes and practices regarding the use of methadone in US State and federal prisons. *J Urban Health.* 2005; 82:411–419. [PubMed: 15917502]
- Rieckmann T, Kovas AE, Rutkowski BA. Adoption of medications in substance abuse treatment: Priorities and strategies of single state authorities. *J Psychoactive Drugs.* 2010; (Supplement 6): 227–238. [PubMed: 21138199]
- Rieckmann TR, Kovas AE, Fussell HE, et al. Implementation of evidence-based practices for treatment of alcohol and drug disorders: The role of the state authority. *J Behav Health Serv Res.* 2009; 36:407–419. [PubMed: 18543111]
- Roman PM, Abraham AJ, Knudsen HK. Using medication-assisted treatment for substance use disorders: Evidence of barriers and facilitators of implementation. *Addict Behav.* 2011; 36:584–589. [PubMed: 21377275]
- Roman PM, Abraham AJ, Rothrauff TC, et al. A longitudinal study of organizational formation, innovation adoption, and dissemination activities within the National Drug Abuse Treatment Clinical Trials Network. *J Subst Abuse Treat.* 2010; 38:S44–52. [PubMed: 20307795]
- Royston P. Multiple imputation of missing values. *Stata J.* 2004; 4:227–241.
- Royston P. Multiple imputation of missing values: Update. *Stata J.* 2005; 5:188–201.
- Royston P. Multiple imputation of missing values: Update of ice. *Stata J.* 2005; 5:527–536.
- SAMHSA. [Accessed April 23, 2012] Treatment Episode Data Set (TEDS): 1992-2000 National Admissions to Substance Abuse Treatment Services. 2002. http://www.dasis.samhsa.gov/teds00/TEDS_2k_index.htm
- Savage SA, Abraham AJ, Knudsen HK, et al. Timing of buprenorphine adoption by privately funded substance abuse treatment programs: The role of institutional and resource-based interorganizational linkages. *J Subst Abuse Treat.* 2012; 42:16–24. [PubMed: 21831565]
- Saxon AJ, McCarty D. Challenges in the adoption of new pharmacotherapeutics for addiction to alcohol and other drugs. *Pharmacol Ther.* 2005; 108:119–128. [PubMed: 16055196]
- Smith-Rohrberg D, Bruce RD, Altice FL. Review of corrections-based therapy for opiate-dependent patients: Implications for buprenorphine treatment among correctional populations. *J Drug Issues.* 2004; 34:451–480.
- Stewart MT, Horgan CM. Health services and financing of treatment. *Alcohol Res Health.* 2011; 33:389–394.
- Terry-McElrath YM, Chiqui JF, McBride DC. Factors related to Medicaid payment acceptance at outpatient substance abuse treatment programs. *Health Serv Res.* 2011; 46:632–653. [PubMed: 21105870]
- Thomas CP, Garnick DW, Horgan WM, et al. Advancing performance measure for use of medications in substance abuse treatment. *J Subst Abuse Treat.* 2011; 40:35–43. [PubMed: 20934836]
- Thomas CP, Reif S, Haq S, et al. Use of buprenorphine for addiction treatment: Perspectives of addiction specialists and general psychiatrists. *Psychiatr Serv.* 2008; 59:909–916. [PubMed: 18678689]
- Wallack SS, Thomas CP, Martin TC, et al. Substance abuse treatment organizations as mediators of social policy: Slowing the adoption of a congressionally approved medication. *J Behav Health Serv Res.* 2010; 37:64–78. [PubMed: 18668369]

TABLE 1**Descriptive Statistics of Organizational Characteristics of Treatment Programs within the Clinical Trials Network**

Variable	Mean (SD) or % (N)
Implementation of medication-assisted treatment (MAT) for opioid use disorders (OUDs)	9.56 (24.06)
<i>Financial Factors</i>	
% past-fiscal year revenues from Medicaid	18.99 (25.17)
% past-fiscal year revenues from private insurance	12.81 (22.26)
% past-fiscal year revenues from criminal justice system	4.12 (13.95)
% past-fiscal year revenues from Federal block grant	11.18 (22.00)
% past-fiscal year revenues from state government (non-block grant)	23.06 (31.97)
% past-fiscal year revenues from county government	9.06 (17.69)
<i>Structural Characteristics</i>	
Hospital-based	12.0% (18)
For-profit	8.6% (13)
Outpatient-only	51.0% (76)
<i>Workforce Characteristics</i>	
Number of physicians on staff	0.96 (1.42)
% counselors with Master's-level degree or higher	46.90 (36.04)
% counselors personally in recovery	39.05 (34.04)

Note: Descriptive statistics reflect available data prior to imputation; sample sizes ranged from 137 to 152 treatment centers.

TABLE 2

Unadjusted Negative Binomial Regression Models of Financial Factors and Implementation of Medication-Assisted Treatment for Opioid Use Disorders

	Unadjusted b (Robust SE)
% past-fiscal year revenues from Medicaid	.024 (.011) *
% past-fiscal year revenues from private insurance	.024 (.014) +
% past-fiscal year revenues from criminal justice system	-.061 (.024) *
% past-fiscal year revenues from Federal block grant	-.007 (.011)
% past-fiscal year revenues from state government (i.e., not the Federal block grant)	-.015 (.012)
% past-fiscal year revenues from county government	-.023 (.011) *

Notes: b = unstandardized coefficient. SE = standard error. Models adjust for the nesting of treatment programs within 70 organizations.

+ p<.10,

* p<.05 (two-tailed tests).

TABLE 3

Multivariate Negative Binomial Regression Models of Implementation of Medication-Assisted Treatment for Opioid Use Disorders

	Model 1 b (Robust SE)	Model 2 b (Robust SE)	Model 3 b (Robust SE)
% past-fiscal year revenues from Medicaid	.028 (.014) [*]	—	—
% past-fiscal year revenues from criminal justice system	—	−.047 (.023) [*]	—
% past-fiscal year revenues from county government	—	—	−.025 (.145) ⁺
Hospital-based (vs. non-hospital)	1.600 (.620) [*]	1.975 (.757) ^{**}	2.615 (.973) ^{**}
For-profit (vs. non-profit)	−2.321 (.911) [*]	−1.730 (.889) ⁺	−2.290 (1.095) [*]
Outpatient-only (vs. not outpatient-only)	−1.225 (.940)	−.559 (.700)	−.260 (.817)
Number of physicians on staff	.404 (.235) ⁺	.437 (.205) [*]	.518 (.244) [*]
% counselors with Master's-level degree or higher	.028 (.012) [*]	.028 (.011) [*]	.021 (.013) ⁺
% counselors personally in recovery	−.000 (.009)	.001 (.009)	.001 (.009)
Constant	−.158 (.957)	.131 (.863)	.226 (.869)
Alpha	13.992 (3.658)	14.410 (3.810)	14.403 (3.808)

Notes: b = unstandardized coefficient. SE = standard error. Models adjust for the nesting of treatment programs within 70 organizations.

⁺ p<.10,

^{*} p<.05,

^{**} p<.01