Genome Sequence of *Exiguobacterium antarcticum* B7, Isolated from a Biofilm in Ginger Lake, King George Island, Antarctica

Adriana Ribeiro Carneiro,^a Rommel Thiago Jucá Ramos,^a Hivana Dall'Agnol,^{a,c} Anne Cybelle Pinto,^b Siomar de Castro Soares,^b Anderson Rodrigues Santos,^b Luis Carlos Guimarães,^b Sintia Silva Almeida,^b Rafael Azevedo Baraúna,^a Diego Assis das Graças,^a Luciano Chaves Franco,^a Amjad Ali,^b Syed Shah Hassan,^b Catarina Isabel P. Nunes,^c Maria Silvanira Barbosa,^a Karina Kelly Fiaux,^b Flávia Figueira Aburjaile,^b Eudes Guilherme Vieira Barbosa,^b Syeda Marriam Bakhtiar,^b Daniella Vilela,^d Felipe Nóbrega,^d Adriana Lopes dos Santos,^d Marta Sofia P. Carepo,^c Vasco Azevedo,^b Maria Paula Cruz Schneider,^a Vivian Helena Pellizari,^d and Artur Silva^a

Institute of Biological Sciences, Federal University of Pará, Belém, PA, Brazil^a; Institute of Biological Sciences, Federal University of Minas Gerais, Belo Horizonte, MG, Brazil^b; REQUIMTE/CQFB, Department of Chemistry at the Faculty of Science and Technology, New University of Lisbon, Caparica, Portugal^c; and Laboratory of Ecology of Microorganisms, IO, University of São Paulo, São Paulo, Brazil^d

Exiguobacterium antarcticum is a psychotropic bacterium isolated for the first time from microbial mats of Lake Fryxell in Antarctica. Many organisms of the genus *Exiguobacterium* are extremophiles and have properties of biotechnological interest, e.g., the capacity to adapt to cold, which make this genus a target for discovering new enzymes, such as lipases and proteases, in addition to improving our understanding of the mechanisms of adaptation and survival at low temperatures. This study presents the genome of *E. antarcticum* B7, isolated from a biofilm sample of Ginger Lake on King George Island, Antarctic peninsula.

The genus *Exiguobacterium* consists of many species isolated from diverse habitats over a wide temperature range (−12 to 55°C), such as glacial ice, hot springs, plant rhizospheres, Siberian permafrost, tropical soils, and temperate soils (11). These species' adaptation to cold requires cellular modifications that may have possible industrial applications. Thermal acclimation proteins and enzymes, such as proteases, lipases, and cellulases, can be used as follows: in the food industry as food additives and flavoring agents, as biosensors, in environmental bioremediation, and in the pharmaceutical industry, among others (5, 9).

Exiguobacterium antarcticum is a Gram-positive bacterium that is mobile by peritrichous flagella but does not form spores and exhibits bright orange, convex colonies with varied sizes and shapes in bacillary morphology when grown on tryptone soy agar (TSA) at 25°C (3). This organism was isolated for the first time from microbial mats of Lake Fryxell, Antarctica (1) and has been named strain H2^T (DSM 14480^T; EMBL accession no. AJ297437). Moreover, studies of the biodiversity and biogeography of the genus *Exiguobacterium*, such as analysis by pulsed-field gel electrophoresis (PFGE), have contributed to classifying this species and estimating that the genome of strain H2^T is approximately 2,550,800 bp according to the PFGE results (14).

Thus, we present the first complete genome of *E. antarcticum* B7, isolated from a microbial biofilm at Ginger Lake, located on King George Island, Antarctic peninsula. This strain was subjected to two sequence analyses using two methods to construct genomic libraries on the SOLiD platform (mate paired and fragments), which generated 89,195,902 reads 25 bp in size and 71,239,854 reads 50 bp in size, respectively. The quality control of the raw data was assessed using the Quality assessment program (10), and a quality filter with Phred 20 was applied.

The filtered and corrected data set totaled 61,981,355 and 39,811,567 reads for the mate-paired and fragment libraries, corresponding to genomic coverages of 499× and 642×, respectively, considering the 3.1-Mb size of the *Exiguobacterium sibiricum* 255-15 (CP001022) reference genome.

To assemble the data, we adopted a *de novo* approach using the Velvet (16) and Edena programs (6). The best assembly results with Velvet and Edena were grouped for a total of 21,878 contigs, which were ordered and aligned using the reference genome through CLC Genomics Workbench 4.7.1. A scaffold with 1,187 contiguous sequences, totaling 2,665,352 bases, was thus obtained. The remaining gaps were closed by recursive alignments with unmapped contigs and short reads not used in the *de novo* assembly, followed by manual strand editing (2, 13).

For the functional genome annotation, open reading frame (ORF) prediction was obtained using the Glimmer program (<http://www.cbcb.umd.edu/software/glimmer/>). The RNAmmer program (7) was used to predict rRNAs. The tRNAscan-SE program (8) was used to identify tRNAs, and the RFAM database (4) was searched to predict other noncoding RNAs (ncRNAs). Domains and protein families were predicted with InterProScan software (15), and the annotations were manually curated in the Artemis program (12).

The *E. antarcticum* B7 genome has a size of 2,815,863 bp, a GC content of 47.48%, 2,772 coding sequences (CDSs), 9 rRNA operons, 66 tRNAs, 52 ncRNAs, and 76 pseudogenes.

Nucleotide sequence accession number. The genome sequence obtained in this study has been deposited in the GenBank database under accession number CP003063.1.

ACKNOWLEDGMENTS

M.P.C.S., A.S., V.A., A.R.C., S.D.C.S., S.S.A., A.S., F.F.A., and E.G.V.B. were supported by the National Council for Scientific and Technological Development (Conselho Nacional de Desenvolvimento Científico e Tec-

Received 27 September 2012 Accepted 1 October 2012

Address correspondence to Artur Silva, asilva@ufpa.br.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

doi:10.1128/JB.01791-12

nológico—CNPq). R.T.J.R., A.R.S., L.C.G., R.A.B., D.A.D.G., H.D., and A.A. were supported by the Brazilian Federal Agency for the Support and Evaluation of Graduate Education (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior—CAPES). This work was part of the Paraense Network of Genomics and Proteomics (Rede Paraense de Genômica e Proteômica), supported by the Paraense Amazonia Foundation (Fundação Amazônia Paraense—FAPESPA), Amazon Center of Excellence in Genomics of Microorganisms (Núcleo Amazônico de Excelência em Genômica de Microorganismos)—Centers of Excellence Support Program (Programa de apoio a Núcleo de Excelência) Pronex/CNPq/FAPESPA, the National Program for Academic Cooperation (Programa Nacional de Cooperação Acadêmica) PROCAD/CAPES, the Studies and Projects Funding Agency (Financiadora de Estudos e Projetos—FINEP), and the Minas Gerais Research Fund (Fundação de Amparo à Pesquisa do estado de Minas Gerais—FAPEMIG). This study is also part of a CAPES-FCT (Science and Technology Fund—Fundação para a Ciência e a Tecnologia) international cooperation with the Department of Chemistry (Departamento de Química) of the New University of Lisbon Faculty of Sciences and Technology (Faculdade de Ciências e Tecnologia da Universidade Nova de Lisboa).

REFERENCES

1. Brambilla E, Hippe H, Hagelstein A, Tindall BJ, Stackebrandt E. 2001. 16S rDNA diversity of cultured and uncultured prokaryotes of a mat sample from Lake Fryxell, McMurdo Dry Valleys, Antarctica. *Extremophiles* 5:23–33.
2. Cerdeira LT, et al. 2011. Rapid hybrid de novo assembly of a microbial genome using only short reads: *Corynebacterium pseudotuberculosis* I19 as a case study. *J. Microbiol. Methods* 86:218–223.
3. Fruhling A, Schumann P, Hippe H, Straubler B, Stackebrandt E. 2002. *Exiguobacterium undae* sp. nov. and *Exiguobacterium antarcticum* sp. nov. *Int. J. Syst. Evol. Microbiol.* 52:1171–1176.
4. Gardner PP, et al. 2011. Rfam: Wikipedia, clans and the “decimal” release. *Nucleic Acids Res.* 39:D141–D145.
5. Gounot AM. 1991. Bacterial life at low temperature: physiological aspects and biotechnological implications. *J. Appl. Bacteriol.* 71:386–397.
6. Hernandez D, Francois P, Farinelli L, Osteras M, Schrenzel J. 2008. De novo bacterial genome sequencing: millions of very short reads assembled on a desktop computer. *Genome Res.* 18:802–809.
7. Lagesen K, et al. 2007. RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res.* 35:3100–3108.
8. Lowe TM, Eddy SR. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res.* 25:955–964.
9. Margesin R, Feller G. 2010. Biotechnological applications of psychrophiles. *Environ. Technol.* 31:835–844.
10. Ramos RT, et al. 2011. Analysis of quality raw data of second generation sequencers with Quality Assessment Software. *BMC Res. Notes* 4:130.
11. Rodrigues DF, Tiedje JM. 2007. Multi-locus real-time PCR for quantitation of bacteria in the environment reveals *Exiguobacterium* to be prevalent in permafrost. *FEMS Microbiol. Ecol.* 59:489–499.
12. Rutherford K, et al. 2000. Artemis: sequence visualization and annotation. *Bioinformatics* 16:944–945.
13. Tsai IJ, Otto TD, Berriman M. 2010. Improving draft assemblies by iterative mapping and assembly of short reads to eliminate gaps. *Genome Biol.* 11:R41.
14. Vishnivetskaya TA, Kathariou S, Tiedje JM. 2009. The *Exiguobacterium* genus: biodiversity and biogeography. *Extremophiles* 13:541–555.
15. Zdobnov EM, Apweiler R. 2001. InterProScan—an integration platform for the signature-recognition methods in InterPro. *Bioinformatics* 17:847–848.
16. Zerbino DR, Birney E. 2008. Velvet: algorithms for de novo short read assembly using de Bruijn graphs. *Genome Res.* 18:821–829.