Published in final edited form as:

Org Lett. 2012 November 16; 14(22): 5808–5810. doi:10.1021/ol3024665.

Biomimetic Synthesis and Biological Evaluation of Aplidiopsamine A

Joseph D. Panarese and Craig W. Lindsley

Departments of Chemistry and Pharmacology, Vanderbilt University, Nashville, Tennessee 37232.

Abstract

The first total synthesis of Aplidiopsamine A, a rare 3*H*-pyrrolo[2,3-*c*]quinolone alkaloid from the *Aplidiopsis confluata*, has been achieved following the proposed biosynthesis. This biomimetic synthesis requires only 5 steps and proceeds in 20.8% overall yield. Biological evaluation across large panels of discrete molecular targets identified that Aplidiopsamine A is a highly selective PDE4 inhibitor, a target for numerous CNS disorders.

The tricyclic 3*H*-pyrrolo[2,3-*c*]quinoline ring system is a rare moiety found only in a few marine natural products.^{1,2} The first members of this structural class were the marinoquinolines A-F (**1-5**), isolated from multiple species of bacteria and found to possess antibacterial, antifungal and acetylcholinesterase activities (Figure 1).^{1,2} Several synthetic approaches to marinoquinolines A-C and E (**1-3**, **5**) have been reported, relying on either a key Morgen-Walls or Pictet-Spengler reaction.^{3,4} Recently, aplidiopsamine A (**6**) was isolated from the Australian ascidian, *Aplidiopsis confluata*, and represents the first example of the 3*H*-pyrrolo[2,3-*c*]quinoline moiety linked to an adenine via a methylene bridge, a rare non glycoside adenine conjugate. Thus, **6** represents only the second report of a marine organism producing the 3*H*-pyrrolo[2,3-*c*]quinoline ring system; importantly, **6** was found to be a potent antimalarial agent, without toxicity to healthy cells, further providing support for synthesis.⁵

Carroll and co-workers developed a convergent biosynthetic proposal for **6** from adenine **7** and tryptophan **10** (Scheme 1). Here, **9** is generated from the reaction of adenine and hydroxyacetic acid. Pyrrole **11**, itself a natural product isolated from *R. thailandica*, was postulated to be a dead-end product in the biosynthesis of pyrrolnitrin from tryptophan. Condensation of **9** and **11** would ultimately lead to aplidiopsamine A, via a Bischler-Napieralski-like cyclization and dehydration sequence.⁵

Correspondence to: Craig W. Lindsley.

craig.lindsley@vanderbilt.edu.

Supporting Information Available: Experimental procedures, characterization data, and ¹H and ¹³C NMR spectra for all new compounds. This material is available free of charge via the internet at <http://pubs.acs.org>.

The authors declare no competing financial interest.

Therefore, we elected to pursue a biomimetic approach for the first total synthesis of aplidiopsamine A, and attempt the condensation of a suitable synthetic surrogate of **9** and pyrrole **11** to access **12**. Interestingly, Correia synthesized **11**, en route to marinoquinolines A-C and E, from advanced materials in five synthetic steps in 47.5% overall yield.⁴ We envisioned a more expedited route, and were gratified that a two step sequence from commercial reagents produced **11**. In the event, aniline **14** smoothly underwent a Suzuki coupling with boronate ester **13** to deliver TIPS protected pyrrole **15**, which upon basic hydrolysis generated **11** in 96% yield over the two steps (Scheme 2).⁶ Acylation with α -bromo acetyl bromide provided **16** in 86% yield. Selective *N*-alkylation at the 9- versus 7-position of adenine to afford **12** was achieved via a hard deprotonation with NaH in 56% yield. Softer approaches with cesium carbonate resulted in 2:1 regioisomeric mixtures and poor conversion. With **12** in hand, we were now poised to perform the biomimetic condensation via a Bischler-Napieralski-type cyclization. Classical variants of this cyclization/dehydration sequence⁷ employ TFA, POCl₃ or other Lewis acids; however, in the presence of the adenine moiety, these conditions either failed, led to intractable gums or poor conversion (less than 10% yields). Ultimately, we found that 4 M HCl/dioxanes under microwave irradiation (130 °C, 10 min) facilitated the reaction sequence to deliver, for the first time, aplidiopsamine A in 45% yield.⁶ Overall, the biomimetic synthesis of **6** required five steps and proceeded in 20.8% overall yield, an ideal route to prepare unnatural analogs. The synthetic **6** exhibited physical and spectroscopic data identical to that of the natural aplidiopsamine A.^{5,6}

With large quantities of **6** in hand, we elected to further profile **6** against a larger panel of discrete molecular targets of therapeutic significance beyond antimalarial activity, as **6** was shown to not be cytotoxic. Indeed, we have previously elucidated intriguing activities for a number of marine alkaloids at CNS targets with unprecedented selectivities amongst highly conserved receptor families.⁸⁻¹¹ As **6** possesses the basic pharmacophore (H-bond donor/acceptor triad) of many known ATP-competitive kinase inhibitors,¹² we profiled synthetic **6** in a KINOMEScan panel against 97 kinases (both wild-type and mutants) at a 10 μ M concentration.^{6,13} Despite the presence of a known pharmacophore,¹² **6** was uniformly inactive; however, this can also be viewed as possessing very clean ancillary pharmacology. In parallel, we also profiled **6** in a Lead Profiling screen (a radio ligand binding panel at a 10 μ M concentration) against 68 GPCRs, ion channels and transporters.¹⁴ Once again, **6** displayed very clean ancillary pharmacology, possessing activity at only two targets: 5-HT_{2B} (62% @ 10 μ M) and PDE4/Rolipram (74% @ 10 μ M). When full concentration-response-curves (CRCs) were obtained for **6**, the 5-HT_{2B} K_i was weak at ~ 10 μ M, but **6** proved to be a moderately potent PDE4 inhibitor (K_i = 1.2 μ M, IC₅₀ = 3.3 μ M).^{6,14} PDE4 is a high profile target for antidepressant, antipsychotic and neuroprotective drug development; moreover, **6** represents a fundamentally new PDE4 inhibitor chemotype.^{15,16}

With the identification of a therapeutically relevant target for **6**, the need emerged to develop rapid chemistry to enable the synthesis of unnatural analogs, and ideally, from a common intermediate. Application of our optimized cyclization/dehydration conditions with the advanced α -bromo intermediate **16** led directly to the 3*H*-pyrrolo[2,3-*c*]quinoline derivative **17**, wherein the reaction conditions with HCl generated the corresponding chloromethyl congener. Notably, high reaction temperatures typically employed in Bischler-Napieralski reactions were avoided as the condensation was found to proceed at room temperature. Not only did **17** prove to be an ideal common intermediate for the synthesis of unnatural analogs of **6**, compounds **18a-18c**, in high yields (82-96%), but also for the synthesis of marinoquinoline A (**1**) when the nucleophile was a hydride source, LiAlH₄. This approach affords the natural product **1** in only 5 steps and in 52% overall yield, representing a highly expeditious route.^{3,4}

With unnatural analogs of **6** in hand, we evaluated them against both rat PDE4 (Rolipram) and human PDE4 to discern structure-activity-relationships (SAR). As shown in Table 1, several of the unnatural analogs possessed PDE4 activity, but were weaker than **6** (rat K_s 4.6 to 25 μ M and IC_{50} s 13 to 71 μ M). All proved to be weaker functional antagonists on human PDE4 than rat. Interestingly, marinoquinoline A (**1**), lacking the amino methyl moiety at C9, was inactive as was the Bischler-Napieralski acyclic substrate **12**, devoid of the 3*H*-pyrrolo[2,3-*c*]quinoline ring system. Overall, this suggests that the tricyclic 3*H*-pyrrolo[2,3-*c*]quinoline ring system with a C9 amino methyl moiety is essential for PDE4 activity, and this chemotype represents a fundamentally new PDE4 pharmacophore.

In summary, we have completed the first total synthesis of aplidiopsamine A (**6**) following the proposed biosynthetic pathway in five steps and with a 20.8% overall yield. Extensive biological profiling identified **6** as a moderately potent ligand for, and inhibitor of, PDE4, an important target for CNS drug discovery. These data further argue for natural products as a source of novel chemotypes to initiate drug discovery efforts. Furthermore, we developed chemistry to rapidly access unnatural analogs of **6**, and in the course synthesized marinoquinoline A (**1**) in only five steps and in 52% overall yield. SAR within this novel class of PDE4 ligands demonstrated that both the tricyclic 3*H*-pyrrolo[2,3-*c*]quinoline ring system and a C9 amino methyl moiety is required for PDE4 activity. Further refinements, DMPK analysis and additional unnatural analog synthesis are underway and will be reported in due course.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported, in part, by the Department of Pharmacology and William K. Warren, Jr.. Funding for the NMR instrumentation was provided in part by a grant from NIH (S10 RR019022).

References

1. Sangoi Y, Sakuleko O, Yuenyongsawad S, Kanjana-opas A, Ingkaninan K, Plubrukan A, Suwanborirux K. *Mar. Drugs*. 2008; 6:578–586. [PubMed: 19172195]
2. Okanya PW, Mohr KI, Gerth K, Jansen R, Muller R. *J. Nat. Prod.* 2011; 74:603–608. [PubMed: 21456549]
3. Ni L, Li Z, Wu F, Xu J, Wu X, Kong L, Yao H. *Tetrahedron Lett.* 2012; 53:1271–1274.
4. Schwalm CS, Correia CRD. *Tetrahedron Lett.* 2012; 53:4836–4840.
5. Carroll AJ, Duffy S, Avery VM. *J. Org. Chem.* 2010; 75:8291–8294. [PubMed: 21043526]
6. See Supporting Information for full details.
7. a Kennedy JP, Breininger ML, Lindsley CW. *Tetrahedron Lett.* 2009; 50:7067–7069. b Capilla AS, Romero M, Pujol MD, Caignard DH, Renard P. *Tetrahedron*. 2001; 57:8297–8303.
8. Kennedy JP, Brogan JT, Lindsley CW. *J. Nat. Prod.* 2008; 71:1783–1788. [PubMed: 18800848]
9. Aldrich LN, Stoops SL, Crews BC, Marnett LM, Lindsley CW. *Bioorg. Med. Chem. Lett.* 2010; 20:5207–5211. [PubMed: 20655217]
10. Brogan JT, Stoops SL, Crews BC, Marnett LM, Lindsley CW. *ACS Chem. Neurosci.* 2011; 2:633–639. [PubMed: 22247792]
11. Brogan JT, Stoops SL, Lindsley CW. *ACS Chem. Neurosci.* 2012; 3:658–664. [PubMed: 23019492]
12. Lewis JA, Lebois EP, Lindsley CW. *Curr. Opin. Chem. Biol.* 2008; 12:269–279. [PubMed: 18342020]
13. For information of the KIONMEscan panel assay, see: www.discoverx.com

14. For information on the Lead Profiling screen see: www.ricerca.com.
15. Pages L, Gavalda A, Lehner MD. Expert Opin. Ther. Pat. 2009; 19:1501–1519. [PubMed: 19832118]
16. Burgin AB, Magnusson OT, Witte P, Bjornsson JM, Thorsteindottir S, Hage T, Kiselyov AS, Stewart LJ, Gurnet ME. Nat. Biotech. 2010; 28:63–70.

Figure 1. Structures of the marinoquinolines A-F (**1-5**) and aplidiopsamine A (**6**), marine natural products possessing the rare 3*H*-pyrrolo[2,3-*c*]quinoline ring system (shown in red).

Scheme 1.
Proposed biosynthesis of Aplidiopsamine (6).

Scheme 2.
Biomimetic total synthesis of Aplidiopsamine (**6**).

Scheme 3.

Synthesis of Marinoquinoline A (**1**) and unnatural analogs of **6** from a common intermediate.

Table 1Phosphodiesterase PDE4 activities of **6** and unnatural analogs.

Cmpd	rat Rolipram ^a		human PDE4 ^b
	K _i (μM)	IC ₅₀ (μM)	IC ₅₀ (μM)
6	1.2	3.3	6.1
1	>100	>100	>100
18a	25.6	71.7	31.8
18b	4.7	13.1	72.9
18c	5.0	14.0	>100
12	>100	>100	>100

^aWistar rat brain, 1.8 nM [³H] rolipram, 1% DMSO, incubation for 60 min at 4 °C in 50 mM Tris-HCl, pH 7.4.^bHuman U937 cells, 1.10 μM [³H]cAMP+cAMP, incubation for 20 min at 25 °C, 50 mM Tris-HCl, pH 7.5, 5 mM MgCl₂, quantitation of [³H]adenosine.¹⁴