Published in final edited form as:

Psychiatry Res. 2012 December 30; 200(2-3): 173–176. doi:10.1016/j.psychres.2012.06.014.

Neurocognitive Predictors of Source Monitoring in Schizophrenia

Mohammed K Shakeel* and Nancy M Docherty

Schizophrenia Research Lab, Department of Psychology, Kent State University, Ohio, USA

Abstract

Rationale—Source monitoring is a metacognitive process involved in making judgments about the origin of information by recruiting cognitive processes. Deficits in source monitoring have been linked to positive symptoms of schizophrenia. We investigated whether certain neurocognitive functions—specifically attention, working memory, and organizational sequencing—were associated with source monitoring in a sample of schizophrenia patients.

Methods—Attention (Auditory Continuous Performance Test), organizational sequencing (Trail Making Test B-A), working memory (Digits Backward), and internal source monitoring (SM) were assessed in 45 outpatients diagnosed with schizophrenia or schizoaffective disorder.

Results—Standard multiple regression analysis showed attention, working memory and organizational sequencing together predicted source monitoring. Organizational sequencing was the only significant individual predictor, with better organizational sequencing ability being associated with better source monitoring. Hierarchical regression analysis showed that working memory by itself did not result in a significant predictive model of source monitoring, but adding organizational sequencing led to a significant change from the working memory model and resulted in a significant overall model, accounting for 26% of the variance in source monitoring.

Conclusions—Neurocognitive functions were associated with source monitoring in schizophrenia. Organizational sequencing, which requires an awareness of self-generated actions, predicted source monitoring performance even after controlling for working memory.

Keywords

Source Monitoring; Neurocognition; Schizophrenia; Attention; Working memory; Organizational Sequencing

1. Introduction

A fundamental idea underlying source monitoring theory is that memories, images, and feelings do not come with abstract labels specifying their source for us; rather, we attribute information in memory to particular origins or sources using heuristic and systematic decision processes, based on evaluation of cues such as contextual and sensory/perceptual information, cognitive operations and semantic detail (Johnson et al., 1993; Johnson and

© 2012 Elsevier Ireland Ltd. All rights reserved.

*Corresponding Author: Schizophrenia Research Lab, Department of Psychology, Kent State University, P.O. Box 5190, Kent, Ohio 44242-0001, USA. Tel.: +1-330-672-7671. mshakeel@kent.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Raye, 2000; Lindsay, 2008). Source monitoring may involve discriminating between externally derived sources of information—external source monitoring (e.g. determining which of two people made a statement); between internally generated sources—internal source monitoring (e.g. determining whether one said, or only thought about, a statement); or discriminating internal from external sources—reality monitoring (e.g. determining whether one thought of, or heard, the statement) (see Johnson et al., 1993). Cognitive functions can affect the evaluative processes involved in making these judgments (e.g. Hashtroudi et al., 1994; also see Lindsay, 2008).

Source monitoring deficits have been observed in schizophrenia (see Mitchell and Johnson, 2009). More specifically, studies have shown that schizophrenia patients with symptoms like delusions and hallucinations have greater source monitoring deficits than patients without these symptoms and healthy control subjects (Brunelin et al., 2007; Keefe et al., 2002). It has therefore been theorized that poor source monitoring may underlie some of the positive symptoms in schizophrenia. For instance, an inability to monitor self-generated thoughts and actions can result in the conclusion that they come from an external source, which may form the basis for hallucinations and some kinds of delusions (Brebion et al., 2000; Brunelin et al., 2007; Keefe et al., 2002).

The current study investigated whether specific facets of neurocognitive functioning are associated with source monitoring in schizophrenia. Lindsay (2008) suggested that manipulations that impair encoding or retrieval of source-specifying information can affect source monitoring. Decreased attention, for instance, may interfere with binding of perceptual detail with content and reduce source accuracy (Hashtroudi et al., 1994), while working memory and organizational sequencing could be related esp. to systematic processing of source information. Working memory may help in deliberate reasoning about the event with respect to additional information, such as discovering inconsistencies among memories or evaluating the plausibility of a target memory, while organizational sequencing may facilitate source judgments by helping in the evaluation of supporting memories, such as remembering other events which took place before or after the target memory (for discussion of systematic processes, see Johnson and Raye, 2000). We hypothesized that attention, working memory and organizational sequencing would predict source monitoring performance in schizophrenia patients.

Although internal source monitoring (Franck et al., 2000; Henquet et al., 2005; Nienow and Docherty, 2004) and reality monitoring deficits (Anselmetti et al., 2007; Brebion et al., 2000; Vinogradov et al., 1997) have been observed in schizophrenia, external source monitoring deficits have generally not been found, unlike in bipolar disorder (Fisher et al., 2008; Harvey, 1985). Also, source monitoring deficits in schizophrenia are usually characterized by patients having a greater tendency to misidentify self-generated thoughts as actions (Franck et al., 2000; Henquet et al., 2005; but see Franck et al., 2001), and to attribute self-generated information to external sources (Anselmetti et al., 2007; Brebion et al., 2000; Vinogradov et al., 1997), which has led some authors to propose that source monitoring in schizophrenia may be characterized by a specific inability to identify and process self generated information (Fisher et al., 2008; Keefe et al., 2002). Although tests of both organizational sequencing and working memory involve self-generation of information, tests that measure the former place greater demands on sequencing and temporal structuring of responses (Wolwer & Gaebel, 2002), which requires greater discrimination of responses that have been implemented (actions) from those held in mind during task performance. We therefore hypothesized that performance on an organizational sequencing task would predict source monitoring even after controlling for general working memory.

2. Methods

2.1. Sample

The sample consisted of 45 outpatients with a DSM-IV diagnosis of schizophrenia or schizoaffective disorder who were part of a larger project examining emotion, neurocognition, and communication disturbance (see Docherty et al., 2003). The Schedule for Affective Disorders and Schizophrenia—Lifetime Version (Spitzer and Endicott, 1978), adapted slightly for use with DSM-IV criteria, was used to arrive at the DSM diagnosis. A clinical psychologist who had acceptable inter-rater reliability with another research group ($k = .88$) arrived at the diagnoses based on information gathered from the interview and clinical records. Patients were excluded if they reported a history suggestive of possible organic brain damage or met DSM-IV criteria for substance abuse. All patients (except one) were receiving psychiatric medications at time of the study.

The study was carried out with due approval of the Kent State University Institutional Review Board, and informed consent was obtained from all participants prior to enrollment in the present study, thus conforming to the ethical standards laid down in the 1964 Declaration of Helsinki. Table 1 shows descriptive characteristics of the sample.

2.2. Measures

Attention, working memory and organizational sequencing tests were administered at the initial sessions, and a source monitoring task was administered in a follow-up session nine months later.

2.2.1. Source Monitoring task—An internal source monitoring (SM) task we previously developed (see Nienow and Docherty, 2004) was used for the present study. Briefly, participants have to generate single word responses to 16 incomplete statements. The statements are ones for which there is only one appropriate response (e.g. “The first month of the year is _____”). On half the trials the participants only think of the answer to themselves, and on the other half they say the answer out loud. Immediately after this, they are given a source recognition sheet with all the responses they said, thought, and with eight new words (24 words total), and they have to identify whether they had said or thought each word on the list, or if the word was new. The task was piloted on 39 college students, who showed 99% agreement on the words they used to complete the statements. The total correct score was used for the present analysis.

2.2.2. Attention—The continuous performance test is a measure of sustained attention (see Lezak, 2004). We administered the auditory continuous performance test (CPT-A; see Strub and Black, 1988) in which the participant listens to a 7 minute audiotape presentation of a quasi-random series of letters read at 1 second intervals, and has to respond every time the target letter 'A' is heard. The total correct score was used for the present analysis.

2.2.3. Working Memory—The Digits Backward (Digits-B), part of the Wechsler Adult Intelligence Scale—Revised (Wechsler, 1981), is a measure of working memory (Garlinghouse et al., 2010). The participant has to listen to a set of numbers and repeat it in the reverse order (see Lezak et al., 2004). The total correct score was used for the present analysis.

2.2.4. Organizational sequencing—We used part B of the Trail Making Test (Reitan and Davidson, 1974) as a measure of organizational sequencing ability. The participant has to connect letters and numbers in alternating order (1-A-2-B-3-C...etc.), as quickly as possible. We followed Reitan's (1958) revised procedure which requires the experimenter to

point out errors as they occur, resulting in the participant always completing the test without errors so scoring can be based on time alone. To arrive at a measure of organizational sequencing after controlling for psychomotor speed, score on part A (which only involves connecting numbers (1-2-3-4...)) was subtracted from part B. The resulting score (Trails B-A) was used.

2.3 Statistical Analysis

Statistical analyses were carried out using SPSS 16.0 (www.spss.com). Correlations were calculated between all variables. Standard multiple regression analysis was used to determine whether measures of attention, working memory and organizational sequencing together predicted source monitoring. Hierarchical multiple regression was used to determine whether organizational sequencing predicted source monitoring after controlling for working memory. A significance level of $p < .05$ (two-tailed) was employed for all analyses.

3. Results

3.1. Preliminary Analysis

Complete data were available for 39 participants for the standard regression and 45 participants for the hierarchical regression analysis. No multivariate outliers were found among the cases using $p < .001$ criterion for Mahalanobis distance. After CPT-A and Trails B-A were log-transformed to reduce skewness and improve homoscedasticity, all assumptions for multiple linear regression analysis were met. Means and standard deviations (non-transformed) for the variables are presented in Table 1, and correlations between them (transformed) are presented in Table 2.

3.2. Hypothesis tests

A standard multiple regression was performed with SM as criterion, and CPT-A, Digits-B, and Trails B-A as predictor variables. Table 3 shows regression coefficients for the analysis. R for the model was significant [$R = 0.53$; $F(3, 35) = 4.65$, $p < .01$] and indicates that 28% of the variance in SM was predicted by CPT-A, Digits-B, and Trails B-A ($R^2 = .28$). Trails B-A was the only significant individual predictor in this model [$t(35) = -3.42$, $p < .01$], with one standard deviation decrease in time taken for Trails B-A being associated with over half a standard deviation increase in SM performance ($\beta = -.53$).

Hierarchical regression was used to determine whether Trails B-A predicted SM after controlling for Digits-B. Table 4 shows regression coefficients for each step of the analysis. In Block 1, R for the model was not significant [$R = 0.19$; $F(1, 43) = 1.55$, $p > .05$] when Digits-B was the only predictor of SM. In Block 2, when Trails B-A was added as the predictor, it significantly improved the model R [$F\text{-change}(1, 42) = 12.47$, $p = .001$] and the overall model also became significant [$R = 0.51$; $F(2, 42) = 7.22$, $p = .01$], accounting for 26% of the variance in SM ($R^2 = .26$).

4. Discussion

In the present study we looked at cognitive predictors of source monitoring performance and found that attention, working memory and organizational sequencing together predicted source monitoring in schizophrenia patients. Organizational sequencing was a significant individual predictor in this model, while working memory and attention were not. Attention has been shown in non-psychiatric populations to be associated with source monitoring (Hashtroudi et al., 1994; Zaragoza and Lane, 1998) and a ceiling effect on the attention task (for 25% of the sample) may have partly contributed to its not being a significant individual

predictor in the present study, although our findings are generally in agreement with other studies which have not found a relation between attention and source monitoring deficits in schizophrenia (Ferchiou et al. 2010 ; but also see Brebion et al. 1996).

Organizational sequencing predicted source monitoring even after controlling for working memory.

The Trail Making Test is a complex measure which assesses psychomotor speed (Lezak et al., 2004), organizational sequencing (Levitt et al., 1995), and working memory (Garlinghouse et al., 2010), while digit span backward is a specific measure of working memory (Lezak et al., 2004). We controlled for psychomotor speed by subtracting Trails A from B, and controlled for the working memory by regressing out digit backward, and still found that Trails B-A predicted source monitoring. Although the digit-span backward test is also a measure of organization to some extent (Garlinghouse et al., 2010; Partington, 1949), the trail making test requires a much longer, ongoing response than the digit span task (participants took an average of 127.56 seconds ($SD = 51.49$) to complete Trail Making Test-B). Functionally, the organizational sequencing task requires the participant to not only hold the letter-number sequence in mind (which constitutes the working memory component) but also to actively monitor and distinguish the part of the sequence they have completed drawing (actions) from those they have not yet completed as they go along (see Levitt et al., 1995). Source monitoring is a metacognitive process that requires higher level integration of several basic cognitive processes (see e.g. Hashtroudi et al., 1994; Lindsay, 2008). We suggest that the cognitive mechanisms underlying the sense of awareness about past and ongoing self-generated events in organizational sequencing may also contribute to an individual's ability to retrospectively distinguish actions they have carried out ('said' words) from those they have only imagined ('thought' words) on the internal source monitoring task.

One limitation of our study is that the source-monitoring test provides three options ('said', 'heard', and 'new') simultaneously, which makes it more susceptible to response bias in patients (see Achim & Weiss, 2008). One way of arriving at a more accurate measure of discrimination would be to first provide participants with an old/new recognition test, and then examine source-monitoring accuracy (said/heard) for the items correctly recognized as old. Another limitation of this study is the time difference (9 months) between collection of data on neurocognitive variables, and on the source monitoring task. Previous studies have shown that source monitoring deficits in schizophrenia are stable and trait-like, and tend to persist over long periods (e.g., 8 months (Harvey et al., 1990), and 2 years (Vinogradov et al., 1997)) despite changes in medication and clinical status. Several studies also have shown that cognitive functions [including attention (Irani et al., 2012), working memory (Irani et al., 2012) and trail making test performance (Heaton et al., 2001)], tend to be stable over long periods of time (Heaton et al., 2001) and are not associated with duration of illness (Gold et al., 1999). Still, we cannot rule out that our results might have been affected to some degree by the fact that all the neurocognitive and metacognitive processes were not assessed simultaneously. If so, the effect would have been to diminish the associations.

Recent studies (Subramaniam et al., 2012) have shown that cognitive remediation programs that succeed in improving cognitive functions can also improve source monitoring as well as real-world functioning. There has been a gradual shift in focus of schizophrenia research from the psychotic symptoms, to the more enduring cognitive impairments (Green, 2010). Our findings suggest that neurocognitive deficits merit attention not only due to their association with functional disability (Green, 1996) but also because they may indirectly contribute to psychotic symptoms due to their influence on metacognitive processes.

Acknowledgments

The authors thank Dr. Maria S. Zaragoza for her comments on the manuscript.

Funding for this study was provided by NIMH Grants 1R15MH57151 AND 5R01-MH58783 and by Kent State University

References

- Achim AM, Weiss AP. No evidence for a differential deficit of reality monitoring in schizophrenia: A meta-analysis of the associative memory literature. *Cognitive Neuropsychiatry*. 2008; 13:369–384. [PubMed: 18781492]
- Anselmetti S, Cavallaro R, Bechi M, Angelone SM, Ermoli E, Cocchi F, Smeraldi E. Psychopathological and neuropsychological correlates of source monitoring impairment in schizophrenia. *Psychiatry Res*. 2007; 150:51–59. [PubMed: 17289157]
- Brébion G, Smith MJ, Gorman JM, Amador X. Reality monitoring failure in schizophrenia: The role of selective attention. *Schizophr. Res*. 1996; 22:173–180. [PubMed: 8958602]
- Brebion G, Amador X, David A, Malaspina D, Sharif Z, Gorman JM. Positive symptomatology and source-monitoring failure in schizophrenia--an analysis of symptom-specific effects. *Psychiatry Res*. 2000; 95:119–131. [PubMed: 10963798]
- Brunelin J, d'Amato T, Brun P, Bediou B, Kallel L, Senn M, Poulet E, Saoud M. Impaired verbal source monitoring in schizophrenia: an intermediate trait vulnerability marker? *Schizophr Res*. 2007; 89:287–292. [PubMed: 17029909]
- Docherty NM, Cohen AS, Nienow TM, Dinzeo TJ, Dangelmaier RE. Stability of formal thought disorder and referential communication disturbances in schizophrenia. *J. Abnorm. Psychol*. 2003; 112:469–475. [PubMed: 12943025]
- Ferchiou A, Schürhoff F, Bulzacka E, Leboyer M, Szöke A. Selective Attention Impairment in Schizophrenia: Can it Explain Source Monitoring Failure? *J. Nerv. Ment. Dis*. 2010; 198:779. [PubMed: 20921871]
- Fisher M, McCoy K, Poole JH, Vinogradov S. Self and other in schizophrenia: a cognitive neuroscience perspective. *Am. J. Psychiatry*. 2008; 165:1465–1472. [PubMed: 18708487]
- Franck N, Farrer C, Georgieff N, Marie-Cardine M, Dalery J, d'Amato T, Jeannerod M. Defective recognition of one's own actions in patients with schizophrenia. *Am. J. Psychiatry*. 2001; 158:454–459. [PubMed: 11229988]
- Franck N, Rouby P, Daprati E, Dalery J, Marie-Cardine M, Georgieff N. Confusion between silent and overt reading in schizophrenia. *Schizophr. Res*. 2000; 41:357–364. [PubMed: 10708345]
- Garlinghouse MA, Roth RM, Isquith PK, Flashman LA, Saykin AJ. Subjective rating of working memory is associated with frontal lobe volume in schizophrenia. *Schizophr. Res*. 2010; 120:71–75. [PubMed: 20303715]
- Gold S, Arndt S, Nopoulos P, O'Leary DS, Andreasen NC. Longitudinal study of cognitive function in first-episode and recent-onset schizophrenia. *Am. J. Psychiatry*. 1999; 156:1342–1348. [PubMed: 10484943]
- Green, M. Cognition and Daily Functioning in schizophrenia, in *Anonymous Neuropsychology of Everyday Functioning*. The Guilford Press; 2010. p. 441–456.
- Green MF. What are the functional consequences of neurocognitive deficits in schizophrenia? *Am. J. Psychiatry*. 1996; 153:321–330. [PubMed: 8610818]
- Harvey PD. Reality monitoring in mania and schizophrenia: The association of thought disorder and performance. *J. Nerv. Ment. Dis*. 1985; 173:67–73. [PubMed: 3968548]
- Harvey PD, Docherty NM, Serper MR, Rasmussen M. Cognitive deficits and thought disorder: II. An 8-month followup study. *Schizophr. Bull*. 1990; 16:147–156. [PubMed: 2333475]
- Hashtroudi S, Johnson MK, Vnek N, Ferguson SA. Aging and the effects of affective and factual focus on source monitoring and recall. *Psychol. Aging*. 1994; 9:160–170. [PubMed: 8185863]

- Heaton RK, Gladsjo JA, Palmer BW, Kuck J, Marcotte TD, Jeste DV. Stability and course of neuropsychological deficits in schizophrenia. *Arch. Gen. Psychiatry*. 2001; 58:24. [PubMed: 11146755]
- Henquet C, Krabbendam L, Dautzenberg J, Jolles J, Merckelbach H. Confusing thoughts and speech: source monitoring and psychosis. *Psychiatry Res*. 2005; 133:57–63. [PubMed: 15698677]
- Irani F, Brensinger CM, Richard J, Calkins ME, Moberg PJ, Bilker W, Gur RE, Gur RC. Computerized Neurocognitive Test Performance in Schizophrenia: A Lifespan Analysis. *American Journal of Geriatric Psych*. 2012; 20:41.
- Johnson MK, Raye CL. Cognitive and brain mechanisms of false memories and beliefs. *Memory, brain, and belief*. 2000:35–86.
- Johnson MK, Hashtroudi S, Lindsay DS. Source monitoring. *Psychol. Bull.* 1993; 114:3–28. [PubMed: 8346328]
- Keefe RS, Arnold MC, Bayen UJ, McEvoy JP, Wilson WH. Source-monitoring deficits for self-generated stimuli in schizophrenia: multinomial modeling of data from three sources. *Schizophr. Res.* 2002; 57:51–67. [PubMed: 12165376]
- Levitt, JJ.; Nestor, PG.; Karapellou, ME.; Law, S.; O'Donnell, BF.; McCarley, RW. Spatial working memory in schizophrenia: cognitive correlates; Paper presented at the Annual Meeting of the American Psychiatric Association; Miami, FL.. 1995. p. 20-25.
- Lezak, MD. *Neuropsychological Assessment*. USA: Oxford University Press; 2004.
- Lindsay DS. Source monitoring. *Cognitive psychology of memory*. 2008; 2:325–348.
- Mitchell KJ, Johnson MK. Source monitoring 15 years later: what have we learned from fMRI about the neural mechanisms of source memory? *Psychol. Bull.* 2009; 135:638–677. [PubMed: 19586165]
- Mitchell KJ, Johnson MK, Raye CL, Greene EJ. Prefrontal cortex activity associated with source monitoring in a working memory task. *J. Cogn. Neurosci.* 2004; 16:921–934. [PubMed: 15298780]
- Nienow TM, Docherty NM. Internal source monitoring and thought disorder in schizophrenia. *J. Nerv. Ment. Dis.* 2004; 192:696–700. [PubMed: 15457113]
- Partington JE, Leiter RG. Partington's Pathways Test. *Psychological Service Center Journal*. 1949; 1:11–20.
- Reitan, RM.; Davison, LA. *Clinical Neuropsychology: Current Status and Applications*. VH Winston & Sons; 1974.
- Reitan R. Validity of the Trail Making Test as an indicator of organic brain damage. *Percept. Mot. Skills*. 1958; 8:271–276.
- Spitzer, RL.; Endicott, J. *Schedule for affective disorders and schizophrenia-lifetime version*. New York: New York State Psychiatric Institute; 1978.
- Strub, RL.; Black, FW. *Neurobehavioral Disorders: A Clinical Approach*. Philadelphia: Davis; 1988.
- Subramaniam K, Luks TL, Fisher M, Simpson GV, Nagarajan S, Vinogradov S. Computerized Cognitive Training Restores Neural Activity within the Reality Monitoring Network in Schizophrenia. *Neuron*. 2012; 73:842–853. [PubMed: 22365555]
- Vinogradov S, Willis-Shore J, Poole JH, Marten E, Ober BA, Shenaut GK. Clinical and neurocognitive aspects of source monitoring errors in schizophrenia. *Am. J. Psychiatry*. 1997; 154:1530–1537. [PubMed: 9356560]
- Wechsler D. *WAIS-R Manual: Wechsler Adult Intelligence Scale-Revised*. Psychological corp. 1981
- Wolwer W, Gaebel W. Impaired Trail-Making Test-B performance in patients with acute schizophrenia is related to inefficient sequencing of planning and acting. *J. Psychiatr. Res.* 2002; 36:407–416. [PubMed: 12393310]
- Zaragoza M, Lane S. Processing resources and eyewitness suggestibility. *Legal and Criminological Psychology*. 1998; 3:305–320.

Table 1

Descriptive information for sample (n=45).

Mean age in years (std.deviation)	35.66 (8.23)
Mean years of Education	12.41 (1.59)
% Female	48.9
% Caucasian	64.4
% African-American	33.3
Mean age at first psychiatric treatment	20.31 (5.97)
% Schizophrenia	68.9
% Schizoaffective disorder	31.1
Mean GAF score (initial session)	50.22 (11.99)
Mean GAF score (follow-up session)	53.82 (9.46)
Mean score CPT-A ^a	36.18 (3.89)
Mean score Digits Span Backward	5.78 (2.16)
Mean score TMT B-A (Time taken)	107.24 (49.05)
Mean score Source Monitoring	15.02 (2.89)

^a
n=39

GAF = Global Assessment of Functioning Scale

CPT-A = Auditory Continuous Performance test

TMT = Trail-Making Test

Table 2

Correlations among variables.

	SM	CPT-A (log)	Digits-B
CPT-A (log)	-.12		
Digits-B	.21	-.34 [*]	
Trails B-A (log)	-.53 ^{**}	.07	-.38 ^{**}

^{*}
 $p < .01$ (2-tailed)

^{**}
 $p < .01$ (2-tailed)

CPT-A = Auditory Continuous Performance test

Digits-B = Digit Span Backward

SM = Source Monitoring

TMT = Trail-Making Test

Table 3

Standard multiple regression of CPT-A, Digits-B, and Trails B-A on source monitoring (n=39).

	B	SE	β	pr^2
Intercept	30.12	5.12		
CPT-A (log)	-.67	1.12	-.09	-.10
Digits-B	-0.03	0.22	-.02	-.02
Trails B-A (log)	-7.46**	2.18	-.53	-.50
R	.53**			
R ²	.28			

**
 $p < .01$

CPT-A = Auditory Continuous Performance test

Digits-B = Digit Span Backward

TMT = Trail-Making Test

Table 4

Hierarchical Regression Analysis of Digits-B and Trails B-A on source monitoring (n=45).

	B	SE	β	pr^2
<i>Block 1</i>				
Intercept	13.58	1.23		
Digits-B	0.25	0.20	.19	.19
R	.19			
<i>Block 2</i>				
Intercept	29.91	4.75		
Digits-B	-.001	0.19	-.00	-.00
Trails B-A (log)	-7.49 **	2.12	-.51	-.48
R	.51 **			
R ²	.27			

**
 $p < .01$

Digits-B = Digit Span Backward

TMT = Trail-Making Test