Published in final edited form as:

Tetrahedron Lett. 2012 September 12; 53(37): 4921–4924. doi:10.1016/j.tetlet.2012.06.116.

Microwave-assisted Hantzsch thiazole synthesis of *N*-phenyl-4-(6-phenylimidazo[2,1-*b*]thiazol-5-yl)thiazol-2-amines from the reaction of 2-chloro-1-(6-phenylimidazo[2,1-*b*]thiazol-5-yl)ethanones and thioureas

Sukanta Kamila, Kimberly Mendoza, and Edward R. Biehl*

Department of Chemistry, Southern Methodist University, Dallas, Texas-75275, USA

Abstract

N-Phenyl-4-(6-phenylimidazo[2,1-*b*]thiazol-5-yl)thiazol-2-amines (**6a-q**) have been synthesized by the Hantzsch thiazole reaction of 2-chloro-1-(6-phenylimidazo[2,1-*b*]thiazol-5-yl)ethanones (**4a-e**) with suitably substituted thioureas using microwave heating. The ethanones (**4a-e**) were prepared by the reaction of 6-phenylimidazo[2,1-*b*]thiazoles (**3a-e**) with chloroacetylchloride in refluxing 1,4-dioxane whereas the thiazoles (**3a-e**) were synthesized by the reaction of 2-bromo-1-phenylethanones (**2a-e**) with thiazol-2-amine in refluxing acetone.

Keywords

microwave irradiation; (6-phenylimidazo[2; 1-*b*]thiazol-5-yl)thiazol-2-amine 2-chloro-1-(6-phenylimidazo[2; 1-*b*]thiazol-5-yl)ethanones; 6-phenylimidazo[2; 1-*b*]thiazoles

1. Introduction

Many natural products and biologically active compounds containing imidazo[2,1-*b*]thiazole moieties have been synthesized and shown to exhibit potent biological activity.¹ As shown in Figure 1, tetramisole (**A**) is a strong anthelmintic in the treatment of many nematodes.² Thieno[3,2-*d*]pyrimidinone (**B**)³ and 5,6-diarylimidazo[2,1-*b*][1,3]thiazoles (**C**),⁴ are excellent antibacterial agents and C-2 aryl-substituted pilicides [thiazole ring-fused 2-pyridones (**D**)]⁵ exhibit anti-coccidial behavior. ¹¹C-labeled imidazo[2,1-*b*]benzothiazole (**E**)⁶ has been shown to be a superb fluoroprobe in PET analysis of Alzheimer's disease. Milne et al.⁷ have prepared compound (**F**) which possesses an imidazo[2,1-*b*]benzothiazole ring. Compound **F**, although unrelated in structure to resveratrol, showed a 1000-fold greater affinity toward SIRT1 than resveratrol. These workers also demonstrated that compound **F** binds to the SIRT1 enzyme–peptide substrate complex at an allosteric amino terminal site to the catalytic domain resulting in lower Michaelis constant for acetylated substrates. Compound **G** (Figure 1) inhibits binding of radio labeled TARC (Thymus and Activation Regulated Chemokine) and MDC (Macrophage-derived Chemokine) to CCR4 receptors on the surface of CEM⁸ and also inhibits the in vitro migration of CEM cells mediated by

© 2012 Elsevier Ltd. All rights reserved.

*Corresponding author: E. R. Biehl; ebiehl@smu.edu; Phone 214-768-1280; Fax 214-768-4089.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

TARC.⁸ Subsequently another research group identified hydrazone derivatives of imidazo[2,1-*b*]benzothiazoles as a potent antitumor agent.⁹ It also been reported that imidazo[2,1-*b*]benzothiazole system acts as a scaffold endowing dihydropyridines with selective cardiodepressant activities.¹⁰

For the past few years, our group has, also, been preparing biologically important heterocycles using microwave irradiation^{11a-g} Of particular importance to this report, is our preparation of novel bis (2-thioxothiazolidin-4-one) derivatives,^{11a} several of which possess strong neuroprotecting activities against Alzheimer's and Parkinson's diseases. We have now extended our research to the microwave-assisted synthesis of potentially biologically active *N*-phenyl-4-(6-phenylimidazo[2,1-*b*]thiazol-5-yl)thiazol-2 amine derivatives (**6a-p**) in which imidazo[2,1-*b*]thiazole core moieties are attached to a variety of 2-aminothiazoles, and report the results herein. We were attracted to the Hantzsch synthesis since it has been one of the methods of choice for preparing aminothiazoles.¹² Most of these syntheses involve extended conventional heating and give aminothiazoles in mediocre yields. For example, *N*-thiazoyl α -amino acids were prepared in yields generally ranging from 18–55% in a one-pot, two-step, 7–15 hours Hantzsch reaction.¹³ There are only a few reports on microwave-assisted Hantzsch reactions.^{14a-d} Thus this study should add to the increasing importance of microwave-assisted reactions in heterocyclic chemistry.¹⁵

2. Results and Discussion

Scheme 1 outlines the synthesis of **6a-p**. Optimum conditions for carrying out the Hantzsch microwave-assisted reactions were ascertained by carrying out a series of reactions of 2-chloro-1-(6-phenylimidazo[2,1-*b*]thiazol-5-yl)ethanones (**4a**) with *N*-phenylthiourea. The results, which are summarized in Table 1, showed that maximum yield of **6a** (95%) were obtained by heating at 90 °C for 30 min in methanol.

The results for the reactions of **6a-p** are shown in Table 2. Compounds **6a-p** were obtained in 89–95% yields and most of them are powder-like solids. They are insoluble in usual organic solvents such as, dichloromethane, ethyl acetate, THF, ethanol or methanol but soluble in DMF or DMSO. The proposed structures of **6a-p** were confirmed, in part, by the presence of C=N signals around δ 164 ppm (indicative of an 2-aminothiazole ring) in the ¹³C NMR spectra. The ¹H NMR spectra of these compounds showed a broad singlet (equivalent to 1H) around δ 11 ppm which corresponds to NH functionality. The starting 2-chloro-1-(6-phenylimidazo[2,1-*b*]thiazol-5-yl)ethanones (**4a-e**) were prepared by refluxing 2.5 equivalent of chloroacetylchloride and 1 equivalent of 6-phenylimidazo[2,1-*b*]thiazoles (**3a-e**) in refluxing 1,4-dioxane. Although we tried other solvents, 1,4-dioxane proved to be superior. The structures of **4a-e** were confirmed by the presence of C=O signals around δ 180 ppm and the CH₂ signal around δ 47 ppm. The ¹H NMR spectra of **4a-e** showed characteristic singlets (equivalent to 2H) around δ 4 ppm which corresponds to -CH₂ functionality. The starting 6-phenylimidazo[2,1-*b*]thiazoles (**3a-e**) were synthesized according to a literature¹⁶ procedure. All compounds were characterized by ¹H NMR, ¹³C NMR, DEPT-135 and HRMS analysis.

Interestingly, the same reactions carried out under conventional reflux conditions using methanol as a solvent gave **6a-p** in lower yields (Table 2), required longer reaction time (8 h) and/or the products required rigorous purification. However, the microwave reactions provided **6a-p** in higher yields in faster reaction times (usually less than 30 min) and the pure products were obtained by simple washing of the crude products with cold ethanol. Our method compares favorably to that reported for the microwave-assisted synthesis of functionalized simple 2-aminothiazoles^{14a} with respect to yield and reaction time. However

a higher temperature was needed in our procedure presumably due to the more complex nature of the phenylimidazo[2,1-*b*]thiazol-5-yl)thiazol-2-amine products.

In conclusion, we have successfully developed an easy practical access to novel series of *N*-phenyl-4-(6-phenylimidazo[2,1-*b*]thiazol-5-yl)thiazol-2-amine derivatives. Thus, the mild reaction condition, easy work up procedure, good to excellent yields, and readily available starting materials make this reaction an attractive method for the preparation of titled compounds. To the best of our knowledge, there has been no reported example of synthesis of this type of biologically important molecule. Efforts directed toward the synthesis of other important drug molecule with imidazo[2,1-*b*]thiazol-5-yl)thiazol moieties by microwave irradiation are ongoing in our laboratory. Also work is in progress to obtain biological activity (antibacterial, antifungal, anticancer, antitumor and neuroprotective kinase inhibitory activity) of these important compounds. Results in these studies will be presented in due course.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors are grateful to NIH (IRC2NS064950) for generous financial support.

References and notes

1. (a) Budriesi R, Ioan P, Leoni A, Pedemonte N, Locatelli A, Micucci M, Chiarini A, Galiotta LJV. *J Med Chem.* 2011; 54:3885. [PubMed: 21568323] (b) Guzeldemirci NU, Kucukbasmaci O. *Eur J med Chem.* 2010; 45:63. [PubMed: 19939519] (c) Metaye T, Millet C, Kraimps JL, Saunier B, Barbier J, Begon F. *Biochem Pharmacol.* 1992; 43:1507. [PubMed: 1314603]
2. Raeymaekers AHM, Allewijn FTN, JV, Demoen PJA, VanOffenwert TTT, Janssen PAJ. *J Med Chem.* 1966; 9:545. [PubMed: 5968020]
3. Carpenter, AJ.; Cooper, JP.; Handlon, AL.; Hertzog, DL.; Hyman, CE.; Guo, YC.; Speake, JD.; Witty, DR. WO03033476. 2003.
4. Scribner A, Meitz S, Fisher M, Wyvratt M, Leavitt P, Liberator P, Gurnett A, Brown C, Mathew J, Thompson D, Schmatz D, Biftu T. *Bioorg Med Chem Lett.* 2008; 18:5263. [PubMed: 18789686]
5. Chorell E, Pinkner JS, Phan G, Edvinsson S, Buelens F, Remaut H, Waksman G, Haultgren SJ, Almqvist F. *J Med Chem.* 2010; 53:5690. [PubMed: 20586493]
6. Yousefi BH, Manook A, Drzezga A, Reutern BV, Schwaiger M, Wester HJ, Henriksen G. *J Med Chem.* 2011; 54:949. [PubMed: 21275403]
7. Milne JC, Lambert PD, Schenk S, Carney D, Smith JJ, Gangne DJ, Jin L, Boss O, Perni RB, Vu CB, Bemis JE, Xie R, Disch JS, Ng PY, Nunes JJ, Lynch AV, Yang HY, Galonek H, Israelian K, Choy W, Iffland A, Lavu S, Medvedik O, Sinclair DA, Olefsky JM, Jirousek MR, Elliott PJ, Westphal CH. *Nature.* 2007; 450:712. [PubMed: 18046409]
8. Wang X, Xu F, Xu Q, Mahmud H, Houze J, Zhu L, Akerman M, Tonn G, Tang L, McMaster BE, Dairaghi DJ, Schall TJ, Collins TL, Medina JC. *Bioorg Med Chem Lett.* 2006; 16:2800. [PubMed: 16497499]
9. Andreani A, Burnelli S, Granaola M, Leoni A, Locatelli A, Morigi R, Rambaldi M, Varoli L, Calonghi N, Cappadone C, Farruggia G, Zini M, Stefanelli C, Masotti L, Radin NS, Shoemaker RH. *J Med Chem.* 2008; 51:809. [PubMed: 18251494]
10. Budriesi R, Ioan P, Locatelli A, Cosconati S, Leoni A, Ugenti MP, Andreani A, Toro RD, Bedini A, Spampinato S, Marinelli L, Novellino E, Chiarini A. *J Med Chem.* 2008; 51:1592. [PubMed: 18303827]
11. See, for example: Kamila S, Biehl ER. *Tetrahedron Lett.* 2012 In press. Kamila S, Ankati H, Harry E, Biehl ER. *Tetrahedron Lett.* 2012; 53:2195. Kamila S, Ankati H, Biehl ER. *Tetrahedron Lett.*

- 2011; 52:4375. Kamila S, Ankati H, Mendoza K, Biehl ER. *Open Org Chem J.* 2011; 5:127. Kamila S, Ankati H, Biehl ER. *Molecules.* 2011; 16:5527. [PubMed: 21716174] Kamila S, Ankati H, Biehl ER. *Arkivoc.* 2011; 9:94. Kamila S, Biehl ER. *J Heterocycl Chem.* 2007; 44:407. Kamila S, Koh B, Khan O, Zhang H, Biehl ER. *J Heterocycl Chem.* 2006; 43:1641. Kamila S, Koh B, Biehl ER. *J Het Chem.* 2006; 43:1609. Kamila S, Zhang H, Biehl ER. *Heterocycles.* 2005; 65:2119.
12. Hantzsch A, Weber JH. *Ber Dtsch Chem Ges.* 1887; 20:3118.
13. Rudolph J. *Tetrahedron.* 2006; 56:3161.
14. See, for example: Kabalka GW, Mereddy AR. *Tetrahedron Lett.* 2006; 47:517. Leadbeater NE. *Chem Commun.* 2010; 46:6693. Gaikwad SA, Patil AA, Deshmukh MB. Phosphorus, Sulfur, Silicon Relat Elem. 2010; 185:103. Bougrin K, Loupy A, Soufiaoui M. *J Photo Chem Photobio C: Photo Chem Reviews.* 2005; 6:139.
15. See, For example: *Microwaves in Organic Synthesis.* Loupy A. Wiley-VCH Weinheim 2002. *Controlled Microwave Heating in Modern Organic Chemistry.* *Angew Chem Intern Ed.* 2004; 4:6250. Varma R. *Solvent-Free Synthesis of Heterocyclic Compounds Using Microwaves.* *J Heterocycl Chem.* 1999; 36:1565.
16. Chekotylo OA. *Synthesis.* 2010; 10:1692.

Figure 1.
Imidazo[2,1-*b*]thiazoles and 2-aminothiazoles

Scheme 1. Schematic representation for the synthesis of compounds **6a-p**.

Table 1 Screening of solvents, reaction time, and temperature for synthesis of **6a**

Condition	Temp (°C)	Time (min)	Yield (%)
No solvent	90–120	15	trace
Ethanol	90–120	15	79
Ethanol	90–120	30	85
Methanol	90	15	71
Methanol	90	30	95
Acetonitrile	90–100	15–30	55
Water	90–120	30–45	trace
THF	70–100	30–45	trace
n-Butanol	90–110	30–45	trace
DME	90–110	30–45	trace
Sulfonate	90–130	30–45	trace
Benzene/Toluene	90–110	30–45	trace
DMF	90–110	30–45	trace

Reaction condition: microwave irradiation using **4a**: **5a** (1:1 equiv) and 1 mmol of **4a**, solvent (2 mL) under 250 psi pressure in a capped specially designed microwave test tube.

Table 2

The synthesis of *N*-phenyl-4-(6-phenylimidazo[2,1-*b*]thiazol-5-yl)thiazol-2-amines (**6a-p**) by the microwave-assisted Hantzsch reaction of ethanone derivatives (**4a-e**) with substituted thioureas (**5a-d**).

Entry				Yield ^{a,b}	Yield ^{a,c}
1	4a , R ₁ = H	5a , R ₂ = R ₃ = R ₄ = H	6a , R ₁ = R ₂ = R ₃ = R ₄ = H	98	65
2	4b , R ₁ = OCH ₃	5a , R ₂ = R ₃ = R ₄ = H	6b , R ₁ = OCH ₃ , R ₂ = R ₃ = R ₄ = H	96	45
3	4c , R ₁ = CH ₃	5a , R ₂ = R ₃ = R ₄ = H	6c , R ₁ = CH ₃ , R ₂ = R ₃ = R ₄ = H	92	69
4	4d , R ₁ = CN	5a , R ₂ = R ₃ = R ₄ = H	6d , R ₁ = CN, R ₂ = R ₃ = R ₄ = H	90	60
5	4e , R ₁ = Br	5a , R ₂ = R ₃ = R ₄ = H	6e , R ₁ = Br, R ₂ = R ₃ = R ₄ = H	95	78
6	4a , R ₁ = H	5b , R ₂ = R ₄ = H, R ₃ = Cl	6f , R ₁ = H, R ₂ = R ₄ = H, R ₃ = Cl	93	74
7	4a , R ₁ = H	5c , R ₂ = R ₄ = Cl, R ₃ = H	6g , R ₁ = H, R ₂ = R ₄ = Cl, R ₃ = H	99	88
8	4a , R ₁ = H	5d , R ₂ = R ₄ = H, R ₃ = OCH ₃	6h , R ₁ = H, R ₂ = R ₄ = H, R ₃ = OCH ₃	90	53
9	4c , R ₁ = CH ₃	5d , R ₂ = R ₄ = H, R ₃ = OCH ₃	6i , R ₁ = CH ₃ , R ₂ = R ₄ = H, R ₃ = OCH ₃	89	47
10	4e , R ₁ = Br	5c , R ₂ = R ₄ = Cl, R ₃ = H	6j , R ₁ = Br, R ₂ = R ₄ = Cl, R ₃ = H	98	67

Entry	4	5	6	Yield ^{a,b}	Yield ^{d,e}
11	4	5	6	93	81
12				96	80
13				89	55
14				91	58
15				98	67
16				96	70

^a Isolated yield, all compounds were characterized by ¹H NMR, ¹³C NMR, IR and HRMS analysis.

^b MW, 30 min, 90° C. 1 mmol:0.6 mmol **4**:**5**:2 mL MeOH

^c Conventional heating 8 h at 90°C; 0.1mmol:0.6mmol **4:5:2** mL MeOH d. 3.6mmol:23.4mmol/10 mL MeOH