Published in final edited form as:

Curr Opin HIV AIDS. 2012 November ; 7(6): 514–519. doi:10.1097/COH.0b013e3283582d30.

Oral antiretroviral chemoprophylaxis: current status

Jared Baeten, MD, PhD and Connie Celum, MD, MPH

Departments of Global Health, Medicine, and Epidemiology University of Washington

Abstract

Purpose of review—Pre-exposure prophylaxis (PrEP), in which HIV uninfected persons with ongoing HIV risk use oral antiretroviral medications as chemoprophylaxis against sexual HIV acquisition, is a promising new HIV prevention strategy.

Recent findings—During the past two years, proof-of-concept that PrEP protects against sexual HIV acquisition has been demonstrated in three clinical trials, conducted among men who have sex with men and heterosexual men and women. These trials used daily oral tenofovir disoproxil fumarate, alone or co-formulated with emtricitabine. The degree of HIV protection in these trials was strongly related to the level of adherence to PrEP. Two additional clinical trials, both among heterosexual women, did not demonstrate HIV protection with PrEP, with low adherence to daily use of PrEP the leading hypothesis for lack of efficacy; adherence and biologic mechanisms for lack of efficacy in these trial populations are being evaluated.

Summary—Oral chemoprophylaxis, using tenofovir and combination emtricitabine/tenofovir, is effective for prevention of sexual HIV transmission. Next steps in the field include rigorous evaluation of uptake and adherence to PrEP in implementation settings.

Keywords

HIV prevention; pre-exposure prophylaxis; sexual HIV-1 transmission; tenofovir; emtricitabine

Introduction

Antiretroviral medications have the potential to be used for HIV prevention as treatment to reduce the infectiousness of HIV infected persons (1) or, for HIV-uninfected persons, as chemoprophylaxis after a recognized high-risk exposure (i.e., post-exposure prophylaxis (PEP)) or on an ongoing basis as pre-exposure prophylaxis (PrEP) for persons with repeated HIV exposures (2). While many clinicians have experience prescribing antiretroviral treatment and PEP, antiretroviral PrEP is a new HIV prevention strategy. This review will focus on the rationale and evidence for oral antiretrovirals as PrEP for HIV prevention, hypotheses to explain areas of uncertainty in the available data, and next steps for the field.

Chemoprophylaxis for HIV prevention: rationale

The oral nucleotide reverse transcriptase inhibitor tenofovir disoproxil fumarate (TDF), either alone or combination with the nucleoside reverse transcriptase inhibitor emtricitabine (FTC), has been most intensively studied as PrEP. TDF and FTC/TDF have important biologic qualities that made these agents attractive as potential PrEP medications: potent

Corresponding author: Jared Baeten, MD, PhD International Clinical Research Center Department of Global Health University of Washington, Box 359927 325 Ninth Avenue Seattle, WA 98104 206 520-3808 jbaeten@uw.edu.

Conflict of Interest Gilead Sciences provided study drug for the Partners PrEP Study but did not provide funding to the authors or the institution.

antiretroviral activity, including activity against all HIV subtypes; rapid onset of activity after ingestion; early action in HIV's lifecycle, which could be important for blocking initial infection; and once-daily dosing with few drug interactions. TDF and FTC/TDF are widely used as part of combination antiretroviral therapy regimens for treatment of HIV infection, and a substantial and reassuring safety and tolerability profile has been established. TDF is administered once-daily for HIV treatment, at a dose of 300 mg (as branded Viread®), and FTC/TDF also includes 200 mg of FTC (co-formulated FTC/TDF is sold branded Truvada®); these standard doses were chosen for studies of PrEP.

Animal model studies and evidence from studies of prevention of mother-to-child HIV transmission provided additional rationale for evaluating oral PrEP for HIV prevention. Non-human primate studies found that daily or intermittent PrEP using TDF and FTC/TDF given prior to SIV/SHIV systemic and mucosal challenge provided high protection (70-100%), in a dose-dependent manner (3-7). There was some evidence of greater HIV protection using FTC/TDF compared to TDF alone, suggesting that combination PrEP could provide greater benefit than from a single agent. Additional evidence that PrEP may be effective for HIV prevention comes from studies of infant prophylaxis using oral antiretrovirals to reduce HIV risk from ongoing HIV exposure through breastmilk (8).

Efficacy trials of daily oral PrEP for HIV prevention

Six randomized, double-blind, placebo-controlled clinical trials of oral PrEP, prescribed for use once-daily, are completed or ongoing (Table 1, listed in chronological order of reporting of their results). These trials were designed to rigorously evaluate the safety and efficacy of oral PrEP for HIV prevention, and they mandated intensive clinical and counseling procedures, including monthly study visits with HIV serologic testing to identify breakthrough infections early and minimize exposure to PrEP during acute HIV infection (generally with point-of-care rapid tests) and clinical evaluation, regular laboratory safety testing, and individualized medication adherence counseling. PrEP was delivered in a context of a package of HIV prevention services, including HIV and risk-reduction counseling, screening and treatment for sexually transmitted infections, free provision of condoms, and other services.

The iPrEx study enrolled 2499 HIV seronegative men who have sex with men and transgender women from Brazil, Ecuador, Peru, South Africa, and Thailand, with the majority from the South American sites and 9% from the US (9). The trial demonstrated that daily oral FTC/TDF reduced HIV acquisition risk by 44% (95% CI 15-63%, $p=0.005$). Subgroup analyses indicated higher efficacy associated with higher adherence to the study medication: 73% among those with 90% adherence as measured by counts of unused pills. The relationship between detection of tenofovir or emtricitabine in blood samples (a biomarker of adherence) and HIV protection was assessed as well: only 8% of seroconverters had detectable study drug at the visit closest to seroconversion, compared with 54% of a matched subset of non-seroconverters. Having detectable drug was strongly associated with substantially lower HIV risk (relative risk reduction 92%, 95% CI 40-99%, $p<0.001$).

The FEM-PrEP study enrolled 2021 high-risk HIV uninfected women from Kenya, South Africa, and Tanzania. The study was stopped due to futility by its Independent Data Monitoring Committee in April 2011 (efficacy estimate 6%, 95% CI -52-41%, $p=0.8$) (10). Like iPrEx, FEM-PrEP tested archived blood samples from the trial for the presence of the study medication, and this testing suggested that adherence to daily oral FTC/TDF was very low in FEM-PrEP: only 26% of non-seroconverting controls had consistent tenofovir levels detected in plasma (and only 15% of seroconverters as well). The study team concluded that

study drug adherence was too low to assess the efficacy of FTC/TDF PrEP for HIV prevention in the study population.

The TDF2 study enrolled 1200 heterosexual HIV uninfected men and women in Botswana (90% <30 years of age). The study demonstrated that FTC/TDF PrEP had 63% efficacy (95% CI 22-83%, $p=0.01$) for HIV protection compared to placebo (11). Among those known to be receiving study product at the time of seroconversion (i.e., censoring follow-up time for those who had been lost to follow-up or had study product held for other reasons), efficacy was 78% (95% CI 41-94, $p=0.005$). FTC/TDF appeared to provide protection for both men (overall: 80%, $p=0.03$; subgroup receiving medication: 82%, $p=0.06$) and women (overall: 49%, $p=0.1$; subgroup receiving medication: 76%, $p=0.02$), although the relatively small size of the study limited statistical significance for these subgroup analyses.

The Partners PrEP Study enrolled 4758 HIV uninfected men and women from Kenya and Uganda who were at risk of HIV because of having a known HIV-infected partner who was not yet eligible for antiretroviral according to the national guidelines of those two countries (16). In July 2011, the study's Data Safety Monitoring Board recommended that the placebo arm be discontinued because the study crossed a pre-defined stopping boundary for demonstrating PrEP efficacy for HIV protection. TDF efficacy was 67% (95% CI 44-81, $p<0.0001$) and FTC/TDF efficacy was 75% (95% CI 55-87, $p<0.0001$); the difference between TDF and FTC/TDF was not statistically significant ($p=0.23$). Both TDF and FTC/TDF significantly reduced HIV risk for both men and women (12): for TDF 63% ($p=0.01$) for men and 71% ($p=0.002$) for women and for FTC/TDF 84% ($p<0.001$) for men and 66% ($p=0.005$) for women, and these degrees of HIV protection for women and men were statistically comparable. Adherence to study drug was measured by multiple means – pill counts of unused study medication, electronic pill cap monitoring, and home visits for unannounced pill counts (17) – and was very high. Tenofovir was detected in 82% of blood samples from a randomly-selected subpopulation of non-seroconverters (confirming high adherence); detection was less frequent (31%) in those who acquired HIV. Detection of tenofovir was associated with substantial HIV protection (86%, $p<0.001$ for the TDF arm and 90%, $p=0.002$ for the FTC/TDF arm) (18).

The VOICE trial is an ongoing five-arm study of daily oral or topical PrEP (i.e., oral TDF, oral FTC/TDF, oral placebo, vaginal tenofovir gel, vaginal placebo gel) among HIV uninfected African women. The Data Safety Monitoring Board for the VOICE trial recommended discontinuation of the oral TDF arm in September 2011 (13) due to inability to demonstrate efficacy; the FTC/TDF and placebo arms are ongoing and will report results in early 2013. Lastly, the ongoing Bangkok Tenofovir Study is testing daily oral TDF PrEP among HIV uninfected 2413 injection drug users in Thailand; results are expected in late 2012(15).

Additional outcomes from PrEP clinical trials: safety, resistance, sexual behavior

Trials have found that oral TDF and FTC/TDF appear to be well-tolerated among HIV-uninfected persons, with the rate of both serious and mild adverse events generally balanced between those receiving PrEP and those receiving placebo. In both iPrEx and Partners PrEP, gastrointestinal side effects (e.g., nausea, diarrhea) occurred more commonly in those assigned active PrEP, although these symptoms were present only in a minority of subjects (~10% or less), were mild in severity, and were generally limited to the first month after initiation of the medication. A modest (1%) reduction in bone mineral density was observed in the iPrEx study and in an earlier phase II study of TDF PrEP in men who have sex with men (19); decline in bone mineral density is a known side effect of TDF when used for HIV

treatment and has not been associated with increased risk of fracture. Oral TDF has been associated with renal complications in HIV-infected persons, particularly proximal tubular dysfunction with or without reduced glomerular filtration, but PrEP clinical trials have not found increased risk of renal complications in HIV uninfected persons. Finally, data from Partners PrEP (20) and from the Antiretroviral Pregnancy Registry (21) suggest that use of TDF and FTC/TDF in early pregnancy is not associated with increased rates of birth defects, although more data are needed to fully assess the safety of these medications through pregnancy.

Antiretroviral resistance has been rare in PrEP trials and limited to those with seronegative acute infection at the time of randomization: 2/2 subjects in iPrEx (both M184I/V mutations), 2/8 subjects in Partners PrEP (one K65R and one M184V mutation), and 1/1 subject in TDF2 (K65R and M184V). Five cases of M184V resistance were observed in FEM-PrEP, one in the placebo arm and three potentially transmitted and not acquired on PrEP. Adherence to pre-exposure prophylaxis, protection against HIV infection, and antiretroviral resistance appear to be tightly interwoven: low adherence provides little HIV protection but little risk of resistance if infection is acquired, whereas high adherence potentially blocks most transmissions. Whether there is a window of modest PrEP use that confers diminished HIV protection but sufficient drug pressure to select for resistance is unknown. Clearly, HIV testing must be done to minimize PrEP use by those already infected.

Finally, the question increased sexual risk-taking accompanying PrEP use has been explored in iPrEx and Partners PrEP, where self-reported condom use increased during the studies and sexually transmitted infection rates fell, to an equivalent degree across both active PrEP and placebo arms. In iPrEx, one important correlate of detection of tenofovir in blood was unprotected anal receptive sex – suggesting that those at highest risk of HIV were more, and not less, likely to use the medication.

Understanding the divergent results of efficacy trials of daily oral PrEP

The iPrEx, TDF2, and Partners PrEP trials provide clear evidence that daily oral PrEP, using TDF and FTC/TDF, reduces the risk of sexual HIV acquisition. The lack of HIV protection in the FEM-PrEP (testing FTC/TDF) and VOICE (testing TDF) trials, as well as the wide range of efficacy estimates in iPrEx, TDF2, and Partners PrEP suggests that there are important factors that influence PrEP efficacy.

Biological hypotheses to explain divergent PrEP trial results have focused on trying to explain the lack of efficacy for PrEP in women at high-risk who were enrolled in FEM-PrEP and VOICE. Pharmacokinetics studies have found that oral dosing of TDF achieves higher concentrations (by a factor of 10-fold) in rectal tissue compared to cervicovaginal tissue (22; 23), leading some to hypothesize that oral PrEP could be less effective in women, whose primary exposure is through vaginal sex, compared to men who have sex with men, whose primary risk is through receptive anal sex (of note, no studies of tenofovir concentrations in relevant mucosal tissue for heterosexual men have been done). Arguing against this hypothesis are gender-specific subgroup results from Partners PrEP and TDF2, which found that PrEP provides high protection against HIV for women – equivalent to that seen for heterosexual men. Others have hypothesized that important co-factors for HIV acquisition – including mucosal inflammation due to sexually transmitted infections or other conditions, contact with partners with acute HIV infection, and intravaginal practices and use of hormonal contraception (24; 25) – that likely were common in the FEM-PrEP and VOICE populations might interact with PrEP in some way as to influence HIV protection efficacy.

While this hypothesis is intriguing, there are no data at this time to explain how these factors would blunt or even completely override HIV protection provided by PrEP.

The strongest hypothesis to explain divergent results across PrEP trials is differences in adherence to daily oral PrEP. Adherence to antiretroviral therapy is key to its efficacy for HIV treatment, and thus it is reasonable to expect that adherence would be critical to the efficacy of antiretroviral PrEP. Moreover, of course, for individuals who did not take PrEP at all, no protection would be expected. Most subjects in FEM-PrEP (70%) perceived themselves to have little or no chance of acquiring HIV, which could explain low PrEP use in that trial. Missed doses and missed visits to collect PrEP study medication undoubtedly diminished efficacy in PrEP trials, and there appears to be a strong dose-response relationship between PrEP use and HIV protection in PrEP trials (Table 2). Thus, efficacy estimates from the clinical trials are underestimates of the true biologic efficacy of PrEP for preventing HIV infection – and the protection estimates when tenofovir was present in blood from the iPrEx and Partners PrEP trials may thus more closely reflect the true biologic efficacy. Additional analyses in the FEM-PrEP and VOICE studies will be necessary to understand PrEP use, predictors of use, and HIV protection for African women at high risk of infection.

Next steps for PrEP

In May 2012, the Antiviral Drugs Advisory Committee to the US Food and Drug Administration recommended that a formal label indication for HIV prevention be made for branded FTC/TDF (Truvada®) – a landmark for HIV prevention. Guidelines from WHO and CDC are in development and will help guide use of PrEP in clinical settings.

The critical next step question for PrEP will be whether implementation outside of rigorous clinical trials can be feasibly done. Several PrEP trials have provided active PrEP to their study participants (iPrEx, Partners PrEP, TDF2), to fulfill promises of access to effective products for placebo-arm participants and to understand adherence and sexual behavior in the absence of placebo. In addition, demonstration projects of PrEP are planned, in diverse populations and geographic settings. A follow-on study to Partners PrEP will evaluate staged use of PrEP and antiretroviral therapy in Kenyan and Ugandan HIV serodiscordant couples (26), an important next step to bring together the benefits of treatment and PrEP for HIV prevention. Most demonstration projects are planning less frequent and less-intensive visits than were done in the clinical trials. Operations research needs to be done to understand delivery of PrEP to those at highest risk (26-28), evaluate delivery models of PrEP, and motivate and monitor PrEP adherence.

The potential for less-than-daily dosing of oral FTC/TDF is being evaluated in pharmacokinetics studies (29) and in a recently-initiated trial among men who have sex with men in France and Canada (IPERGAY) (30). For most persons, PrEP should be envisioned as a time-limited prevention strategy, for periods (months to a few years) of highest behavioral risk – for example during periods when attempting to conceive (31). In this way, time-limited PrEP is an important contrast to use of antiretrovirals as treatment, which is necessarily life-long.

Conclusions

Daily oral TDF and FTC/TDF PrEP are effective strategies for HIV prevention. Analyses from ongoing and completed PrEP trials will provide a more complete understanding of effectiveness in different populations, and reasons for divergent efficacy estimates. Successful HIV prevention on a population scale will need to incorporate multiple, evidence-based biologic and behavioral strategies to achieve maximal benefits, including

behavior change, HIV testing, male circumcision, antiretroviral treatment for HIV-infected persons, and PrEP.

Acknowledgments

The authors have received research grant support from the US National Institutes of Health and from the Bill & Melinda Gates Foundation for research related to pre-exposure prophylaxis and are the protocol chairs of the Partners PrEP Study.

Funding support: US National Institutes of Health (R01 MH095507) and the Bill & Melinda Gates Foundation (grant 47674).

References

- ** 1. Cohen MS, Chen YQ, McCauley M, Gamble T, Hosseinipour MC, et al. Prevention of HIV-1 infection with early antiretroviral therapy. *New England Journal of Medicine*. 2011; 365:493–505. [PubMed: 21767103] This randomized clinical trial provided definitive evidence that use of antiretroviral treatment by HIV infected persons results in substantial reduction in infectiousness and risk of HIV transmission to sexual partner (96% relative risk reduction, mediated through high adherence and viral suppression). Use of antiretroviral medications for treatment of HIV infected persons to reduce infectiousness and pre-exposure prophylaxis for HIV uninfected persons to reduce susceptibility are complementary prevention strategies.
2. Cohen MS, Gay C, Kashuba AD, Blower S, Paxton L. Narrative review: antiretroviral therapy to prevent the sexual transmission of HIV-1. *Annals of Internal Medicine*. 2007; 146:591–601. [PubMed: 17438318]
3. Tsai CC, Follis KE, Sabo A, Beck TW, Grant RF, et al. Prevention of SIV infection in macaques by (R)-9-(2-phosphonylmethoxypropyl)adenine. *Science*. 1995; 270:1197–9. [PubMed: 7502044]
4. Garcia-Lerma JG, Otten RA, Qari SH, Jackson E, Cong ME, et al. Prevention of rectal SHIV transmission in macaques by daily or intermittent prophylaxis with emtricitabine and tenofovir. *PLoS Medicine*. 2008; 5:e28. [PubMed: 18254653]
5. Garcia-Lerma JG, Cong ME, Mitchell J, Youngpairoj AS, Zheng Q, et al. Intermittent prophylaxis with oral truvada protects macaques from rectal SHIV infection. *Science Translational Medicine*. 2010; 2:14ra4.
6. Parikh UM, Dobard C, Sharma S, Cong ME, Jia H, et al. Complete protection from repeated vaginal simian-human immunodeficiency virus exposures in macaques by a topical gel containing tenofovir alone or with emtricitabine. *Journal of Virology*. 2009; 83:10358–65. [PubMed: 19656878]
7. Subbarao S, Otten RA, Ramos A, Kim C, Jackson E, et al. Chemoprophylaxis with tenofovir disoproxil fumarate provided partial protection against infection with simian human immunodeficiency virus in macaques given multiple virus challenges. *Journal of Infectious Diseases*. 2006; 194:904–11. [PubMed: 16960777]
8. Chasela CS, Hudgens MG, Jamieson DJ, Kayira D, Hosseinipour MC, et al. Maternal or infant antiretroviral drugs to reduce HIV-1 transmission. *New England Journal of Medicine*. 2010; 362:2271–81. [PubMed: 20554982]
- ** 9. Grant RM, Lama JR, Anderson PL, McMahan V, Liu AY, et al. Preexposure chemoprophylaxis for HIV prevention in men who have sex with men. *New England Journal of Medicine*. 2010; 363:2587–99. [PubMed: 21091279] This pivotal randomized clinical trial demonstrated that oral pre-exposure prophylaxis, using daily oral emtricitabine/tenofovir, was efficacious for prevention of HIV acquisition in men who have sex with men.
- ** 10. Van Damme L, Corneli A, Ahmed K, Agot K, Lombaard J, et al. Preexposure prophylaxis for HIV infection among African women. *New England Journal of Medicine*. 2012 Published online ahead of press July 11, 2012. This randomized clinical trial among high-risk African women failed to demonstrate HIV protection with oral pre-exposure prophylaxis. The study investigators hypothesized that poor adherence to the study medication explained the results.
- ** 11. Thigpen MC, Kebaabetswe PM, Paxton LA, Smith DK, Rose CE, et al. Antiretroviral preexposure prophylaxis for heterosexual HIV transmission in Botswana. *New England Journal of Medicine*. 2012; 11:2012. Published online ahead of press July. This randomized clinical trial

among young men and women from Botswana provides supportive evidence for oral pre-exposure prophylaxis for HIV prevention in heterosexuals.

- ** 12. Baeten JM, Donnell D, Ndase P, Mugo NR, Campbell JD, et al. Antiretroviral prophylaxis for HIV prevention in heterosexual men and women. *New England Journal of Medicine*. 2012. Published online ahead of press July 11, 2012. This pivotal randomized clinical trial demonstrated that oral pre-exposure prophylaxis, using daily oral emtricitabine/tenofovir and daily oral tenofovir alone, was efficacious for prevention of HIV acquisition in both heterosexual men and women.
13. National Institute of Allergy and Infectious Diseases (NIAID). NIH modifies 'VOICE' HIV prevention study in women. 2011. <http://www.nih.gov/news/health/sep2011/niaid-28.htm>
14. National Institute of Allergy and Infectious Diseases (NIAID). NIH discontinues tenofovir vaginal gel in 'Voice' HIV prevention study. 2011. <http://www.nih.gov/news/health/nov2011/niaid-25.htm>
15. Martin M, Vanichseni S, Suntharasamai P, Sangkum U, Chuachoowong R, et al. Enrollment characteristics and risk behaviors of injection drug users participating in the Bangkok Tenofovir Study, Thailand. *PLoS ONE*. 2011; 6:e25127. [PubMed: 21969870]
16. Mujugira A, Baeten JM, Donnell D, Ndase P, Mugo NR, et al. Characteristics of HIV-1 serodiscordant couples enrolled in a clinical trial of antiretroviral pre-exposure prophylaxis for HIV-1 prevention. *PLoS ONE*. 2011; 6:e25828. [PubMed: 21998703]
17. Bangsberg, D.; Haberer, J.; Psaros, C.; Baeten, J.; Katabira, E., et al. High adherence and high effectiveness observed in HIV discordant couples: Partners PrEP Study, adherence monitoring and counseling substudy. 19th Conference on Retroviruses and Opportunistic Infections (CROI); Seattle, WA. 2012. Abstract 1067
18. Donnell, D.; Baeten, J.; Hendrix, C.; Bumpus, N.; Bangsberg, D., et al. Tenofovir disoproxil fumarate drug levels indicate PrEP use is strongly correlated with HIV-1 protective effects: Kenya and Uganda. 19th Conference on Retroviruses and Opportunistic Infections (CROI); Seattle, WA. 2012. Abstract 30
19. Liu, A.; Vittinghoff, E.; Irby, R.; Mulligan, K.; Sellmeyer, D., et al. BMD loss in HIV- men participating in a TDF PrEP clinical trial in San Francisco. 18th Conference on Retroviruses and Opportunistic Infections (CROI); Boston, MA. 2011. Abstract 93
20. Mugo, N.; Celum, C.; Donnell, D.; Campbell, J.; Bukusi, E., et al. Pregnancy incidence and birth outcomes in a clinical trial of PrEP: Uganda and Kenya. 19th Conference on Retroviruses and Opportunistic Infections; Seattle, USA. 2012. Abstract 1060
21. Antiretroviral Pregnancy Registry Interim Report, 1 January 1989 through 31 July 2011. 2011. http://www.apregistry.com/forms/interim_report.pdf
22. Anderson, P.; Meditz, A.; Zheng, J-H.; Predhomme, J.; Klein, B., et al. Cellular pharmacology of tenofovir and emtricitabine in blood, rectal, and cervical cells from HIV- volunteers. 19th Conference on Retroviruses and Opportunistic Infections (CROI); Seattle, WA. 2012. Abstract 587
23. Patterson KB, Prince HA, Kraft E, Jenkins AJ, Shaheen NJ, et al. Penetration of tenofovir and emtricitabine in mucosal tissues: implications for prevention of HIV-1 transmission. *Sci Transl Med*. 2011; 3:112re4.
24. Heffron R, Donnell D, Rees H, Celum C, Mugo N, et al. Use of hormonal contraceptives and risk of HIV-1 transmission: a prospective cohort study. *Lancet Infectious Diseases*. 2012; 12:19–26. [PubMed: 21975269]
25. Low N, Chersich MF, Schmidlin K, Egger M, Francis SC, et al. Intravaginal practices, bacterial vaginosis, and HIV infection in women: individual participant data meta-analysis. *PLoS Medicine*. 2011; 8:e1000416. [PubMed: 21358808]
- * 26. Hallett TB, Baeten JM, Heffron R, Barnabas R, de Bruyn G, et al. Optimal uses of antiretrovirals for prevention in HIV-1 serodiscordant heterosexual couples in South Africa: a modelling study. *PLoS Medicine*. 2011; 8:e1001123. [PubMed: 22110407] This mathematical modeling study explores relative effectiveness and cost-effectiveness of different strategies for using antiretroviral treatment and pre-exposure prophylaxis to reduce HIV risk in HIV serodiscordant African couples.

- * 27. Juusola JL, Brandeau ML, Owens DK, Bendavid E. The cost-effectiveness of preexposure prophylaxis for HIV prevention in the United States in men who have sex with men. *Annals of Internal Medicine*. 2012; 156:541–50. [PubMed: 22508731] This study analyzed the cost-effectiveness of pre-exposure prophylaxis for use for HIV prevention among men who have sex with men in the US.
- * 28. Walensky RP, Park JE, Wood R, Freedberg KA, Scott CA, et al. The cost-effectiveness of pre-exposure prophylaxis for HIV infection in South African women. *Clinical Infectious Diseases*. 2012; 54:1504–13. [PubMed: 22474224] This important study analyzed the cost-effectiveness of pre-exposure prophylaxis for use for HIV prevention among heterosexual South African women.
- 29. HIV Prevention Trials Network. HPTN 067 - The ADAPT Study: A phase II, randomized, open-label, pharmacokinetic and behavioral study of the use of intermittent oral emtricitabine/tenofovir disoproxil fumarate pre-exposure prophylaxis (PrEP). 2011. http://www.hptn.org/research_studies/hptn067.asp
- 30. The IPERGAY Study. 2012. <http://www.ipergay.fr/>
- 31. Matthews LT, Baeten JM, Celum C, Bangsberg DR. Periconception pre-exposure prophylaxis to prevent HIV transmission: benefits, risks, and challenges to implementation. *AIDS*. 2010; 24:1975–82. [PubMed: 20679759]

Key bullet points

- Proof-of-concept has been demonstrated oral pre-exposure prophylaxis (PrEP), using daily oral tenofovir/emtricitabine, protects against sexual HIV acquisition in three clinical trials, conducted among men who have sex with men and heterosexual men and women.
- The degree of HIV protection in clinical trials of PrEP was strongly related to pill-taking adherence, and one clinical trial among heterosexual women did not demonstrate HIV protection with PrEP, with low adherence to daily use of PrEP the leading hypothesis for lack of efficacy.
- Next steps in the field include rigorous evaluation of uptake and adherence to PrEP in implementation settings.

Table 1

Current status of efficacy trials of daily oral PrEP for HIV prevention

Study (location)	Population	Design and PrEP Agent	Status	Relative reduction in HIV incidence due to PrEP	Reference
iPrEx (Brazil, Ecuador, Peru, South Africa, Thailand, US)	2499 men who have sex with men and transgender women	1:1 randomization to FTC/TDF or Placebo	Completed. FTC/TDF effective for HIV prevention.	FTC/TDF: 44% (95% CI 15-63%, p=0.005)	(9)
FEM-PrEP (Kenya, South Africa, Tanzania)	2120 women	1:1 randomization to FTC/TDF or Placebo	Completed. Trial stopped early for lack of efficacy for HIV prevention.	FTC/TDF: 6% (95% CI -52-41%, p=0.8)	(10)
TDF2 Study (Botswana)	1219 heterosexual men and women	1:1 randomization to FTC/TDF or Placebo	Completed. FTC/TDF effective for HIV prevention.	FTC/TDF: 63% (95% CI 22-83%, p=0.01)	(11)
Partners PrEP Study (Kenya, Uganda)	4758 heterosexual men and women with known HIV infected partners (HIV serodiscordant couples)	1:1:1 randomization to TDF, FTC/TDF, or Placebo	Ongoing. TDF and FTC/TDF effective for HIV prevention. Trial continued with blinded follow-up to compared TDF to FTC/TDF.	TDF: 67% (95% CI 44-81%, p<0.0001) FTC/TDF: 75% (95% CI 55-87%, p<0.0001)	(12)
VOICE (South Africa, Uganda, Zimbabwe)	3021 women (additional women randomized to tenofovir gel topical PrEP or placebo)	1:1:1 randomization to TDF, FTC/TDF, or Placebo	Ongoing. Oral TDF arm stopped early for lack of efficacy for HIV prevention. FTC/TDF arm continued.	TDF: No efficacy for HIV prevention. Efficacy estimate not yet available. FTC/TDF: Results expected early 2013.	(13; 14)
Bangkok Tenofovir Study (Thailand)	2413 injection drug users	1:1 randomization to TDF or Placebo	Ongoing.	TDF: Results expected late 2013.	(15)

Table 2

Dose-response relationship between adherence and efficacy for HIV prevention in trials of daily oral FTC/TDF PrEP

Study	Population	HIV protection estimate (randomized comparison versus placebo)	Frequency of detection of study medication in blood samples of non-seroconverters	HIV protection estimate (as related to high adherence)
Partners PrEP Study	Heterosexual men and women with known HIV infected partners (HIV serodiscordant couples)	75%	81%	90% in subjects with detectable tenofovir levels
TDF2	Young heterosexual men and women	63%	79%	78% excluding follow-up periods when subjects had no PrEP refills for >30 days
iPrEx	Men who have sex with men and transgender women	44%	51%	92% in subjects with detectable tenofovir levels
FEM-PrEP	Women	6%	35-38% at a single visit, 26% at two consecutive visits	Investigators concluded that use of PrEP was too low to evaluate efficacy.
VOICE	Women	Trial still ongoing	Not yet available	Not yet available