Published in final edited form as:

Cell Mol Life Sci. 2009 October ; 66(19): 3149–3160. doi:10.1007/s00018-009-0068-5.

Faithful after break-up: suppression of chromosomal translocations

Sang Eun Lee and

Department of Molecular Medicine, Institute of Biotechnology, University of Texas Health Science Center at San Antonio, 15355 Lambda Drive, San Antonio, TX 78245, USA

Kyungjae Myung

Genome Instability Section, Genetics and Molecular Biology Branch, National Human Genome Research Institute, National Institutes of Health, Bethesda, MD 20892, USA

Sang Eun Lee: lees4@uthscsa.edu; Kyungjae Myung: kmyung@mail.nih.gov

Abstract

Chromosome integrity in response to chemically or radiation-induced chromosome breaks and the perturbation of ongoing replication forks relies on multiple DNA repair mechanisms. However, repair of these lesions may lead to unwanted chromosome rearrangement if not properly executed or regulated. As these types of chromosomal alterations threaten the cell's and the organism's very own survival, multiple systems are developed to avoid or at least limit break-induced chromosomal rearrangements. In this review, we highlight cellular strategies for repressing DNA break-induced chromosomal translocations in multiple model systems including yeast, mouse, and human. These pathways select proper homologous templates or broken DNA ends for the faithful repair of DNA breaks to avoid undesirable chromosomal translocations.

Keywords

Chromosomal translocation; Double strand break; DNA repair; Homologous recombination; End joining

Introduction

Chromosomal translocation is a type of genome rearrangements that results in the fusion of two different chromosomes [1]. Specific chromosomal translocations are intimately associated certain types of leukemias and childhood sarcomas [2, 3], and the cause of which can be traced to the fusion of two unrelated genes that produces new fusion proteins with oncogenic properties or the fusion with regulatory elements which lead to abnormal gene expression in proto-oncogenes [4]. Identifying and characterizing chromosomal translocations facilitates diagnosis, and could provide guidelines for a therapeutic treatment course designed to target the oncogenic proteins generated by a specific chromosomal translocation [1, 4]. In many instances, treatments designed for specific chromosomal translocations have been highly effective and have improved the success of the treatment. Puzzlingly, despite its incredible clinical value, little is known about the molecular mechanism that leads to chromosomal translocations. However, accumulated evidence

suggests a role for DNA double-strand breaks (DSBs) as initiating lesions for chromosomal translocations.

Chromosomes are inherently fragile. Their integrity is constantly challenged by multiple agents that can damage either their phosphate backbone or nucleotide bases. The resulting DNA lesions, if not repaired properly, severely compromise the genetic integrity of the cell and endanger the survival of cells and ultimately organisms. Among these lesions, DNA DSBs pose a distinctly difficult challenge to an actively growing cell because even a single unrepaired DSB is sufficient to cause a cell's demise [5]. Naturally, all living cells devote substantial resources to detect and repair DNA DSBs efficiently. The two major mechanisms for repairing DSBs are homologous recombination (HR), wherein the break is repaired by copying across the lesion using homologous sequence as a template, and non-homologous end joining (NHEJ), the rejoining of DNA ends by ligation [6, 7]. A third, less well-defined mechanism, known as microhomology-mediated end joining (MMEJ) that is distinct from HR and Ku-dependent NHEJ, has been recently described [8]. We refer readers to these reviews [9, 10] to learn more about the genetics and molecular steps of DSB repair pathways.

The failure to repair DSBs is terminal and can lead to cell death. Additionally, if DSBs are misrepaired, they can lead to gross chromosomal aberrations and rearrangements [11–14]. To multi-cellular organisms, chromosomal rearrangements are highly dangerous, because even if the population of cells carrying chromosomal rearrangements may not face death, oncogenic rearrangements can cause cellular transformation that effectively decimates an organism's long-term viability. Distinct pathways have evolved to ensure that repair of DNA breaks does not result in chromosomal rearrangements. Recent work in the budding yeast and mouse model systems has begun to characterize the mechanisms of a few specific pathways that suppress chromosomal translocations during DSB repair. In this review, we discuss the recent progress in the field in identifying the molecular and genetic components of these pathways.

Role of DNA double-strand breaks in chromosomal translocations

Before we discuss particular pathways that suppress DSB-induced chromosomal translocation, we briefly outline available evidence, from various models and organisms, demonstrating the role of DSBs and the improper repair of such lesions in the formation of chromosomal translocations. First, the elevated level of chromosomal translocations was readily detected in mice deficient in DNA DSB repair such as Ku70 or Ku80 [11, 13]. Additionally, yeasts deficient in restarting collapsed replication forks rely on DSB repair mechanisms for survival [15] and elevate gross chromosomal rearrangements up to ~ 10,000-fold [16]. In yeast, a single DSB could lead to a ~ 1,000-fold induction in the gross chromosomal rearrangement rate [14]. In humans, genomic fragile sites are also associated with chromosomal rearrangements and translocations [17–19]. Clinically, chromosomal translocations and associated leukemias are frequent among nuclear bomb survivors, who suffer from radiation-induced DSBs [20]. Certain chromosomal translocations are also tightly linked to V(D)J recombination or class switch recombination events, two programmed gene rearrangements mediated by DNA DSB repair [21]. Finally, therapy-induced chromosomal translocations and leukemias stem from the treatment with chemotherapeutics or radiation that induces DNA strand breaks [22].

To form DSB-induced chromosomal translocations, non-homologous chromosomes have to interact and engage in the repair process. In fact, DSB repair frequently entails a situation in which more than a single homologous template or multiple ends are available for strand invasion or ligation during HR or NHEJ, respectively. Judicious selection of the proper HR

templates or ends for ligation is crucial because illegitimate selection may generate chromosomal rearrangements and oncogenic transformation [7]. Several basic principles as to how a cell selects certain homologous templates and ligation partners for repair have recently emerged and have shed light on the molecular and biochemical attributes of chromosomal translocation suppression during DSB repair. Next, we will describe a few strategies that cells employ to repress HR-induced chromosomal translocations.

Repression of recombination between templates other than sister chromatids

HR is an important DSB repair pathway and is evolutionarily conserved from bacteria to human. Depending on the type of repair products and associated genes and gene products, HR can be further classified into three distinct categories: gene conversion, break-induced replication (BIR), and single-strand annealing (SSA) (Fig. 1) [23]. Substantial evidence links HR, in particular, BIR and SSA, to DSB-induced chromosomal translocations in yeast and mammals [7]. For instance, gross chromosomal rearrangements in yeast strains deficient in the intra-S phase DNA damage checkpoint partially depend on HR [16, 24]. Reduced expression of Pol alpha or Pol delta, polymerases involved in DNA replication, also leads to chromosomal rearrangements featuring nonreciprocal translocations congruent with their origin by BIR [17, 25]. Repetitive DNA elements such as Alu, LINE, and SINE sequences in metazoan genomes can promote excessive chromosomal rearrangements including chromosomal translocations by SSA. Telomere fusions during the telomeric crisis also rely on the SSA mechanism [26]. Such fusions will contribute to additional complex gross chromosomal rearrangements [27].

In principle, HR between sequences other than sister chromatids, specifically between ectopic sequences located on non-homologous chromosomes, can yield reciprocal or non-reciprocal chromosomal translocations. However, the paucity of HR-mediated chromosome translocations suggests that distinct mechanisms must exist to suppress aberrant repair and/or promote the faithful repair of DNA breaks (Table 1). Even if the detailed suppression mechanisms of these HR-induced chromosomal translocation need further elucidation, in this review, we will outline the two most prominent mechanisms: (1) restricting the sources of available homologous sequences flanking the breaks to sister chromatids, and (2) controlling the resolution of recombination intermediates to yield non-crossovers.

Bolstering proximity between sister chromatids

Sister chromatids that form after successful DNA replication are ideal substrates for HR due to their clonal identity. Recombination between sister chromatids does not carry the risk of mutagenesis or gross chromosomal rearrangements [28, 29]. The physical proximity between sister chromatids serves a key role in their preferential use as HR templates. However, proximity installed during replicative DNA synthesis alone may not be sufficient to ensure the use of sister chromatids as homologous templates. As a result, additional devices must be in place to promote the use of sister chromatids or alternatively to discourage the use of other potential templates (e.g., homologous chromosomes or ectopic sequences) during recombination. Nevertheless, uncovering the molecular basis underlying preferential usage of sister chromatids for recombination and the gene products that enforce this regulation faces significant challenges as there are few assays available to date to detect such recombinations genetically. Therefore, our knowledge in this process is substantially limited and relies heavily on those studies that detect unequal chromatid exchange events or cytological detection of recombination products in mammalian cells [30, 31]. Recently, a new assay that induces a site-specific DSB in one of the sister chromatids has been developed and has begun to shed light on the unique genetic features of equal chromatid exchange [29, 32].

One of the surprising discoveries from such studies was that sister chromatid recombination is encouraged by further enhancing the physical proximity between chromatids via the recruitment of the cohesion complex at or near DNA breaks (Fig. 2) [33]. Cohesin is a multi-subunit, ring-shaped protein complex composed of Smc1p, Smc3p, Scc1p, and Scc3p [34–36]. The cohesin ring is typically loaded onto DNA by the action of the Scc2p/Scc4p complex in the late G1 and early S phase of the cell cycle to tether sister chromatids together following replication [37]. In addition, upon DNA damage, cohesins are recruited to DSBs via their association with phosphorylated H2AX. Cohesins are then rapidly loaded onto the DNA in the vicinity of the DSBs to promote sister chromatid cohesion in a Chk1p- and Eco1p-dependent manner [38–40]. In support of this model, H2AX mutant replacing the serine residue that is subjected to damage-induced phosphorylation to alanine, or the cohesin mutants that have lost their ability to catalyze sister chromatid association upon DNA damage, have dramatically reduced sister chromatid recombination upon DNA damage [39] and exhibit an elevated frequencies of chromosomal translocations [41, 42]. Recently, yet another pair of SMC-like proteins, Smc5p/Smc6p, have also been implicated in recombination between sister chromatids, but how Smc5p/Smc6p stimulates recombination between sister chromatids is not clearly understood [43]. Additionally, the ATP-dependent chromatin remodeling RSC complex facilitates the loading of cohesin at DNA breaks and thereby promotes sister chromatid recombination ([44] SEL; manuscript in preparation).

Recombine when sister chromatids are available

Sister chromatids are produced during S phase and become available for recombination at S/G2. Reflecting the temporal availability of sister chromatids, it is logical that cells couple HR to the cell cycle and limit the recombination process primarily to the S/G2 phase. This optimizes the use of sister chromatids as recombination templates. To accomplish this goal, cells put one of the key initial steps of HR, single-stranded DNA (ssDNA) formation, under the influence of cell cycle control [45–47]. Efficient end resection and the formation of recombinogenic ssDNA at DNA breaks are catalyzed by nucleolytic degradation of 5' ends, which requires a combination of at least four nucleases (Dna2p, Exo1p, Mre11p, and Sae2p) and a helicase (Sgs1p) [48–50]. Multiple studies show that 5' end resection takes place at S/G2 even if limited end resection can still be detected at G1 [45, 46, 51, 52]. Central to the cell cycle-dependent 5' end processing lies Sae2p and its mammalian counterpart, CtIP, that undergo Cdk-dependent phosphorylation constituting the key regulatory circuit for activation of 5' end resection at S/G2 [47]. However, one or more Cdk1 targets besides Sae2p/CtIP likely exist to regulate 5' end resection, because deletion of *SAE2* alone does not fully eliminate end resection at G2. Notably, Mre11p/Rad50p/Xrs2p and Sae2p are all responsible for the initial 5' end processing but not for the sustained end resection that depends on Dna2p/Sgs1p and Exo1p [49, 50, 53]. Therefore, Dna2p, Sgs1p, and Exo1p are potential targets for Cdk1-dependent phosphorylation and cell cycle-dependent resection control. In this context, it is notable that Dna2p has been previously identified as a target of Cdk1-dependent phosphorylation in yeast [54].

End resection at G1 phase is further inhibited by the Ku and Lig4p core NHEJ proteins [51, 55, 56]. Consequently, deletion of NHEJ proteins in mice dramatically elevates recombination, in particular, non-reciprocal chromosome translocations [11, 13, 57, 58]. Ku and/or Lig4 complex has been suggested to inhibit end resection by promoting recurrent joining of DNA ends or by sterically protecting DNA ends from nuclease cleavage [51, 55, 56, 59].

Sister chromatids are identical to each other from top to bottom. In contrast, allelic sequences between homologous chromosomes can be divergent, and the extent of homology between ectopic homologous sequences is likely limited to regions immediately flanking the

DSB. When yeast cells were provided with two homologous templates; one homologous to the broken ends, and the other to sequences distant from the break, they elected the sequences distant from the broken ends preferentially to engage in recombination [60]. The results suggest that cells take into account the length and the sequence divergence of homologous partner when they select homologous partner and the preferential usage of longer stretches of homology emphasizes the use of sister chromatids as the ideal recombination templates. It also explains why a cell catalyzes substantial end resection (1–2 kb) prior to initiating recombination even if the recombination only requires a few hundred base pairs of single-stranded DNA [60–62]. Lastly, damage-induced checkpoints also stimulate recombination between sister chromatids and thus repress HR-mediated chromosomal translocations in a yet-to-be identified manner [31].

Resolution to remain faithful

Following strand pairing/exchange and repair synthesis, Holliday junction recombination intermediates should be resolved to generate either the patched or crossover recombination products (Fig. 1) [63]. As the crossover between ectopic homologies yields chromosome translocations, molecular mechanisms that modulate the extent of crossover product formation and the helicases and nucleases associated with this regulation have crucial importance in suppressing DSB-induced chromosome translocations [7]. Studies suggest that the Mus81p-Mms4p (Eme1) complex plays a role in Holliday junction resolution in metazoan and fission yeast systems [64, 65]. In line with this premise, Mus81p-Mms4p (Eme1) cleaves Holliday junction-mimicking molecules in vitro [64–67]. Deletion of Mus81p-Mms4p also leads to defects in meiosis, a hallmark of recombination deficiency [68–70]. However, Mus81p-Mms4p (Eme1) may not be the major Holliday junction resolvase because inactivation of Mus81p-Mms4p only causes a modest reduction in crossover product formation in meiosis [70]. Furthermore, the cleavage pattern of this nuclease is substantially different from that of the *E. coli* Holliday resolution enzyme, which cleaves Holliday junctions to yield symmetrical re-ligatable ends [71]. Recently, Gen1 (Yen1p in *S. cerevisiae*) [72] has been identified as a putative novel Holliday junction resolvase in human cells. Functional redundancy likely exists between these two enzyme complexes and potentially others. For example, nucleases such as Slx1p and Slx4p have been suggested, but not characterized, for the roles in resolving the Holliday junction intermediates during mitotic and/or meiotic recombination.

The Srs2p and Blm helicases have been implicated in repressing crossover product formation by either (1) stimulating strand displacement after initial pairing and DNA synthesis, or (2) dissolving recombination intermediates in conjunction with topoisomerase III and Blap75 proteins, respectively [73]. Both in yeast and mammals, the strand unwinding activity of Blm/Top3/Blap75 have been con-firmed biochemically using DNA substrates mimicking a double Holliday junction recombination intermediate [74, 75]. In agreement with its role in crossover suppression, inactivation of Blm leads to a dramatic increase in sister chromatid exchange that may be attributed to elevated crossover formation between sister chromatids in Bloom patients' cells [76]. Inactivation of Sgs1p also yields high levels of gross chromosomal rearrangements with elevated chromosome translocations [79, 24]. Besides its role in Holliday junction dissolution, Sgs1p also suppresses chromosome translocations between homeologous sequences along with a few mismatch repair factors [24, 77–79]. Indeed, inactivation of Blm leads to a high incidence of cancers, in particular, leukemia and lymphomas [80]. Recently, Mph1p/Fml1, the FANCM ortholog in budding and fission yeasts, respectively, as well as Rad54p, have been implicated in the branch migration step during HR [81, 82]. Deletion of *MPH1* or *FML1* substantially increases crossover product formation during mitotic recombination. Biochemically, Mph1p or Fml1 disassembles the D-loop via its branch migrating activity [83, 84]. Interestingly, excess

Mph1p or its helicase-deficient derivative interferes with Rad51p-dependent recombination and elevates gross chromosomal rearrangements [85].

Promiscuous joining of DNA breaks produces chromosome translocations

The role of NHEJ in the origin of clinically relevant chromosome translocations has been suggested for some time. Lack of sufficient lengths of homology at the breakpoint junctions opposes the involvement of homologous recombination in most chromosome translocations [4]. Comprehensive analysis of 358 spontaneous genome rearrangements in yeast selected by the loss of marker genes at the end of one arm in chromosome V indicated that they were generated primarily by NHEJ [86]. Furthermore, oncogenic chromosome translocations are often linked to V(D)J recombination or class switch recombination, two programmed gene rearrangements supported by NHEJ [21]. Experimentally, two simultaneous DNA DSBs induced on two different chromosomes are sufficient to form reciprocal chromosome translocations in yeast and mice [87, 88].

Even if NHEJ plays a major role in the formation of disease-causing chromosome translocations, evidence indicates that NHEJ-mediated chromosome translocations are extremely rare and thus cells possess the mechanisms to limit the promiscuous joining of broken DNA ends [89]. Often, NHEJ has been dubbed as the error-prone mechanism of DSB repair, as the process inherently carries the risk of mutagenesis at the repair junctions due to inevitable end processing prior to ligation [90]. Paradoxically, however, NHEJ is far more accurate than anticipated and very competent in restoring chromosome integrity. Overall, the fidelity of NHEJ is truly remarkable if one considers that multiple DNA breaks are produced throughout lymphocyte development and in most cases they are all repaired properly without chromosomal rearrangements.

In principle, three parameters can greatly influence the frequency of NHEJ-mediated chromosome translocations: the frequency of DSB production, the stability of the connection between broken DNA ends, and the mobility of broken DNA ends (Table 2). NHEJ-mediated chromosome translocations require two or more breaks to be generated at any given time and to be compatible for joining each other. Thereby, any conditions that elevate the frequency of DNA breaks will likely improve the chance of illegitimate end joining such as chromosome translocations. Even if DNA lesions that initiate NHEJ-mediated chromosome translocations are as yet unknown, a number of translocation break points feature topoisomerase II cleavage sites, fragile sites to impede ongoing replication forks, and potential recognition sequences for RAG protein cleavage [1, 91]. Additionally, CpG dinucleotides are disproportionately represented in the annotated breakpoints in blood cancers [92, 93]. Therefore, genes that can decrease the formation of these unscheduled/scheduled breaks are likely candidates for suppressing translocations, and mutations in these genes may increase the occurrence of blood cell malignancies.

In budding yeast, broken ends are tethered together and remained proximal to each other in an Mre11p/Rad50p/Xrs2p-, Sae2p-, and ATM-dependent manner even after several hours of DNA cleavage [53, 88, 94, 95]. End tethering also requires Rad52p during G2 [95]. The precise nature of end tethering remains unknown, but is likely mediated through various protein-protein interactions and is distinct from the annealing of imperfect repeat sequences flanking DNA breaks (as in SSA), since it occurs in G1 where DNA end resection is markedly decreased and is independent of Rad52p [88]. Gene mutations that disrupt end tethering (i.e., *sae2* and *tel1* mutants) dramatically elevate the frequency of chromosome translocations in a model system that induces two contemporaneous DNA breaks. This suggests that tethering inhibits the ends from engaging in promiscuous end joining and thus reduces the risk of NHEJ-mediated chromosome translocations [88].

Substantial controversy exists regarding the mobility of broken chromosomes and its role in the formation of chromosomal translocations. Live-cell microscopy suggests that radiation-induced DNA DSBs in general exhibit confined movement and larger motions are rare [96]. The fixed position of ultrasoft X-ray-induced DNA lesions detected in human fibroblasts is also consistent with the limited mobility of DNA breaks [97]. These observations support the hypothesis known as “contact first”, where the damaged chromatin should be proximal to each other prior to DNA damage to form chromosome translocations [98]. However, the efficient formation of chromosomal translocations from breaks on any two different chromosomes in yeast and mouse ES cells challenges such a model [87, 88]. Furthermore, analysis of over 300 chromosomal translocation events in yeast did not uncover prevalent chromosome translocations between distinct chromosome pairs [86], which would be expected if certain chromosomes are always near one another. Chemical- or radiation-induced breaks, dysfunctional telomeres, and activation-induced cytidine deaminase (AID)-induced DNA breaks in immunoglobulin class switch recombination recruit multiple proteins that function in DNA damage signaling and repair [99, 100] and may modulate break mobility. 53BP1, one of the factors rapidly recruited to these lesions, has recently been implicated in sustaining the mobility of DNA ends and thus stimulates long-range end joining and NHEJ-dependent chromosome translocations during V(D)J joining and telomere fusions [101, 102]. Despite evidence for long-range end joining, typically DNA ends that are not joined immediately with adjacent partners are quickly processed into recombination intermediates so that they are no longer available for chromosome translocation events [88, 103].

ATM suppresses NHEJ-mediated chromosome translocations

ATM is a central factor for end joining fidelity in yeast and metazoan cells. In *Saccharomyces cerevisiae*, the ATM ortholog Tel1p represses NHEJ-mediated chromosome translocations by tethering broken DNA ends together prior to 5' end processing (Fig. 3) [88, 103]. Tel1p achieves this by modulating multiple DNA damage response factors including Sae2p, H2AX, and Dun1p through its kinase activity [88]. During V(D)J joining, ATM deficiency causes dissociation of RAG-generated double-stranded ends from one another and yields hybrid joints and chromosome translocations [104, 105]. ATM itself has also been proposed to directly hold DNA ends together and prevent the persistence of DNA breaks during V(D)J recombination and class switch recombination [106, 107]. The downstream targets of ATM including H2AX, Mdc1, and 53BP1 also participate in catalyzing the end tethering reaction [108]. Accordingly, a truncation mutation in H2AX that blocks its phosphorylation by ATM and ATR [41, 42, 109] or inactivation of 53BP1 elevates chromosome instability including chromosome translocations [110]. ATM also contributes to faithful end joining as a signal transducer by promoting cell death when end joining is not immediately successful during V(D)J joining [106]. ATM deficiency also elevates chromosomal translocations induced from dysfunctional telomeres [16, 111, 112]. Collectively, ATM suppresses DSB-induced chromosomal translocation in multiple ways.

Does microhomology-mediated end joining generate DSB-induced chromosome translocations?

Microhomology-mediated end joining (MMEJ) is a distinct repair mechanism that uses substantially longer (5–15 bp) microhomologies for annealing and repairing DNA breaks [8]. Due to its reliance on the annealing of microhomology flanking DSB, MMEJ always produces deletions and thus is exceptionally mutagenic. Several lines of evidence strongly suggest that MMEJ can readily create chromosome rearrangements, including chromosome translocations, during DSB repair. First, reciprocal chromosome translocations formed by joining ends between I-*Scel*-induced DSBs and RAG-induced breaks occurs in the absence

of the NHEJ protein Ku [89]. Second, the breakpoint junctions from bladder carcinoma cells exhibit microhomology suggesting that MMEJ is the repair mechanism for chromosome translocations in these cells [113]. Third, deletion of Ku or Lig4 also leads to elevated DSB-induced chromosome translocations that showed typical signs of MMEJ: microhomology [13, 57, 86]. Alternative end joining also imparts in the repair of DSBs induced by reactive oxygen in cells from chronic myelogenous leukemia patient, and thus contributes to genome instability and progression of diseases [114]. Still, the frequency of chromosome translocations mediated by MMEJ has not yet been fully evaluated and thus the contribution of this mechanism to pathological chromosome translocations needs further research.

Perspectives

In this review, we described potential mechanisms to sustain the fidelity of chromosomes during the repair of DNA DSBs. The selection of proper homologous templates and the mode of intermediate resolution (in HR), maintaining proximity between proper ends, the mobility of DNA breaks, and the processing of un-repaired DNA breaks (by NHEJ) represent a few of the strategies to suppress chromosome translocations during DSB repair. Nevertheless, how all these processes are catalyzed and the biochemistry of the genetic components required for suppression of DSB-induced chromosomal translocations, have not yet fully emerged. It is particularly challenging as there are only a few reliable assays to monitor recombination between sister chromatids or address the biochemistry of end tethering events. Many of these repair fidelity maintenance mechanisms are also tightly associated with the chromatin structure at DNA breaks and thus understanding the nature of damaged chromatin is essential. It should also be noted that the fidelity of HR or NHEJ is regulated depending on the cell type and the status of cell differentiation: mitosis versus meiosis, fibroblast versus hematopoietic cells, diploid versus haploid. How cells switch on and/or off these multiple repair fidelity mechanisms is not yet known and needs to be elucidated.

In the past, several factors important for DSB repair regulation have been implicated in tumor suppression. However, it is an open question if the repair fidelity maintenance processes are universally targeted for tumor progression. In particular, the role of alternative end joining and MMEJ in chromosome translocations during DSB repair needs further scrutiny. In the future, we should carefully evaluate the role of individual components of these repair pathways and assess their general significance in tumor avoidance/progression to define a preventative/causal relationship between them and cancer. Such knowledge will lead to potential clinical applications that mediate DSB-induced chromosome translocation suppression for tumor protection and therapeutic intervention.

Acknowledgments

We are grateful to members of the S. Lee and K. Myung's laboratories for helpful discussions. We also thank to D. Walther and D. Daee for editorial help and J. Fekecs (NHGRI) for figure preparation. This work was supported by grants in the National Institutes of Health GM 083010 and the CTRC CDP pilot grants to S.E.L. and the intramural research program of the National Human Genome Research Institute, NIH to K.M. S.E.L. is a Leukemia and Lymphoma Society Scholar.

References

1. Aplan PD. Causes of oncogenic chromosomal translocation. *Trends Genet.* 2006; 22:46–55. [PubMed: 16257470]
2. Rabbitts TH. Chromosomal translocations in human cancer. *Nature.* 1994; 372:143–149. [PubMed: 7969446]

3. Nowell PC, Croce CM. Chromosomal approaches to the molecular basis of neoplasia. *Symp Fundam Cancer Res.* 1986; 39:17–29. [PubMed: 3321306]
4. Zhang Y, Rowley JD. Chromatin structural elements and chromosomal translocations in leukemia. *DNA Repair (Amst).* 2006; 5:1282–1297. [PubMed: 16893685]
5. Lee SE, Moore JK, Holmes A, Umez K, Kolodner RD, Haber JE. *Saccharomyces* Ku70, Mre11/Rad50 and RPA proteins regulate adaptation to G2/M arrest after DNA damage. *Cell.* 1998; 94:399–409. [PubMed: 9708741]
6. Lieber MR. The mechanism of human nonhomologous DNA end joining. *J Biol Chem.* 2008; 283:1–5. [PubMed: 17999957]
7. Haber JE. Transpositions and translocations induced by site-specific double-strand breaks in budding yeast. *DNA Repair (Amst).* 2006; 5:998–1009. [PubMed: 16807137]
8. McVey M, Lee SE. MMEJ repair of double-strand breaks (director's cut): deleted sequences and alternative endings. *Trends Genet.* 2008; 24:529–538. [PubMed: 18809224]
9. Krogh BO, Symington LS. Recombination proteins in yeast. *Annu Rev Genet.* 2004; 38:233–271. [PubMed: 15568977]
10. Daley JM, Palmboos PL, Wu D, Wilson TE. Nonhomologous end joining in yeast. *Annu Rev Genet.* 2005; 39:431–451. [PubMed: 16285867]
11. Difilippantonio MJ, Zhu J, Chen HT, Meffre E, Nussenzweig MC, Max EE, Ried T, Nussenzweig A. DNA repair protein Ku80 suppresses chromosomal aberrations and malignant transformation. *Nature.* 2000; 404:510–514. [PubMed: 10761921]
12. Weinstock DM, Richardson CA, Elliott B, Jasin M. Modeling oncogenic translocations: distinct roles for double-strand break repair pathways in translocation formation in mammalian cells. *DNA Repair (Amst).* 2006; 5:1065–1074. [PubMed: 16815104]
13. Ferguson DO, Sekiguchi JM, Chang S, Frank KM, Gao Y, DePinho RA, Alt FW. The nonhomologous end-joining pathway of DNA repair is required for genomic stability and the suppression of translocations. *Proc Natl Acad Sci USA.* 2000; 97:6630–6633. [PubMed: 10823907]
14. Kolodner RD, Putnam CD, Myung K. Maintenance of genome stability in *Saccharomyces cerevisiae*. *Science.* 2002; 297:552–557. [PubMed: 12142524]
15. Lopes M, Cotta-Ramusino C, Liberi G, Foiani M. Branch migrating sister chromatid junctions form at replication origins through Rad51/Rad52-independent mechanisms. *Mol Cell.* 2003; 12:1499–1510. [PubMed: 14690603]
16. Myung K, Datta A, Kolodner RD. Suppression of spontaneous chromosomal rearrangements by S phase checkpoint functions in *Saccharomyces cerevisiae*. *Cell.* 2001; 104:397–408. [PubMed: 11239397]
17. Lemoine FJ, Degtyareva NP, Lobachev K, Petes TD. Chromosomal translocations in yeast induced by low levels of DNA polymerase a model for chromosome fragile sites. *Cell.* 2005; 120:587–598. [PubMed: 15766523]
18. Schwartz M, Zlotorynski E, Goldberg M, Ozeri E, Rahat A, le Sage C, Chen BP, Chen DJ, Agami R, Kerem B. Homologous recombination and nonhomologous end-joining repair pathways regulate fragile site stability. *Genes Dev.* 2005; 19:2715–2726. [PubMed: 16291645]
19. Admire A, Shanks L, Danzl N, Wang M, Weier U, Stevens W, Hunt E, Weinert T. Cycles of chromosome instability are associated with a fragile site and are increased by defects in DNA replication and checkpoint controls in yeast. *Genes Dev.* 2006; 20:159–173. [PubMed: 16384935]
20. Miller RW. Delayed radiation effects in atomic-bomb survivors. Major observations by the Atomic Bomb Casualty Commission are evaluated. *Science.* 1969; 166:569–574. [PubMed: 4898614]
21. Lieber MR, Yu K, Raghavan SC. Roles of nonhomologous DNA end joining, V(D)J recombination, and class switch recombination in chromosomal translocations. *DNA Repair (Amst).* 2006; 5:1234–1245. [PubMed: 16793349]
22. Sung PA, Libura J, Richardson C. Etoposide and illegitimate DNA double-strand break repair in the generation of MLL translocations: new insights and new questions. *DNA Repair (Amst).* 2006; 5:1109–1118. [PubMed: 16809075]
23. Paques F, Haber JE. Multiple pathways of recombination induced by double-strand breaks in *Saccharomyces cerevisiae*. *Microbiol Mol Biol Rev.* 1999; 63:349–404. [PubMed: 10357855]

24. Schmidt KH, Wu J, Kolodner RD. Control of translocations between highly diverged genes by Sgs1, the *Saccharomyces cerevisiae* homolog of the Bloom's syndrome protein. *Mol Cell Biol.* 2006; 26:5406–5420. [PubMed: 16809776]
25. Lemoine FJ, Degtyareva NP, Kokoska RJ, Petes TD. Reduced levels of DNA polymerase delta induce chromosome fragile site instability in yeast. *Mol Cell Biol.* 2008; 28:5359–5368. [PubMed: 18591249]
26. Wang X, Baumann P. Chromosome fusions following telomere loss are mediated by single-strand annealing. *Mol Cell.* 2008; 31:463–473. [PubMed: 18722173]
27. VanHulle K, Lemoine FJ, Narayanan V, Downing B, Hull K, McCullough C, Bellinger M, Lobachev K, Petes TD, Malkova A. Inverted DNA repeats channel repair of distant double-strand breaks into chromatid fusions and chromosomal rearrangements. *Mol Cell Biol.* 2007; 27:2601–2614. [PubMed: 17242181]
28. Xie A, Puget N, Shim I, Odate S, Jarzyna I, Bassing CH, Alt FW, Scully R. Control of sister chromatid recombination by histone H2AX. *Mol Cell.* 2004; 16:1017–1025. [PubMed: 15610743]
29. Gonzalez-Barrera S, Cortes-Ledesma F, Wellinger RE, Aguilera A. Equal sister chromatid exchange is a major mechanism of double-strand break repair in yeast. *Mol Cell.* 2003; 11:1661–1671. [PubMed: 12820977]
30. Chaganti RS, Schonberg S, German J. A manyfold increase in sister chromatid exchanges in Bloom's syndrome lymphocytes. *Proc Natl Acad Sci USA.* 1974; 71:4508–4512. [PubMed: 4140506]
31. Fasullo M, Bennett T, AhChing P, Koudelik J. The *Saccharomyces cerevisiae* RAD9 checkpoint reduces the DNA damage-associated stimulation of directed translocations. *Mol Cell Biol.* 1998; 18:1190–1200. [PubMed: 9488434]
32. Cortes-Ledesma F, Tous C, Aguilera A. Different genetic requirements for repair of replication-born double-strand breaks by sister-chromatid recombination and break-induced replication. *Nucleic Acids Res.* 2007; 35:6560–6570. [PubMed: 17905819]
33. Strom L, Sjogren C. Chromosome segregation and double-strand break repair—a complex connection. *Curr Opin Cell Biol.* 2007; 19:344–349. [PubMed: 17466504]
34. Gruber S, Haering CH, Nasmyth K. Chromosomal cohesin forms a ring. *Cell.* 2003; 112:765–777. [PubMed: 12654244]
35. Haering CH, Lowe J, Hochwagen A, Nasmyth K. Molecular architecture of SMC proteins and the yeast cohesin complex. *Mol Cell.* 2002; 9:773–788. [PubMed: 11983169]
36. Haering CH, Farcas AM, Arumugam P, Metson J, Nasmyth K. The cohesin ring concatenates sister DNA molecules. *Nature.* 2008; 454:297–301. [PubMed: 18596691]
37. Gruber S, Arumugam P, Katou Y, Kuglitsch D, Helmhart W, Shirahige K, Nasmyth K. Evidence that loading of cohesin onto chromosomes involves opening of its SMC hinge. *Cell.* 2006; 127:523–537. [PubMed: 17081975]
38. Strom L, Lindroos HB, Shirahige K, Sjogren C. Postreplicative recruitment of cohesin to double-strand breaks is required for DNA repair. *Mol Cell.* 2004; 16:1003–1015. [PubMed: 15610742]
39. Unal E, Arbel-Eden A, Sattler U, Shroff R, Lichten M, Haber JE, Koshland D. DNA damage response pathway uses histone modification to assemble a double-strand break-specific cohesin domain. *Mol Cell.* 2004; 16:991–1002. [PubMed: 15610741]
40. Strom L, Karlsson C, Lindroos HB, Wedahl S, Katou Y, Shirahige K, Sjogren C. Postreplicative formation of cohesion is required for repair and induced by a single DNA break. *Science.* 2007; 317:242–245. [PubMed: 17626884]
41. Celeste A, Petersen S, Romanienko PJ, Fernandez-Capetillo O, Chen HT, Sedelnikova OA, Reina-San-Martin B, Coppola V, Meffre E, Difilippantonio MJ, Redon C, Pilch DR, Oлару A, Eckhaus M, Camerini-Otero RD, Tessarollo L, Livak F, Manova K, Bonner WM, Nussenzweig MC, Nussenzweig A. Genomic instability in mice lacking histone H2AX. *Science.* 2002; 296:922–927. [PubMed: 11934988]
42. Bassing CH, Suh H, Ferguson DO, Chua KF, Manis J, Eckersdorff M, Gleason M, Bronson R, Lee C, Alt FW. Histone H2AX: a dosage-dependent suppressor of oncogenic translocations and tumors. *Cell.* 2003; 114:359–370. [PubMed: 12914700]

43. De Piccoli G, Cortes-Ledesma F, Ira G, Torres-Rosell J, Uhle S, Farmer S, Hwang JY, Machin F, Ceschia A, McAleenan A, Cordon-Preciado V, Clemente-Blanco A, Vilella-Mitjana F, Ullal P, Jarmuz A, Leitao B, Bressan D, Dotiwala F, Papusha A, Zhao X, Myung K, Haber JE, Aguilera A, Aragon L. Smc5-Smc6 mediate DNA double-strand-break repair by promoting sister-chromatid recombination. *Nat Cell Biol.* 2006; 8:1032–1034. [PubMed: 16892052]
44. Liang B, Qiu J, Ratnakumar K, Laurent BC. RSC functions as an early double-strand-break sensor in the cell's response to DNA damage. *Curr Biol.* 2007; 17:1432–1437. [PubMed: 17689960]
45. Ira G, Pellicoli A, Balijja A, Wang X, Fiorani S, Carotenuto W, Liberi G, Bressan D, Wan L, Hollingsworth NM, Haber JE, Foiani M. DNA end resection, homologous recombination and DNA damage checkpoint activation require CDK1. *Nature.* 2004; 431:1011–1017. [PubMed: 15496928]
46. Aylon Y, Liefshitz B, Kupiec M. The CDK regulates repair of double-strand breaks by homologous recombination during the cell cycle. *EMBO J.* 2004; 23:4868–4875. [PubMed: 15549137]
47. Huertas P, Cortes-Ledesma F, Sartori AA, Aguilera A, Jackson SP. CDK targets Sae2 to control DNA-end resection and homologous recombination. *Nature.* 2008; 455:689–692. [PubMed: 18716619]
48. Gravel S, Chapman JR, Magill C, Jackson SP. DNA helicases Sgs1 and BLM promote DNA double-strand break resection. *Genes Dev.* 2008; 22:2767–2772. [PubMed: 18923075]
49. Mimitou EP, Symington LS. Sae2, Exo1 and Sgs1 collaborate in DNA double-strand break processing. *Nature.* 2008; 455:770–774. [PubMed: 18806779]
50. Zhu Z, Chung WH, Shim EY, Lee SE, Ira G. Sgs1 helicase and two nucleases Dna2 and Exo1 resect DNA double-strand break ends. *Cell.* 2008; 134:981–994. [PubMed: 18805091]
51. Zhang Y, Hefferin ML, Chen L, Shim EY, Tseng HM, Kwon Y, Sung P, Lee SE, Tomkinson AE. Role of Dnl4-Lif1 in nonhomologous end-joining repair complex assembly and suppression of homologous recombination. *Nat Struct Mol Biol.* 2007; 14:639–646. [PubMed: 17589524]
52. Barlow JH, Lisby M, Rothstein R. Differential regulation of the cellular response to DNA double-strand breaks in G1. *Mol Cell.* 2008; 30:73–85. [PubMed: 18406328]
53. Clerici M, Mantiero D, Lucchini G, Longhese MP. The *Saccharomyces cerevisiae* Sae2 protein promotes resection and bridging of double strand break ends. *J Biol Chem.* 2005; 280:38631–38638. [PubMed: 16162495]
54. Ubersax JA, Woodbury EL, Quang PN, Paraz M, Blethrow JD, Shah K, Shokat KM, Morgan DO. Targets of the cyclin-dependent kinase Cdk1. *Nature.* 2003; 425:859–864. [PubMed: 14574415]
55. Zierhut C, Diffley JF. Break dosage, cell cycle stage and DNA replication influence DNA double strand break response. *EMBO J.* 2008; 27:1875–1885. [PubMed: 18511906]
56. Clerici M, Mantiero D, Guerini I, Lucchini G, Longhese MP. The Yku70–Yku80 complex contributes to regulate double-strand break processing and checkpoint activation during the cell cycle. *EMBO Rep.* 2008; 9:810–818. [PubMed: 18600234]
57. Gao Y, Ferguson DO, Xie W, Manis JP, Sekiguchi J, Frank KM, Chaudhuri J, Horner J, DePinho RA, Alt FW. Interplay of p53 and DNA-repair protein XRCC4 in tumorigenesis, genomic stability and development. *Nature.* 2000; 404:897–900. [PubMed: 10786799]
58. Myung K, Ghosh G, Fattah FJ, Li G, Kim H, Dutia A, Pak E, Smith S, Hendrickson EA. Regulation of telomere length and suppression of genomic instability in human somatic cells by Ku86. *Mol Cell Biol.* 2004; 24:5050–5059. [PubMed: 15143195]
59. Wasko BM, Holland CL, Resnick MA, Lewis LK. Inhibition of DNA double-strand break repair by the Ku heterodimer in mrx mutants of *Saccharomyces cerevisiae*. *DNA Repair (Amst).* 2009; 8:162–169. [PubMed: 18992851]
60. Inbar O, Kupiec M. Homology search and choice of homologous partner during mitotic recombination. *Mol Cell Biol.* 1999; 19:4134–4142. [PubMed: 10330153]
61. Budd ME, Campbell JL. Interplay of Mre11 nuclease with Dna2 plus Sgs1 in Rad51-dependent recombinational repair. *PLoS ONE.* 2009; 4:e4267. [PubMed: 19165339]
62. Sugawara N, Haber JE. Characterization of double-strand break-induced recombination: homology requirements and single-stranded DNA formation. *Mol Cell Biol.* 1992; 12:563–575. [PubMed: 1732731]

63. Heyer WD. Recombination: Holliday junction resolution and crossover formation. *Curr Biol*. 2004; 14:R56–R58. [PubMed: 14738748]
64. Chen XB, Melchionna R, Denis CM, Gaillard PH, Blasina A, Van de Weyer I, Boddy MN, Russell P, Vialard J, McGowan CH. Human Mus81-associated endonuclease cleaves Holliday junctions in vitro. *Mol Cell*. 2001; 8:1117–1127. [PubMed: 11741546]
65. Boddy MN, Gaillard PH, McDonald WH, Shanahan P, Yates JR 3rd, Russell P. Mus81-Eme1 are essential components of a Holliday junction resolvase. *Cell*. 2001; 107:537–548. [PubMed: 11719193]
66. Gaskell LJ, Osman F, Gilbert RJ, Whitby MC. Mus81 cleavage of Holliday junctions: a failsafe for processing meiotic recombination intermediates? *EMBO J*. 2007; 26:1891–1901. [PubMed: 17363897]
67. Kaliraman V, Mullen JR, Fricke WM, Bastin-Shanower SA, Brill SJ. Functional overlap between Sgs1-Top3 and the Mms4-Mus81 endonuclease. *Genes Dev*. 2001; 15:2730–2740. [PubMed: 11641278]
68. Oh SD, Lao JP, Taylor AF, Smith GR, Hunter N. RecQ helicase, Sgs1, and XPF family endonuclease, Mus81-Mms4, resolve aberrant joint molecules during meiotic recombination. *Mol Cell*. 2008; 31:324–336. [PubMed: 18691965]
69. Jessop L, Lichten M. Mus81/Mms4 endonuclease and Sgs1 helicase collaborate to ensure proper recombination intermediate metabolism during meiosis. *Mol Cell*. 2008; 31:313–323. [PubMed: 18691964]
70. de los Santos T, Hunter N, Lee C, Larkin B, Loidl J, Hollingsworth NM. The Mus81/Mms4 endonuclease acts independently of double-Holliday junction resolution to promote a distinct subset of crossovers during meiosis in budding yeast. *Genetics*. 2003; 164:81–94. [PubMed: 12750322]
71. Haber JE, Heyer WD. The fuss about Mus81. *Cell*. 2001; 107:551–554. [PubMed: 11733053]
72. Ip SC, Rass U, Blanco MG, Flynn HR, Skehel JM, West SC. Identification of Holliday junction resolvases from humans and yeast. *Nature*. 2008; 456:357–361. [PubMed: 19020614]
73. Ira G, Malkova A, Liberi G, Foiani M, Haber JE. Srs2 and Sgs1-Top3 suppress crossovers during double-strand break repair in yeast. *Cell*. 2003; 115:401–411. [PubMed: 14622595]
74. Wu L, Hickson ID. The Bloom's syndrome helicase suppresses crossing over during homologous recombination. *Nature*. 2003; 426:870–874. [PubMed: 14685245]
75. Chen CF, Brill SJ. Binding and activation of DNA topoisomerase III by the Rmi1 subunit. *J Biol Chem*. 2007; 282:28971–28979. [PubMed: 17693398]
76. Ellis NA, Lennon DJ, Proytcheva M, Alhadeff B, Henderson EE, German J. Somatic intragenic recombination within the mutated locus BLM can correct the high sister-chromatid exchange phenotype of Bloom syndrome cells. *Am J Hum Genet*. 1995; 57:1019–1027. [PubMed: 7485150]
77. Spell RM, Jinks-Robertson S. Examination of the roles of Sgs1 and Srs2 helicases in the enforcement of recombination fidelity in *Saccharomyces cerevisiae*. *Genetics*. 2004; 168:1855–1865. [PubMed: 15611162]
78. Sugawara N, Goldfarb T, Studamire B, Alani E, Haber JE. Heteroduplex rejection during single-strand annealing requires Sgs1 helicase and mismatch repair proteins Msh2 and Msh6 but not Pms1. *Proc Natl Acad Sci USA*. 2004; 101:9315–9320. [PubMed: 15199178]
79. Myung K, Datta A, Chen C, Kolodner RD. SGS1, the *Saccharomyces cerevisiae* homologue of BLM and WRN, suppresses genome instability and homeologous recombination. *Nat Genet*. 2001; 27:113–116. [PubMed: 11138010]
80. Bohr VA. Rising from the RecQ-age: the role of human RecQ helicases in genome maintenance. *Trends Biochem Sci*. 2008; 33:609–620. [PubMed: 18926708]
81. Bugreev DV, Mazina OM, Mazin AV. Rad54 protein promotes branch migration of Holliday junctions. *Nature*. 2006; 442:590–593. [PubMed: 16862129]
82. Gari K, Decaillet C, Stasiak AZ, Stasiak A, Constantinou A. The Fanconi anemia protein FANCM can promote branch migration of Holliday junctions and replication forks. *Mol Cell*. 2008; 29:141–148. [PubMed: 18206976]

83. Prakash R, Satory D, Dray E, Papusha A, Scheller J, Kramer W, Krejci L, Klein H, Haber JE, Sung P, Ira G. Yeast Mph1 helicase dissociates Rad51-made D-loops: implications for crossover control in mitotic recombination. *Genes Dev.* 2009; 23:67–79. [PubMed: 19136626]
84. Sun W, Nandi S, Osman F, Ahn JS, Jakovleska J, Lorenz A, Whitby MC. The FANCM ortholog Fml1 promotes recombination at stalled replication forks and limits crossing over during DNA double-strand break repair. *Mol Cell.* 2008; 32:118–128. [PubMed: 18851838]
85. Banerjee S, Smith S, Oum JH, Liaw HJ, Hwang JY, Sikdar N, Motegi A, Lee SE, Myung K. Mph1p promotes gross chromosomal rearrangement through partial inhibition of homologous recombination. *J Cell Biol.* 2008; 181:1083–1093. [PubMed: 18591428]
86. Putnam CD, Pennaneach V, Kolodner RD. *Saccharomyces cerevisiae* as a model system to define the chromosomal instability phenotype. *Mol Cell Biol.* 2005; 25:7226–7238. [PubMed: 16055731]
87. Richardson C, Jasin M. Frequent chromosomal translocations induced by DNA double-strand breaks. *Nature.* 2000; 405:697–700. [PubMed: 10864328]
88. Lee K, Zhang Y, Lee SE. *Saccharomyces cerevisiae* ATM orthologue suppresses break-induced chromosome translocations. *Nature.* 2008; 454:543–546. [PubMed: 18650924]
89. Weinstock DM, Brunet E, Jasin M. Formation of NHEJ-derived reciprocal chromosomal translocations does not require Ku70. *Nat Cell Biol.* 2007; 9:978–981. [PubMed: 17643113]
90. Gu J, Lieber MR. Mechanistic flexibility as a conserved theme across 3 billion years of nonhomologous DNA end-joining. *Genes Dev.* 2008; 22:411–415. [PubMed: 18281457]
91. Raghavan SC, Swanson PC, Wu X, Hsieh CL, Lieber MR. A non-B-DNA structure at the Bcl-2 major breakpoint region is cleaved by the RAG complex. *Nature.* 2004; 428:88–93. [PubMed: 14999286]
92. Ramiro AR, Jankovic M, Callen E, Difilippantonio S, Chen HT, McBride KM, Eisenreich TR, Chen J, Dickins RA, Lowe SW, Nussenzweig A, Nussenzweig MC. Role of genomic instability and p53 in AID-induced c-myc-Igh translocations. *Nature.* 2006; 440:105–109. [PubMed: 16400328]
93. Tsai AG, Lu H, Raghavan SC, Muschen M, Hsieh CL, Lieber MR. Human chromosomal translocations at CpG sites and a theoretical basis for their lineage and stage specificity. *Cell.* 2008; 135:1130–1142. [PubMed: 19070581]
94. Lobachev K, Vitriol E, Stemple J, Resnick MA, Bloom K. Chromosome fragmentation after induction of a double-strand break is an active process prevented by the RMX repair complex. *Curr Biol.* 2004; 14:2107–2112. [PubMed: 15589152]
95. Kaye JA, Melo JA, Cheung SK, Vaze MB, Haber JE, Toczyski DP. DNA breaks promote genomic instability by impeding proper chromosome segregation. *Curr Biol.* 2004; 14:2096–2106. [PubMed: 15589151]
96. Jakob B, Splinter J, Durante M, Taucher-Scholz G. Live cell microscopy analysis of radiation-induced DNA double-strand break motion. *Proc Natl Acad Sci USA.* 2009; 106(9):3172–3177. [PubMed: 19221031]
97. Nelms BE, Maser RS, MacKay JF, Lagally MG, Petrini JH. In situ visualization of DNA double-strand break repair in human fibroblasts. *Science.* 1998; 280:590–592. [PubMed: 9554850]
98. Soutoglou E, Dorn JF, Sengupta K, Jasin M, Nussenzweig A, Ried T, Danuser G, Misteli T. Positional stability of single double-strand breaks in mammalian cells. *Nat Cell Biol.* 2007; 9:675–682. [PubMed: 17486118]
99. Franco S, Alt FW, Manis JP. Pathways that suppress programmed DNA breaks from progressing to chromosomal breaks and translocations. *DNA Repair (Amst).* 2006; 5:1030–1041. [PubMed: 16934538]
100. Bailey SM, Murnane JP. Telomeres, chromosome instability and cancer. *Nucleic Acids Res.* 2006; 34:2408–2417. [PubMed: 16682448]
101. Dimitrova N, Chen YC, Spector DL, de Lange T. 53BP1 promotes non-homologous end joining of telomeres by increasing chromatin mobility. *Nature.* 2008; 456:524–528. [PubMed: 18931659]
102. Difilippantonio S, Gapud E, Wong N, Huang CY, Mahowald G, Chen HT, Kruhlak MJ, Callen E, Livak F, Nussenzweig MC, Sleckman BP, Nussenzweig A. 53BP1 facilitates long-range DNA end-joining during V(D)J recombination. *Nature.* 2008; 456:529–533. [PubMed: 18931658]

103. Mantiero D, Clerici M, Lucchini G, Longhese MP. Dual role for *Saccharomyces cerevisiae* Tel1 in the checkpoint response to double-strand breaks. *EMBO Rep.* 2007; 8:380–387. [PubMed: 17347674]
104. Bredemeyer AL, Sharma GG, Huang CY, Helmink BA, Walker LM, Khor KC, Nuskey B, Sullivan KE, Pandita TK, Bassing CH, Sleckman BP. ATM stabilizes DNA double-strand-break complexes during V(D)J recombination. *Nature.* 2006; 442:466–470. [PubMed: 16799570]
105. Bredemeyer AL, Huang CY, Walker LM, Bassing CH, Sleckman BP. Aberrant V(D)J recombination in ataxia telangiectasia mutated-deficient lymphocytes is dependent on nonhomologous DNA end joining. *J Immunol.* 2008; 181:2620–2625. [PubMed: 18684952]
106. Callen E, Jankovic M, Difilippantonio S, Daniel JA, Chen HT, Celeste A, Pellegrini M, McBride K, Wangsa D, Bredemeyer AL, Sleckman BP, Ried T, Nussenzweig M, Nussenzweig A. ATM prevents the persistence and propagation of chromosome breaks in lymphocytes. *Cell.* 2007; 130:63–75. [PubMed: 17599403]
107. Reina-San-Martin B, Chen HT, Nussenzweig A, Nussenzweig MC. ATM is required for efficient recombination between immunoglobulin switch regions. *J Exp Med.* 2004; 200:1103–1110. [PubMed: 15520243]
108. Bassing CH, Alt FW. H2AX may function as an anchor to hold broken chromosomal DNA ends in close proximity. *Cell Cycle.* 2004; 3:149–153. [PubMed: 14712078]
109. Franco S, Gostissa M, Zha S, Lombard DB, Murphy MM, Zarrin AA, Yan C, Tepsuporn S, Morales JC, Adams MM, Lou Z, Bassing CH, Manis JP, Chen J, Carpenter PB, Alt FW. H2AX prevents DNA breaks from progressing to chromosome breaks and translocations. *Mol Cell.* 2006; 21:201–214. [PubMed: 16427010]
110. Ward IM, Difilippantonio S, Minn K, Mueller MD, Molina JR, Yu X, Frisk CS, Ried T, Nussenzweig A, Chen J. 53BP1 cooperates with p53 and functions as a haploinsufficient tumor suppressor in mice. *Mol Cell Biol.* 2005; 25:10079–10086. [PubMed: 16260621]
111. Chan SW, Blackburn EH. Telomerase and ATM/Tel1p protect telomeres from nonhomologous end joining. *Mol Cell.* 2003; 11:1379–1387. [PubMed: 12769860]
112. Karlseder J, Broccoli D, Dai Y, Hardy S, de Lange T. p53- and ATM-dependent apoptosis induced by telomeres lacking TRF2. *Science.* 1999; 283:1321–1325. [PubMed: 10037601]
113. Bentley J, Diggle CP, Harnden P, Knowles MA, Kiltie AE. DNA double strand break repair in human bladder cancer is error prone and involves microhomology-associated end-joining. *Nucleic Acids Res.* 2004; 32:5249–5259. [PubMed: 15466592]
114. Sallmyr A, Tomkinson AE, Rassool FV. Up-regulation of WRN and DNA ligase IIIalpha in chronic myeloid leukemia: consequences for the repair of DNA double-strand breaks. *Blood.* 2008; 112:1413–1423. [PubMed: 18524993]
115. Heidinger-Pauli JM, Unal E, Guacci V, Koshland D. The kleisin subunit of cohesin dictates damage-induced cohesion. *Mol Cell.* 2008; 31:47–56. [PubMed: 18614046]
116. Hwang JY, Smith S, Ceschia A, Torres-Rosell J, Aragon L, Myung K. Smc5-Smc6 complex suppresses gross chromosomal rearrangements mediated by break-induced replications. *DNA Repair (Amst).* 2008; 7:1426–1436. [PubMed: 18585101]
117. Smith S, Gupta A, Kolodner RD, Myung K. Suppression of gross chromosomal rearrangements by the multiple functions of the Mre11-Rad50-Xrs2 complex in *Saccharomyces cerevisiae*. *DNA Repair (Amst).* 2005; 4:606–617. [PubMed: 15811632]
118. Lengsfeld BM, Rattray AJ, Bhaskara V, Ghirlando R, Paull TT. Sae2 is an endonuclease that processes hairpin DNA cooperatively with the Mre11/Rad50/Xrs2 complex. *Mol Cell.* 2007; 28:638–651. [PubMed: 18042458]
119. Barber LJ, Youds JL, Ward JD, McIlwraith MJ, O'Neil NJ, Petalcorin MI, Martin JS, Collis SJ, Cantor SB, Auclair M, Tissenbaum H, West SC, Rose AM, Boulton SJ. RTEL1 maintains genomic stability by suppressing homologous recombination. *Cell.* 2008; 135:261–271. [PubMed: 18957201]
120. Chiolo I, Saponaro M, Baryshnikova A, Kim JH, Seo YS, Liberi G. The human F-Box DNA helicase FBH1 faces *Saccharomyces cerevisiae* Srs2 and postreplication repair pathway roles. *Mol Cell Biol.* 2007; 27:7439–7450. [PubMed: 17724085]

121. Dronkert ML, Beverloo HB, Johnson RD, Hoeijmakers JH, Jasin M, Kanaar R. Mouse RAD54 affects DNA double-strand break repair and sister chromatid exchange. *Mol Cell Biol.* 2000; 20:3147–3156. [PubMed: 10757799]
122. Osman F, Dixon J, Doe CL, Whitby MC. Generating crossovers by resolution of nicked Holliday junctions: a role for Mus81-Eme1 in meiosis. *Mol Cell.* 2003; 12:761–774. [PubMed: 14527420]
123. Blais V, Gao H, Elwell CA, Boddy MN, Gaillard PH, Russell P, McGowan CH. RNA interference inhibition of Mus81 reduces mitotic recombination in human cells. *Mol Biol Cell.* 2004; 15:552–562. [PubMed: 14617801]
124. Stewart GS, Wang B, Bignell CR, Taylor AM, Elledge SJ. MDC1 is a mediator of the mammalian DNA damage checkpoint. *Nature.* 2003; 421:961–966. [PubMed: 12607005]
125. Lou Z, Minter-Dykhouse K, Franco S, Gostissa M, Rivera MA, Celeste A, Manis JP, van Deursen J, Nussenzweig A, Paull TT, Alt FW, Chen J. MDC1 maintains genomic stability by participating in the amplification of ATM-dependent DNA damage signals. *Mol Cell.* 2006; 21:187–200. [PubMed: 16427009]
126. Chen C, Kolodner RD. Gross chromosomal rearrangements in *Saccharomyces cerevisiae* replication and recombination defective mutants. *Nat Genet.* 1999; 23:81–85. [PubMed: 10471504]
127. Chen C, Umezu K, Kolodner RD. Chromosomal rearrangements occur in *S. cerevisiae* rfa1 mutator mutants due to mutagenic lesions processed by double-strand-break repair. *Mol Cell.* 1998; 2:9–22. [PubMed: 9702187]

Fig. 1.

Mechanisms of homologous recombination-mediated chromosome translocations. DNA double-strand breaks (DSBs) are processed by nucleolytic degradation of the 5' strand and the resulting 3' single-stranded DNA either invades a homologous template such as a sister chromatid, a homologous chromosome, or an ectopic region of homology (in BIR and gene conversion), or the flanking repeats anneal to form annealed intermediates (in SSA). In gene conversion, DNA synthesis displaces the D-loop and captures the 2nd end to yield a double Holliday junction (dHJ). Resolution by a nuclease may produce chromosome translocations (crossover product). Alternatively, 3' ssDNA ends that invade and initiate DNA synthesis is dissociated from the donor, re-anneals with the other 3' ssDNA tails to yield non-crossover products (synthesis dependent strand annealing, SDSA). In BIR, lack of homology in the 2nd end leads to the assembly of a replication fork by the invading strand and repair synthesis continues to the end of the chromosome, generating a nonreciprocal chromosome translocation. In SSA, annealing between repetitive sequences on different chromosomes is further processed by 3' flap removal and gap synthesis to yield chromosome translocations

Fig. 2.

DNA damage-induced cohesin loading at DNA breaks stimulates recombination between sister chromatids. Sister chromatid cohesion is established by loading cohesin complex at cohesion attachment regions (CAR, shown in scattered at chromosome arm) during S phase. Upon DNA break induction, cohesin is rapidly recruited to the ~50-kb flanking region adjacent to DNA breaks (shown in *arrows*). Break-induced cohesin recruitment depends on Mre11p/Rad50p/Xrs2p complex that binds immediately next to DNA breaks (~2 kb from each side of a DNA break) and phosphorylated H2AX that covers the region co-occupying with cohesin. Cohesin accumulation reinforces the proximity of sister chromatid to further stimulate recombination between sister chromatids

Fig. 3.

ATM plays a central role in suppressing NHEJ-mediated chromosome translocations. ATM suppresses chromosome translocations by facilitating end tethering and degradation of the 5' strand to yield recombinogenic 3' single stranded DNAs. ATM facilitates end tethering and resection by phosphorylation of Mre11/Rad50/Xrs2-CtIP and H2AX. Phosphorylation of H2AX and methylation of H3 lysine 73 by the methylase Dot1 lead to the recruitment of 53BP1 and modulate the mobility of DNA ends for long-range end joining. Cells that fail to repair broken chromosomes are eliminated by ATM-dependent apoptosis

Table 1

Genes and their biochemical activities that potentially suppress recombination-induced chromosome translocations

Name ^a	Function	Biochemical activity	Reference
Smc1 (Smc1p), Smc3 (Smc3p), Scc1 (Scc1p), Scc3 (Scc3p) (Scc2p), Scc4 (Scc4p)	Tether sister chromatids; promote sister chromatid recombination	Sister chromatid cohesion	[38, 39]
Chk1 (Chk1p)	Regulate damage-induced sister chromatid cohesion	Protein kinase	[115]
(Smc5p/Smc6p)	Promote sister chromatid recombination; suppress gross chromosomal rearrangements (GCRs)	SMC like proteins; Sumo E3 ligase	[43, 116]
(RSC)	Loading of cohesin upon DNA damage	ATP-dependent chromatin remodeler	[44]
(Rad9p)	n/a	Checkpoint mediator	[31]
Cdk1 (Cdc28p)	Promote 5' end processing	Protein kinase	[45, 46]
γH2AX (γH2Ap)	Sister chromatid cohesion; promote 5' end processing	Recruitment/structural	[39, 41, 42]
Mre11 (Mre11p), Rad50 (Rad50p), Nbs1 (Xrs2p)	Promote 5' end processing; sister chromatid cohesion; GCR suppressors	Nuclease; ATPase complex	[16, 29, 39, 117]
CtIP (Sae2p)	Promote 5' end processing	5' exonuclease	[47, 49, 118]
Blm (Sgs1p)	Promote 5' end processing; dissolution of dHJ/D-loop; suppress GCRs	3' helicase	[24, 48–50, 74]
RTEL? hFBH1? (Srs2p)	Displace Rad51 filament	3' helicase	[73, 119, 120]
FANCM (Mph1p)	Disrupt D-loop; branch migration	3' helicase	[82, 83]
Rad54 (Rad54p)	Branch migration; sister recombination	ATPase	[81, 121]
Mus81 (Mus81p)/Eme1 (Mms4p)	Resolution of dHJ/D-loop	Structure-specific endonuclease	[65, 122, 123]
Gen1 (Yen1p)	Resolution of dHJ	Structure-specific endonuclease	[72]
Ku70/80 (Yku70p/Yku80p)	Inhibit end resection/recombination	DNA end binding	[51, 55, 56]

^aNames of genes and their *Saccharomyces cerevisiae* orthologs are shown in parentheses

Table 2

Genes and their biochemical activities that potentially suppress end joining-induced chromosome translocations

Name ^a	Function	Biochemical activity	Reference
ATM (Tel1p)	End tethering; 5' end resection; apoptosis; end motility	Protein kinase	[88, 101, 102, 104, 106]
Mre11 (Mre11p), Rad50 (Rad50p), Nbs1 (Xrs2p)	Promote 5' end processing; end tethering; suppress GCRs	Nuclease; ATPase complex	[16, 29, 39, 88, 94, 95, 117]
CtIP (Sae2p)	Promote 5' end processing; end tethering	5' exonuclease	[47, 49, 53, 94, 118]
γ H2AX (γ H2Ap)	End tethering; promote 5' end processing	Recruitment/structural	[39, 41, 42, 88]
(Dun1p)	Promote 5' end processing; suppress GCRs	Protein kinase	[16, 88]
53BP1 (Rad9p)	Increase end mobility	Checkpoint mediator	[101, 102]
Mdc1	Establish γ H2AX	Mediator	[124, 125]
p19/Arf	Cell cycle control; cell death	Protein kinase inhibitor	[92]
p53	Checkpoint arrest; cell death	Transcription factor	[92]
Fen1 (Rad27p) ^b	Lagging strand synthesis; DNA repair; suppress GCRs	5' exonuclease	[126]
Rfa1 (Rfa1p)	DNA replication; DNA repair; suppress GCRs	Single strand DNA binding	[126, 127]
Rad52 (Rad52p)	Recombination; suppress GCRs	Single strand DNA binding; recombination mediator; 2nd end capture	[126]
TLC1 (TLC1p)	Telomere elongation; suppress GCRs	Telomerase	[16]

^aNames of genes and their *Saccharomyces cerevisiae* orthologs are shown in parentheses

^bChromosome translocations form in a Dnl4p-independent manner