Published in final edited form as:

J Hosp Med. 2012 November ; 7(9): 684–689. doi:10.1002/jhm.1974.

Therapeutic hypothermia for cardiac arrest: Real-world utilization trends and hospital mortality

Anupam B. Jena, M.D., Ph.D.^A, John A. Romley, Ph.D.^B, Christopher Newton-Cheh, M.D., MPH.^C, and Peter Noseworthy, M.D.^D

^ADepartment of Health Care Policy, Harvard Medical School, 180 Longwood Avenue, Boston, MA 02115; and Department of Medicine, Massachusetts General Hospital; and National Bureau of Economic Research, Cambridge, MA; Tel: 617-432-8322; jena.anupam@mgh.harvard.edu.

^BLeonard D. Schaeffer Center for Health Policy and Economics, University of Southern California, Los Angeles, CA 90089-0626. Tel: 213-821-7965; Fax: 213-740-3460; romley@sppd.usc.edu.

^CCardiology Division, Massachusetts General Hospital, Harvard Medical School, Cardiovascular Research Center and Center for Human Genetic Research, Massachusetts General Hospital, 185 Cambridge St, CPZN 5.242, Boston, Massachusetts 02114; Tel: 643-3615, Fax: 617-249-0127; cnewtoncheh@partners.org

^DCardiology Division, Massachusetts General Hospital, Harvard Medical School, Center for Human Genetic Research, Cardiovascular Research Center, 185 Cambridge St, CPZN 5.814, Boston, Massachusetts 02114. Tel: 617-643-6328; Fax: 617-507-7766; pnoseworthy@partners.org

Abstract

Background—Therapeutic hypothermia (TH) improves outcomes following cardiac arrest in small clinical trials.

Objective—To study real-world utilization and outcomes in U.S. hospitals.

Design—Retrospective cohort study

Setting—California hospitals

Patients—Patients eligible for therapeutic hypothermia after cardiac arrest.

Interventions—We analyzed all discharges from California (1999-2008) to identify patients eligible for TH after cardiac arrest. Patients were considered eligible for TH if both cardiac arrest and anoxic brain injury were among the administrative diagnoses ($n = 46,833$). Patients undergoing TH ($n = 204$) were identified through billing codes.

Measurements—TH utilization and in hospital mortality

Results—Use of TH increased over the study period with 87.3% (178/204) of TH occurring between 2006 and 2008. Few hospitals appeared to perform TH over the study period (47/419, 11.2%). Utilization of TH was concentrated in a few centers, with the top 3 of 419 centers accounting for 31.4% (64/204) of cases. Patients undergoing TH were younger, less likely to be male, more likely to be treated at teaching centers, and had similar co-morbidities compared to

Corresponding Author: Anupam B. Jena, M.D., Ph.D. Department of Health Care Policy Harvard Medical School 180 Longwood Avenue Boston, MA 02115 Tel: 617-432-8322 jena.anupam@mgh.harvard.edu.

Statement of disclosure: The contents of this manuscript have not been published in any other peer-reviewed media. This manuscript is not under review elsewhere. All authors listed on this manuscript have contributed sufficiently to this project to be included as authors. The authors have no conflict of interest, financial or otherwise.

eligible individuals who did not undergo TH. The adjusted odds ratio for hospital mortality among patients undergoing TH was 0.80 (95% CI 0.60-1.06, $p=0.11$).

Conclusions—TH utilization appears low but implementation is increasing. Case selection and referral biases limit the analysis of the relationship between center TH volume and in hospital mortality.

Keywords

Cardiac arrest; therapeutic hypothermia; cooling

INTRODUCTION

There are over 350,000 cases of out-of-hospital cardiac arrest (OHCA) each year in the United States,(1, 2) and with supportive therapy alone, only a fraction of victims survive to hospital discharge. Rapid intervention including cardiopulmonary resuscitation in the moments following arrest is critical to minimizing neurologic injury, morbidity, and mortality. In 2002, two small randomized controlled trials showed a survival benefit of therapeutic hypothermia (TH) when provided within 1-2 hours after return of circulation following an OHCA and, to date, TH remains one of the few interventions with proven mortality benefit after initial cardiopulmonary resuscitation.(3, 4) Since 2003, TH has been incorporated into the American Heart Association practice guidelines(5-7) and use of TH has steadily increased, but widespread clinical uptake remains low.(8, 9)

The initial studies that evaluated TH were small, with only 189 patients included in the TH arms of the two trials combined. To date, only a few studies have replicated this initial observation in real world settings, with little analysis of outcomes in U.S. centers in particular.(10-13) Accordingly, we aimed to examine the real-world experience with TH in the U.S. using a large administrative claims database of all California hospital admissions to describe utilization trends, hospital mortality, and volume-outcome relationships associated with the intervention.

MATERIALS AND METHODS

Data

We identified all admissions to California hospitals during 1999-2008 based on discharge records from the California Office of Statewide Health Planning and Development. Our study period included cases of TH performed prior to the 2002 major clinical trials since TH was in occasional use prior to the publication of these trials. The data was de-identified and publicly available and therefore exempt from review by the Institutional Review Board. In addition to hospital name, each discharge record included patient age, gender, admission year, International Classification of Disease Version 9 (ICD-9) code for presenting primary and secondary diagnoses, procedure codes, and disposition (discharge to home or rehabilitation, in-hospital death). All California hospitals were included in the registry ($n = 419$). We defined teaching status for each hospital based on membership in the Council of Teaching Hospitals, as reported in the American Hospital Association's Annual Survey ($n = 19$ teaching hospitals).(14)

Setting and Participants

We used discharge diagnoses to identify patients who could be considered eligible for therapeutic hypothermia after cardiac arrest. We classified patients as eligible for therapeutic hypothermia after cardiac arrest based on ICD-9 diagnosis codes that indicated the presence of both cardiac arrest and anoxic brain injury in the administrative diagnoses. Because of

known imprecision in using billing codes to identify patients with cardiac arrest,(15, 16) we broadly defined cardiac arrest to include those patients with ICD-9 codes for cardiac arrest, ventricular fibrillation (VF), or ventricular tachycardia (VT), (supplementary Table 1). We could not distinguish between out-of-hospital and in-hospital cardiac arrest based on administrative diagnoses. To ensure that we included only patients with cardiac arrest complicated by neurologic insult, we required an ICD-9 diagnosis of either anoxic brain injury, coma, or persistent vegetative state. Claims did not allow us to distinguish among initial cardiac arrest rhythms (VF vs. pulseless VT vs. asystole). Patients younger than 18 years of age, and those who were pregnant, suffered traumatic brain injury, intracranial hemorrhage, metastatic cancer, or dementia were excluded.(3, 17) We did not exclude patients based on coagulopathy (which is considered a contraindication to TH) since ICD-9 coding does not allow us to determine the severity of the coagulopathy or whether it was a result of therapeutic hypothermia itself.

We used the ICD-9 procedure code (99.81) for TH to first identify patients who underwent TH from 1999-2008. Since this code also applies to TH used during cardiac and neurosurgery, we examined each of these cases and excluded individuals who underwent cardiac surgery or neurosurgery during the hospitalization. As in our “eligible for TH” definition, we excluded patients younger than 18 years of age, and those who were pregnant, suffered traumatic brain injury, intracranial hemorrhage, metastatic cancer, or dementia. Patients who underwent therapeutic hypothermia but for whom a specific procedure code was not recorded in the discharge abstract – perhaps because the medical institution did not directly bill for the procedure – could not be identified.

Statistical analysis

We used a multivariable logistic model to estimate differences in hospital mortality after cardiac arrest associated with use of therapeutic hypothermia. We conducted two specifications. In our baseline specification, we accounted for case-mix differences between those who underwent TH and those who did not by adjusting for age, gender, year of admission, the number of Charlson-Deyo co-morbidities,(18-20) and hospital teaching status. Because mortality after cardiac arrest in centers that perform therapeutic hypothermia may be different from centers that do not, even for patients who do not undergo the procedure, we included indicators for volume tercile of therapeutic hypothermia cases performed. Volume of therapeutic hypothermia was defined at the hospital level as the total number of cases performed by that hospital from 1999-2008.

In order to explore how hospital teaching status, volume of therapeutic hypothermia procedures (broken into terciles), and year of admission affected the association between hospital mortality after cardiac arrest and therapeutic hypothermia, our baseline logistic model was expanded to include interactions between therapeutic hypothermia and each of these variables. The interaction between therapeutic hypothermia and year explored whether the effectiveness of the procedure changed over time as case-selection, method of therapeutic hypothermia (cold saline vs. commercially available devices) and experience changed in California hospitals. For both specifications, we reported the odds ratio of hospital mortality among patients undergoing therapeutic hypothermia as well as risk-adjusted mortality for both TH and non-TH groups. STATA version 11 (STATA Corp, College Station, Texas) was used for statistical analyses and a two-sided $p = 0.05$ was used.

RESULTS

Descriptive data

Table 1 reports summary statistics for patients with cardiac arrest complicated by neurologic insult (anoxic brain injury, coma, or persistent vegetative state) between 1999 and 2008. Across all years, 204 patients were identified as undergoing TH. In comparison, 105 patients were identified as undergoing TH in 2008 alone. Patients who underwent TH were less likely to be male (30.7% vs. 44.6% male, $p < 0.01$), were younger (63.9 ± 15.0 years vs. 67.3 ± 15.7 years, $p = 0.03$), and had equivalent numbers of Charlson-Deyo co-morbidities (2.5 ± 2.0 diagnoses vs. 2.5 ± 2.0 diagnoses, $p = 0.89$). Therapeutic hypothermia was more commonly employed at teaching hospitals (51/3,709 [1.4%] vs. 153/42,942 [0.4%], $p < 0.01$). There was a trend toward decreased unadjusted mortality among patients who underwent therapeutic hypothermia compared with those who did not (56.9% vs. 62.8%, $p = 0.08$).

Figures 1 and 2 provide additional aggregate statistics on therapeutic hypothermia in California hospitals. Figure 1 plots the number of therapeutic hypothermia cases recorded in the administrative discharge registry between 1999 and 2008. Of the 204 total cases identified during this period, 178 (87.3%) were performed between 2006 and 2008. Figure 2 shows the distribution of TH cases across centers that performed therapeutic hypothermia ($n = 47$ hospitals, 11.3 % of all hospitals). Ten centers accounted for 124/204 (60.7%) of the total patients treated with TH after cardiac arrest; the top three centers accounted for 64 (31.4%) of treated patients. Twenty-seven hospitals were identified as performing therapeutic hypothermia on only one or two patients between 1999 and 2008.

Risk-adjusted mortality

Table 2 presents the odds ratio of factors predicting in-hospital mortality after cardiac arrest complicated by neurologic insult. Factors include use of TH after cardiac arrest, age, gender, year of admission, number of Charlson-Deyo co-morbidities, hospital teaching status, and volume tercile of hospitals that performed therapeutic hypothermia. Overall, patients who were older, male, and had greater co-morbidities were statistically more likely to die after cardiac arrest complicated by neurologic insult. Regardless of whether they underwent TH, patients admitted to hospitals in the highest volume tercile of TH use were more likely to die after cardiac arrest. Adjusting for volume tercile, teaching hospital status was not independently associated with mortality after cardiac arrest. The adjusted odds ratio of mortality among patients undergoing therapeutic hypothermia was 0.80 (95% CI 0.60-1.06, $p = 0.11$). The adjusted probability of inpatient mortality among patients undergoing therapeutic hypothermia was 57.5% (95% CI 50.7 – 64.3%) compared to those who did not 62.8% (95% CI 61.7 – 63.9%), $p = 0.11$.

Figure 3 presents adjusted mortality after cardiac arrest in hospitals that did not perform TH as well as adjusted mortality associated with TH for each volume tercile of hospitals that performed the procedure. Hospital mortality rates among patients not receiving TH after cardiac arrest were slightly higher in hospitals in the high volume tercile of TH (66.3%, 95% CI 63.8 – 68.8%) compared to hospitals in low and moderate volume terciles and to hospitals not performing TH ($p < 0.001$). Hospital mortality rates among low and moderate TH volume centers and in centers not performing TH were similar (62.3%, 61.3%, and 63.4% respectively). Among both the low volume and moderate volume terciles, however, patients who underwent TH after cardiac arrest were significantly less likely to die in hospital compared to those who did not. For patients admitted to hospitals in the low volume tercile, those undergoing therapeutic hypothermia had an adjusted hospital mortality rate of 25.5% (95% CI 3.0 - 47.9%) compared to those who did not undergo TH (adjusted mortality

61.3%, 95% CI 57.4 - 65.1%), $p < 0.001$. In the moderate volume tercile, patients receiving therapeutic hypothermia had an adjusted hospital mortality rate of 31.0% (95% CI 9.2% - 52.8%) compared to 63.4% (95% CI 57.7 - 69.1%), $p < 0.001$, among those not undergoing the procedure. There was no statistically significant difference in adjusted mortality between those who underwent TH and those who did not in hospitals in the highest volume tercile ($p = 0.211$). In addition to examining how volume of therapeutic hypothermia performed by hospitals affected the association between TH and hospital mortality, we also examined whether year of admission and teaching hospital independently modified the association. Neither year of admission nor teaching hospital statistically significantly affected the association between therapeutic hypothermia and hospital mortality after cardiac arrest at the $p < 0.10$ level.

DISCUSSION

In an administrative database of all admissions to California hospitals, we demonstrated that use of TH increased steadily since the publication of the initial clinical trials in 2002. The absolute level of TH utilization in our study undoubtedly represents a significant underestimation of actual TH utilization; however, our study does provide an assessment of the utilization trends over time. The bulk of TH use appears to be performed in a small group of high volume centers and 89% of California hospitals did not perform TH during the study period (as assessed by procedure billing codes). Additionally, within the limitations of a retrospective, administrative claims-based study design, TH appears to be associated with a similar in-hospital mortality rate to that seen in clinical trials.(3, 4) In exploratory analyses, there appears to be a particular benefit of TH in low and moderate volume centers, though these findings should be considered hypothesis-generating.

Despite the body of evidence in favor of TH, utilization in our study and others appears quite low. In a 2005 survey of physicians, 87% of respondents had never used therapeutic hypothermia, citing inadequate data, technical limitations, and lack of incorporation in the ACLS protocol as principal justifications.(8) Other surveys have shown similar results and noted that critical care physicians and those working in large medical centers were more likely to adopt the therapy.(9) Advocates of the therapy have suggested that an explicit hospital-based plan developed by key stakeholders can help facilitate implementation.(24) Accordingly, there is growing interest in developing centers of expertise in highly intensive therapies such as TH. For instance, the New York City Emergency Medical Service has begun to explore a protocol to divert TH candidates to specialized centers.(25, 26) Some favorable results have been reported in individual hospitals and local hospital systems. (27-31)

Our data suggest that TH is associated with an in-hospital mortality rate that is comparable to that reported in the clinical trials. For example, in a 2009 meta-analysis of four clinical trials and one abstract (481 patients in total), TH was associated with a 35% relative mortality benefit as compared to standard post-resuscitation care.(21) It has been estimated that broad TH implementation could save thousands of lives(22) and many authors have advocated for its use and outlined explicit protocols for implementation.(17) Furthermore, TH appears to be cost effective in line with other accepted therapies. Assuming the Hypothermia After Cardiac Arrest (HACA) trial inclusion criteria, even at extreme estimates for costs, the cost-effectiveness of hypothermia remains less than \$100,000 per quality-adjusted life year.(23)

There are important limitations of this study. Our use of administrative claims data certainly underestimates the level of TH utilization since we could only identify cases in which TH was included in the billing codes for the hospitalization. Hospitals may vary in utilization of

this particular billing code for TH in ways that bias our estimated associations. The ICD-9 99.81 code for therapeutic hypothermia was also not developed for post-cardiac arrest TH specifically, so use of the code may actually lag clinical utilization. Although the observed trend in TH utilization is likely mainly due to a true increase in utilization, it is possible that some of the increase observed is due to an increase in utilization of TH procedure billing codes. Our TH utilization estimates should be construed as a lower bound of the actual rates. Additionally, although the estimated real-world mortality benefit of TH may be comparable to that of clinical trials, the equivalence of patients in our sample to those in published randomized trials is uncertain. Similarly, even after adjusting for age, gender, year of admission, co-morbidities, hospital teaching status, and TH volume there are likely many unmeasured variables that influence mortality in both the TH and comparison groups. There are also likely patients included in our comparison group who had both cardiac arrest or ventricular tachycardia and anoxic brain injury, but who were not candidates for TH as the episode of cardiac arrest followed rather than preceded the anoxic brain injury. Since we lack detailed clinical data about the TH cases (initial rhythm, time before return of circulation, pre-existing disease states, etc.), we are unable to match controls directly to cases. Additionally, we lack data to assess neurologic recovery or quality of life after arrest.

The observation that a mortality benefit in our study could be detected only in low and moderate volume centers requires further exploration. Indeed, one might expect that high volume centers may have better outcomes with TH as a result of more robust infrastructure, technical experience, and available resources. Our finding that mortality benefits of TH appear concentrated in centers with low to moderate volume of TH utilization suggest at least one of two possibilities. First, low and moderate volume centers may perform TH in a subset of patients who benefit most from the intervention or alternatively in the most viable cardiac arrest cases (those who may fare well with or without the therapy). Consequently, we may observe relatively favorable outcomes in this group due to this selection bias. Second, high volume centers – despite having more expertise – may also attract patients at higher mortality risk due to referral bias. This would lead us to estimate lower mortality benefits associated with TH in these high volume centers. Indeed, greater observed mortality at high volume centers regardless of TH status suggests that overall acuity is higher at high volume centers. While our inferences are greatly affected by issues of case selection and referral bias, it is also important to consider the possibility that the estimated mortality benefit of TH in higher volume centers is lower because of the selection of patients who do not meet current guidelines for treatment with TH. Distinguishing whether the selection of patients undergoing TH at high volume centers is appropriate or inappropriate based on current guidelines is an important issue that merits further research with datasets with more refined patient clinical information.

In summary, therapeutic hypothermia utilization is low but the rate of implementation has increased since the publication of the initial clinical trials in 2002. The bulk of TH utilization appears limited to a subset of high volume centers and most centers in California appear to have not used the therapy. Real-world in-hospital mortality associated with TH is comparable to that reported in randomized clinical trials.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

None

Grant funding: Dr. Romley received support from NIH grant R03AG031990-A1. Dr. Noseworthy received support from the Max Schaldach Fellowship in Cardiac Pacing and Electrophysiology granted by the Heart Rhythm Society. The design, conduct, analysis, interpretation, and presentation of the data are the responsibility of the investigators, with no involvement from the funding sources.

REFERENCES

1. Zheng ZJ, Croft JB, Giles WH, et al. Sudden cardiac death in the United States, 1989 to 1998. *Circulation*. 2001; 104(18):2158–2163. [PubMed: 11684624]
2. Mathers CD, Loncar D. Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med*. 2006; 3(11):e442. [PubMed: 17132052]
3. Bernard SA, Gray TW, Buist MD, et al. Treatment of comatose survivors of out-of-hospital cardiac arrest with induced hypothermia. *N Engl J Med*. 2002; 346(8):557–563. [PubMed: 11856794]
4. Mild therapeutic hypothermia to improve the neurologic outcome after cardiac arrest. *N Engl J Med*. 2002; 346(8):549–556. [PubMed: 11856793]
5. Nolan JP, Morley PT, Vanden Hoek TL, et al. Therapeutic hypothermia after cardiac arrest: an advisory statement by the advanced life support task force of the International Liaison Committee on Resuscitation. *Circulation*. 2003; 108(1):118–121. [PubMed: 12847056]
6. 2005 American Heart Association Guidelines for Cardiopulmonary Resuscitation and Emergency Cardiovascular Care: Part 7.5: Postresuscitation Support *Circulation*. 2005; 112(Suppl I):IV–84–IV–88.
7. Deakin CD, Morrison LJ, Morley PT, et al. Part 8: Advanced life support: 2010 International Consensus on Cardiopulmonary Resuscitation and Emergency Cardiovascular Care Science with Treatment Recommendations. *Resuscitation*. 2010; 81(Suppl 1):e93–e174. [PubMed: 20956032]
8. Abella BS, Rhee JW, Huang KN, et al. Induced hypothermia is underused after resuscitation from cardiac arrest: a current practice survey. *Resuscitation*. 2005; 64(2):181–186. [PubMed: 15680527]
9. Merchant RM, Soar J, Skrifvars MB, et al. Therapeutic hypothermia utilization among physicians after resuscitation from cardiac arrest. *Crit Care Med*. 2006; 34(7):1935–1940. [PubMed: 16691134]
10. Nielsen N, Sunde K, Hovdenes J, et al. Adverse events and their relation to mortality in out-of-hospital cardiac arrest patients treated with therapeutic hypothermia. *Crit Care Med*. 39(1):57–64. [PubMed: 20959789]
11. Nielsen N, Hovdenes J, Nilsson F, et al. Outcome, timing and adverse events in therapeutic hypothermia after out-of-hospital cardiac arrest. *Acta Anaesthesiol Scand*. 2009; 53(7):926–934. [PubMed: 19549271]
12. Arrich J. Clinical application of mild therapeutic hypothermia after cardiac arrest. *Crit Care Med*. 2007; 35(4):1041–1047. [PubMed: 17334257]
13. Mooney MR, Unger BT, Boland LL, et al. Therapeutic Hypothermia After Out-of-Hospital Cardiac Arrest: Evaluation of a Regional System to Increase Access to Cooling. *Circulation*. 124(2):206–214. [PubMed: 21747066]
14. American Hospital Association. Annual Survey. Health Forum, LLC; Chicago, IL: 2001.
15. Austin PC, Daly PA, Tu JV. A multicenter study of the coding accuracy of hospital discharge administrative data for patients admitted to cardiac care units in Ontario. *Am Heart J*. 2002; 144(2):290–296. [PubMed: 12177647]
16. Goldstein LB. Accuracy of ICD-9-CM coding for the identification of patients with acute ischemic stroke: effect of modifier codes. *Stroke*. 1998; 29(8):1602–1604. [PubMed: 9707200]
17. Holzer M. Targeted temperature management for comatose survivors of cardiac arrest. *N Engl J Med*. 2010; 363(13):1256–1264. [PubMed: 20860507]
18. Charlson ME, Pompei P, Ales KL, et al. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis*. 1987; 40(5):373–383. [PubMed: 3558716]
19. Deyo RA, Cherkin DC, Ciol MA. Adapting a clinical comorbidity index for use with ICD-9-CM administrative databases. *J Clin Epidemiol*. 1992; 45(6):613–619. [PubMed: 1607900]

20. Romano PS, Roos LL, Jollis JG. Adapting a clinical comorbidity index for use with ICD-9-CM administrative data: differing perspectives. *J Clin Epidemiol.* 1993; 46(10):1075–1079. discussion 1081-1090. [PubMed: 8410092]
21. Arrich J, Holzer M, Herkner H, et al. Hypothermia for neuroprotection in adults after cardiopulmonary resuscitation. *Cochrane Database Syst Rev.* 2009; (4):CD004128. [PubMed: 19821320]
22. Holzer M, Behringer W. Therapeutic hypothermia after cardiac arrest and myocardial infarction. *Best Pract Res Clin Anaesthesiol.* 2008; 22(4):711–728. [PubMed: 19137812]
23. Merchant RM, Becker LB, Abella BS, et al. Cost-effectiveness of therapeutic hypothermia after cardiac arrest. *Circ Cardiovasc Qual Outcomes.* 2009; 2(5):421–428. [PubMed: 20031872]
24. Gaieski DF, Fuchs B, Carr BG, et al. Practical implementation of therapeutic hypothermia after cardiac arrest. *Hosp Pract (Minneap).* 2009; 37(1):71–83. [PubMed: 20877174]
25. Zraick K. Paramedics Will Employ New Therapy in Cardiac Arrest Cases. *New York Times.* Aug 3.2010 :A18.
26. Mechem CC, Goodloe JM, Richmond NJ, et al. Resuscitation center designation: recommendations for emergency medical services practices. *Prehosp Emerg Care.* 2010; 14(1):51–61. [PubMed: 19947868]
27. Szumita PM, Baroletti S, Avery KR, et al. Implementation of a Hospital-wide Protocol for Induced Hypothermia Following Successfully Resuscitated Cardiac Arrest. *Crit Pathw Cardiol.* 2010; 9(4): 216–220. [PubMed: 21119341]
28. Kulstad CE, Holt SC, Abrahamsen AA, et al. Therapeutic hypothermia protocol in a community emergency department. *West J Emerg Med.* 2010; 11(4):367–372. [PubMed: 21079711]
29. Walters EL, Morawski K, Dorotta I, et al. Implementation of a Post-Cardiac Arrest Care Bundle Including Therapeutic Hypothermia and Hemodynamic Optimization in Comatose Patients with Return of Spontaneous Circulation After Out-of-Hospital Cardiac Arrest: A Feasibility Study. *Shock.* 2011; 35(4):360–366. [PubMed: 21068697]
30. Hinchey PR, Myers JB, Lewis R, et al. Improved out-of-hospital cardiac arrest survival after the sequential implementation of 2005 AHA guidelines for compressions, ventilations, and induced hypothermia: the Wake County experience. *Ann Emerg Med.* 2010; 56(4):348–357. [PubMed: 20359771]
31. Prior J, Lawhon-Triano M, Fedor D, et al. Community-based application of mild therapeutic hypothermia for survivors of cardiac arrest. *South Med J.* 2010; 103(4):295–300. [PubMed: 20224484]

Figure 1.
Annual cases of therapeutic hypothermia performed after cardiac arrest complicated by neurologic insult, administrative claims data from California hospitals 1999-2008.

Figure 2.

Total cases of therapeutic hypothermia performed after cardiac arrest complicated by neurologic insult, by hospitals performing procedure, administrative claims data from California hospitals 1999-2008.

Figure 3.

Adjusted hospital mortality after cardiac arrest complicated by neurologic insult, stratified by therapeutic hypothermia and volume of hospital.

TABLE 1

Characteristics of patients suffering cardiac arrest complicated by neurologic insult, stratified by therapeutic hypothermia use, California hospitals 1999-2008

	Therapeutic hypothermia	No therapeutic hypothermia	p-value
No. observations	204	46629	
No. cases in teaching hospitals	51	3658	
No. cases in non-teaching hospitals	153	42789	
Age, y	63.9 ± 15.0	67.3 ± 15.7	0.06
Male	30.7	44.6	< 0.01
Hospital mortality, %	56.9	62.8	0.08
<i>Co-morbidities</i>			
No. Charlson-Deyo co-morbidities	2.5 ± 2.0	2.5 ± 2.0	0.89
Coronary artery disease, %	48.0	38.0	< 0.01
Acute myocardial infarction, %	42.6	28.9	< 0.01
Congestive heart failure, %	27.9	35.3	0.03
Hypertension, %	36.3	33.2	0.83
Acute renal failure, %	33.3	26.6	0.03
Diabetes Mellitus, %	30.9	23.0	< 0.01
Chronic obstructive pulmonary disease, %	10.3	19.3	< 0.01

P-values for age, hospital mortality, and no. of Charlson-Deyo co-morbidities reflect 2-sided t-test of continuous variables; p-values for all other variables reflect χ^2 test of categorical data.

TABLE 2

Adjusted odds ratio of hospital mortality after cardiac arrest complicated by neurologic insult in a multivariable regression model, California hospitals 1999-2008

Variable	Odds ratio of hospital mortality (95% CI)	p
No. observations	46651	
Age [†]		
65-69	1.19 (1.12-1.28)	< 0.001
70-74	1.29 (1.20-1.39)	< 0.001
75-79	1.55 (1.44-1.67)	< 0.001
80-84	1.79 (1.65-1.93)	< 0.001
85 and over	2.06 (1.89-2.25)	< 0.001
Male	1.15 (1.10-1.21)	< 0.001
Teaching hospital	1.13 (0.95-1.34)	0.17
No. Charlson-Deyo co-morbidities	1.09 (1.08-1.10)	< 0.001
Year trend	0.98 (0.97-0.99)	< 0.001
Volume tercile among hospitals performing TH ^{††}		
First tercile	0.94 (0.79-1.12)	0.48
Second tercile	1.03 (0.80-1.33)	0.82
Third tercile	1.20 (1.05-1.36)	0.006
Therapeutic hypothermia	0.80 (0.60-1.06)	0.11

[†] Odds ratios compared to individuals aged 60-65.

^{††} Odds ratios compared to hospitals not performing TH