Published in final edited form as:

Angew Chem Int Ed Engl. 2012 January 2; 51(1): 204–208. doi:10.1002/anie.201105801.

Palladium-Catalyzed Allylic Alkylation of Carboxylic Acid Derivatives: *N*-Acyloxazolinones as Ester Enolate Equivalents

Dr. Barry M. Trost, Dr. David J. Michaelis, Julie Charpentier, and Dr. Jiayi Xu

Department of Chemistry, Stanford University, Stanford, CA 94305-5080, Fax: (1+)650-725-0002

Barry M. Trost: bmtrost@stanford.edu

Abstract

A general asymmetric allylic alkylation of ester enolate equivalents at the carboxylic acid oxidation state is reported. *N*-Acylbenzoxazolinone-derived enol carbonates are synthesized in good yield and employed in the palladium-catalyzed alkylation reaction. Good yields (up to 99%) and enantioselectivities (up to 99% ee) are obtained and the imide products are readily converted to a series of carboxylic acid derivatives without loss of enantiopurity. High enantioselectivity in the reaction is achieved by a new strategy for ligand design involving variation of the steric properties of the diarylphosphine moiety.

Keywords

allylic alkylation; asymmetric catalysis; benzoxazolinone; palladium; ester enolate

The asymmetric alkylation of carbonyl compounds is one of the most important methods for generating stereocenters in organic synthesis.¹ The alkylation of ester enolates is particularly useful as the resulting products can be converted to a variety of carboxylic acid derivatives or reduced to alcohols without loss of enantioenrichment at the α -center.² The most common procedures for asymmetric ester enolate alkylations, however, involve the use of stoichiometric chiral auxiliaries.^{3,4} Attempts to render this asymmetric transformation catalytic have achieved only limited success. For example, successful asymmetric alkylations of specialized carboxylic acid derivatives including oxindoles,⁵ azalactones,⁶ and zinc enolates of glycine esters⁷ have been reported. A recent example of an asymmetric Pd-catalyzed Claisen rearrangement of allyl phenylacetate demonstrated an alternative strategy for obtaining enantioenriched ester derivatives.⁸ While these reports demonstrate the continued importance of developing asymmetric alkylations of ester derivatives, a highly general and enantioselective alkylation of simple ester derivatives is still elusive.⁹

As a solution to this unmet need, we and others have recently reported the use of ester enolate surrogates in the asymmetric allylic alkylation (AAA) reaction, including 2-acylimidazoles,¹⁰ *N,N*-dialkyl amides,¹¹ and acylsilanes.¹² However, there are several drawbacks to each of these methods. First, the acylimidazoles and acylsilanes require multiple steps to synthesize from carboxylic acids. Second, subsequent transformations that take advantage of the reactivity of the carboxylic acid functionality are not straightforward. Thus, a general enantioselective method for enolate alkylations of simple ester derivatives that can function directly in subsequent transformations is still elusive.¹³ We report herein

Correspondence to: Barry M. Trost, bmtrost@stanford.edu.

Homepage : <http://www.stanford.edu/group/bmtrost/>

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.

the palladium-catalyzed asymmetric alkylation of *N*-acylbenzoxazolinone-derived enol carbonates, which represents a general asymmetric alkylation of ester enolate equivalents at the carboxylic acid oxidation state.¹⁴ Importantly, the resulting enantioenriched imide products are easily converted to the acid, ester, thioester, amide, or alcohol derivatives under mild conditions without prior activation. In addition, the modularity of our diamino bis(phosphine) ligands allowed for a new strategy for catalyst design wherein the steric properties of the diaryl phosphines was varied to enable high enantioselectivity.

From the outset, our goal was to develop an enantioselective ester enolate alkylation where the products could be easily derivatized under mild conditions to a variety of carboxylic acid derivatives. Thus, our initial studies focused on employing ester surrogates at the *carboxylic acid oxidation state* that are known to hydrolyze under mild conditions (Table 1). One significant advantage of using these activated esters over previously reported systems is the ability to quickly access substrates in one step from any carboxylic acid by amide bond formation. A major challenge in the alkylation of ester derivatives is the propensity of the intermediate enolate to undergo elimination to form a ketene. We believed that the recently developed decarboxylative allylic alkylation methodology^{15, 16} provided a unique way to avoid this problem because the enolate could be trapped as an enol carbonate at low temperature where competing ketene formation would be minimized. The enol carbonate might then be purified and employed in the AAA reaction. In our initial studies, we found that a variety of enol carbonates derived from ester equivalents could indeed be isolated, including *N*-acyl imidazoles, indoles, and oxazolinones (Table 1). In the ensuing decarboxylative asymmetric alkylation reaction using our anthracenediamine-derived bisphosphine ligand **4** (*vide infra*), the best yield and enantioselectivity was observed with *N*-acylbenzoxazolinone-derived substrate **1d**. Further optimization studies, therefore, focused on the *N*-acyloxazolinone derived enol carbonates as substrates for the asymmetric alkylation reaction.

The major challenge with employing the benzoxazolinone-derived enol carbonates was the low yield of the enol forming reaction. Careful analysis of the crude reaction mixture showed that the remainder of the yield was consumed by side reactions resulting from ketene elimination, with benzoxazolinone **3** as the major side product (eq. 1). As a solution to this problem, we found that formation of the enolate in the presence of the chloroformate provided the desired product **1d** in much improved and synthetically

(1)

useful yields (65%). Our optimization studies also demonstrated that the chloroformate acylating agent was necessary for efficient generation of the enol carbonate. This new procedure for enol carbonate formation proved exceptionally general. For instance, a wide range of chloroformate electrophiles could be employed in the process, including those substituted at the terminal (eq. 2, 4) and internal positions of the olefin (eq. 3, 5). The tolerance for substitution on the *N*-acylbenzoxazolinone partner also proved to be very general. Enol carbonates bearing longer alkyl chains (eq. 3), β,β -disubstituted carbons (eq. 4) and a variety of functional groups including primary alkyl chlorides (eq. 5) could all be

isolated in good yield. These enol carbonates are relatively stable and can be stored at low temperature for several months without decomposition.

(2)

(3)

(4)

(5)

Having established that imide-derived enol carbonates could be synthesized in good yield, we then sought to improve the enantioselectivity of the asymmetric alkylation process (Table 2). By varying both the ligand (entries 1–4) and solvent (entries 5–8), we found that the product could be obtained with 75% ee using the stilbene diamine-derived bisphosphine ligand **6a** in THF as solvent. We next varied the structure of the phosphine ligand, believing that increasing the steric bulk around the phosphines could increase the selectivity of the reaction. We were pleased to find that this was indeed the case; *o*-tolyl-substituted bisphosphine **6b** gave the product in higher enantioselectivity (entry 9). This new approach to ligand modification further demonstrates the modularity of our class of enantiopure bisphosphine ligands and the ability to selectively tune both the steric and electronic properties of the ligand to improve selectivity. Other substitution on the phosphine, however, did not lead to an increase in selectivity (entries 10–11). In addition, varying the

structure of the diamine backbone to the 1,2-dicyclohexylethane diamine (**8**) led to decreased yield and enantioselectivity (entry 12).

With optimal conditions in hand, we next explored the substrate scope of the asymmetric alkylation reaction (Table 3). In general, the substituted allyl carbonates proved to be excellent substrates. Specifically, substitution at the internal position of the olefin (entries 2–3), the allylic position (entry 4) or the terminus of the olefin (entry 5) is tolerated and in most cases serves to increase the enantioselectivity of the reaction over the simple allyl electrophile. Where regioselectivity in the alkylation is possible, excellent regiomic ratios are observed in favor of the linear product (entries 5, 9, 13, >10:1 linear:branched selectivity). In addition, the reaction proceeds with high diastereoselectivity when prochiral electrophiles are employed (entry 4, >95:5 d.r.).¹⁷ The AAA reaction also proceeds efficiently with a variety of substituents at R₁. Mono- and disubstitution at the β -position of the enol is tolerated (entries 6, 12). However, β,β -disubstituted enols (entries 12–13) and longer alkyl chains (entry 15) gave a small decrease in the enantioselectivity of product formation. A variety of functional groups are also tolerated in the reaction, including primary alkyl chlorides (entry 14), esters (entry 15), and silyl ethers (entries 16–17). This process is also readily conducted on large scale: when table 3, entry 2 was conducted on 4.0 mmol scale, **2e** was obtained in 66% yield and 97% ee.

A major goal of this research was to develop an efficient asymmetric alkylation of ester enolate equivalents that could be derivatized to any number of carboxylic acid derivatives under mild conditions. We were delighted, therefore, to find that the alkylated products can be functionalized under very mild conditions without significant loss of enantiopurity (Scheme 2).² The benzoxazolinone moiety was cleaved efficiently to generate the corresponding acid (**9**), ester (**10**), and thioester (**11**) by simply treating the imide with the respective lithium anion of the nucleophile. In addition, reduction to the alcohol (**12**) proceeded in good yield. Importantly, little or no racemization of the α -stereocenter of the products was observed for these transformations. The absolute stereochemistry of the alkylation was confirmed by comparison of acid **9** to previously reported materials.¹⁸

Having established that the benzoxazolinone auxiliary was an excellent leaving group for substitution reactions, we next wondered what additional nucleophiles could react with the enantioenriched imide products. We found that simple treatment of product **2s** with a primary amine led to formation of amide product **13** in just 4 hours in high yield (97%) and with complete retention of enantiopurity (eq. 2). This transformation demonstrates the mild reaction conditions under which substitution of the benzoxazolinone can occur, even when compared with Evan's chiral oxazolidinone auxiliary, which generally requires stoichiometric trialkylaluminum additives to effect the analogous transformation.¹⁹ In addition, this direct conversion to amide products is a marked improvement over our previous system¹⁰ where the 2-acylimidazole products had to be converted to the carboxylic acid before subsequent amide bond formation, or racemization would occur.

(6)

The absolute sense of stereochemistry in this decarboxylative alkylation reaction can be rationalized using the wall and flap cartoon model,²⁰ which was developed in our laboratory to predict the selective formation of one enantiomer when the diphenylphosphinobenzoic acid-based chiral ligand scaffolds are employed. Using the (*R,R*)-enantiomer of our stilbene diamine-derived ligand, our working model correctly indicates that selective formation of the (*S*)-enantiomer of the allylation product should occur based on unfavourable steric interactions in the transition state for formation of the corresponding (*R*)-enantiomer (Scheme 2). The selective formation of the (*S*)-enantiomer in this reaction is also in accordance with the structure-based rationale presented by Lloyd-Jones and coworkers.²¹

In summary, we report a general asymmetric allylic alkylation of ester enolate equivalents at the *carboxylic acid oxidation state*. Specifically, a variety of *N*-acylbenzoxazolidinone-derived enol carbonates can be generated in good yields and are excellent substrates for the palladium-catalyzed asymmetric alkylation reaction. For the first time, variation of the steric properties of the phosphine moiety on this class of bis(phosphine) ligands was shown to provide increased enantioselectivity in an allylic alkylation reaction. An important improvement over previous methods is that the activated ester derivatives are easily accessed from readily available carboxylic acids via simple amide bond formation. In addition, the enantioenriched imide products are readily transformed to the corresponding acid, ester, thioester, alcohol, and amide derivatives without need for prior activation or oxidation of the substrate, providing easy access to a variety of highly useful enantioenriched building blocks for organic synthesis.

Experimental Section

General procedure for synthesis of enol carbonates

Into a 25 ml round bottom flask is placed 3-propionylbenzo[d]oxazol-2(3H)-one (0.466 g, 2.43 mmol) and allylchloroformate (0.308g, 2.56 mmol, 1.05 equiv) in 7 ml THF and the flask was cooled to $-78\text{ }^\circ\text{C}$ in a dry ice/acetone bath. To this solution was slowly added a $-78\text{ }^\circ\text{C}$ solution of sodium bis(trimethylsilyl)amide (0.492 g, 2.68 mmol, 1.1 equiv) in 7 ml THF via cannula. The reaction was stirred 1 hr then quenched with PH 7 phosphate buffer. The mixture was poured into a separatory funnel and the aqueous layer extracted with 3×30 ml Et_2O . The combined organics were dried over Na_2SO_4 and filtered through a short plug of silica gel. The solvent was then removed and the product purified on a column of silica gel, eluting with 5:1 hexanes: Et_2O to give 0.434 g (65% yield) of the desired product **1d** as a colorless oil.

General procedure for asymmetric alkylation reaction

Into an oven dried 2 dram vial was placed $\text{Pd}_2(\text{dba})_3 \cdot \text{CHCl}_3$ (5.3 mg, .005 mmol, 0.025 equiv) and Trost Stilbene ligand (*R,R*)-**6b** (9.5 mg, .012 mmol, .06 equiv), and the vial was flushed with argon from a balloon for 5 minutes. Freshly distilled and degassed THF (1 ml) was then added and the solution was sonicated at room temperature for ~20 min until the reaction turned from heterogeneous deep red to homogeneous deep red-orange in color. The

catalyst solution was then transferred via syringe to a degassed 2 dram vial containing **1d** (0.055 g, 0.2 mmol) in 1 ml degassed THF. The reaction was stirred 10 minutes, at which time TLC showed complete consumption of starting material. The solvent was then removed and the product was purified on silica gel with 8:1 the 6:1 hexanes:Et₂O as eluent to give 0.044 g (96% yield) of the desired product with 85% ee.

Acknowledgments

We thank the National Science Foundation and the National Institutes of Health, General Medical Sciences (Grant GM33049), for their generous support of our programs. D.J.M. also thanks the National Institutes of Health for a postdoctoral fellowship (F32GM093467-01).

References

1. Soai, K.; Shibata, T. *Comprehensive Asymmetric Catalysis II*. Jacobsen, EN.; Pfaltz, A.; Yamamoto, H., editors. Springer; New York: 2000. p. 911-922.
2. a) Evans DA, Ennis MD, Mathre DJ. *J Am Chem Soc.* 1982; 104:1737–1739. (b) Evans DA, Britton TC, Ellman JA. *Tetrahedron Lett.* 1987; 28:6141–6144. (b) Damon RE, Coppola GM. *Tetrahedron Lett.* 1990; 31:2849–2852.
3. For reviews see: Seyden-Penne J. *Chiral Auxiliaries and Ligands in Asymmetric Synthesis*. John Wiley & Sons, Inc New York 1995. Evans DA, Helmchen G, Rüping M, Christmann M, Bräse S. *Asymmetric Synthesis—The Essentials*. Wiley-VCH Weinheim 2006. Prabhat A, Qin H. *Tetrahedron.* 2000; 56:917–947. Vicario JL, Badia D, Carrillo L, Reyes E, Etxebarria J. *Curr Org Chem.* 2005; 9:219–235.
4. A recent report of an asymmetric alkylation of arylacetic acids with stoichiometric chiral amine bases also demonstrates the continued interest in this transformation. See: Stivala CE, Zakarian A. *J Am Chem Soc.* 2011; 133:11936–11939. [PubMed: 21744818]
5. a) Trost BM, Frederiksen MU. *Angew Chem.* 2005; 117:312–314. *Angew Chem Int Ed.* 2005; 44:308–310. b) Trost BM, Zhang Y. *J Am Chem Soc.* 2006; 128:4590–4591. [PubMed: 16594693]
6. a) Trost BM, Ariza X. *Angew Chem.* 1997; 109:2749–2751. *Angew Chem Int Ed.* 1997; 36:2635–2637. b) Trost BM, Dogra K. *J Am Chem Soc.* 2002; 124:7256–7257. [PubMed: 12071719]
7. Braun M, Meletis P, Schrader W. *Eur J Org Chem.* 2010:5369–5372. For a review on asymmetric Claisen rearrangements, see: Mikami K, Akiyama K, Hiersemann M, Nubbemeyer U. *The Claisen Rearrangement. Methods and Applications*. Wiley-VCH Weinheim 2007; chapter 2:25–43.
8. Weiß TD, Helmchen G, Kazmaier U. *Chem Commun.* 2002:1270–1271.
9. For reviews on the alkylation of glycine derivatives by chiral phase-transfer catalysis, see: Maruoka K, Ooi T. *Chem Rev.* 2003; 103:3013–3028. [PubMed: 12914490] Jew S, Park H. *Chem Commun.* 2009; 46:7090–7103.
10. Trost BM, Lehr K, Michaelis DJ, Xu J, Buckl AK. *J Am Chem Soc.* 2010; 132:8915–8917. [PubMed: 20550121]
11. Zhang K, Peng Q, Hou X-L, Wu Y-D. *Angew Chem.* 2008; 120:1765–1768. *Angew Chem Int Ed.* 2008; 47:1741–1744.
12. Chen J-P, Ding C-H, Liu W, Hou X-L, Dai L-X. *J Am Chem Soc.* 2010; 132:15493–15495. [PubMed: 20945898]
13. For alkylation of specialized ester derivatives under phase transfer conditions, see: Andrus MB, Christiansen MA, Hicken EJ, Gainer MJ, Bedke DK, Harper KC, Mikkelsen SR, Dodson DS, Harris DT. *Org Lett.* 2007; 9:4865–4868. [PubMed: 17944480] Andrus MB, Harper KC, Christiansen MA, Binkley MA. *Tetrahedron Lett.* 2009; 50:4541–4544.
14. The asymmetric alkylation of dialkyl amides by Wu and coworkers (see Ref. 8) does not represent an asymmetric alkylation of *ester enolate equivalents* due to the difficulty of derivatizing the amide products.
15. a) Tsuji J, Minami I, Shimizu I. *Tetrahedron Lett.* 1983; 24:1793–1796. b) Behenna DC, Stoltz BM. *J Am Chem Soc.* 2004; 126:15044–15045. [PubMed: 15547998] c) Trost BM, Xu J. *J Am Chem*

- Soc. 2005; 127:2846–2847. [PubMed: 15740108] d) Trost BM, Xu J, Schmidt T. J Am Chem Soc. 2010; 131:18343–18357. [PubMed: 19928805]
16. For reviews of transition metal catalyzed decarboxylative allylic alkylations, see: You SL, Dai LX. Angew Chem. 2006; 118:5372–5374. Angew Chem Int Ed. 2006; 45:5246–5248. Braun M, Meier T. Angew Chem. 2006; 118:7106–7109. Angew Chem Int Ed. 2006; 45:6952–6955. Stoltz BM, Mohr JT. Chem Asian J. 2007; 2:1476–1491. [PubMed: 17935094]
17. The relative stereochemistry of the major diastereomer of **2g** was assigned by analogy to similar compounds. See reference 12b.
18. Tannert R, Milroy L-G, Ellinger B, Hu T-S, Arndt H-D, Waldmann H. J Am Chem Soc. 2010; 132:3063–3077. [PubMed: 20148556]
19. For recent examples, see: Evans DA, Sjogren EB, Bartroli J, Dow RL. Tetrahedron Lett. 1986; 27:4957–4960. Evans DA, Rajapakse HA, Stenkamp D. Angew Chem. 2002; 114:4751–4755. Angew Chem Int Ed. 2002; 41:4569–4573.
20. Trost BM, Machacek MR, Aponick A. Acc Chem Res. 2006; 39:747–760. [PubMed: 17042475]
21. Butts CP, Filali E, Lloyd-Jones GC, Norrby P-O, Sale DA, Schramm Y. J Am Chem Soc. 2009; 131:9945–9957. [PubMed: 19435358]

Scheme 1.
Functionalization of chiral acylbenzoxazolidinone **2e**.

Scheme 2.
Transition state model for enantioselectivity.

Table 1

Auxiliary screen of ester enolate equivalents.^[a-c]

1a: 54% 2a: 87% [†] 12% ee	1b: 79% 2b: 71% 30% ee	1c: 83% 2c: 68% 10% ee	1d: 30% 2d: 87% 49% ee

^[a] Reactions performed on 1–2 mmol scale.^[b] Yield are isolated yields.^[c] Ee values determined by chiral HPLC analysis; absolute configuration of **2a–2c** not determined. Reagents and conditions: a) NaHMDS, DME, –78 °C, then allylchloroformate. b) Pd₂dba₃•CHCl₃ (2.5 mol%), **4** (6 mol%), dioxane, rt, 16 h.[†] Run in toluene. DME = 1,2-dimethoxyethane. dba = dibenzylideneacetone. NaHMDS = sodium bis(trimethylsilyl)amide.

Table 2

Optimization of asymmetric alkylation reaction.

entry ^a	ligand	solvent	yield (%) ^b	ee (%) ^c
1	4	Dioxane	87	49
2	5	Dioxane	88	62
3	6a	Dioxane	96	64
4	7	Dioxane	83	60
5	6a	Toluene	99	75
6	6a	CH_2Cl_2	66	63
7	6a	THF	96	75
8	6a	DME	99	73
9	6b	THF	99	85
10 ^d	6c	THF	12	54
11	6d	THF	25	30
12	8	THF	22	60

^aReactions run using 0.2 mmol enol carbonate **1d**, 0.005 mmol $\text{Pd}_2(\text{dba})_3 \cdot \text{CHCl}_3$, and 0.014 mmol ligand at 0.2 M for 10–30 min.^bIsolated yields.^cDetermined by chiral HPLC analysis.^dRun for 12 hrs.

Table 3

Substrate scope for the asymmetric alkylation reaction.

entry ^a	R ₁	R ₂	Yield (%) ^b	ee (%) ^c
1	Me		99	85
		2d		
2	Me		89	97
		2e		
3 ^d	Me		82	82
		2f		
4 ^e	Me		58	99
		2g		
5 ^{d,f}	Me		96	90
		2h		
6 ^g	Et		68	88
		2i		
7 ^h	Et		93	97
		2j		
8 ^h	<i>n</i> -Pr		76	97
		2k		
9 ^f	<i>n</i> -Pr		83	92
		2l

entry ^a	R ₁	R ₂	Yield (%) ^b	ee (%) ^c
10 ^h		 2m	94	95
11	CH ₂ Ph	 2n	73	94
12	cyclohexyl	 2o	77	80
13 ^f	cyclohexyl	 2p	80	83
14 ^h		 2q	66	95
15 ^g		 2r	71	79
16 ^h		 2s	94	95
17 ^d		 2t	72	81

^a See Table 1 for experimental details. Reactions run on 0.1–0.2 mmol scale for 4–24 hrs.

^b Isolated yield.

^c Determined by chiral HPLC analysis.

^d Using ligand **6a**.

^e Diastereomeric ratio of **2g** is >95:5, see reference 14 for relative stereochemistry assignment.

^fLinear : branched ratio of **2h**, **2l**, **2p** is >10:1.

^gRun at 50 °C.

^hRun with 5 mol% Pd₂(dba)₃•CHCl₃ and 14 mol% **6b**.