Published in final edited form as:

Angew Chem Int Ed Engl. 2009 ; 48(4): 767–770. doi:10.1002/anie.200805140.

Formation of the Pyridazine Natural Product Azamerone Involves the Biosynthetic Rearrangement of an Aryl Diazoketone

Jaclyn M. Winter,

Scripps Institution of Oceanography, University of California at San Diego, 9500 Gilman Drive, La Jolla, CA 92093-0204, Fax: (+1) 858-534-1305

Ariane L. Jansma,

Skaggs School of Pharmacy and Pharmaceutical Sciences, University of California at San Diego, 9500 Gilman Drive, La Jolla, CA 92093

Dr. Tracy M. Handel, and

Skaggs School of Pharmacy and Pharmaceutical Sciences, University of California at San Diego, 9500 Gilman Drive, La Jolla, CA 92093

Dr. Bradley S. Moore

Scripps Institution of Oceanography, University of California at San Diego, 9500 Gilman Drive, La Jolla, CA 92093-0204, Fax: (+1) 858-534-1305. Skaggs School of Pharmacy and Pharmaceutical Sciences, University of California at San Diego, 9500 Gilman Drive, La Jolla, CA 92093

Bradley S. Moore: bsmoore@ucsd.edu

All in the family

A variety of feeding experiments with ^{13}C and ^{15}N -labeled molecules established the phthalazinone core of azamerone is derived from the diazo chlorinated meroterpenoid SF2415A3 (see scheme). We propose that an oxidative rearrangement of the aryl diazoketone followed by rearomatization with the dinitrogen group occurs during the biotransformation. This unique biochemistry extends our limited knowledge of the biosynthesis of natural products containing N–N bonds.

Keywords

biosynthesis; marine natural product; nitrogen heterocycle; stable isotopes; *Streptomyces*

Compounds containing N–N bonds have been made synthetically for over a hundred years. However, it wasn't until 1951 that the first naturally occurring compound containing dinitrogen, the azoxy containing toxin macrozamin, was reported.^[1] Since that time, natural products containing hydrazine, nitrosamine, azoxy, diazo and other N–N bonded functional groups have been identified. The enzymes responsible for N–N bond formation could make valuable biocatalysts, but unfortunately, the biosynthesis of this rare structural unit is poorly understood.

Biosynthetic investigations of the azoxy containing antibiotic valanimycin (**1**)^[2] and the antifungal antibiotic pyridazomycin (**2**)^[3a] (Figure 1) have shown that both of these natural

Correspondence to: Bradley S. Moore, bsmoore@ucsd.edu.

Homepage: <http://moorelab.ucsd.edu>

Supporting information for this article (experimental methods and NMR spectra) is available on the WWW under <http://www.angewandte.org> or from the author.

products are derived from the condensation of amino acids. Pyridazomycin^[3b] was the only natural product containing a pyridazine ring until the phthalazinone meroterpenoid azamerone (**3**) was isolated from the marine sediment-derived bacterium *Streptomyces* sp. CNQ-766.^[4] It was first speculated that the pyridazine ring moiety in **3** was also derived from the cyclization of amino acid residues, but identification of co-produced chlorinated meroterpenoids such as A80915C (**4**), 3''-hydroxy-7-methylnapyradiomycin B2 (**5**) and analogs possessing a diazo functional group (**6–8**)^[5] suggest that **3** may rather be derived from a naphthoquinone precursor (Scheme 1). Herein, we report the biosynthesis of **3** from a series of stable isotope tracer experiments, which validate its biosynthetic relationship to the napyradiomycin family of meroterpenoid antibiotics.

The original isolation of **3** from *S.* sp. CNQ-766 generated only ~0.25 mg/L,^[4] which was insufficient for labeling experiments. Different fermentation conditions were thus evaluated to improve production yields that resulted in >30-fold increase in **3** at 8.5 mg/L with M1 medium.^[5a] In addition to exploring the biosynthesis of **3** with stable isotopes, we also examined **4** and the diazo meroterpenoids SF2415A3 (**6**) and A80915D (**7**) in order to probe the biosynthetic relationship of this extended structural family. Feeding experiments with [1,2-¹³C₂]acetate clearly revealed that the naphthoquinone core of **4** was derived from the symmetrical pentaketide 1,3,6,8-tetrahydroxynaphthalene (THN) and that the two isoprenoid units originated via the mevalonate pathway (Scheme 1).^[6] These observations were consistent with previous isotopic experiments in this compound series from *Chainia rubra* MG802-AF1^[7] and from the analysis of the 43-kbp napyradiomycin biosynthetic gene cluster (*nap*) from *Streptomyces aculeolatus* NRRL 18422 and *Streptomyces* sp. CNQ-525.^[8]

¹³C-NMR analysis of [1,2-¹³C₂]acetate-enriched **3** also showed that all carbons except for the C8 methyl group originate from acetate (Figure 2).^[6] The bicyclic phthalazinone unit clearly harbored two distinct ¹³C-labeling patterns of equal proportions, thereby implying that it too derives from the symmetrical THN unit common to the napyradiomycins. Further analysis of L-[methyl-¹³C]methionine-enriched **3** revealed the 26% selective enrichment of C8, confirming that the acetyl methyl group of **3** is methionine-derived.^[6] Importantly, it was observed that if labeled materials were administered to *S.* sp. CNQ-766 during the initial production of **3**, which is roughly five days after inoculation during stationary growth, no enrichment could be detected. Labeled materials had to be added at the time of inoculation in order for them to be efficiently incorporated, thereby suggesting that **3** originates from a methylated intermediate produced earlier in the fermentation process. Altogether these data suggest that **3** is biosynthetically linked to the napyradiomycin family of natural products.

We turned our attention to the biosynthetic origin of the diazo functional group as seen in **6–8** and its relation to **3**. Previous studies on the biosynthesis of the kinamycins with synthetic and natural intermediates revealed that its diazonium group is assembled from the step-wise addition of two nitrogen donors to a ketobenzo[b]fluorene precursor through an aminobenzo[b]fluorene intermediate into kinamycin D (**9**) (Figure 1).^[9] Considering a similar scenario, we administered different forms of ¹⁵N-enriched substrates, including ammonium sulfate, nitrite, and nitrate, and measured their incorporation into **3**, **6** and **7** by MS and NMR (Table 1).^[6] MS analysis revealed that all ¹⁵N-labeled precursors were incorporated to different extents with the oxidized forms having the highest assimilation rates. While [¹⁵N]ammonium sulfate singly enriched **3** and **7** between 7–11%, ¹H-¹⁵N HMBC analysis of **3** showed that both nitrogen atoms were equally labeled. This result was in stark contrast to the enrichment by [¹⁵N]nitrite and [¹⁵N]nitrate, which not only were incorporated at a much greater extent at >70% but only labeled one of the two nitrogen atoms in the natural product. In the case of **7**, ¹⁵N-NMR showed selective enrichment of the

distal diazo nitrogen (-31 ppm).^[9c,10] ¹⁵N-labeling of **3** occurred at -2 ppm, which was assigned to the C6-bound nitrogen N2 in the pyridazine ring based upon the J_{H-N} coupling constants in the ¹H-¹⁵N HMBC experiment.^[6]

Results from the ¹⁵N feeding studies suggested that the napyradiomycin diazo group is also formed in a step-wise manner. Oxidation of the naphthyl ring at C5 facilitates the introduction of the first nitrogen atom by a transamination reaction. The formation of the N–N bond arises from a nucleophilic attack of the aminodihydroquinone intermediate on nitrous acid (Scheme 1). The direct addition of a dinitrogen precursor was also probed with [¹⁵N₂]hydrazine. MS analysis revealed that [¹⁵N₂]hydrazine was only incorporated singly and at a low level into **3** (Table 1). Furthermore, ¹H-¹⁵N HMBC analysis showed that both nitrogen atoms in **3** were labeled, yet unlike the [¹⁵N]ammonium-labeled product, N2 was enriched to a slightly higher extent than N1 suggesting that metabolism of hydrazine in *S. sp.* CNQ-766 results in non-equivalent forms of single nitrogen species.^[6]

The ¹⁵N labeling studies indirectly supported that the diazo group such as in **6** and **7** is a biosynthetic precursor to the pyridazine unit in **3** based on their similar enrichment characteristics. In order to unequivocally link the aryl diazoketones to **3**, we biosynthetically prepared [2-¹⁵N, 9-¹³C]**6** from [¹⁵N]nitrite and L-[methyl-¹³C]methionine and fed the labeled natural product back to *S. sp.* CNQ-766. HPLC-MS inspection of the other napyradiomycin derivatives indicated that **6** was converted into **7** and **3** with retention of both isotopes in the same molar ratio as in the substrate. Isolation and NMR characterization of [2-¹⁵N, 8-¹³C]**6**-enriched **3** showed ¹³C enrichment at C8 and ¹⁵N enrichment at N2,^[6] thereby confirming that not only is **6** a precursor of **3**, but importantly that the diazo N2 atom is biosynthetically equivalent with N2 of the pyridazine group in **3**.

These stable isotope experiments paint a picture uniting **3** with the napyradiomycin family of chlorinated meroterpenoids in which a Baeyer–Villiger-type oxidation of a diazonaphthoquinone such as **7** may initiate the biosynthetic interconversion (Scheme 2). Hydrolysis of the 7-membered heterocyclic intermediate would open the ring to facilitate (1) the 1,2-alkyl shift of the monoterpene subunit and (2) the assembly of the pyridazine ring in which the diazo group forms a new linkage with the C8 diketide. Aromaticity of the ring would be restored by subsequent decarboxylation and dehydration reactions.

The molecular basis for the formation of the diazo group and its rearrangement, however, is presently unresolved even though the napyradiomycin biosynthetic gene cluster from the related *Streptomyces sp.* CNQ-525 has been cloned and sequenced.^[8] Further examination of this strain revealed that like CNQ-766, it too synthesizes **3**. However inspection of the *nap* gene cluster did not reveal an obvious mechanism for diazo synthesis and rearrangement nor did its heterologous expression yield nitrogenated napyradiomycin analogs. The only gene that may encode a diazo biosynthetic enzyme is *napB3*, which codes for a putative aminotransferase and may facilitate the transamination to the aminodihydroquinone intermediate. The *nap* cluster furthermore harbors a number of oxygenases whose functions have not yet been clarified that may be involved in the nitrogen biochemistry of **3**. It is quite likely though that some of the encoding genes are extraneous to the *nap* locus and may reside elsewhere in the *S. sp.* CNQ-525 genome given the expression of the cluster only yielded non-nitrogenated napyradiomycins.^[8]

In conclusion, we established with ¹³C- and ¹⁵N-labeled precursors that azamerone (**3**) is biosynthesized via the *nap* pathway through SF2415A3 (**6**) in which the aryl diazoketone undergoes a novel rearrangement wherein the aromatic ring is oxidatively cleaved to allow for its rearomatization with dinitrogen to give the unique phthalazinone core of **3**. This unprecedented biochemistry extends our limited knowledge of the biosynthesis of natural

products containing N–N bonds and opens the door to exploring and exploiting its molecular basis at the biochemical and genetic levels.

Acknowledgments

This work was supported by the California Sea Grant Program (R/NMP-98 to B.S.M.) and by a Ruth L. Kirschstein National Research Service Award to J.M.W. from the NIH Training Program in Marine Biotechnology (GM067550). We thank Drs W. Fenical, P. R. Jensen, H. C. Kwon and J. Y. Cho (University of California at San Diego, USA) for strain *S. sp.* CNQ-766, standard azamerone, and with R. J. Parry (Rice University, USA), T. F. Molinski, T. A. M. Gulder, and A. A. Mrse for helpful advise.

References

1. a) LaRue TA. Lloydia. 1977; 40:307–321. [PubMed: 142890] b) Langley BW, Lythgoe B, Riggs NV. J Chem Soc. 1951:2309–2316.
2. a) Parry RJ, Li Y, Lii FW. J Am Chem Soc. 1992; 114:10062–10064. b) Garg RP, Gonzalez JM, Parry RJ. J Biol Chem. 2006; 281:26785–26791. [PubMed: 16857674] c) Garg RP, Qian XL, Alemany LB, Moran S, Parry RJ. Proc Nat Acad Sci USA. 2008; 105:6543–6547. [PubMed: 18451033]
3. a) Bockholt H, Beale JM, Rohr J. Angew Chem Int Ed. 1994; 33:1648–1651. b) Grote R, Chen Y, Zeeck A. J Antibiot. 1988; 41:595–601. [PubMed: 3384747]
4. Cho JY, Kwon HC, Williams PG, Jensen PR, Fenical W. Org Lett. 2006; 8:2471–2474. [PubMed: 16737291]
5. a) Fukuda DS, Mynderse JS, Baker PJ, Berry DM, Boeck LD, Counter LD, Ensminger PW, Allen NE, Alborn WE, Hobbs JN. J Antibiot. 1989; 63:623–633. b) Soria-Mercado IE, Prieto-Davo A, Jensen PR, Fenical W. J Nat Prod. 2005; 68:904–910. [PubMed: 15974616] c) Gomi S, Ohuchi S, Sasaki T, Itoh J, Sezaki M. J Antibiot. 1987; 40:740–749. [PubMed: 3610831]
6. See supporting information for full details.
7. Shiomi K, Iinuma H, Naganawa H, Isshiki K, Takeuchi T, Umezawa H. J Antibiot. 1987; 40:1740–1745. [PubMed: 3429338]
8. Winter JM, Moffitt MC, Zazopoulos E, McAlpine JB, Dorrestein PC, Moore BS. J Biol Chem. 2007; 282:16362–16368. [PubMed: 17392281]
9. a) Gould SJ, Melville CR, Cone MC, Chen J, Carney JR. J Org Chem. 1997; 62:320–324. [PubMed: 11671405] b) Gould SJ, Melville CR. Bioorg Med Chem Lett. 1995; 5:51–54. c) Seaton PJ, Gould SJ. J Am Chem Soc. 1988; 110:5912–5914.
10. a) Lippmaa E, Pehk T, Saluvere T, Mägi M. Org Magn Reson. 1973; 5:441–444. b) Glaser R, Horan CJ. J Org Chem. 1995; 60:7518–7528.
11. Biemann, K. Mass Spectrometry: Organic Chemical Applications. McGraw-Hill Book Co; New York: 1962. p. 223–231.

Figure 1.
Selected natural products containing a N–N bond.

Figure 2.

^{13}C -labeling of azamerone (**3**) derived from $[1,2-^{13}\text{C}_2]$ acetate and L-[methyl- ^{13}C]methionine. Two equally independent labeling patterns **3a** and **3b** were evident from the incorporation of $[1,2-^{13}\text{C}_2]$ acetate (bold lines and dots signify double and single enrichments, respectively). The enrichment from L-[methyl- ^{13}C]methionine is shown as a dashed line. The numbering scheme for azamerone has been adopted from Ref. 5c and 8.

Proposed biosynthetic pathway for the diazo chlorinated meroterpenoids **6–8**^[8] and their relation to the chlorinated dihydroquinones such as **4** and **5**. A summation of the two [1,2-¹³C₂]acetate labeling patterns are shown in black (bold lines and dots signify double and single enrichments, respectively). Enrichment from L-[methyl-¹³C]methionine is shown as a dashed line, and enrichment from [¹⁵N]nitrite or [¹⁵N]nitrate is shown in bold font.

Scheme 2.

Proposed oxidative rearrangement of the aryl diazoketone SF2415A3 (**6**) via A80915D (**7**) to azamerone (**3**) in *S. sp.* CNQ-766. Isotopically labeled C and N atoms from $[2\text{-}^{15}\text{N}, 9\text{-}^{13}\text{C}]\textbf{6}$ are shown in bold font.

Table 1

^{15}N NMR data of **3** and **7** and incorporation percentages from nitrogen labeled precursors.

Labeled Precursors	Enrichment of 3 ^[a]		Enrichment of 7 ^[a]	
	Singly/Doubly	$\delta^{15}\text{N}$ ppm ^[b]	Singly/Doubly	$\delta^{15}\text{N}$ ppm ^[c]
$^{15}\text{NO}_2\text{Na}$	77/6	−2	77/−	−31
$^{15}\text{NO}_3\text{Na}$	74/3	−2	79/−	−31
$^{15}\text{NH}_4\text{SO}_4$	11/−	−2 & 13	7/−	− [d]
$^{15}\text{N}_2\text{H}_4\text{SO}_4$	5/0.3	−2 & 13	0/−	− [d]
$[2\text{-}^{15}\text{N}, 9\text{-}^{13}\text{C}]\text{-6}$	11/6	−2	6/4	− [d]

^[a] Enrichment was calculated by m/z ^[11];

^[b] $^1\text{H}\text{-}^{15}\text{N}$ HMBC spectra were measured in CD_3OD at 600 MHz and referenced internally to nitromethane (δ 0 ppm);

^[c] ^{15}N NMR spectra were measured in CD_3CN at 30.4 MHz and referenced externally to nitromethane (δ 0 ppm);

^[d] chemical shift(s) were not measured due to inadequate sample size and low enrichment levels.