A Primary Giant Hydatid Cyst of the Ovary

Abdurrahman Kaya^{1*}, Sibel Yildiz¹, Reşat Özaras¹, Ali Mert¹

¹ Department of Infectious Diseases, Medical School of Cerrahpasa, Istanbul University, Istanbul, Turkey

ARTICLE INFO

Article type:
 Letter to Editor

Article history:
 Received: 18 Jul 2011
 Revised: 01 Oct 2011
 Accepted: 16 Oct 2011

Keywords:
 Cysts
 Ovary

► Please cite this paper as:

Kaya A, Yildiz S, Özaras R, Mert A. A Primary Giant Hydatid Cyst of Ovary. *Iran J Radiol.* 2012;9(3):165-6. DOI: 10.5812/iranjradiol.7955

Copyright ©2012, Tehran University of Medical Sciences and Iranian Society of Radiology. Published by Kowsar Corp. All rights reserved.

Dear Editor,

A 38-year-old female patient was admitted with right lower abdominal pain of 4 months duration. On her clinical examination, right lower abdominal tenderness was noted. Complete blood count including the numbers of eosinophils (80/mm³) was normal. Biochemistry was normal. An abdominal MRI showed an 11 × 9 × 5.5 cm sized cystic lesion in the right ovary surrounded by loculated fluid collection with septations (*Figure 1*). Echinococcal serology was positive (hydatid cyst antibodies by ELISA and indirect hemagglutination test dilutions up to 1/1280 were positive). Pathological examination of the material had a cuticle and germinative membrane of the hydatid cyst and around it chronic nonspecific inflammation leading to the formation of lymphoid follicles was seen. No other organ involvement was detected. Albendazole (10mg/kg/day) was prescribed. The cyst was removed surgically on the 30th day of albendazole treatment. Pathological examination of the cyst was compatible with hydatid cyst. The patient had regular follow-ups, with no recurrence of the cyst. Echinococcal disease is endemic in

Figure 1. Pelvic MRI revealed a septated, lobulated and contrast enhancing cystic mass that is hyperintense on T2 and hypointense on T1 FLAIR sequences.

* Corresponding author: Abdurrahman Kaya, Department of Infectious Diseases, Istanbul, Turkey, Tel: +90-5066113328, Fax: +90-2126320050, E-mail: dr.abdkaya@hotmail.com

many countries (1). It involves multiple organs particularly the liver (50-70%) and the lung (20-30%), while primary involvement of the ovary is very rare (2). The liver and spleen cysts can rupture into the peritoneal cavity which is the way of secondary involvement (3). In our case, there were no cysts in other organs. Although ovarian cysts are usually single, this case showed multilocular cysts the treatment is surgical removal and albendazole is recommended before surgery.

Authors' Contribution

Abdurrahman Kaya followed the patients and wrote the paper. Sibel Yıldız assisted in writing and took the pho-

tos. Reşat Özaras coordinated the follow-up and edited the paper. Ali Mert coordinated the follow-up and edited the paper.

Financial Disclosure

None declared.

References

1. Nakayama DK. Survival in a fetus with sacrococcygeal teratoma and hydrops. *Am J Obstet Gynecol.* 1990;**163**(2):682.
2. Fahy AM, Peters EE. Eccrine tumours of the hand: two cases. *J Hand Surg Br.* 1987;**12**(1):120-2.
3. Sing P, Mushtaq D, Verma N, Mahajan NC. Pelvic hydatidosis mimicking a malignant multicystic ovarian tumor. *Korean J Parasitol.* 2010;**48**(3):263-5.