Published in final edited form as:

Chem Sci. 2013 January 1; 4(1): 398–404. doi:10.1039/C2SC20763J.

Expanding DNAzyme Functionality through Enzyme Cascades with Applications in Single Nucleotide Repair and Tunable DNA-Directed Assembly of Nanomaterials

Yu Xiang^a, Zidong Wang^b, Hang Xing^a, and Yi Lu^{*,a,b}

^aDepartment of Chemistry, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA

^bDepartment of Material Science, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA

Abstract

Many biological functions require two or more enzymes working together in cascades. While many examples of protein and RNA enzyme cascades are known, few enzyme cascades containing solely DNAzymes have been reported. Herein we demonstrate the combination of an 8–17 DNAzyme with RNA cleavage activity and an E47 DNAzyme with DNA ligation activity to achieve a new function of single ribonucleotide repair in DNA while maintaining the integrity of the original DNA sequence, which is difficult for a single DNAzyme to achieve. In addition, this method is applied to modify the sequences of DNA strands immobilized on the surface of nanoparticles to control the DNA-directed assembly selectively and sequentially. Such an approach can be applied to other DNAzymes with different activities to expand the functions of DNAzymes and the scope of their applications.

Introduction

The discovery of deoxyribozymes (DNAzymes) with enzymatic activity in the 1990s^{1, 2} has demonstrated that DNA molecules are not simply inert biopolymers for genetic storage; they can be active catalysts as well.^{3–8} Since then, many DNAzymes have been obtained with catalytic functions such as cleavage,^{2, 9–13} ligation,^{14–16} phosphorylation,¹⁷ adenylation¹⁸ or depurination¹⁹ of nucleic acids, as well as other reactions including porphyrin metallation, C–C bond formation, nucleopeptide linkage formation, oxygen transfer and thymine dimer repair.^{20–26} Because DNAzymes are facile to synthesize and more stable than protein and RNA enzymes, they have been widely used in applications such as nanomaterial assembly,^{27, 28} biosensing,^{29–31} logical computing,³² nanomachine engineering,³³ antiviral or gene therapy,³⁴ and *in vitro* RNA manipulation.³⁵ Despite these successes, the application of DNAzymes is limited by the narrower range of catalytic functionality compared to protein enzymes. One possible approach to addressing this issue would be to combine enzymes with different reactivities to form a cascade of successive enzymatic reactions, which together create new functionality. Indeed, many such examples exist in biology, since nearly all important biological functions, such as the pathways involved in DNA repair and protein synthesis, require a cascade of multiple protein enzymes to carry out their full function. In contrast, little has been reported about the use of DNAzyme cascades to realize enhanced functionality. Such a strategy could expand the functionality of DNAzymes to a

level more on par with protein and RNA enzymes, which should greatly increase the range of possible applications.

One such application is single nucleotide repair, i.e., excision of a misincorporated ribonucleotide in single-stranded DNA and subsequent insertion of the corresponding deoxyribonucleotide at the excision site. The misincorporation of ribonucleotides into DNA strands can occur from exposure to external oxidizing agents or ionizing radiation,³⁶ or spontaneously during DNA replication.³⁷ misincorporation of ribonucleotide can distort the structure of DNA,³⁸ reduce its stability,³⁹ and interfere with the normal interaction between DNA and DNA polymerases.⁴⁰ In fact, the overexpression of DNA polymerases that are prone to ribonucleotide misincorporation has been linked to many cancers, including ovarian, prostate, breast and colon cancers.⁴¹ In nature, protein enzymes such as RNase H and FEN-1 can efficiently excise misincorporated ribonucleotides in DNA by cleaving the DNA at the ribonucleotide site and then restore the correct deoxyribonucleotides by DNA polymerases,^{42, 43} which is an example of an enzyme cascade. It would be interesting to find out if a similar function could be achieved through DNAzyme cascades.

Another potential application is in tuning the properties of DNA-functionalized nanomaterials. For example, DNA-functionalized gold nanoparticles²⁷ have emerged as an attractive platform for biosensing,^{32, 44–50} nanomedicine,⁴⁵ and as building blocks for controlled nanoassemblies.^{51–55} Although much research has been focused on the surface modification of gold nanoparticles with DNA for various applications, there are still limited methods to modify the sequences of DNA already immobilized on gold nanoparticles in order to make the properties of the DNA-modified nanomaterials tunable after fabrication. The use of DNAzymes is a promising approach for DNA modification on nanomaterials⁵⁶ due to the excellent stability of DNAzymes and their smaller size compared to protein enzymes, which should minimize steric effects between the enzyme and the DNA to avoid reduction in reaction efficiency. However, it is still very challenging to modify a specific DNA sequence on multiple-DNA-functionalized nanomaterials to tune their functions in a selective and sequential fashion.

Herein, we demonstrate a cascade of two DNAzymes with RNA cleavage and DNA ligation activities, respectively, in order to carry out single nucleotide repair or selective sequence modification of DNA. In a one-pot reaction, a single misincorporated ribonucleotide in a DNA strand was converted to the corresponding deoxyribonucleotide while maintaining sequence integrity. Furthermore, the sequences of DNA strands immobilized on multiple functional nanoparticles were successfully modified in order to control and alter the DNA-directed assembly of nanoparticles in a stepwise and selective fashion.

Results and Discussion

To demonstrate that single nucleotide repair in DNA can be achieved by the cascade of two DNAzymes, we used a 26-nt DNA strand (**O1**) containing a misincorporated cytidine (rC) ribonucleotide as an example. The goal was to convert the rC in **O1** into a deoxycytidine (C), as seen in **O4** (Figure 1a), while maintaining the integrity of the DNA sequence. The DNAzymes 17E_{m1} (Figure 1a, blue) with RNA cleavage activity^{2, 57–59} and E47 (Figure 1a, red) with DNA ligation activity^{14, 60} were chosen as the cascade pair in this study. The 17E_{m1} DNAzyme catalyzes the hydrolysis of the 3' phosphodiester linkage of the internal rC in the DNA strand when metal ion cofactors such as Pb²⁺ and Zn²⁺ are present (Figure S1a in Supplementary Information). On the other hand, the E47 DNAzyme can induce the catalytic ligation of the 5' –OH of the DNA substrate with another 3'- phosphorylated DNA strand (activated by imidazole)¹⁴ in the presence of Cu²⁺ or Zn²⁺ as the metal cofactor (Figure S1b in Supplementary Information). Therefore, by sequential cleavage and ligation

reactions catalyzed by these two DNAzymes on **O1** containing rC, **O1** could first be cleaved at the 3' phosphodiester of the rC by 17E_{m1} and then undergo ligation at the cleavage site with another 3'-phosphorylated DNA strand of identical sequence (except with deoxyribonucleotide C in place of ribonucleotide rC) by E47. The product **O4** has a sequence identical to the starting strand **O1**, with the rC replaced with C.

Initially, 3'-fluorescein-labeled **O1** was treated with 17E_{m1} to form DNA duplex **O1**-17E_{m1} via 18 matched base pairs (9 on each binding arm). In the presence of Pb²⁺, **O1** was efficiently cleaved by 17E_{m1} into fragments **O2** and **Oc**, resulting in the dehybridization of the duplex because the melting temperature of the duplex between 17E_{m1} and **O2** or between 17E_{m1} and **Oc** is below room temperature. The fluorescence image after polyacrylamide gel electrophoresis (PAGE) suggested the complete cleavage of **O1** and formation of **O2** (Figure 2a, lane 1 and 2 for **O1** and **O2**, respectively), while **Oc** was not visible on the gel due to the lack of a fluorescein label. The cleavage reaction product **O2** was also confirmed by the result from matrix-assisted laser desorption ionization-time of flight (MALDI-TOF) mass spectrum (Table 1 and Figure S2 in Supplementary Information). Control experiments using DNAzyme of a different sequence (17E_{m2}) or without Pb²⁺ showed negligible cleavage of the substrate **O1** (Figure S3 in Supplementary Information) due to the specificity of the DNAzyme and the essential role of the metal ion cofactor.^{2, 57-59} Subsequently, without any purification of **O2** from the mixture solution after the previous cleavage step, E47 and 3'-phosphorylated **O3** (imidazole-activated) were added into the solution to generate another DNA complex **O2**-**O3**-E47, which gave **O4** as the product after the E47-catalyzed ligation reaction in the presence of Cu²⁺ (Figure 1a).^{14, 60} The formation of **O4** was confirmed by both fluorescent PAGE (Figure 2a, the upper band of lane 3) and MALDI-TOF MS (Table 1 and Figure S2 in Supplementary Information), while some unreacted **O2** was also observed on the gel (Figure 2a, the lower band of lane 3). Here **O3** was invisible due to the lack of a fluorescein label. Considerably lower levels of ligation between **O2** and **O3** were observed if either E47 or Cu²⁺ was absent (Figure S3 in Supplementary Information). Note that the reactions catalyzed by the DNAzyme cascade were achieved through a one-pot reaction without isolation and purification of the intermediate **O2**.

To provide further confirmation of the above successful conversion of rC to C in from **O1** to **O4**, a longer **O3+8A** (**O3** extended by A₈ at 5') was used in place of **O3** (Figure 1a). Under the same conditions, a longer product **O4+8A** was obtained (Figure 2a, the upper band of lane 4) with slower gel migration compared to **O4** (Figure 2a, the upper band of lane 3), suggesting that the ligation reaction occurred mostly between **O2** and imidazole-activated **O3**, and not between **O2** and un-activated **Oc** (**Oc** is the product from the previous cleavage reaction of **O1** and 17E_{m1}), in which case a band with the same migration as **O4** would have been observed. The presence of C rather than rC in the product **O4** was supported by the lower molecular weight of **O4** in the MALDI-TOF mass spectrum compared to that of **O1** (Table 1), as well as the increased resistance to hydrolysis of **O4** even in the presence of Pb²⁺ and 17E_{m1} (Figure 2a, lane 5), which can catalyze the cleavage of a substrate containing an internal ribonucleotide linkage (**O1**),^{2, 57} but not a substrate containing entirely deoxyribonucleotides (**O4**).

In addition to the single nucleotide repair functionality, it is also possible to use this methodology to edit the sequence of a DNA strand, which was used to convert the DNA strand **O5** into **O4** using the same cascade and conditions as before (Figure 1b and Figure S4 in Supplementary Information). The product **O4** was confirmed by PAGE (Figure 2b) and MALDI-TOF MS (Table 1 and Figure S2 in Supplementary Information) and found to be identical to that obtained from the method in Figure 1a.

Encouraged by the above results, we applied this method to modify the sequence of DNA immobilized on gold nanoparticles²⁷ (AuNPs) to control the DNA-directed assembly of the AuNPs in a selective manner. DNA-functionalized gold AuNPs have been used in a variety of applications due to both their unique properties and the sequence-dependent hybridization of ssDNA immobilized on the AuNPs for controlled assembly.^{51–55} As shown in Figure 3, when AuNPs are functionalized by complementary DNAs, the AuNPs can assemble into an “aggregated” state via DNA hybridization, which shows red-shifted and broadened absorption spectra compared to AuNPs functionalized by non-complementary DNAs. By modifying the sequences of DNA on the AuNPs, the assembly of the particles can be effectively controlled. Although methods for fabrication of DNA-functionalized AuNPs have been developed,²⁷ there are still limited methods to modify the DNA sequences already immobilized on AuNPs in order to tune their functions. It is even more challenging to modify a specific DNA sequence on multiply-functionalized AuNPs with different DNA sequences on each nanoparticle. Selective modification can allow each different function of the AuNP to be controlled in a selective fashion for potential applications.

AuNPs of 13 nm diameter were functionalized with DNA molecules via 3'-end thiols and used for this study. The formation of AuNP assembly was confirmed by TEM images (Figure S5 in Supplementary Information) and characterized as large changes in absorption spectra²⁷ (A_{700}/A_{532} changed from < 0.15 to > 0.50 as illustrated in Figure 5, and Table S1 in Supplementary Information). As shown in Figure 4, **O6**-functionalized AuNPs (red) were found to be able to form aggregates with **O9**- functionalized AuNPs (blue) via DNA-directed assembly through 12 complementary base pairs (Figure 5, Figure S5 and Table S1), but not with **O10**-functionalized AuNPs (purple), because of the 4 mismatched base pairs in the middle of the binding arm (Figure 5, Figure S5 and Table S1). After treated with $17E_{m2}$ and Pb^{2+} , the **O6** on the surface of AuNPs was cleaved and converted to **O8**, which could not hybridize with either **O9** or **O10** efficiently. Thus no DNA-directed assembly was observed between the resulting **O8**- functionalized AuNPs and either **O9**- or **O10**-functionalized AuNPs (Figure 5, Figure S5 and Table S1). However, after a subsequent ligation reaction catalyzed by E47 in the presence of imidazole-activated **O3** and Cu^{2+} , **O8** on the surface of AuNPs could be extended to **O7**, making the AuNPs capable of assembling with **O10**-, but not **O9**-functionalized AuNPs (Figure 5, Figure S5 and Table S1).

Interestingly, the product, **O7**-functionalized AuNPs showed inverse characteristics in the formation of DNA-directed assembly with **O10**- and **O9**-functionalized AuNPs, compared to the original **O6**-functionalized AuNPs. TEM images of **O6**- functionalized AuNPs, either with or without treatment with $17E_{m2}$ /E47, mixed with **O10**-functionalized AuNPs, are displayed in the inset of Figure 4. It clearly demonstrates that the ability to edit and replace DNA on AuNPs allows for exquisite programmable control over the assembly of nanoparticles.

Taking advantage of the specificity of DNAzyme to its nucleic acid substrates by complementary base pairing in the binding arms, selective modification of DNA sequences on surface of multiple functional AuNPs was also achieved in this work. As depicted in Figure 6, **O6** (red) and **O11** (blue) bi-functional AuNPs could be modified by $17E_{m1}$, $17E_{m2}$ and E47 selectively and sequentially.. As shown in Figure 6, AuNPs capable of forming DNA-directed assembly with both (A, E), either (B, C, F), or neither (D) of the **O9**- and **O10**-functionalized AuNPs could be obtained (Figure 6 and Table 2S in Supplementary Information) by monitoring the significant increase of A_{700}/A_{532} as indication of assembly formation. Such a result, which is challenging to achieve by other techniques, can be used for the construction of tunable nanoassemblies for various applications.

Conclusions

In summary, by putting together a cascade of two DNAzymes with cleaving and ligating activities, we have generated a new functionality for effective DNA modification. This function was applied in the conversion of a single misincorporated ribonucleotide into the corresponding deoxyribonucleotide in DNA and the modification of DNA sequences on the surface of gold nanoparticles to modify and control their self-assembly through DNA hybridization. The results suggest that combining DNAzymes with different catalytic activities may achieve more interesting functions and thus broaden the applications of DNAzymes.

Experimental Section

Materials

All DNA samples were purchased from Integrated DNA Technologies Inc. (Coralville, IA). Substrate and enzyme strands of the DNAzyme were purified by HPLC; thiol-modified and phosphorylated DNAs underwent standard desalting. Other chemicals were purchased from Sigma-Aldrich Inc.

The sequences of DNA used in this work were as follows:

Cleaving DNAzyme 17E_{m1}: 5'-ATAGTGTCT CCGAGCCGGTCGAA
ATAGGATG-3'

Cleaving DNAzyme 17E_{m2}: 5'-ATAGTGTCT CCGAGCCGGTCGAA
CTTGATG-3'

Ligating DNAzyme E47: 5'-
CGGATAGTGTCTTTCGCTAGACCATGTGACGCATGGTGAGATGCTT-3'

Fluorescein-labeled substrate O1: 3'-FAM-
GCCTATCACAAAGGrATATCTCTACGAA-5'

3'-phosphorylated O3: 3'-PO₃-CGAACTCTACGAA-5'

3'-phosphorylated O3+8A: 3'-PO₃-CGAACTCTACGAAAAAAAAA-5'

Fluorescein-labeled substrate O5: 3'-FAM-
GCCTATCACAAAGGrCGAACTCTACGAA-5'

3'-thiol-modified O6: 3'-HS-
AAAAAAGCCTATCACAAAGGrATATCTCTACGAA-5'

3'-thiol-modified O9: 3'-HS-AAAAAAGAGATATCCTT-5'

3'-thiol-modified O10: 3'-HS-AAAAAAGAGTTCGCCTT-5'

3'-thiol-modified O11: 3'-HS-
AAAAAAGCCTATCACAAAGGrCGAACTCTACGAA-5'

Experimental procedures

DNAzyme (17E_{m1} or 17E_{m2})-catalyzed cleavage of fluorescein-labelled substrate O1 or O5—The 3'-Fluorescein-labeled substrate DNA **O1** or **O5** (1 μM) was mixed with DNAzyme 17E_{m1} or 17E_{m2} (1 μM) and Pb(NO₃)₂ (5 μM) in 25 mM HEPES (pH 7.0) containing 100 mM NaCl. The mixture was incubated at room temperature for 1 h (for **O1**) or 3 h (for **O5**) to achieve complete cleavage (> 95 %) and subsequently analyzed by PAGE and MALDI-TOF MS.

Imidazole-activation of O3 and O3+8A—In a 0.5 mL centrifuge tube, 20 μ L 100 μ M **O3** or **O3+8A** with 3'-phosphorylation were incubated with 2.5 μ L of 1 M imidazole (pH 6.0, adjusted by adding concentrated HCl), and 4.5 μ L of 1 M EDC•HCl at room temperature for 1 hour. The mixture was then purified with a PD-10 (Amersham Biosciences) desalting column, and the first fraction with strong absorption at 260 nm was collected (about 0.5~1.5 mL range). The DNA concentration of the eluted fraction was determined by monitoring the absorbance at 260 nm. The imidazole-activated **O3** and **O3+8A** were used right away, since they were not very stable at room temperature.

DNAzyme (E47)-catalyzed ligation of O2—The solution containing cleaved substrate from 17E treatment was further mixed with the ligation DNAzyme E47 (2 μ M), $\text{Cu}(\text{NO}_3)_2$ (20 μ M) and imidazole-activated **O3** or **O3+8A** (3 μ M) in 25 mM HEPES (pH 7.0) containing 300 mM NaCl, and allowed to incubate at room temperature for 4 h before analysis by PAGE and MALDI-TOF MS.

PAGE and MALDI-TOF MS—3'-fluorescein-labeled DNA samples (1 μ M) were mixed in a 10:1 ratio with a stop solution containing 100 mM EDTA, to quench the cleavage and ligation reactions. This solution was mixed 1:1 with glycerol and transferred to a denaturing 20 % acrylamide gel. The resulting gel was documented by a fluorescence image scanner (model FLA-3000G, Fuji). DNA isolation from the gel was performed by soaking the cut-out gel band in 1 mL Millipore water for 1 day, then centrifuging at 10 000 rpm for 15 minutes. Afterwards, the fluorescent supernatant was concentrated to about 20 μ L and purified by PD-10 column. The fraction with fluorescence at 520 nm was collected (total volume approximately 0.5~1.5 mL), concentrated to 20 μ L, and then analyzed by MALDI-TOF MS.

Nanoparticle preparation and functionalization (O6, O9, O10 and O11)—The preparation of 13 nm gold nanoparticles was as reported.²⁷ DNAs with 3'-thiol-modifications (**O6**, **O9**, **O10** and **O11**) were activated by incubation with TCEP. Typically, 9 μ L of 1 mM DNA in Millipore water were incubated with 1 μ L of 20 mM freshly prepared TCEP and 1 μ L 500 mM NaAc-HAc buffer (pH 5.2) at room temperature for 1 hour. The mixture was then directly added into 3 mL of citrate-capped 13 nm AuNPs (about 3 nM). After incubation for 16 hours at room temperature, 0.3 mL of buffer containing 1 M NaCl and 100 mM of Tris-acetate (pH 8.2) was added by drop to the nanoparticle solution under stir. After incubation for 24 hours, the nanoparticles were centrifuged at 13 000 rpm for 15 minutes. The supernatant was removed and nanoparticles were re-dispersed in buffer containing 100 mM NaCl and 25 mM Tris-acetate (pH 8.2). This centrifugation process was repeated once more to remove free DNA in solution. For nanoparticles functionalized with both **O6** and **O11** in sequential tuning experiments, the molar ratio of **O6** and **O11** in loading was 4:1, with a total loading amount of 25 μ L of 1 mM DNA. Other procedures were the same as above.

DNAzyme (17E_{m1} or 17E_{m2})-catalyzed cleavage of DNA on gold nanoparticles (O6 and O11)—DNA-functionalized gold nanoparticles were centrifuged at 13 000 rpm for 15 minutes. The supernatant was removed and nanoparticles were re-dispersed in buffer containing 100 mM NaCl and 25 mM HEPES (pH 7.0) to prepare a 3 nM solution. Then DNAzyme 17E_{m1} or 17E_{m2} (3 μ M) and $\text{Pb}(\text{NO}_3)_2$ (4 μ M) were added to the solution. The mixture was incubated at room temperature for 10 hours, then centrifuged at 13 000 rpm for 15 minutes. The supernatant was removed and nanoparticles were re-dispersed in 1 mM HEPES (pH 7.0) without NaCl for 15 min to denature DNA hybridization. This centrifugation process was repeated once to remove free DNA in solution. At last, gold

nanoparticles were re-dispersed in 100 mM NaCl and 25 mM HEPES (pH 7.0) to prepare a 3 nM solution.

DNAzyme (E47)-catalyzed ligation of DNA on gold nanoparticles (O8)—The above solution was mixed with E47 (3 μ M), Cu(NO₃)₂ (20 μ M) and imidazole-activated **O3** (4 μ M) in 25 mM HEPES (pH 7.0) containing 150 mM NaCl. Then the mixture was incubation at room temperature for 16 hours. After that, nanoparticles were centrifuged at 13 000 rpm for 15 minutes. The supernatant was removed and nanoparticles were re-dispersed in 1 mM HEPES (pH 7.0) without NaCl for 15 min to denature DNA hybridization. This centrifugation process was repeated twice to remove free DNA in solution. Finally, nanoparticles were redispersed in 100 mM NaCl and 25 mM HEPES (pH 7.0) to a final concentration of 3 nM.

Formation of gold nanoparticle assemblies—DNA-functionalized gold nanoparticles were washed and redispersed in 100 mM NaCl and 25 mM HEPES (pH 7.0) to prepare 3 nM solutions as mentioned above. Equal volumes of **O6-**, **O7-** or **O8-** functionalized AuNPs (3 nM), **O9-** or **O10-** functionalized gold nanoparticles (3 nM), and 1 M NaCl were mixed. After vortexing, the resulting solution was incubated at room temperature for at least 1 hour to form nanoparticle assemblies.

TEM—Gold nanoparticles were analyzed using a JEOL 2010LaB6 transmission electron microscope (TEM) operated at 200 kV. Samples were prepared by putting a drop of nanoparticle solutions onto a carbon-coated copper TEM grid (Ted Pella).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We wish to thank the US National Institute of Health (ES016865) and Department of Energy (DE-FG02-08ER64568) for financial support.

Notes and references

1. Robertson DL, Joyce GF. Nature. 1990; 344:467–468. [PubMed: 1690861]
2. Breaker RR, Joyce GF. Chem. Biol. 1994; 1:223–229. [PubMed: 9383394]
3. Sen D, Geyer CR. Curr. Opin. Chem. Biol. 1998; 2:680–687. [PubMed: 9914188]
4. Li YF, Breaker RR. Curr. Opin. Struc. Biol. 1999; 9:315–323.
5. Lu Y. Chem. Eur. J. 2002; 8:4588–4596. [PubMed: 12362396]
6. Breaker RR. Nature. 2004; 432:838–845. [PubMed: 15602549]
7. Schlosser K, Li YF. Chem. Biol. 2009; 16:311–322. [PubMed: 19318212]
8. Silverman SK. Acc. Chem. Res. 2009; 42:1521–1531. [PubMed: 19572701]
9. Breaker RR, Joyce GF. Chem. Biol. 1995; 2:655–660. [PubMed: 9383471]
10. Carmi N, Balkhi SR, Breaker RR. Proc. Nat. Acad. Sci. U.S.A. 1998; 95:2233–2237.
11. Feldman AR, Sen D. J. Mol. Biol. 2001; 313:283–294. [PubMed: 11800557]
12. Liu JW, Brown AK, Meng XL, Cropek DM, Istok JD, Watson DB, Lu Y. Proc. Nat. Acad. Sci. U.S.A. 2007; 104:2056–2061.
13. Chandra M, Sachdeva A, Silverman SK. Nat. Chem. Biol. 2009; 5:718–720. [PubMed: 19684594]
14. Cuenoud B, Szostak JW. Nature. 1995; 375:611–614. [PubMed: 7791880]
15. Sreedhara A, Li YF, Breaker RR. J. Am. Chem. Soc. 2004; 126:3454–3460. [PubMed: 15025472]

16. Purtha WE, Coppins RL, Smalley MK, Silverman SK. *J. Am. Chem. Soc.* 2005; 127:13124–13125. [PubMed: 16173722]
17. Li YF, Breaker RR. *Proc. Nat. Acad. Sci. U.S.A.* 1999; 96:2746–2751.
18. Li YF, Liu Y, Breaker RR. *Biochemistry.* 2000; 39:3106–3114. [PubMed: 10715132]
19. Sheppard TL, Ordoukhanian P, Joyce GF. *Proc. Nat. Acad. Sci. U.S.A.* 2000; 97:7802–7807.
20. Li YF, Sen D. *Nat. Struct. Biol.* 1996; 3:743–747. [PubMed: 8784345]
21. Sen D, Poon LCH. *Crit. Rev. Biochem. Mol.* 2011; 46:478–492.
22. Chandra M, Silverman SK. *J. Am. Chem. Soc.* 2008; 130:2936–2937. [PubMed: 18271591]
23. Pradeepkumar PI, Hobartner C, Baum DA, Silverman SK. *Angew. Chem., Int. Ed.* 2008; 47:1753–1757.
24. Travascio P, Li YF, Sen D. *Chem. Biol.* 1998; 5:505–517. [PubMed: 9751647]
25. Poon LCH, Methot SP, Morabi-Pazooki W, Pio F, Bennet AJ, Sen D. *J. Am. Chem. Soc.* 2011; 133:1877–1884. [PubMed: 21265562]
26. Chinnapen DJF, Sen D. *Proc. Natl. Acad. Sci. U. S. A.* 2004; 101:65–69. [PubMed: 14691255]
27. Mirkin CA, Letsinger RL, Mucic RC, Storhoff JJ. *Nature.* 1996; 382:607–609. [PubMed: 8757129]
28. Liu JW, Lu Y. *J. Am. Chem. Soc.* 2003; 125:6642–6643. [PubMed: 12769568]
29. Li, Y.; Lu, Y. *Functional Nucleic Acids for Sensing and Other Analytical Applications.* New York: Springer; 2009.
30. Liu JW, Cao ZH, Lu Y. *Chem. Rev.* 2009; 109:1948–1998. [PubMed: 19301873]
31. Ali MM, Aguirre SD, Lazim H, Li YF. *Angew. Chem., Int. Ed.* 2011; 50:3751–3754.
32. Willner I, Shlyahovsky B, Zayats M, Willner B. *Chem. Soc. Rev.* 2008; 37:1153–1165. [PubMed: 18497928]
33. Chen Y, Wang MS, Mao CD. *Angew. Chem., Int. Ed.* 2004; 43:3554–3557.
34. Peracchi A. *Rev. Med. Virol.* 2004; 14:47–64. [PubMed: 14716691]
35. Pyle AM, Chu VT, Jankowsky E, Boudvillain H. *Method Enzymol.* 2000; 317:140–146.
36. Isildar M, Schuchmann MN, Schultefrohlinde D, Vonsonntag C. *Int. J. Radiat. Biol.* 1981; 40:347–354.
37. Eder PS, Walder RY, Walder JA. *Biochimie.* 1993; 75:123–126. [PubMed: 8389211]
38. Horton NC, Finzel BC. *J. Mol. Biol.* 1996; 264:521–533. [PubMed: 8969302]
39. Nakano S, Kanzaki T, Sugimoto N. *J. Am. Chem. Soc.* 2004; 126:1088–1095. [PubMed: 14746477]
40. Joyce CM. *Proc. Nat. Acad. Sci. U.S.A.* 1997; 94:1619–1622.
41. Scanlon KJ, Kashanisabet M, Miyachi H. *Cancer Invest.* 1989; 7:581–587. [PubMed: 2698255]
42. Rydberg B, Game J. *Proc. Nat. Acad. Sci. U.S.A.* 2002; 99:16654–16659.
43. Kim N, Huang SYN, Williams JS, Li YC, Clark AB, Cho JE, Kunkel TA, Pommier Y, Jinks-Robertson S. *Science.* 2011; 332:1561–1564. [PubMed: 21700875]
44. Park SJ, Taton TA, Mirkin CA. *Science.* 2002; 295:1503–1506. [PubMed: 11859188]
45. Rosi NL, Mirkin CA. *Chem. Rev.* 2005; 105:1547–1562. [PubMed: 15826019]
46. Thaxton CS, Georganopoulou DG, Mirkin CA. *Clin. Chim. Acta.* 2006; 363:120–126. [PubMed: 16214124]
47. Wang ZD, Lu Y. *J. Mater. Chem.* 2009; 19:1788–1798.
48. Wang H, Yang RH, Yang L, Tan WH. *Acs Nano.* 2009; 3:2451–2460. [PubMed: 19658387]
49. Xu W, Xue XJ, Li TH, Zeng HQ, Liu XG. *Angew. Chem., Int. Ed.* 2009; 48:6849–6852.
50. Wang F, Han Y, Lim CS, Lu YH, Wang J, Xu J, Chen HY, Zhang C, Hong MH, Liu XG. *Nature.* 2010; 463:1061–1065. [PubMed: 20182508]
51. Park SY, Lytton-Jean AKR, Lee B, Weigand S, Schatz GC, Mirkin CA. *Nature.* 2008; 451:553–556. [PubMed: 18235497]
52. Sharma J, Chhabra R, Cheng A, Brownell J, Liu Y, Yan H. *Science.* 2009; 323:112–116. [PubMed: 19119229]
53. Wang Y, Chen G, Yang MX, Silber G, Xing SX, Tan LH, Wang F, Feng YH, Liu XG, Li SZ, Chen HY. *Nat. Commun.* 2010; 1

54. Pinheiro AV, Han DR, Shih WM, Yan H. *Nat. Nanotechnol.* 2011; 6:763–772. [PubMed: 22056726]
55. Zhao Z, Jacovetty EL, Liu Y, Yan H. *Angew. Chem., Int. Ed.* 2011; 50:2041–2044.
56. Zhao WA, Lam JCF, Chiuman W, Brook MA, Li YF. *Small.* 2008; 4:810–816. [PubMed: 18537135]
57. Li J, Lu Y. *J. Am. Chem. Soc.* 2000; 122:10466–10467.
58. Okumoto Y, Tanabe Y, Sugimoto N. *Biochemistry.* 2003; 42:2158–2165. [PubMed: 12590605]
59. Nakano S, Karimata HT, Kitagawa Y, Sugimoto N. *J. Am. Chem. Soc.* 2009; 131:16881–16888. [PubMed: 19874030]
60. Liu JW, Lu Y. *Chem. Commun.* 2007:4872–4874.

Figure 1.

(a) Conversion of a single ribonucleotide (rC) in a DNA strand **O1** to a deoxyribonucleotide (C) by the cascade of DNAzymes $17E_{m1}$ and E47: **O1** is cleaved by $17E_{m1}$ to afford products of **O2** and **Oc**; **O2** is then ligated with **O3** (activated by imidazole) to form **O4** by E47. (b) Sequence modification of a DNA strand **O5** to **O4** through a similar protocol by the cascade of DNAzymes $17E_{m2}$ and E47.

Figure 2.

(a) Fluorescent PAGE (20% denaturing gel) images of the transformation from **O1** to **O4** by DNAzymes 17E_{m1} and E47. Lanes in (a): **1, O1**; **2, 1** after cleavage by 17E_{m1} to yield **O2** and **Oc** in the presence of Pb²⁺; **3, 2** after ligation to yield **O4** by E47 in the presence of **O3** and Cu²⁺; **4, 2** after ligation to yield **O4+8A** by E47 in the presence of **O3+8A** and Cu²⁺; **5, O4** in the presence of Pb²⁺ and 17E_{m1}. (b) Fluorescent PAGE images of the transformation from **O5** to **O4** by 17E_{m2} and E47: Lanes in (b): **1, O5**; **2, 1** after cleavage by 17E_{m2} to yield **O2** in the presence of Pb²⁺; **3, 2** after ligation to yield **O4** by E47 in the presence of **O3** and Cu²⁺; **4, 2** after ligation to yield **O4+8A** by E47 in the presence of **O3+8A** and Cu²⁺; **5, O4** in the presence of Pb²⁺ and 17E_{m2}.

Figure 3.

Assembly of two types of DNA-functionalized gold nanoparticles. If the sequences of the two DNAs are not complementary, the gold nanoparticles are in a “dispersed” state and exhibit a red color with a sharp absorption band peaked around 532 nm (left). In contrast, the assembly of the gold nanoparticles with complementary DNAs causes the formation of an “aggregated” state with a broad absorption band around 600 nm (right).

Figure 4.

Controlling the assembly of DNA-functionalized gold nanoparticles via cascade-mediated modification of the DNA sequences. The solid and dashed lines indicate the successful and unsuccessful formation of aggregates, respectively. The inset shows the assembly of AuNPs modified with the complementary strands **O6** and **O9** (top) and the lack of assembly between AuNPs modified with noncomplementary strands **O9** and **O7** (obtained by treating **O6**-functionalized AuNPs with 17E_{m2} and E47) (bottom).

Figure 5. Absorption spectra of **O6-**, **O7-** or **O8-**functionalized AuNPs in the presence of **O9-** and **O10-**functionalized AuNPs, respectively. The red-shift of the peak indicates the formation of AuNP aggregations due to the hybridization of complementary DNAs on the AuNPs.²⁷For the ratios of absorbance (A_{700}/A_{532}), see Table S1 in Supplementary Information.

Figure 6.

(a) Scheme showing stepwise modification of DNA sequences on multiply-functional gold nanoparticles by the collaboration of 17E_{m1} or 17E_{m2} and E47. (b) DNA sequences of O6~O8 and O11.

Figure 7. Absorption spectra of DNA-functionalized gold nanoparticles **A-F** (Figure 6a) in the presence of **O9**- and **O10**-functionalized AuNPs, respectively. The red-shift of the peak indicates the formation of AuNP aggregations due to the hybridization of complementary DNAs on the AuNPs.²⁷ For the ratios of absorbance (A_{700}/A_{532}), see Table S2 in Supplementary Information.

Table 1

Measured and calculated molecular weight (m/z) in MALDI-TOF mass spectra of 3'-fluorescein-labeled DNAs (**O1**, **O2**, **O4** and **O5**). For full spectra, see Figure S2 in Supplementary Information.

DNA	O1	O2	O4	O5
Measured	8831.6	4824.1	8812.3	8819.1
Calculated	8831.9	4826.3	8815.9	8821.9