Published in final edited form as:

Epidemiology. 2013 January ; 24(1): 122–128. doi:10.1097/EDE.0b013e318276c98a.

Hypothetical mid-life interventions in women and risk of type 2 diabetes

Goodarz Danaei^{1,2}, An Pan³, Frank B Hu^{2,3,4}, and Miguel A Hernán^{2,5}

¹Department of Global Health and Population, Harvard School of Public Health, Boston, MA

²Department of Epidemiology, Harvard School of Public Health, Boston, MA

³Department of Nutrition, Harvard School of Public Health, Boston, MA

⁴Channing Laboratory, Harvard Medical School and Brigham and Women's Hospital, Boston MA

⁵Harvard-MIT Division of Health Sciences and Technology, Cambridge, MA

Abstract

Background—Randomized trials have examined short-term effects of lifestyle interventions for diabetes prevention only among high-risk individuals. Prospective studies have examined the associations between lifestyle factors and diabetes in healthy populations but have not characterized the intervention. We estimated long-term effects of “hypothetical” lifestyle interventions on diabetes in a prospective study of healthy women, using the parametric g-formula.

Methods—Using data from the Nurses' Health Study, we followed 76,402 women from 1984 to 2008. We estimated the risk of type 2 diabetes under 8 “hypothetical” interventions: quitting smoking, losing weight by 5% every 2 years if overweight/obese, exercising at least 30 minutes a day, eating less than 3 servings a week of red meat, eating at least 2 servings a day of whole grain, drinking 2 or more cups of coffee a day, drinking 5 or more grams of alcohol a day and drinking less than 1 serving of soda a week.

Results—The 24-year risk of diabetes was 9.6% under no intervention and 4.3% when all interventions were imposed (55% lower risk [95% confidence interval= 47% to 63%]). The most effective interventions were weight loss (24% lower risk), physical activity (19%) and moderate alcohol use (19%). Overweight/obese women would benefit the most, with 10.8 percentage points reduction in 24-year risk of diabetes. The validity of these estimates relies on absence of unmeasured confounding, measurement error, and model misspecification.

Conclusion—A combination of dietary and non-dietary lifestyle modifications, begun in mid-life or later in relatively healthy women, could have prevented at least half of the cases of type 2 diabetes in this cohort of US women.

Address correspondence to: Goodarz Danaei, Assistant Professor of Global Health, Harvard School of Public Health, 665 Huntington Avenue, Building 1, 1107, Boston MA 02115 USA, gdanaei@hsph.harvard.edu, Tel: +1 617 432 5722, Fax: +1 617 432 6733.

Conflicts of Interest: Authors have no conflicts of interest to declare.

SDC Supplemental digital content is available through direct URL citations in the HTML and PDF versions of this article (www.epidem.com). This content is not peer-reviewed or copy-edited; it is the sole responsibility of the author.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Diabetes is a major cause of death and disability worldwide,^{1,2} and its prevalence has increased substantially in most regions of the world in the last three decades.³ Complications of diabetes put a major economic burden on health systems in both developed and developing countries.⁴⁻⁶ Primary prevention of type 2 diabetes, which constitutes more than 90% of cases, is a major concern for health systems worldwide.

Several randomized trials have examined the effect of physical activity, smoking cessation, weight loss and healthy diet on the incidence of type 2 diabetes in high-risk participants over relatively short periods of about 3 years.⁷⁻¹⁵ Overall, these randomized trials reported about 50% reduction in the incidence of type 2 diabetes after intensive lifestyle modifications.¹⁶⁻¹⁸ Prospective observational studies have mostly examined the long-term association between lifestyle and type 2 diabetes incidence¹⁹⁻²² in relatively healthy populations, but these studies did not specify the corresponding interventions or the time of their initiation.

Devising an informed policy for diabetes prevention requires estimating the effect of lifestyle interventions initiated in mid-life or later (as randomized trials did) over long periods and in relatively healthy populations (as observational studies did). We applied the parametric g-formula to data from the observational Nurses' Health Study to estimate the 24-year risk of type 2 diabetes under various hypothetical lifestyle interventions that start in mid-life or later.

Methods

Study population

The Nurses' Health Study is a prospective cohort study that started in 1976 by enrolling 121,700 US female nurses aged 30 to 55 years. A questionnaire was mailed to collect data on sociodemographic, lifestyle and dietary factors and on history of diseases and treatments. Biennial questionnaires have been mailed since then to update information on risk factors and disease incidence. More details on this cohort are available elsewhere.²³

We used the 1984 questionnaire as baseline, because a detailed 131-item food frequency questionnaire (FFQ) was distributed in that year. Women were excluded from our analysis if they had a diagnosis of diabetes before 1984 or did not have information on diabetes diagnosis date. In addition, we excluded participants who left more than 70 items blank on the baseline FFQ, or who reported unusual total energy intakes (i.e., energy intake <500 or >3500 kcal/day). We also excluded participants without baseline information on date of birth, body weight and height, smoking status, physical activity, and dietary variables, and those who had cancer or cardiovascular disease at baseline (see eFigure 1 for a flowchart of participant selection). After these exclusions, 76,402 women were available for the analysis. Women were followed until the occurrence of type 2 diabetes, death, incomplete follow-up (i.e. not returning a questionnaire), or administrative end of follow-up in June 2008, whichever happened first.

Diet was measured using a semi-quantitative FFQ in 1984, 1986 and every 4 years afterwards. The FFQ asked about the usual intake of various food items, including alcoholic drinks, during the past 12 months. The reproducibility and validity of the questionnaire have been shown elsewhere.²⁴ Dietary data recorded in the 1980 questionnaire was used to adjust for pre-baseline diet. Physical activity was reported in 1980, 1982, 1986, 1988 and every 4 years thereafter, using a validated questionnaire²⁵ on type, frequency and intensity of each activity. We summed the duration of moderate or vigorous activities per week (i.e. requiring at least 3 Metabolic Equivalent of Task [MET] scores/hour, including brisk walking). Body weight was self-reported in each biennial questionnaire and height was self-reported in 1976.

We truncated the values of dietary risk factors, body mass index (BMI) and physical activity in each period at the 99th percentile to prevent implausible values from affecting our analyses. Sensitivity analyses using various thresholds based on expert knowledge, or setting values above the 99th percentile to missing and carrying the last observed value forward, did not change the results materially.

Diabetes ascertainment

Details of diabetes ascertainment has been described elsewhere.²⁶ Briefly, a supplementary questionnaire was mailed to participants who reported a diagnosis of diabetes. A case of type 2 diabetes was considered confirmed if, according to the National Diabetes Data Group criteria,²⁷ at least one of the following was reported on the supplementary questionnaire: 1) one or more classic symptoms plus fasting plasma glucose levels of ≥ 7.8 mmol/L or random plasma glucose levels of ≥ 11.1 mmol/L; 2) at least two elevated plasma glucose concentrations on different occasions (fasting plasma glucose ≥ 7.8 mmol/L, random plasma glucose levels of ≥ 11.1 mmol/L, or concentrations of ≥ 11.1 mmol/L after two hours or more shown by oral glucose tolerance testing) in the absence of symptoms; or 3) treatment with insulin or oral hypoglycaemic agents. The diagnostic criteria changed in June 1998: according to the American Diabetes Association criteria,²⁸ a fasting plasma glucose of 7.0 mmol/L instead of 7.8 mmol/L was considered the threshold for the diagnosis of diabetes. Only confirmed cases were included in the analysis. We excluded cases designated as gestational or secondary diabetes. The validity of the supplementary questionnaire has been previously documented by reviewing medical records and assessing undiagnosed diabetes in a random sample of women.^{26,27,29} Deaths were identified by reports from next of kin or postal authorities, or by searching the National Death Index. At least 98% of deaths among the study participants were identified.³⁰

Hypothetical lifestyle interventions

We considered 8 hypothetical interventions and their combinations, based on the evidence from both randomized trials and observational studies on their potential effect for diabetes prevention:^{19,20,31-35} quitting smoking, losing 5% of BMI every 2 years if overweight or obese (defined as BMI ≥ 25 kg/m²), exercising at least 30 minutes a day (moderate or vigorous), eating at least 2 servings of whole grain per day, drinking at least 2 cups of coffee per day, drinking at least 5 grams of alcohol per day, eating at most 3 servings of red meat per week (including unprocessed and processed), and drinking at most 1 serving of soda per week. All interventions started at baseline in 1984 and continued until the end of follow-up. Except for the weight loss intervention, which imposed a gradual decline in body weight, all other interventions imposed a minimum or maximum threshold on the level of a risk factor. For these interventions, values beyond the threshold were set to the threshold level.

The intensities of the interventions were chosen to match current public health guidelines (e.g. Center of Disease Control and Prevention guidelines for physical activity, World Health Organization definition of overweight and obesity) or were selected to reflect feasible public health interventions. We estimated the effects of more intensive interventions for weight loss and physical activity in sensitivity analyses.

Statistical analysis

We used the parametric g-formula,³⁶ a generalization of standardization for time-varying exposures and confounders, to estimate the 24-year risks of type 2 diabetes under the selected lifestyle interventions. The parametric g-formula has been previously used to estimate the effect of multiple lifestyle interventions on the risk of coronary heart disease.³⁷ If all time-varying confounders have been correctly measured and modeled at all time-points, the g-formula can be used to consistently estimate the standardized risk of type 2

diabetes under hypothetical interventions. We used a Kaplan-Meier estimator that incorporates censoring by death and loss to follow-up.³⁸ For the exact formula see electronic appendix eText.

A simplified description of the process of estimating risks using parametric g-formula is as follows: we start by fitting regression models for all potential confounders and for the disease outcome using data on the entire study population. We then use these models to simulate the risk of the disease under various interventions in five steps: (1) take the observed joint distribution of covariates at baseline; (2) estimate the joint distribution of time-varying covariates at the next time-point using the parametric models; (3) “intervene” by setting the values of some covariates to values determined by the hypothetical interventions; (4) estimate the predicted probability of the outcome using these new values; (5) repeat steps 2 through 4 for the entire duration of follow-up to estimate the predicted risk of the disease under the selected interventions. See papers by Taubman et al³⁹ and Young et al⁴⁰ for a more detailed description.

More formally, the standardized cumulative risk estimated by the g-formula is a weighted average of the risks of type 2 diabetes conditional on the specified intervention values and the observed confounder history. The weights are the probability density functions of the time-varying confounders, which are estimated via parametric regression models. We approximated the weighted average by using a Monte Carlo simulation of 10,000 individuals with the baseline values of covariates sampled from their empirical distribution. The values of time-varying covariates for each 2-year interval were drawn from the distribution estimated via the regression models after setting the values of lifestyle factors to those specified by the interventions.

To increase comparability, the models included calendar year for each period and the following potential baseline confounders: age, history of diabetes in first-degree relatives, smoking and oral contraceptive use prior to 1980, marital status, education, husband’s education, employment, and stress in daily life and work, as well as pre-baseline values of the variables corresponding to the 8 selected intervention (i.e. BMI, smoking, exercise, and intake of meat, whole grain, coffee, alcohol and soda). These latter adjustments enabled us to estimate the effect of changes in lifestyle in middle-aged women as opposed to the life-long effect of a healthy lifestyle.

We modeled the distribution of 20 time-varying covariates: multivitamin use, aspirin use, statin use, post-menopausal hormone use, smoking, physical activity, soda intake, coffee intake, red meat intake, whole grain intake, alcohol use, BMI, high blood pressure, high serum cholesterol, coronary heart disease, stroke, angina or coronary artery bypass graft, cancer, menopause and osteoporosis (see eTable 1).

We compared the estimated risks under various interventions with the risk under no intervention to calculate the population risk ratio and population risk difference. The population attributable risk is one minus the population risk ratio. To estimate the 95% confidence intervals (CIs), we used non-parametric bootstrapping with 500 samples.⁴¹ We also computed the proportion of women who were hypothetically intervened on in any period and the average proportion of women intervened on in each 2-year period. The latter measures overall adherence in the observed data to the hypothetical intervention among those following the intervention up until the previous period. We examined the possibility of effect modification by conducting the analysis separately in subsets of the study population defined at baseline according to age, BMI, and family history of diabetes. All analyses were conducted using SAS 9.2 (Cary, NC). The SAS macro and its documentation are available at <http://www.hsph.harvard.edu/causal/software>.

Results

Table 1 shows the baseline characteristics of the 76,402 women who met all eligibility criteria in 1984. Mean age was 49.9 years and mean BMI was 24.4 kg/m²; 25% of women were current smokers and 19% had a family history of diabetes. During 24 years (1.4 million person-years) of follow-up, there were 6,044 cases of type 2 diabetes and 8,260 deaths. 17,690 participants did not reach the end of follow-up (i.e. did not return a questionnaire). The models collectively performed very well in estimating the risk-factor distributions under no intervention. For example, the mean difference between the observed and simulated number of cigarettes smoked per day was less than 0.4 cigarettes during follow-up; similarly, the point estimate of the ratio of the observed to simulated BMI was never smaller than 0.99 or larger than 1.00 (see eFigures 2-9). More importantly, the simulated 24-year risk of diabetes was the same as the observed risk at 9.6%, which is a necessary condition for no model misspecification. The coefficients of the models used in the simulations are presented in eTable 2.

Table 2 shows the 24-year risk of diabetes under various hypothetical lifestyle interventions. Among the non-dietary lifestyle interventions, weight loss was estimated to reduce the risk by 24% (95% CI= 22% to 26%), exercise by 19% (6% to 30%), and quitting smoking by 0% (-1% to 2%) compared with no lifestyle intervention. Among the selected dietary interventions, drinking at least 5 grams of alcohol a day was estimated to reduce the risk by 19% (12% to 23%), eating less than 3 servings of red meat per week by 8% (5% to 11%), and drinking at least 2 cups of coffee a day by 3% (0% to 6%). The mean alcohol intake under the hypothetical intervention “drink at least 5 grams of alcohol a day” ranged from 9.2 to 11.7 grams a day during follow-up, which is equivalent to 1 drink a day.

We estimated that the 24-year risk of type 2 diabetes would be reduced by 39% (29% to 47%) under the 3 non-dietary interventions, by 29% (21% to 35%) under the 5 dietary interventions, and by 55% (47% to 63%) under all 8 interventions. The estimated 24-year risk of type 2 diabetes under all interventions was 4.3% (3.6% to 5.1%). Of all participants, 25% maintained a BMI of 25 kg/m², fewer than 11% followed each of the dietary interventions, and 0% followed all 8 interventions for the whole duration of follow-up (9% followed all 8 interventions at some point during follow-up).

Table 3 presents the results of the analyses for more extreme weight-loss interventions. We estimated that reducing BMI at 5% per 2-year period down to 23 kg/m² would reduce the risk of diabetes by 53% (51% to 56%), while reducing BMI at a faster pace of 10% every 2 years would reduce the risk by 60% (57% to 63%). Combining this latter intervention with the 7 other lifestyle interventions would reduce the risk of type 2 diabetes by 72%. We estimated a risk reduction of 16% (8% to 22%) under a hypothetical intervention in which participants exercise for 30 minutes/day if they have a normal BMI (i.e., BMI < 25 kg/m²) and 1 hour/day if they are overweight or obese (i.e., BMI ≥ 25 kg/m²). A more intensive intervention to engage in at least 1 hour/day of moderate or vigorous activity, regardless of BMI, would reduce the risk of type 2 diabetes by 15% (7% to 22%).

To evaluate the sensitivity of the estimates to our analytic decisions, we conducted analyses that varied the ordering of the variables measured in each questionnaire, estimated censoring due to incomplete follow-up, considered a different intervention on alcohol where women would drink 5 to 10 grams/day of alcohol, and included women who did not return 1 or 2 questionnaires by carrying their last reported values forward. The estimates of relative risks and risk differences did not change materially.

The effects of lifestyle interventions were stronger in younger women (< 50 years old) and those who were overweight or obese at baseline, in both the risk ratio and risk difference

scales (Table 4). For all 8 interventions combined, the 24-year risk of type 2 diabetes would be reduced by 10.8 percentage points in women who were overweight or obese compared with only 1.9 percentage points in women who had a BMI of $< 25 \text{ kg/m}^2$ at baseline (Table 4). Women with a family history of diabetes were also estimated to benefit more from these interventions: the risk difference for all 8 interventions in this group was 7.3% versus 4.8% in those without a family history of diabetes.

Discussion

Our results suggest that, in this cohort of US women, 55% of cases of type 2 diabetes could have been prevented by a combination of dietary and non-dietary lifestyle modifications. Our estimates are particularly relevant for health policy because they quantify the 24-year impact of lifestyle interventions that start in mid-life or later in relatively healthy women.

A beneficial effect of lifestyle modification on diabetes risk had been previously found in several randomized trials. However, these studies were designed to evaluate a short-term effect (over approximately a 3-year period) in overweight participants with impaired glucose tolerance. Both the Diabetes Prevention Program⁷ and the Diabetes Prevention Study⁸ found that a combined intervention on diet and physical activity reduced diabetes risk by 58% compared with general guidance or written advice. The risk of diabetes in the control group was 29% at 3 years in the Diabetes Prevention Program and 23% at 4 years in the Diabetes Prevention Study, which are much higher than our 9.6% risk at 24 years in a population of healthy US women.

Our estimates may not be generalizable to other populations with different distributions of risk factors, as the g-formula standardizes the risk to the distribution of risk factors in the particular population under study. For example, we estimated no reduction in risk of type 2 diabetes if all women had quit smoking but only 25% of women in our population were smokers. When we compared the risk had everyone been “forced” to smoke 20 cigarettes a day to the risk had everyone quit smoking, the estimated population risk ratio was 1.1. Also, the magnitude of our estimates is specific to the set of interventions that was considered. Though our results are generally consistent with previous analyses of prospective studies,^{34,35,42,43} differences do exist because previous studies did not specify the time of initiation of the lifestyle interventions and considered more extreme weight loss.¹⁹⁻²² For example, a previous analysis of the Nurses’ Health Study cohort classified women as low-risk if they met 5 criteria (BMI $< 25 \text{ kg/m}^2$, diet high in cereal fiber and polyunsaturated fat and low in trans-fat and glycemic load, at least half an hour/day of moderate-to-vigorous physical activity, no current smoking, at least a half-serving of an alcoholic beverage/day) between 1986 and 1996. Compared with the rest of the cohort, women in the low-risk group had an average two-year relative risk of diabetes of 0.09 (95% CI= 0.05 to 0.17). The authors estimated that some hypothetical intervention (different from the ones we considered here) on the above 5 factors could have prevented 91% of diabetes cases.

Our analysis has several strengths. The Nurses’ Health Study cohort has collected detailed data on usual dietary intakes, physical activity, body weight and smoking every 2-4 years, and diabetes diagnosis was validated. We included 24 years of follow-up and had a large number of cases, which allowed meaningful subgroup analyses. By applying the parametric g-formula^{39,44} we could estimate the effect of hypothetical interventions starting in middle-age or later while (i) appropriately adjusting for time-varying confounders affected by prior exposure, (ii) generating adjusted estimates of absolute risk and population attributable risk, and (iii) estimating effects of interventions individually and in various combinations.

Like other analyses of observational data, the validity of our results relies on the key assumptions of no residual confounding, no measurement error, and no model misspecification. The possibility of residual confounding cannot be logically excluded despite adjustment for many potential confounders. A certain degree of measurement error is expected for lifestyle variables and may have contributed to bias. We had to rely on self-reported weight and height; a validation study on a small sample of the study population showed a 0.96 correlation between self-reported and measured weight.⁴⁵ Though the prediction of our models under the hypothetical interventions cannot be directly evaluated (0% of women followed all interventions during the entire follow-up), our models provided accurate predictions under no intervention, a necessary condition for no model misspecification.

We tried to simplify the public health interpretation of our estimates by comparing interventions with well-defined start times and exposure values, by using food items rather than dietary scores¹⁹ or the Healthy Eating Index,⁴⁶ and by including total caloric intake in the models so that our hypothetical interventions imply replacing the selected food item with other foods that are usually taken as a substitute in the study population. However, the interpretation of our estimates for weight loss and exercise is complicated because there are multiple versions of the interventions. For example, participants may lose weight by reducing their caloric intake or by using weight loss medications; similarly, the selected duration and level of exercise may be achieved by increasing frequency or duration of different types of activities.⁴⁷

In summary, our results suggest that 55% of cases of diabetes that occurred during 24 years of follow-up in this large prospective study of US women could have been prevented by 8 lifestyle interventions initiated in mid-life or later. The most effective intervention in this population was losing weight, followed by eating a healthy diet and engaging in regular moderate or vigorous physical activity.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We are grateful to Roger Logan and Jessica Young for their technical assistance, to Walter Willett and Meir Stampfer for their comments on a previous draft, and to the Channing Laboratory, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, and to all the women enrolled in the Nurses' Health Study.

Source of Funding: This work was funded through National Institute of Health (NIH) grants HL080644, DK58845, CA87969.

References

1. Lopez AD, Mathers CD, Ezzati M, Jamison DT, Murray CJ. Global and regional burden of disease and risk factors, 2001: systematic analysis of population health data. *Lancet*. 2006; 367(9524): 1747–1757. [PubMed: 16731270]
2. Danaei G, Lawes CM, Vander HS, Murray CJ, Ezzati M. Global and regional mortality from ischaemic heart disease and stroke attributable to higher-than-optimum blood glucose concentration: comparative risk assessment. *Lancet*. 2006; 368(9548):1651–1659. [PubMed: 17098083]
3. Danaei G, Finucane MM, Lu Y, Singh GM, Cowan MJ, Paciorek CJ, Lin JK, Farzadfar F, Khang YH, Stevens GA, Rao M, Ali MK, Riley LM, Robinson CA, Ezzati M. National, regional, and global trends in fasting plasma glucose and diabetes prevalence since 1980: systematic analysis of

- health examination surveys and epidemiological studies with 370 country-years and 2.7 million participants. *Lancet*. 2011; 378(9785):31–40. [PubMed: 21705069]
4. Abegunde DO, Mathers CD, Adam T, Ortegon M, Strong K. The burden and costs of chronic diseases in low-income and middle-income countries. *Lancet*. 2007; 370(9603):1929–38. [PubMed: 18063029]
 5. Bjork S, Kapur A, King H, Nair J, Ramachandran A. Global policy: aspects of diabetes in India. *Health Policy*. 2003; 66(1):61–72. [PubMed: 14499166]
 6. MacLeod KM, Tooke JE. Direct and indirect costs of cardiovascular and cerebrovascular complications of type II diabetes. *Pharmacoeconomics*. 1995; 8(Suppl 1):46–51. [PubMed: 10159003]
 7. Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, Nathan DM. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *The New England Journal of Medicine*. 2002; 346(6):393–403. [PubMed: 11832527]
 8. Tuomilehto J, Lindstrom J, Eriksson JG, Valle TT, Hamalainen H, Ilanne-Parikka P, Keinanen-Kiukkaanniemi S, Laakso M, Louheranta A, Rastas M, Salminen V, Uusitupa M. Prevention of type 2 diabetes mellitus by changes in lifestyle among subjects with impaired glucose tolerance. *N Engl J Med*. 2001; 344(18):1343–1350. [PubMed: 11333990]
 9. Pan XR, Li GW, Hu YH, Wang JX, Yang WY, An ZX, Hu ZX, Lin J, Xiao JZ, Cao HB, Liu PA, Jiang XG, Jiang YY, Wang JP, Zheng H, Zhang H, Bennett PH, Howard BV. Effects of diet and exercise in preventing NIDDM in people with impaired glucose tolerance. The Da Qing IGT and Diabetes Study. *Diabetes Care*. 1997; 20(4):537–544. [PubMed: 9096977]
 10. Dyson PA, Hammersley MS, Morris RJ, Holman RR, Turner RC. The Fasting Hyperglycaemia Study: II. Randomized controlled trial of reinforced healthy-living advice in subjects with increased but not diabetic fasting plasma glucose. *Metabolism*. 1997; 46(12 Suppl 1):50–5. [PubMed: 9439560]
 11. Kosaka K, Noda M, Kuzuya T. Prevention of type 2 diabetes by lifestyle intervention: a Japanese trial in IGT males. *Diabetes Res Clin Pract*. 2005; 67(2):152–62. [PubMed: 15649575]
 12. Ramachandran A, Snehalatha C, Mary S, Mukesh B, Bhaskar AD, Vijay V. The Indian Diabetes Prevention Programme shows that lifestyle modification and metformin prevent type 2 diabetes in Asian Indian subjects with impaired glucose tolerance (IDPP-1). *Diabetologia*. 2006; 49(2):289–97. [PubMed: 16391903]
 13. Bo S, Ciccone G, Baldi C, Benini L, Dusio F, Forastiere G, Lucia C, Nuti C, Durazzo M, Cassader M, Gentile L, Pagano G. Effectiveness of a lifestyle intervention on metabolic syndrome. A randomized controlled trial. *J Gen Intern Med*. 2007; 22(12):1695–703. [PubMed: 17922167]
 14. Roumen C, Corpeleijn E, Feskens EJ, Mensink M, Saris WH, Blaak EE. Impact of 3-year lifestyle intervention on postprandial glucose metabolism: the SLIM study. *Diabet Med*. 2008; 25(5):597–605. [PubMed: 18445174]
 15. Lindahl B, Nilsson TK, Borch-Johnsen K, Roder ME, Soderberg S, Widman L, Johnson O, Hallmans G, Jansson JH. A randomized lifestyle intervention with 5-year follow-up in subjects with impaired glucose tolerance: pronounced short-term impact but long-term adherence problems. *Scand J Public Health*. 2009; 37(4):434–42. [PubMed: 19181821]
 16. Yamaoka K, Tango T. Efficacy of lifestyle education to prevent type 2 diabetes: a meta-analysis of randomized controlled trials. *Diabetes Care*. 2005; 28(11):2780–6. [PubMed: 16249558]
 17. Orozco LJ, Buchleitner AM, Gimenez-Perez G, Roque IFM, Richter B, Mauricio D. Exercise or exercise and diet for preventing type 2 diabetes mellitus. *Cochrane Database Syst Rev*. 2008; (3):CD003054. [PubMed: 18646086]
 18. Gillies CL, Abrams KR, Lambert PC, Cooper NJ, Sutton AJ, Hsu RT, Khunti K. Pharmacological and lifestyle interventions to prevent or delay type 2 diabetes in people with impaired glucose tolerance: systematic review and meta-analysis. *BMJ*. 2007; 334(7588):299. [PubMed: 17237299]
 19. Hu FB, Manson JE, Stampfer MJ, Colditz G, Liu S, Solomon CG, Willett WC. Diet, lifestyle, and the risk of type 2 diabetes mellitus in women. *N Engl J Med*. 2001; 345(11):790–797. [PubMed: 11556298]

20. Mozaffarian D, Kamineni A, Carnethon M, Djousse L, Mukamal KJ, Siscovick D. Lifestyle risk factors and new-onset diabetes mellitus in older adults: the cardiovascular health study. *Arch Intern Med.* 2009; 169(8):798–807. [PubMed: 19398692]
21. Laaksonen MA, Knekt P, Rissanen H, Harkanen T, Virtala E, Marniemi J, Aromaa A, Heliovaara M, Reunanen A. The relative importance of modifiable potential risk factors of type 2 diabetes: a meta-analysis of two cohorts. *Eur J Epidemiol.* 2009; 25(2):115–24. [PubMed: 20012885]
22. Steinbrecher A, Morimoto Y, Heak S, Ollberding NJ, Geller KS, Grandinetti A, Kolonel LN, Maskarinec G. The preventable proportion of type 2 diabetes by ethnicity: the multiethnic cohort. *Ann Epidemiol.* 2011; 21(7):526–35. [PubMed: 21497517]
23. Colditz GA, Manson JE, Hankinson SE. The Nurses' Health Study: 20-year contribution to the understanding of health among women. *J Womens Health.* 1997; 6(1):49–62. [PubMed: 9065374]
24. Willett WC, Sampson L, Stampfer MJ, Rosner B, Bain C, Witschi J, Hennekens CH, Speizer FE. Reproducibility and validity of a semiquantitative food frequency questionnaire. *Am J Epidemiol.* 1985; 122(1):51–65. [PubMed: 4014201]
25. Hu FB, Sigal RJ, Rich-Edwards JW, Colditz GA, Solomon CG, Willett WC, Speizer FE, Manson JE. Walking compared with vigorous physical activity and risk of type 2 diabetes in women: a prospective study. *JAMA.* 1999; 282(15):1433–1439. [PubMed: 10535433]
26. Manson JE, Rimm EB, Stampfer MJ, Colditz GA, Willett WC, Krolewski AS, Rosner B, Hennekens CH, Speizer FE. Physical activity and incidence of non-insulin-dependent diabetes mellitus in women. *Lancet.* 1991; 338(8770):774–778. [PubMed: 1681160]
27. Field AE, Coakley EH, Must A, Spadano JL, Laird N, Dietz WH, Rimm E, Colditz GA. Impact of overweight on the risk of developing common chronic diseases during a 10-year period. *Arch Intern Med.* 2001; 161(13):1581–6. [PubMed: 11434789]
28. Report of the expert committee on the diagnosis and classification of diabetes mellitus. *Diabetes Care.* 1997; 20(7):1183–1197. [PubMed: 9203460]
29. Hu FB, Leitzmann MF, Stampfer MJ, Colditz GA, Willett WC, Rimm EB. Physical activity and television watching in relation to risk for type 2 diabetes mellitus in men. *Arch Intern Med.* 2001; 161(12):1542–1548. [PubMed: 11427103]
30. Dellavalle RP, Drake A, Graber M, Heilig LF, Hester EJ, Johnson KR, McNealy K, Schilling L. Statins and fibrates for preventing melanoma. *Cochrane Database Syst Rev.* 2005; (4):CD003697. [PubMed: 16235336]
31. Knowler WC, Fowler SE, Hamman RF, Christophi CA, Hoffman HJ, Brenneman AT, Brown-Friday JO, Goldberg R, Venditti E, Nathan DM. 10-year follow-up of diabetes incidence and weight loss in the Diabetes Prevention Program Outcomes Study. *Lancet.* 2009; 374(9702):1677–86. [PubMed: 19878986]
32. Lindstrom J, Louheranta A, Mannelin M, Rastas M, Salminen V, Eriksson J, Uusitupa M, Tuomilehto J. The Finnish Diabetes Prevention Study (DPS): Lifestyle intervention and 3-year results on diet and physical activity. *Diabetes Care.* 2003; 26(12):3230–6. [PubMed: 14633807]
33. Malik VS, Popkin BM, Bray GA, Despres JP, Willett WC, Hu FB. Sugar-sweetened beverages and risk of metabolic syndrome and type 2 diabetes: a meta-analysis. *Diabetes Care.* 33(11):2477–83. [PubMed: 20693348]
34. Huxley R, Lee CM, Barzi F, Timmermeister L, Czernichow S, Perkovic V, Grobbee DE, Batty D, Woodward M. Coffee, decaffeinated coffee, and tea consumption in relation to incident type 2 diabetes mellitus: a systematic review with meta-analysis. *Arch Intern Med.* 2009; 169(22):2053–63. [PubMed: 20008687]
35. Freeman SR, Drake AL, Heilig LF, Graber M, McNealy K, Schilling LM, Dellavalle RP. Statins, fibrates, and melanoma risk: a systematic review and meta-analysis. *J Natl Cancer Inst.* 2006; 98(21):1538–46. [PubMed: 17077356]
36. Robins, JM.; Hernán, MA. Estimation of the causal effects of time-varying exposures. In: Fitzmaurice, G.; Davidian, M.; Verbeke, G.; Monenberghs, G., editors. *Longitudinal data analysis. Handbooks of modern statistical methods.* Chapman & Hall/CRC; Boca Raton: 2009. p. 553-599.
37. Taubman SL, Robins JM, Mittleman MA, Hernan MA. Intervening on risk factors for coronary heart disease: an application of the parametric g-formula. *Int J Epidemiol.* 2009; 38(6):1599–611. [PubMed: 19389875]

38. Gooley TA, Leisenring W, Crowley J, Storer BE. Estimation of failure probabilities in the presence of competing risks: new representations of old estimators. *Stat Med.* 1999; 18(6):695–706. [PubMed: 10204198]
39. Taubman SL, Robins JM, Mittleman MA, Hernán MA. Intervening on risk factors for coronary heart disease: an application of the parametric g-formula. *Int J Epidemiol.* 2009
40. Young JG, Cain LE, Robins JM, O'Reilly E, Hernán MA. Comparative effectiveness of dynamic treatment regimes: an application of the parametric g-formula. *Statistics in Biosciences.* 2011; 3:119–143.
41. Efron B, Tibshirani R. Bootstrap Methods for Standard Errors, Confidence Intervals, and Other Measures of Statistical Accuracy. *Statistical Science.* 1986; 1(1):54–75.
42. de Koning L, Malik VS, Rimm EB, Willett WC, Hu FB. Sugar-sweetened and artificially sweetened beverage consumption and risk of type 2 diabetes in men. *Am J Clin Nutr.* 2011; 93(6): 1321–7. [PubMed: 21430119]
43. Aune D, Ursin G, Veierod MB. Meat consumption and the risk of type 2 diabetes: a systematic review and meta-analysis of cohort studies. *Diabetologia.* 2009; 52(11):2277–87. [PubMed: 19662376]
44. Robins JM. A new approach to causal inference in mortality studies with a sustained exposure period-application to control of the healthy worker survivor effect. *Mathematical modelling.* 1986; 7:1393–1512.
45. Willett W, Stampfer MJ, Bain C, Lipnick R, Speizer FE, Rosner B, Cramer D, Hennekens CH. Cigarette smoking, relative weight, and menopause. *Am J Epidemiol.* 1983; 117(6):651–8. [PubMed: 6859020]
46. Fung TT, McCullough M, van Dam RM, Hu FB. A prospective study of overall diet quality and risk of type 2 diabetes in women. *Diabetes Care.* 2007; 30(7):1753–7. [PubMed: 17429059]
47. Hernán MA, VanderWeele TJ. Compound treatments and transportability of causal inference. *Epidemiology.* 2011; 22(3):368–77. [PubMed: 21399502]

Table 1Baseline characteristics^a of the eligible 76,402 women, Nurses' Health Study 1984

Characteristic	
Age at baseline (years); mean (SD)	49.9 (7.2)
Married	91.4
College education	25.6
Husband's education:	
not married or husband less than high-school education	29.1
High-school	32.2
College	21.0
Graduate school	17.8
Employment:	
Nursing education	13.5
Operating room nurse	4.0
Room nurse	2.7
In-patient nurse	20.1
Out-patient nurse	7.4
Other nursing	26.4
Non-nursing	6.3
Homemaker	19.8
High stress in work or life	67.3
Family history of diabetes	18.6
Oral contraceptive use in 1980	50.4
Menopause	45.4
Postmenopausal hormones	12.0
Diet	
Red meat (servings/day); mean (SD)	1.2 (0.8)
Coffee intake (cup/day); mean (SD)	2.4 (1.8)
Whole grain intake (serving/day); mean (SD)	1.2 (1.2)
Soda (serving/day); mean (SD)	0.9 (1.0)
Alcohol (gram/day); mean (SD)	6.8 (10.4)
Calories per day (Kcal); mean (SD)	1718 (530)
Cigarette smoking	
Smoking history in 1980	56.2
Current smoker	25.0
Cigarettes per day (among smokers); mean (SD)	20 (11)
Multivitamin use	37.5
Activity (hours/week of moderate or vigorous); mean (SD)	2.3 (2.2)
BMI (kg/m ²); mean (SD)	
At age 18	21.3 (2.8)
At baseline	24.4 (1.2)
High cholesterol	7.4

Characteristic	
High blood pressure	19.6
Aspirin use	
Less than daily	57.0
Daily or more	9.3
Osteoporosis	2.6

^a% unless otherwise indicated.

Table 2
Diabetes risk under hypothetical lifestyle interventions, Nurses' Health Study, 1984-2008

No.	Intervention	24-year risk of diabetes (%) ^a (95% CI)	Population risk ratio ^b (95% CI)	Population risk difference (%) ^b (95% CI)	Cumulative percent intervened on	Average percent intervened on
0	Natural course, no intervention ^c	9.6 (9.0 to 10.2)	1.00	0	0	0
1	Quit smoking	9.6 (9.0 to 10.1)	1.00 (0.98 to 1.01)	0.01 (-0.1 to 0.2)	28	3
2	Exercise at least 30 min/day	7.8 (6.7 to 9.1)	0.81 (0.70 to 0.94)	1.8 (0.6 to 2.8)	100	71
3	Lose 5% of BMI if BMI >25 kg/m ²	7.3 (6.8 to 7.8)	0.76 (0.74 to 0.78)	2.3 (2.1 to 2.5)	75	57
4	Eat at most 3 servings/week red or processed meat	8.8 (8.2 to 9.4)	0.92 (0.89 to 0.95)	0.8 (0.5 to 1.0)	98	56
5	Drink at least 2 cups/day coffee	9.3 (8.7 to 9.8)	0.97 (0.94 to 1.00)	0.3 (0.0 to 0.6)	96	51
6	Eat at least 2 servings/day whole grain	9.6 (8.9 to 10.2)	0.99 (0.96 to 1.04)	0.05 (-0.3 to 0.4)	99	59
7	Drink at most 1 serving/week soda	9.5 (8.7 to 10.0)	0.98 (0.94 to 1.02)	0.2 (-0.2 to 0.6)	95	55
8	Drink at least 5 grams/day of alcohol	7.8 (7.2 to 8.6)	0.81 (0.77 to 0.88)	1.8 (1.2 to 2.3)	90	47
9	All lifestyle interventions (numbers 1-3)	5.9 (4.9 to 6.9)	0.61 (0.53 to 0.71)	3.7 (2.8 to 4.6)	100	81
10	All dietary interventions (numbers 4-8)	6.9 (6.1 to 7.7)	0.71 (0.65 to 0.79)	2.8 (2.0 to 3.4)	100	97
11	All interventions (numbers 1-8)	4.3 (3.6 to 5.1)	0.45 (0.37 to 0.53)	5.3 (4.4 to 6.1)	100	91

^aThere were 6,044 cases of diabetes among 76,402 women after 1,391,216 person-years of follow-up. The observed risk was 9.6%.

^bIn addition to the 8 intervention variables, we modeled 13 other covariates in the analysis: calendar year, multivitamin use, aspirin use, statin use, postmenopausal hormone use, high blood pressure, high serum cholesterol, stroke, myocardial infarction, cancer, menopause and osteoporosis (see Table S1 for more detail on covariates and models). All models included lagged values of time-varying covariates plus baseline non-time varying variables: age, history of diabetes in first-degree relatives, marital status, education, husband's education, employment, stress in daily life and work, smoking and oral contraceptive use prior to 1980.

^cReference category.

Table 3
Effect of hypothetical weight-loss interventions on diabetes incidence in the Nurses' Health Study 1984-2008

No.	Intervention	24-year risk of diabetes (%) ^a (95% CI)	Population risk ratio ^b (95% CI)	Population risk difference (%) ^b (95% CI)	Cumulative percent intervened on	Average percent intervened on
0	Natural course, no intervention ^c	9.6 (9.0 to 10.2)	1.00	0	0	0
1	Maintain BMI < 25 kg/m ²	6.7 (6.3 to 7.2)	0.70 (0.68 to 0.72)	2.9 (2.6 to 3.1)	75	52
2	Lose 5% of BMI if BMI > 25 kg/m ²	7.3 (6.8 to 7.8)	0.76 (0.74 to 0.78)	2.3 (2.1 to 2.5)	75	57
3	Lose 5% of BMI if BMI > 23 kg/m ²	4.5 (4.1 to 4.8)	0.47 (0.44 to 0.49)	5.1 (4.7 to 5.5)	91	64
4	Lose 10% of BMI if BMI > 23 kg/m ²	3.8 (3.5 to 4.2)	0.40 (0.37 to 0.43)	5.8 (5.3 to 6.2)	91	58

^aThere were 6,044 cases of diabetes among 76,402 women after 1,391,216 person-years of follow-up. The observed risk was 9.6%.

^bIn addition to the 8 intervention variables, we modeled 13 other covariates in the analysis: calendar year, multivitamin use, aspirin use, statin use, postmenopausal hormone use, high blood pressure, high serum cholesterol, stroke, myocardial infarction, cancer, menopause and osteoporosis (see Table S1 for more detail on covariates and models). All models included lagged values of time-varying covariates plus baseline non-time varying variables: age, history of diabetes in first-degree relatives, marital status, education, husband's education, employment, stress in daily life and work, smoking and oral contraceptive use prior to 1980.

^cReference category.

Table 4

Effect of joint hypothetical interventions in sub-groups defined at baseline by (A) body mass index, (B) age and (C) family history of diabetes *

Intervention	Population risk ratio (95% CI)		Population risk difference (95% CI)		Test for heterogeneity
(A)	BMI at baseline	Test for heterogeneity	BMI at baseline	Test for heterogeneity	
	$<25^a$	25^b	$<25^a$	25^b	
Quit smoking, lose weight if BMI 25, exercise 30 min/day	0.76 (0.60 to 0.99)	0.56 (0.46 to 0.68)	1.1 (0.1 to 1.7)	8.1 (5.6 to 9.9)	P<0.001
Dietary interventions ^c	0.70 (0.60 to 0.83)	0.71 (0.62 to 0.80)	1.3 (0.7 to 1.9)	5.3 (3.6 to 6.8)	P<0.001
All interventions	0.56 (0.43 to 0.71)	0.41 (0.32 to 0.52)	1.9 (1.2 to 2.6)	10.8 (8.8 to 12.4)	P<0.001
(B)	Age at baseline		Age at baseline		
	$<50^d$	50^e	$<50^d$	50^e	
Quit smoking, lose weight if BMI 25, exercise 30 min/day	0.49 (0.40 to 0.66)	0.74 (0.58 to 0.91)	4.6 (3.4 to 5.8)	2.6 (0.8 to 4.0)	P=0.06
Dietary interventions ^c	0.65 (0.56 to 0.78)	0.77 (0.67 to 0.87)	3.1 (2.1 to 4.1)	2.3 (1.3 to 3.2)	P=0.27
All interventions	0.33 (0.26 to 0.50)	0.57 (0.43 to 0.73)	5.9 (4.9 to 7.0)	4.3 (2.6 to 5.6)	P=0.07
(C)	Family history		Family history		
	No ^f	Yes ^g	No ^f	Yes ^g	
Quit smoking, lose weight if BMI 25, exercise 30 min/day	0.56 (0.46 to 0.68)	0.70 (0.55 to 0.89)	3.5 (2.5 to 4.4)	4.9 (1.8 to 7.5)	P=0.36
Dietary interventions ^c	0.69 (0.62 to 0.79)	0.76 (0.66 to 0.92)	2.5 (1.7 to 3.0)	4.0 (1.4 to 5.8)	P=0.19
All interventions	0.41 (0.32 to 0.51)	0.53 (0.41 to 0.72)	4.8 (3.9 to 5.4)	7.3 (4.7 to 9.9)	P=0.07

* In addition to the 8 intervention variables, we modeled 13 other covariates in these analyses: calendar year, multivitamin use, aspirin use, statin use, post-menopausal hormone use, high blood pressure, high serum cholesterol, stroke, myocardial infarction, cancer, menopause and osteoporosis (see Table S1 for more detail on covariates and models). All models included lagged values of time-varying covariates plus baseline non-time varying variables: age, history of diabetes in first-degree relatives, marital status, education, husband's education, employment, stress in daily life and work, smoking and oral contraceptive use prior to 1980.

^aThe 24-year risk of type 2 diabetes under no intervention was 4.4% (95% CI 3.5, 5.2).

\$watermark-text

\$watermark-text

\$watermark-text

- ^bThe 24-year risk of type 2 diabetes under no intervention was 18.3% (95% CI 17.0, 19.2).
- ^cEat <3 servings/week of red or processed meat and 2 servings/day of whole grain; drink 2 cups/day of coffee and 5 grams/day of alcohol and <1 serving/week of soda
- ^dThe 24-year risk of type 2 diabetes under no intervention was 8.9% (95% CI 8.2, 11.5).
- ^eThe 24-year risk of type 2 diabetes under no intervention was 10.0% (95% CI 9.1, 11.6).
- ^fThe 24-year risk of type 2 diabetes under no intervention was 8.1% (95% CI 7.4, 8.6).
- ^gThe 24-year risk of type 2 diabetes under no intervention was 16.6% (95% CI 15.3, 18.0).