Published in final edited form as:

Am J Primatol. 2013 February ; 75(2): 107–115. doi:10.1002/ajp.22089.

The association of intergroup encounters, dominance status, and fecal androgen and glucocorticoid profiles in wild male white-faced capuchins (*Cebus capucinus*)

Valérie A. M. Schoof and Katharine M. Jack

Department of Anthropology, Tulane University, New Orleans, LA 70118

Abstract

Androgens play a role in male reproductive competition, frequently via aggression, while glucocorticoids are associated with the stress response. However, the relationships of these hormones with different sources of competition (intra-versus intergroup) and dominance status are highly variable. Here we consider the fecal androgen (fA) and glucocorticoid (fGC) profiles of alpha and subordinate male *Cebus capucinus* in the context of intergroup competition during a rare period of low intragroup competition (i.e. all females were either pregnant or lactating). Intergroup encounters (IGEs) are a long-term reproductive strategy in male white-faced capuchins, enabling them to assess the composition of neighboring groups. IGEs pose a threat to resident males as these can result in injury or death, loss of dominance rank, group eviction, and group takeovers that are frequently associated with infanticide. From February to July 2007, fecal samples were collected from eight males in three groups of white-faced capuchins in the Santa Rosa Sector of the Área de Conservación Guanacaste, Costa Rica. IGE rate was positively associated with both fA and fGC levels, indicating that IGEs are perceived as reproductive challenges by resident males, and may be associated with elevated metabolic costs. Alpha males sire the majority of group offspring and, accordingly, the threat of IGEs to both future (via rank loss or eviction) and current (via infanticide) reproductive success is greater than for subordinate males. Consistent with this observation, alpha males had higher fA and fGC levels than subordinate males. Given that all females were either pregnant or lactating and pronounced overt intragroup competition was absent, we interpret the difference in hormone profiles of alpha and subordinate males as being primarily associated with variation in the perceived threats of IGEs to dominant versus subordinate males. Future studies should focus on the interaction of intra- and intergroup competition by examining hormone levels in the presence of periovulatory females.

Keywords

Endocrinology; reproductive competition; infanticide; dominance; reproductive skew

INTRODUCTION

Androgens, including testosterone, are a group of steroid hormones that facilitate reproductive competition [Wingfield et al., 1990], maintain reproductive physiology, develop secondary sexual characteristics, and inhibit paternal behavior in males [reviewed in Fernandez-Duque et al., 2009]. Androgens also have deleterious effects such as immunosuppression and compromised survivability [reviewed in Muehlenbein, 2009], though these costs may be outweighed by the benefits of increased competitive ability

during periods of elevated reproductive competition. Glucocorticoids (GCs) are associated with the stress response, which is adaptive in the short term because it focuses energetic resources to areas relevant to immediate survival needs (e.g. energy release) and temporarily suppresses nonvital functions (e.g. growth, reproduction) [Sapolsky, 2004]. However, long-term or chronic expression of the stress response, as observed by high GC levels over extended periods, can be detrimental to health [reviewed in Sapolsky, 2004].

As yet, no single hormonal profile exists that reflects either dominant or subordinate status [Creel, 2001], suggesting that many species-specific or population-based social and environmental factors influence the socioendocrinology of male dominance relationships. Research has primarily focused on the many factors that influence male intragroup competition, and thus the relationship between dominance and hormones. These factors may include breeding seasonality [Bales et al., 2006; Brockman et al., 1998; Lynch et al., 2002; Ostner et al., 2002, 2008, 2011] and the presence of periovulatory females [Vandenberg, 1969; Muller & Wrangham, 2004a; Setchell et al., 2008, 2010], dominance style [Strier et al., 1999; Muehlenbein et al., 2004; Muller & Wrangham 2004a] and group stability [Sapolsky, 1993; Brockman et al., 2001], relative allostatic load or metabolic expenditure [Goymann & Wingfield, 2004; Muller & Wrangham, 2004b; Sapolsky, 2004], as well as kinship, the frequency of being subjected to stress, the ability to avoid overt competition (i.e. via evasion) and the availability of social partners [Abbott, et al. 2003; Sapolsky, 2004].

In addition to the role of intragroup reproductive competition (i.e. internal source) on hormone levels in multi-male species, the influence of dispersing males, intergroup encounters (IGEs), and group takeovers (i.e. external sources) must also be considered as these may be the greater source of reproductive competition in some species [Strier, 1994; van Hooft, 2000]. Nonresident males can pose a threat to resident male reproductive success via copulations with group females, and via IGEs and takeovers which may result in loss of rank, group eviction, infanticide, injury, or death [Hrdy, 1977; Brockman et al., 2001, 2009; Fedigan & Jack, 2004; van Schaik et al., 2004; Bergman et al., 2005; Cristóbal-Azkarate et al., 2006; Ostner et al., 2008; Van Belle et al., 2009b, 2010]. These external sources of reproductive competition may exert a strong influence on male hormone levels. Variation in the androgen profiles of males may be associated with differences in group composition [*Cercopithecus aethiops*: Whitten & Turner, 2004; *Alouatta pigra*: Rangel-Negrin et al., 2011], while androgen elevations are frequently associated with challenges from extragroup males [*Propithecus verreauxi*: Brockman et al., 1998; *Callithrix kuhlii*: Ross et al., 2004; *Colobus vellerosus*: Teichroeb & Sicotte, 2008; *Alouatta palliata*: Cristóbal-Azkarate et al., 2006; but see *Eulemur fulvus rufus*: Ostner et al., 2002, 2008; *Saguinus mystax*: Huck et al., 2005; *A. pigra*: Van Belle et al., 2009a], immigration and emigration [*Papio cynocephalus*: Alberts et al., 1992; *P. verreauxi*: Brockman et al., 1998, 2001], and infanticide risk [*E. fulvus rufus*: Ostner et al., 2002, 2008; *C. aethiops*: Whitten & Turner, 2004].

Participation in group defense is likely energetically expensive, such that an increase in metabolic load could be associated with increased GC levels, but evidence of this is equivocal. For example, while male chacma baboons experience elevated GC levels after changes in male group membership [i.e. via immigrations, *P. hamadryas ursinus*: Bergman et al., 2005], male moustached tamarins, mantled howlers, black howlers, and captive male marmosets do not show an increase in GCs after encounters with extragroup males [*C. kuhlii*: Ross et al., 2004; *S. mystax*: Huck et al., 2005; *A. palliata*: Cristóbal-Azkarate et al., 2007; *A. pigra*: Van Belle et al., 2009a]. It may be that GC levels are associated more with actual changes rather than threats of changes in group membership. It also appears that group males maintain elevated GCs as an anticipatory response to the risk of takeover and killing of infants by extragroup males [*P. verreauxi*: Brockman et al., 2009; *E. fulvus rufus*: Ostner et al., 2008].

In this study, we examine the association between fecal androgen (fA) and fecal glucocorticoid (fGC) levels and intergroup reproductive competition in alpha and subordinate male white-faced capuchins residing in the Santa Rosa Sector, Costa Rica. By chance our study took place during an unusually lengthy non-reproductive period during which all group females were either pregnant or lactating. This provided us with the rare opportunity to examine male hormone profiles in the absence of periovulatory females and during a time when male intragroup reproductive competition is expected to be low, though perhaps not completely absent. White-faced capuchins are female philopatric and reside in groups comprised of related adult females, multiple immigrant subadult and adult males, and their offspring. Intragroup male relationships are characterized by affiliative interactions and low agonism in the study population [Jack, 2003], and cooperation in group defense against predators and extragroup males [Jack 2003, Perry 1998]. While intragroup relationships among coresident males are tolerant, at the intergroup level, severe and sometimes lethal male-male agonism occurs in the context of IGEs and group takeovers [Perry, 1996; Fedigan & Jack, 2004]. During IGEs, male white-faced capuchins are the primary participants, while females only rarely participate and generally flee from encounter locations with immature group members [pers. obs.; Perry, 1996; Rose & Fedigan, 1995; Rose 1998; but see Crofoot, 2007]. Intergroup encounters in white-faced capuchins occur throughout the year, although they are slightly more frequent in the dry season [Perry, 1996].

Overall, IGEs play an important role in male long-term reproductive strategies, especially in terms of assessing the composition of neighboring groups for dispersal opportunities [Perry, 1996; Fedigan & Jack, 2004; Crofoot, 2007]. Adult male white-faced capuchins usually immigrate into groups by aggressively and cooperatively evicting resident males [Fedigan & Jack, 2004] and immigration is the primary way in which they increase their dominance rank [Jack & Fedigan, 2004]. In our long-term study groups, most takeovers (14 of 15) occur during the dry season months of January to April, occur approximately every four years [Fedigan & Jack, 2004], and result in the eviction of resident males, infanticides, and the occasional deaths or disappearances of both males and females [Rose, 1998; Fedigan, 2003; see also Gros-Louis et al., 2003 and Perry, 1995 for similar observations at Lomas Barbudal, Costa Rica]. While infanticides are most common after takeovers [Fedigan & Jack, 2004], they can also occur during IGEs [Vogel & Fuentes-Jiménez, 2006].

In species where dominant males sire the majority of offspring, the negative impact of lost future reproductive opportunities due to decreased rank or eviction may mean that dominant individuals are disproportionately affected by threats from extragroup males. The additional threat of infanticide, which would result in loss of current offspring, may be especially important. For example, dominant male sifakas, who have priority of access to females [Brockman et al., 1998], appear to respond to challenges from extragroup males and immigration events with elevated testosterone [Brockman et al., 2001] but not fGC [Brockman et al., 2009]. Evidence of overt intragroup male reproductive competition in white-faced capuchins is rare [but see Perry, 1998] and alpha males appear to obtain only a slightly disproportionate number of copulations [Rose, 1998]. However, genetic studies demonstrate that alpha males sire the majority of offspring [Jack & Fedigan, 2006; Muniz et al., 2010]. Since alpha males have the highest reproductive success, they have the most to lose during IGEs and group takeovers in terms of future reproductive opportunities due to eviction or rank loss, and current reproductive success due to infanticide. This may explain why alpha males are the most vigilant [Rose & Fedigan, 1995] and the most involved in group defense during IGEs and takeover attempts [Perry, 1996].

Given that IGEs (and, by association, takeovers) may result in loss of rank, group eviction, infanticide, injury, or death, we predict that male fA and fGC levels will be positively associated with IGE rates. We further predict that alpha males will have higher fA than

subordinate males because of the disproportional threat of IGEs to alpha male reproductive success. We also predict that alpha males will have higher fGC levels than subordinates because of the presumed higher metabolic costs associated with elevated vigilance and high level of participation in IGEs.

METHODS

Study site and subjects

Our study took place in the Santa Rosa Sector of the Área de Conservación de Guanacaste (formerly known as Santa Rosa National Park) from February through July 2007. Located in the northwestern part of Costa Rica, the Santa Rosa Sector is an approximately 108 km² protected area made up of a mosaic of dry deciduous, semi-evergreen and riparian forest fragments interspersed with regenerating pasture [for additional information on the field site, see Fedigan & Jack, 2012]. Research complied with protocols approved by the Área de Conservación de Guanacaste and the Tulane University IACUC and adhered to the legal requirements of Costa Rica and the United States as well as the American Society of Primatologists principles for the ethical treatment of primates.

We collected hormone data on all male white-faced capuchins (*Cebus capucinus*) aged > 6 years (N = 8) residing in three habituated social groups. Over 940 contact hours were logged in the dry season (EX: 280h, GN: 316h, LV: 344h) and 255 hours in the wet season (EX: 80h, GN: 87h, LV: 88h). With the exception of the only subadult male in our sample (MZ in LV), all males were immigrants and estimated to be ~10 years of age. As in this and other capuchin species [Perry, 1998; Fragaszy et al., 2004], we were unable to determine male dominance rank below the alpha level due to the rarity of agonistic interactions between male dyads. However, alpha status was clearly identifiable based on physical appearance (i.e. larger brows, jaws, and overall size, and greater piloerection) and behavior.

About half of all IGEs involve chasing and/or physical contact, while the remaining IGEs were characterized by avoidance, fleeing, vocal and ritualized physical threats, or persistent vigilance (Childers, 2008). Alpha males are the most active participants, and participate in the majority of IGEs [Rose & Fedigan, 1995; Perry, 1996], and in groups with multiple males, most subordinates participate, though their participation and intensity is more variable (pers. obs.). We used handheld digital voice recorders to record *ad libitum* data on the date and time of the encounter, and which groups were involved. Intergroup encounters are rare events, and were too infrequently observed during our study period to calculate reliable monthly IGE rates. Therefore, we calculated each group's IGE rate in the dry and the early wet seasons by dividing the number of known IGEs by the total contact hours for each group (Table 1). The dry season receives very little, if any, rainfall, and its end is marked by almost daily rainfall beginning in mid-May. Based on rainfall data, the last day of the dry season was May 17, 2007, which we used as the cutoff point between the dry and early wet season.

Collection and analyses of fecal samples

We collected morning fecal samples from all study males at least once per week (N = 194). Sample collection, storage, and preliminary field extraction methods followed Carnegie et al. [2011b], using Prevail C18 Maxi-Clean 300mg SPE cartridges (Alltech Associates, Inc., Deerfield, IL). Samples were transported and analyzed at the Wisconsin National Primate Research Center (WNPRC) by VAMS. Hormones were washed using 1 mL of 5% methanol, eluted from the SPE cartridges using 2 mL of 100% methanol, dried in a hot water bath, reconstituted in 1 mL of 100% ethanol, and stored in a refrigerator. Samples were subsequently analyzed by in-house enzyme immunoassay (EIA) for fA [see Ginther et

al., 2001] and fGC [Carnegie et al., 2011b], both of which are competitive binding assays that use antibodies raised in rabbits (fA antibody R156, fGC antibody R4866, Munro, University of California, Davis). For fA, sample volumes of 5–25 μ L of the ethanol solution were dried and resuspended in the assay solution, and 1/3 was aliquoted into each well. For fGC, 50 μ L of the ethanol solution was used. All samples and standards (1–250 pg, N = 8, Δ 4-androsten-17 β -ol-3-one for fA, and 11 β ,17 α ,21-Trihydroxypregn-4-ene-3,20-dione for fGC, Sigma-Aldrich, St. Louis, MO) were run in duplicate. Concentrations of fA and fGC were measured in ng/g.

Mean intra- and inter-assay coefficients of variation (CV) for fA were 5.5 ng/g and 17.9 ng/g respectively for the low pool, and 7.4 ng/g and 12.4 ng/g respectively for the high pool. Mean intra- and inter-assay CVs for fGC were 6.2 ng/g and 14.0 ng/g respectively for the low pool, and 2.7 ng/g and 6.0 ng/g respectively for the high pool. Mean percent accuracy (the degree to which the assay actually measures the steroid in a known pool) was high for both hormones (fA: 109.05 ± 2.79 , N = 8; fGC: 115.48 ± 2.27 , N = 6). Additionally, the percent bound values from the pooled samples paralleled the percent bound values of the standard curve for both hormones, and there were no differences in their slopes (fA: $t(28) = 0.841$, $P > 0.05$, N = 8; fGC: $t(21) = 0.066$, $P > 0.05$, N = 7).

The validity of the androgen assay is supported by the observation that adult males have higher fecal testosterone than juvenile males [Jack et al., 2011] and fA increased in a subordinate male who became the alpha male [Schoof et al., 2011]. We did not conduct an ACTH challenge in our study population to test the biological validity of the fGC response, because this would interfere with the naturalistic approach of the research conducted at this field site. However, previous research in this population of *Cebus capucinus* indicates that unhabituated individuals respond to observer presence with high fGC levels than habituated individuals [Jack et al., 2008], females respond to a presumably stressful situation (change in male group composition) with an increase in fGC level [Carnegie et al., 2011b], and a case study indicates that a male had an increase in fGC after he became alpha male [Schoof et al., 2011]. ACTH challenge in *Cebus apella* results in a cortisol increase [Torres-Farfan et al. 2009].

Statistical Analysis

We used repeated measures generalized linear mixed models (gamma probability distribution, log link function) to analyze the association between IGE rate, male dominance status (alpha versus subordinate), and hormone levels. Specifically, each model was fitted with the hormone (fA or fGC) as the response variable, fecal sample order (for each male) as the repeated variable, dominance status and IGE rate as fixed effects, and male nested within social group as a random factor. We used a first-order autoregressive (AR1) covariance structure for the repeated measure. A two-tailed t-test revealed no collinearity between the categorical predictor (dominance status) and the continuous predictor (IGE rate) ($t = 0.955$, $df = 192$, $P = 0.341$). We used models permitting variation in degrees of freedom across tests (recommended for small sample sizes), and used robust estimation to handle violations of model assumptions about normality and homogeneity of variance of the residuals. All analyses were run in IBM SPSS Statistics for Windows, version 19.0.

RESULTS

Androgens

Intergroup encounter rate was positively and significantly associated with fA levels ($F = 21.534$, $P < 0.001$, $df = 1, 191$; Figure 1a), and alpha males had significantly higher fA levels than subordinate males ($F = 13.769$, $P < 0.001$, $df = 1, 191$; Figure 2a).

Glucocorticoids

Intergroup encounter rate was positively and significantly associated with fGC levels ($F = 5.693$, $P = 0.018$, $df = 1, 191$; Figure 1b), and alpha males had significantly higher fGC levels than subordinate males ($F = 4.441$, $P = 0.036$, $df = 1, 191$; Figure 2b). Figure 2b indicates that alpha males had higher fGC levels in the two larger groups with multiple subordinates (GN and LV), but not in the smaller group with a single subordinate (EX).

DISCUSSION

White-faced capuchins are classified as moderate seasonal breeders [for classification see van Schaik et al., 1999]; however, sexual behavior, copulations, conceptions and births can and do occur throughout the year [Carnegie et al., 2011a]. During our study period all females were either pregnant or lactating (confirmed by progesterone and estradiol assays, unpublished data), and although courtship behavior and copulations (mounts with thrusting) were observed, these were clearly not associated with conceptions. The timing of our study allowed us to examine the influence of intergroup competition on male androgen and glucocorticoid levels without the possible confound of pronounced intragroup competition, since we can exclude the possibility that males were competing for immediate access to periovulatory females.

Male fA levels were positively and significantly associated with IGE rate. Similar androgen increases have been observed in other species after challenges from extragroup males or bisexual groups [*P. verreauxi*: Brockman et al., 1998; *C. kuhlii*: Ross et al., 2004; *A. palliata*: Cristóbal-Azkarate et al., 2006; *C. vellerosus*: Teichroeb & Sicotte, 2008; *A. pigra*: Rangel-Negrín et al., 2011, but see *E. f. rufus*: Ostner et al., 2008; *S. mystax*: Huck et al., 2005; *A. pigra*: Van Belle et al., 2009a]. Black howler males (*A. pigra*) living in uni-male groups have higher androgen levels than males living in multi-male groups [Rangel-Negrín et al., 2011]. Given that these males have an increased probability of being challenged for group membership, elevated androgen levels may represent an anticipatory response to reproductive challenges from extragroup males [Rangel-Negrín et al., 2011]. In white-faced capuchins, all males face the threat of group eviction since most takeovers result in the complete replacement of resident males [Fedigan & Jack, 2004]. The positive relationship between fA and IGE rates in white-faced capuchins indicates that resident males perceive IGEs as a form of reproductive competition.

Male fGC levels were also significantly associated with IGE rate. Several studies examining challenges by extragroup males did not observe an increase in fGC levels [*C. kuhlii*: Ross et al., 2004; *S. mystax*: Huck et al., 2005; *A. palliata*: Cristóbal-Azkarate et al., 2007; *A. pigra*: Van Belle et al., 2009a]. However, previous research on male baboons has demonstrated that immigration events are associated with elevated GC levels, indicating that these are stressful events [*P. h. ursinus*: Bergman et al., 2005]. To our knowledge, however, this is the first study to report a significant relationship between fGC and the rate of intergroup encounters.

In addition to the threats of IGEs mentioned above, infanticide in white-faced capuchins is frequent after group takeovers [Perry, 1996; Rose, 1998; Fedigan, 2003; Gros-Louis et al., 2003] and occurs occasionally during IGEs [Vogel & Fuentes-Jiménez, 2006]. In our study, we calculated IGE rates by season; the observed association between IGE rates and both fA and fGC may be tied to the greater risk of male takeovers (and therefore infanticides) during the dry season [Fedigan & Jack, 2004], which also coincides with the birth peak [Carnegie et al., 2011a]. Reported elevations in fA and fGC levels during the birth season in Verreaux's sifakas and red-fronted lemurs have been linked to the predictable risk of infanticide in these species [*P. verreauxi*: Brockman et al., 2001, 2009; *E. f. rufus*: Ostner et al., 2002, 2008].

Given that alpha male white-faced capuchins are the most active participants during IGEs and takeover attempts and that they direct most of their vigilance towards non-resident males [Rose & Fedigan, 2005; pers. obs.], we argue that it is not surprising to find that they display significantly higher fA and fGC levels than subordinate males. The observation that alpha males in all three social groups had higher fA than their subordinates supports our contention that IGEs pose a disproportional threat to the reproductive success of alpha males. We also predicted that alpha males would have higher fGC levels than subordinate males because of the presumed higher metabolic costs associated with elevated vigilance and high level of participation in IGEs. Predicting elevated fGC levels in alpha males is also consistent with the notion that perceived infanticide risk may disproportionately affect them since they sire the majority of offspring [Jack & Fedigan, 2006; Muniz et al., 2010]. Indeed, the two social groups in which alpha males did have higher fGC levels had five and four infants (GN, LV, see Figure 2b), respectively, born during the first four months of the study, which corresponded with the dry season. The third group (EX) had only one infant under the age of 6 months at the start of the study period, and no births occurred during the remainder of the study. Therefore, lower perceived risk of infanticide may account for both lower fGC and fA in the alpha male in EX group compared to the alpha males in the other two groups.

We propose that the difference between alpha and subordinate male fA and fGC during our study is driven primarily by the greater threat of IGEs and takeovers to alpha males, rather than by high intragroup male competition, since no periovulatory females were present. However, chimpanzee and grey-cheeked mangabey males display a similar pattern, where dominant males maintain higher androgens and GC levels than subordinates even in the absence of periovulatory females [*P. troglodytes*: Muehlenbein et al., 2004; Muller & Wrangham, 2004a, b; *L. albigena*: Arlet et al., 2009, 2011]. In these species, dominance status is achieved and maintained aggressively, so the association between persistently elevated androgens levels, status, and aggression are not surprising. Muller & Wrangham [2004a] propose that since estrous females are rare, males maintain high androgens levels due to the unpredictability of challenges from rivals for access to these females. Furthermore, they suggest that the positive association between dominance status and GC levels are due to relatively higher energetic demands [Muller & Wrangham, 2004b]. Given that white-faced capuchin births occur throughout the year, females ovulate rarely because of a long interbirth interval (2.2 years), and most (if not all) ovulatory cycles result in conceptions [data not shown; Carnegie et al., 2011a]; therefore, the presence of ovulatory females in this species is also unpredictable. However, previous research shows that within group male-male aggression is extremely rare and that dominance is not aggressively maintained in the studied population [Jack, 2003; but see Perry, 1998 for observations in a different population], making it unlikely that alpha male capuchins maintain higher fA levels due to the unpredictability of aggressive challenges by coresident males. Indeed, in the present study, some males displayed no agonism towards their coresident males, which is why we were unable to assign ranks below the alpha level.

In contrast to chimpanzees and mangabeys, black howler males have low intragroup aggression. Nonetheless, Van Belle et al. [2009a] found that “central” black howler males have higher fA and fGC than “noncentral” males, even in the absence of periovulatory females, a pattern similar to that reported here for white-faced capuchins. Van Belle et al. [2009a] propose that coresident males compete year round, like chimpanzees and mangabeys, but that they do so covertly by fostering relationships with group females. They propose that the costs of this form of competition may account for higher fA and fGC levels observed in “central” compared to “noncentral” males. As with savannah baboons [*P. cynocephalus*: Gesquiere et al., 2011], costs associated with being an alpha or “central” male may exist even during times when intragroup reproductive competition appears to be low. Previous research has indicated that white-faced capuchin alpha males also generally spend

more time affiliating with females than do subordinate males [Perry, 1997; Jack, 2003; Carnegie et al., 2005]. Therefore, we cannot entirely rule out the possibility that covert intragroup competition occurs even in the absence of periovulatory females. To further explore the influence of intragroup reproductive competition on male hormone levels, future work should focus on an examination of male fA and fGC levels during periods when females are ovulating, as well as on male-female interactions.

Acknowledgments

Research complied with protocols approved by the Área de Conservación de Guanacaste (ACG) and the Tulane University IACUC, and adhered to the legal requirements of Costa Rica and the United States. A. Childers, Z. King, and C. Sheller assisted in field data collection, and J. Addicott provided data management consulting. D. Wittwer and T. Ziegler provided valuable training and guidance for hormone analyses [the NIH provides partial support to the Wisconsin National Primate Research Center (RR000167) for the cost of analysis]. We sincerely thank them all. Research support and conference funding was available from J. Armand Bombardier Internationalist Fellowship (VAMS), Fonds Québécois de la Recherche sur la Nature et les Technologies Master's Research Scholarship (VAMS), Tulane University's Graduate Student Studies Association (VAMS), Provost's office (VAMS), Research Enhancement Fund (KMJ), Department of Anthropology (KMJ) and Stone Center for Latin American Studies (VAMS and KMJ).

References

- Abbott DH, Keverne EB, Bercovitch FB, Shively CA, Mendoza SP, Saltzman W, Snowdon CT, Ziegler TE, Banjevic M, Garland T Jr, Sapolsky RM. Are subordinates always stressed? A comparative analysis of rank differences in cortisol levels among primates. *Hormones and Behavior*. 2003; 43:67–82. [PubMed: 12614636]
- Alberts SC, Sapolsky RM, Altmann J. Behavioral, endocrine, and immunological correlates of immigration by an aggressive male into a natural primate group. *Hormones and Behavior*. 1992; 26:167–178. [PubMed: 1612563]
- Arlet ME, Grote MN, Molleman F, Isbell L, Carey JR. Reproductive tactics influence cortisol levels in individual male gray-cheeked mangabeys (*Lophocebus albigena*). *Hormones and Behavior*. 2009; 55:210–216. [PubMed: 18996388]
- Arlet ME, Kaasik A, Molleman F, Isbell L, Carey JR, Mänd R. Social factors increase fecal testosterone in wild male gray-cheeked mangabeys (*Lophocebus albigena*). *Hormones and Behavior*. 2011; 59:605–611. [PubMed: 21376722]
- Baker AJ, Dietz JM, Kleiman DG. Behavioural evidence for monopolization of paternity in multi-male groups of golden lion tamarins. *Animal Behaviour*. 1993; 46:1091–1101.
- Bales KL, French JA, McWilliams J, Lake RA, Dietz JM. Effects of social status, age, and season on androgen and cortisol levels in wild male golden lion tamarins (*Leontopithecus rosalia*). *Hormones and Behavior*. 2006; 49:88–95. [PubMed: 15978593]
- Bergman TJ, Beehner JC, Cheney DL, Seyfarth RM, Whitten PL. Correlates of stress in free-ranging male chacma baboons, *Papio hamadryas ursinus*. *Animal Behaviour*. 2005; 70:703–713.
- Brockman DK, Whitten PL, Richard AF, Schneider A. Reproduction in free-ranging male *Propithecus verreauxi*: The hormonal correlated of mating and aggression. *American Journal of Physical Anthropology*. 1998; 105:137–151. [PubMed: 9511910]
- Brockman DK, Whitten PL, Richard AF, Benanader B. Birth season testosterone levels in male Verreaux's sifaka, *Propithecus verreauxi*: Insights into socio-demographic factors mediating seasonal testicular function. *Behavioral Ecology and Sociobiology*. 2001; 49:117–127.
- Brockman DK, Cobden AK, Whitten PL. Birth season glucocorticoids are related to the presence of infants in sifaka (*Propithecus verreauxi*). *Proceedings of the Royal Society B: Biological Sciences*. 2009; 276:1855–1863.
- Carnegie SD, Fedigan LM, Ziegler TE. Behavioral indicators of ovarian phase in white-faced capuchins (*Cebus capucinus*). *American Journal of Primatology*. 2005; 67:51–68. [PubMed: 16163719]

- Carnegie SD, Fedigan LM, Melin AD. Reproductive seasonality in female capuchins (*Cebus capucinus*) in Santa Rosa (Area de Conservación Guanacaste), Costa Rica. *International Journal of Primatology*. 2011a; 32:3–31.
- Carnegie SD, Fedigan LM, Ziegler TE. Social and environmental factors affecting fecal glucocorticoids in wild, female white-faced capuchins (*Cebus capucinus*). *American Journal of Primatology*. 2011b; 73:1–9.
- Childers, A. Unpublished Master's Thesis. Tulane University; New Orleans, Louisiana, USA: 2008. Spatial ecology of Costa Rica white-faced capuchins: Socioecological and cognitive implications.
- Cristóbal-Azkarate J, Chavira R, Boeck L, Rodríguez-Luna E, Veà JJ. Testosterone levels of free-ranging resident mantled howler monkey males in relation to the number and density of solitary males: A test of the challenge hypothesis. *Hormones and Behavior*. 2006; 49:261–267. [PubMed: 16182297]
- Cristóbal-Azkarate J, Chavira R, Boeck L, Rodríguez-Luna E, Veà JJ. Glucocorticoid levels in free ranking resident mantled howlers: a study of coping strategies. *American Journal of Primatology*. 2007; 69:866–876. [PubMed: 17358001]
- Crofoot MC. Mating and feeding competition in white-faced capuchins (*Cebus capucinus*): the importance of short- and long-term strategies. *Behaviour*. 2007; 144:1473–1495.
- Fedigan LM. Impact of male takeovers on infant deaths, births and conceptions in *Cebus capucinus* at Santa Rosa, Costa Rica. *International Journal of Primatology*. 2003; 24:723–741.
- Fedigan LM, Jack KM. The demographic and reproductive context of male replacements in *Cebus capucinus*. *Behaviour*. 2004; 141:755–775.
- Fedigan, LM.; Jack, KM. Tracking monkeys in Santa Rosa: Lessons from a regenerating tropical dry forest. In: Kappeler, P.; Watts, D., editors. *Long-term Field Studies of Primates*. Springer Press; 2012. p. 165-184.
- Fernandez-Duque E, Valeggia CR, Mendoza SP. The biology of paternal care in human and nonhuman primates. *Annual Review of Anthropology*. 2009; 38:115–130.
- Fragaszy, DM.; Visalberghi, E.; Fedigan, LM. *Cebus*. Cambridge: Cambridge University Press; 2004. The complete capuchin: The biology of the genus; p. 339
- Gesquiere LR, Lear NH, Simao MCM, Onyango PO, Alberts SC, Altmann J. Life at the top: Rank and stress in wild male baboons. *Science*. 2011; 333:357–360. [PubMed: 21764751]
- Ginther AJ, Ziegler TE, Snowdon CT. Reproductive biology of captive male cotton-top tamarin monkeys as a function of social environment. *Animal Behaviour*. 2001; 61:65–78. [PubMed: 11170697]
- Gros-Louis J, Perry S, Manson JH. Violent coalitionary attacks and intraspecific killing in wild white-faced capuchin monkeys (*Cebus capucinus*). *Primates*. 2003; 44(4):341–346. [PubMed: 12910384]
- Hrdy SB. Infanticide as a primate reproductive strategy. *American Scientist*. 1977; 65:40–49. [PubMed: 402873]
- Jack KM. Explaining variation in affiliative relationships among male white-faced capuchins (*Cebus capucinus*). *Folia Primatologica*. 2003; 74:1–16.
- Jack KM, Fedigan LM. Male dispersal patterns in white-faced capuchins, *Cebus capucinus*. Part 2: Patterns and causes of secondary dispersal. *Animal Behaviour*. 2004; 67:771–782.
- Jack, KM.; Fedigan, LM. Why be alpha male? Dominance and reproductive success in wild white-faced capuchins (*Cebus capucinus*). In: Estrada, A.; Garber, P.; Luecke, L.; Pavelka, MSM., editors. *New Perspectives in the Study of Mesoamerican Primates: Distribution, Ecology, Behavior and Conservation*. New York: Springer; 2006. p. 367-386.
- Jack KM, Lenz BB, Healan E, Rudman S, Schoof VAM, Fedigan LM. The effects of observer presence on the behavior of *Cebus capucinus* in Costa Rica. *American Journal of Primatology*. 2008; 70:490–494. [PubMed: 18076061]
- Jack KM, Schoof VA, Sheller CR, Rich CI, Klingelhofer PP. The influence of age and dominance status on fecal testosterone, dihydrotestosterone, and cortisol in male white-faced capuchin monkeys (*Cebus capucinus*) in Santa Rosa National Park, Costa Rica [Abstract]. *American Journal of Primatology*. 2011; 73 (suppl 1):61.

- Lynch JW, Ziegler TE, Strier KB. Individual and seasonal variation in fecal testosterone and cortisol levels of wild male tufted capuchin monkeys, *Cebus apella nigrilus*. *Hormones and Behavior*. 2002; 41:275–287. [PubMed: 11971661]
- Muehlenbein, MP. The application of endocrine measures in primate parasite ecology. In: Huffman, MA.; Chapman, CA., editors. *Primate parasite ecology: The dynamics and study of host-parasite relationships*. Cambridge: Cambridge University Press; 2009. p. 63-81.
- Muehlenbein MP, Watts DP, Whitten PL. Dominance rank and fecal testosterone levels in adult male chimpanzees (*Pan troglodytes schweinfurthii*) at Ngogo, Kibale National Park, Uganda. *American Journal of Primatology*. 2004; 64:71–82. [PubMed: 15356859]
- Muller MN, Wrangham RW. Dominance, aggression and testosterone in wild chimpanzees: A test of the 'challenge hypothesis'. *Animal Behaviour*. 2004a; 67:113–123.
- Muller MN, Wrangham RW. Dominance, cortisol and stress in wild chimpanzees (*Pan troglodytes schweinfurthii*). *Behavioral Ecology and Sociobiology*. 2004a,b; 55:332–340.
- Muniz L, Perry S, Manson JH, Gilkenson H, Gros-Louis J, Vigilant L. Male dominance and reproductive success in wild white-faced capuchins (*Cebus capucinus*) at Lomas Barbudal, Costa Rica. *American Journal of Primatology*. 2010; 72:1118–1130. [PubMed: 20717998]
- Ostner J, Kappeler PM, Heistermann M. Seasonal variation and social correlates of androgen excretion in male redfronted lemurs (*Eulemur fulvus rufus*). *Behavioral Ecology and Sociobiology*. 2002; 52:485–495.
- Ostner J, Kappeler P, Heistermann M. Androgen and glucocorticoids levels reflect seasonally occurring social challenges in male redfronted lemurs (*Eulemur fulvus rufus*). *Behavioral Ecology and Sociobiology*. 2008; 62:627–638. [PubMed: 19816530]
- Ostner J, Heistermann M, Schülke O. Male competition and its hormonal correlates in Assamese macaques (*Macaca assamensis*). *Hormones and Behavior*. 2011; 59:105–113. [PubMed: 21056566]
- Perry S. Intergroup encounters in wild white-faced capuchins (*Cebus capucinus*). *International Journal of Primatology*. 1996; 17:309–330.
- Perry S. Male-female social relationships in wild white-faced capuchins (*Cebus capucinus*). *Behaviour*. 1997; 134:477–510.
- Perry S. Male-male social relationships in wild white-faced capuchins, *Cebus capucinus*. *Behaviour*. 1998; 135:139–172.
- Rangel-Negrín A, Dias PAD, Chavira R, Canales-Espinosa D. Social modulation of testosterone levels in male black howlers (*Alouatta pigra*). *Hormones and Behavior*. 2011; 59:159–166. [PubMed: 21081132]
- Rose, LM. PhD Dissertation. Washington University; St. Louis MI: University Microfilms International; 1998. Behavioral ecology of white-faced capuchins (*Cebus capucinus*) in Costa Rica.
- Rose LM, Fedigan LM. Vigilance in white-faced capuchins, *Cebus capucinus*, in Costa Rica. *Animal Behaviour*. 1995; 49:63–70.
- Ross CN, French JA, Patera KJ. Intensity of aggressive interactions modulates testosterone in male marmosets. *Physiology & Behavior*. 2004; 83:437–445. [PubMed: 15581666]
- Sapolsky, RM. The physiology of dominance in stable versus unstable social hierarchies. In: Mason, WA.; Mendoza, SP., editors. *Primate social conflict*. Albany: SUNY Press; 1993. p. 171-204.
- Sapolsky RM. Social status and health in humans and other animals. *Annual Review of Anthropology*. 2004; 33:393–418.
- Schoof VAM, Jack KM, Carnegie SD. Rise to power: A case study of male fecal androgen and cortisol levels before and after a non-aggressive rank change in a group of wild white-faced capuchins (*Cebus capucinus*). *Folia Primatologica*. 2011; 82:299–307.
- Setchell JM, Smith T, Wickings EJ, Knapp LA. Social correlates of testosterone and ornamentation in male mandrills. *Hormones and Behavior*. 2008; 54:365–372. [PubMed: 18582885]
- Setchell JM, Smith T, Wickings EJ, Knapp LA. Stress, social behavior, and secondary sexual traits in a male primate. *Hormones and Behavior*. 2010; 58:720–728. [PubMed: 20688067]
- Strier KB. Myth of the typical primate. *Yearbook of Physical Anthropology*. 1994; 37:233–271.

- Strier KB, Ziegler TE, Wittwer DJ. Seasonal and social correlates of fecal testosterone and cortisol levels in wild male muriquis (*Brachyteles arachnoides*). *Hormones and Behavior*. 1999; 35:125–134. [PubMed: 10202120]
- Strier KB, Lynch JW, Ziegler TE. Hormonal changes during the mating and conception seasons of wild northern muriquis (*Brachyteles arachnoides hypoxanthus*). *American Journal of Primatology*. 2003; 61:85–99. [PubMed: 14582130]
- Teichroeb JA, Sicotte P. Social correlates of fecal testosterone in male ursine colobus monkeys (*Colobus vellerosus*): The effect of male reproductive competition in aseasonal breeders. *Hormones and Behavior*. 2008; 54:417–423. [PubMed: 18555251]
- Torres-Farfan C, Abarzua-Catalan L, Valenzuela FJ, Mendez N, Richter HG, Valenzuela GJ, Serón-Ferré M. Cryptochrome 2 expression level is critical for adrenocorticotropin stimulation of cortisol production in the capuchin monkey adrenal. *Endocrinology*. 2009; 150(6):2717–2722. [PubMed: 19246533]
- Van Belle S, Estrada A, Ziegler TE, Strier KB. Social and hormonal mechanisms underlying male reproductive strategies in black howler monkeys (*Alouatta pigra*). *Hormones and Behavior*. 2009a; 56:355–363. [PubMed: 19712675]
- Van Belle S, Estrada A, Ziegler TE, Strier KB. Sexual behavior across ovarian cycles in wild black howler monkeys (*Alouatta pigra*): Male mate guarding and female mate choice. *American Journal of Primatology*. 2009b; 71:153–164. [PubMed: 19016487]
- Van Belle S, Kulp AE, Thiessen-Bock R, Garcia M, Estrada A. Observed infanticides following a male immigration event in black howler monkeys, *Alouatta pigra*, at Palenque National Park, Mexico. *Primates*. 2010; 41:279–284. [PubMed: 20665226]
- van Hooff, JARAM. Relationships among non-human primate males: A deductive framework. In: Kappeler, PM., editor. *Primate males: Causes and consequences of variation in group composition*. Cambridge: Cambridge University Press; 2000. p. 183–191.
- van Schaik, CP.; van Noordwijk, MA.; Nunn, CL. Sex and social evolution in primates. In: Lee, P., editor. *Comparative primate socioecology*. Cambridge: Cambridge University Press; 1999. p. 204–231.
- van Schaik, CP.; Pradham, GR.; van Noordwijk, MA. Mating conflict in primates: Infanticide, sexual harassment and female sexuality. In: Kappeler, PM.; van Schaik, CP., editors. *Sexual selection in primates: New and comparative perspectives*. Cambridge: Cambridge University Press; 2004. p. 131–150.
- Vandenberg JG. Endocrine coordination in monkeys: Male sexual responses to the female. *Physiology and Behavior*. 1969; 4:261–264.
- Vogel ER, Fuentes-Jiménez A. Rescue behavior in white-faced capuchin monkeys during an intergroup attack: support for the Infanticide Avoidance Hypothesis. *American Journal of Primatology*. 2006; 68:1012–1016. [PubMed: 16892414]
- Wingfield JC, Hegner RE, Dufty AMJ, Ball GF. The “Challenge Hypothesis”: Theoretical implications for patterns of testosterone secretion, mating systems, and breeding strategies. *American Naturalist*. 1990; 136:829–846.

Figure 1a.

Figure 1b.

Figure 1.

Log of individual male mean a) fecal androgen and b) fecal glucocorticoid level and seasonal intergroup encounter (IGE) rate (encounters/hour) for each of three social groups

Figure 2a.

Figure 2b.

Figure 2. Mean male a) fecal androgen and b) fecal glucocorticoid levels (ng/g) across three social groups. Alpha males are indicated with solid bars, and subordinates are indicated by open bars.

Table 1

Rates of intergroup encounters (events/hour) in the dry season and wet season for each of the study groups.

Social Group	Dry Season			Early Wet Season		
	IGE	Contact Hours	IGE rate	IGE	Contact Hours	IGE Rate
EX	9	280h37m	0.043	2	80h43m	0.025
GN	8	316h28m	0.029	1	87h47m	0.011
LV	10	344h03m	0.039	3	88h41m	0.034