Published in final edited form as:

Antivir Ther. 2012 ; 17(8): 1483–1493. doi:10.3851/IMP2492.

Antiretroviral-based HIV-1 Prevention: Antiretroviral Treatment and Pre-Exposure Prophylaxis

Connie Celum, MD, MPH and Jared Baeten, MD, PhD

Departments of Global Health, Medicine, and Epidemiology, University of Washington

Abstract

Antiretroviral-based HIV-1 prevention strategies – including antiretroviral treatment (ART) to reduce the infectiousness of HIV-1 infected persons and oral and topical pre-exposure prophylaxis (PrEP) for uninfected persons to prevent HIV-1 acquisition – are the most promising new approaches for decreasing HIV-1 spread. Observational studies among HIV-1 serodiscordant couples have associated ART initiation with a reduction in HIV-1 transmission risk of 80–92%, and a recent randomized trial demonstrated that earlier initiation of ART (i.e., at CD4 counts between 350 and 550 cells/mm³), in the context of virologic monitoring and adherence support, resulted in a 96% reduction in HIV-1 transmission. A number of ongoing and recently-completed clinical trials have assessed the efficacy of PrEP for HIV-1 prevention as peri-coitally administered or daily-administered 1% tenofovir gel and daily oral tenofovir and combination emtricitabine/tenofovir. Completed studies have demonstrated HIV-1 protection efficacies ranging from 39% to 75%. However, two trials in African women have shown no HIV-1 protection with PrEP; the reasons for lack of efficacy in those trials are being investigated. Adherence is likely key to efficacy of antiretrovirals for HIV-1 prevention, both as ART and PrEP. Critical unanswered questions for successful delivery of antiretroviral-based HIV-1 prevention include how to target ART and PrEP to realize maximum population benefits, whether HIV-1 infected persons at earlier stages of infection would accept ART to reduce their risk for transmitting HIV-1 and highest-risk HIV-1 negative persons would use PrEP, and whether high adherence could be sustained to achieve high effectiveness.

Introduction

Antiretrovirals markedly increase the survival of HIV-1 infected persons and are the cornerstone of strategies to prevent vertical HIV-1 transmission. During the past 10 years, a growing scientific and advocacy interest in antiretroviral-based strategies for prevention of sexual HIV-1 transmission has developed, and antiretroviral-based HIV-1 prevention interventions are now among the most promising strategies for dramatically reducing HIV-1 spread. Antiretrovirals have the potential to be used for HIV-1 prevention as antiretroviral treatment (ART) to reduce the infectiousness of HIV-1 infected persons and oral or topical pre-exposure prophylaxis (PrEP) for uninfected persons with repeated and ongoing HIV-1 exposure. The past two years have seen significant new advances in HIV-1 prevention as a result of antiretroviral-based prevention strategies, both ART and PrEP, and policymakers are considering implications of trials of ART and PrEP in terms of resources and programmatic implementation. This review will focus on the rationale and evidence for ART and PrEP for HIV-1 prevention, and remaining questions and uncertainties.

Corresponding author: Connie Celum, MD, MPH, International Clinical Research Center, Department of Global Health, University of Washington, Box 359927, 325 Ninth Avenue, Seattle, WA 98104, 206 520-3825, ccelum@uw.edu.

The authors report no conflict of interest.

Evidence for ART for HIV-1 Prevention: Observational Studies

The primary determinant of risk of HIV-1 transmission is the concentration of HIV-1 in plasma [1, 2] and genital secretions [3]. ART reduces HIV-1 plasma concentrations to undetectable levels within 6 months of initiation in the majority of persons [4, 5] and seminal and cervicovaginal HIV-1 concentrations are also reduced to undetectable levels in most persons on ART [6–10]. Use of peripartum ART is responsible for the remarkable success in virtually eliminating mother-to-child HIV-1 transmission in resource-rich settings [11].

It was hypothesized that the substantial reduction in HIV-1 quantity in plasma and genital compartments in persons on suppressive ART would translate into markedly reduced risk of HIV-1 transmission to sexual partners [1, 12, 13]. A meta-analysis of data from 11 cohorts among 5021 heterosexual HIV-1 serodiscordant couples with 1098 person-years follow-up found only five cases of HIV-1 transmission to sexual partners from HIV-1 infected persons receiving ART, consistent with a transmission rate between 0.19 and 1.09 per 100 person-years [14] (Table 1).

Among 3381 HIV-1 serodiscordant couples in the Partners in Prevention HSV/HIV Transmission Study, HIV-1 transmission risk was reduced by 92% (95% CI 43–100%, $p=0.004$) among the 349 couples in which the HIV-1 infected partners initiated ART during follow-up (1 transmission event in 273 person-years of follow-up), compared to those who did not start ART (102 transmission events in 4558 person-years of follow-up, Table 1) [15]. As in the meta-analysis [14], the rate of HIV-1 transmission was very low (<0.5% per year) after ART initiation. The single HIV-1 transmission event observed after ART initiation occurred less than four months after ART was begun, and thus it is probable that transmission occurred prior to complete HIV-1 suppression as a result of ART. An important aspect of these observational data is that HIV-1 infected partners in the Partners in Prevention HSV/HIV Transmission Study received ART outside of research clinic settings, thus reflecting the community standard of care for ART delivery, adherence counseling, and clinical monitoring. Viral load monitoring was not standard practice in the research study or in clinical settings in the study communities; retrospective testing of a single plasma sample obtained a median of 7.3 months after ART initiation (interquartile range 3.4–12.1 months) demonstrated that 70% of HIV-1 infected persons achieved virologic suppression.

In the Partners in Prevention HSV/HIV Transmission Study, HIV-1 transmissions occurred across the range of CD4 count strata. HIV-1 incidence among couples where the HIV-1 infected partners had $CD4 < 200$ cells/mm³ was >8% per year in the absence of ART, emphasizing that the greatest priority for ART provision for both HIV-1 treatment and prevention is for HIV-1 infected persons with lower CD4 counts. For HIV-1 infected partners who had CD4 counts >200 cells/mm³, 70% of transmissions were from those who had plasma HIV-1 RNA levels 50,000 copies/mL, even though they accounted for only ~30% of follow-up time [15]. These data suggest that high viral load (e.g., 50,000 copies/mL) may be a useful laboratory criterion to augment CD4 counts for determining ART eligibility since it would identify individuals at higher risk of HIV-1 transmission who would benefit from earlier initiation of ART (e.g., at $CD4 > 350$ cells/mm³, the current WHO standard).

Definitive evidence for ART for HIV-1 Prevention: HPTN 052

In 2011, the observational data associating ART initiation with substantial reduction in HIV-1 risk were confirmed by HIV Prevention Trials Network (HPTN) 052, a randomized trial among 1763 HIV-1 serodiscordant heterosexual couples [16]. At the time of enrollment into the trial, all HIV-1 infected partners had a CD4 count between 350 and 550 cells/mm³

(thus not meeting international guidelines for ART provision). HIV-1 infected partners were randomly assigned to immediately initiate ART or wait to initiate ART until their CD4 count fell to 250 cells/mm^3 , consistent with standard practice at the time. In April 2011, the trial's independent Data and Safety Monitoring Board (DSMB) recommended public report of the study results because HIV-1 prevention benefits of earlier ART initiation had become clearly demonstrated (i.e., the statistical calculations for the study crossed a pre-specified efficacy threshold for early discontinuation of the delayed ART arm). Of 39 HIV-1 transmissions observed in the study, 28 were virologically-linked within the study partnership: 27 in the delayed ART arm and only 1 in the immediate ART arm, a 96% reduction in HIV-1 risk that was highly statistically significant (RR 0.04, 95% 0.01–0.27, $p < 0.001$) [16]. The single transmission in the immediate ART arm was observed soon after ART initiation, a similar situation to what was observed in the Partners in Prevention HSV/HIV Transmission Study observational analysis. Notably, HPTN 052 conducted quarterly viral load monitoring with intensive follow-up and adherence counseling for those on ART who were not achieving viral suppression, and as a result they achieved 89% of participants on ART with viral suppression within 3 months and 97% at 24 months [16].

Public Health Implications of ART for HIV-1 Prevention

Scientific opinion [13] and mathematical modeling [17, 18] have stimulated great interest in the potential of ART to substantially reduce population HIV-1 incidence. Universal ART – administered through universal HIV-1 testing, linkage to care, immediate access to ART regardless of CD4, and optimal adherence (together called the ‘Test and Treat’ concept) – has been predicted to virtually eliminate incident HIV-1 infections, over a period of decades [17]. These mathematical models focused on hyperendemic settings with uncertain generalizability to more concentrated epidemics in more resource-rich settings. In addition, the mathematical models that indicate HIV-1 elimination is feasible require almost universal coverage, of both frequent HIV-1 testing and ART provision, as well as very high ART adherence and consequently, rates of virologic suppression. Mathematical models using potentially more realistic assumptions have suggested that the impact of treatment on HIV-1 incidence could be more modest, although still highly beneficial [18]. Uncertainties about ‘Test and Treat’ include: 1) how to address barriers in the ‘cascade’ of steps between learning one is HIV-1 infected and achieving sustained viral suppression with ART [19, 20], 2) the willingness of HIV-1 infected persons to initiate ART at higher CD4 counts when they are asymptomatic and sustain high adherence levels and viral suppression in order to durably reduce HIV-1 transmission, 3) availability of resources to greatly increase and sustain ART coverage, and 4) the ultimate impact of increasing ART provision on population HIV-1 incidence, the key measure of public health prevention benefits. Evaluation of strategies to achieve high knowledge of HIV-1 serostatus, uptake of ART and retention in care is being evaluated in a feasibility study in two U.S. cities (Bronx and Washington DC) through HPTN 065 [21]. Several large trials of the population level impact of Test and Treat strategies have been initiated in Africa, including studies in Botswana, Tanzania, South Africa, and Zambia [22].

While these approaches are being evaluated, an initial focus of earlier ART provision might be HIV-1 serodiscordant couples – a smaller target population and for whom the clinical trial data [16] and mathematical modeling [23] indicate a substantial impact. WHO guidelines about testing, counseling and antiretrovirals for HIV-1 prevention in HIV-1 serodiscordant couples will be released in 2012 [24]. Considering the available evidence and mathematical modeling data, it is clear that ART has the potential to be a highly effective HIV-1 prevention strategy if implementation challenges can be met and high coverage achieved. the results of community trials and demonstration projects of ‘Test and Treat’ interventions are eagerly awaited.

Rationale for Topical and Oral PrEP for HIV-1 prevention

The rationale for prevention of sexual HIV-1 acquisition with PrEP stems from efficacy of antiretrovirals for the prevention of mother-to-child transmission of HIV-1, first demonstrated with peripartum zidovudine [25]. Recent studies have shown that post-natal antiretrovirals, provided to infants who have ongoing exposure to HIV-1 through breastmilk, can substantially reduce HIV-1 risk [26]. Infant studies provide compelling analogous evidence that antiretroviral prophylaxis could be highly efficacious for preventing infection in the context of known and ongoing HIV-1 exposure [27].

Animal studies also provided evidence to suggest that antiretrovirals could be used for prevention of HIV-1 acquisition. Macaque SHIV challenge studies [28] and more recently, humanized mouse HIV-1 challenge studies [29], have tested topical tenofovir gel, oral tenofovir disoproxil fumarate (TDF), and oral combination emtricitabine (FTC)/TDF. The choices of the first candidate ARVs for prevention, TDF and FTC/TDF, were based on their early action in the HIV-1 life cycle, high potency, high genital tract levels [30, 31], excellent safety and tolerability, and low incidence of resistance (e.g., low levels of resistance to tenofovir in short-term monotherapy studies) [32]. Animal model studies have tested drug dosing and delivery routes that reflect oral dosing in humans and repeat low dose mucosal virus challenges to mimic sexual exposure to HIV-1. Overall, these studies indicated high levels of protection from topical tenofovir gel [33] and daily oral dosing of TDF and FTC/TDF, with potentially greater protection from combination FTC/TDF than TDF alone [28, 34], as well as efficacy when FTC/TDF was dosed intermittently (3 days before and 2 hours after rectal viral exposure) [35]. Animal studies also are being used to identify potential new PrEP candidates, including topical maraviroc [36] and raltegravir [37].

Human Efficacy Trials of Topical and Oral PrEP

Proof-of-concept clinical trials of the safety and efficacy of topical tenofovir vaginal gel and oral TDF and oral FTC/TDF as PrEP have been conducted in African women at risk of heterosexual HIV acquisition through six studies: 1) CAPRISA 004 (CAPRISA is the Center for AIDS Programme of Research in South Africa), 2) FEM-PrEP, 3) Partners PrEP, 4) TDF2, and 5) VOICE (Vaginal and Oral Interventions to Control the Epidemic study) and 6) FACTS 001 (FACTS is the Follow-on African Consortium for Tenofovir Studies). Two of these studies enrolled African heterosexual men (Partners PrEP, TDF2). One study enrolled men who have sex with men (MSM) from the Americas, Thailand, and South Africa (iPrEx represents the Spanish acronym for Chemoprophylaxis for HIV Prevention in men). One ongoing study enrolled injection drug users (Bangkok Tenofovir Study) (Table 2). These studies are summarized below in chronological order of their reporting of efficacy findings.

CAPRISA 004 enrolled 889 HIV-1 uninfected women, ages 18–44 in Durban and rural KwaZulu-Natal, South Africa, who were randomized to peri-coital use of 1% tenofovir gel or placebo, dosed by the BAT24 strategy (BAT24 = for before and after sex, not to exceed two doses in 24 hours). The event-driven BAT24 strategy was developed to mimic dosing of nevirapine for prevention of mother-to-child HIV-1 transmission as it is used for peripartum prophylaxis. In CAPRISA 004, tenofovir gel significantly reduced the risk of HIV-1 acquisition by 39% (95% CI 6–60%, $p=0.017$) [38]. Notably, HIV-1 incidence was 9.1 per 100 women-years in the placebo arm, an extraordinarily high rate. No antiretroviral resistance to tenofovir (i.e., the K65R mutation) was detected using standard population sequencing in HIV-1 samples obtained from HIV-1 seroconverters. In subgroup analyses, efficacy was estimated to be 54% in women who reported >80% adherence to gel use with sex acts in the prior month ($p=0.025$). In a case-control analysis of cervicovaginal tenofovir

levels among HIV-1 seroconverters and matched controls, women with levels >1000ng/mL had 74% lower risk of HIV-1 infection than those with <1000 ng/mL [39], supporting evidence of an adherence-efficacy relationship. An unexpected 'bonus' of CAPRISA 004 was a 51% reduction in HSV-2 incidence among women assigned to the tenofovir gel arm, which is consistent with *in vitro* studies that have found that the high mucosal concentrations of tenofovir diphosphate, the active metabolite, achieved with vaginal application have direct anti-herpetic activity [40]. Concentrations of tenofovir diphosphate in vaginal tissues are approximately 100-fold higher with 1% tenofovir gel dosing compared to oral dosing using TDF and topical tenofovir gel results in little systemic absorption [41].

The iPrEx study enrolled 2499 HIV-1 seronegative MSM and transgender women from North and South America, South Africa and Thailand in a randomized placebo-controlled trial of daily oral FTC/TDF. The majority of men were enrolled from the South American sites. The iPrEx study demonstrated that FTC/TDF reduced HIV-1 acquisition risk by 44% (95% CI 15–63%, $p=0.005$). Subgroup analyses indicated higher efficacy (73%) among those with 90% adherence [42]. Importantly, plasma and intracellular drug levels from a subset of study visits demonstrated that only 8% of seroconverters and 54% non-seroconverters had detectable tenofovir or emtricitabine; having detectable drug was strongly associated with substantially lower risk of acquiring HIV-1 (OR 12.9, 95% CI 1.7–99.3). The low level of detection of drug in the non-seroconverters indicates that overall adherence in the study population was only moderate; correlates of higher adherence included age >25 years, participants from US sites, and recent unprotected anal receptive sex [43]. Resistance to emtricitabine (i.e., M184I/V mutations) was detected in two participants randomized to FTC/TDF who had seronegative acute HIV-1 infection at the time of randomization. For these two individuals, emtricitabine resistance was no longer detectable, even by ultrasensitive resistance testing, six months after PrEP discontinuation [44]. Although M184V resistance was rare in the breakthrough infections, retrospective drug level testing in seroconverters indicated that men became infected during periods of low exposure to FTC/TDF [45]. Some experts remain concerned about the potential for undetected archived resistance in PrEP breakthrough infections to compromise response to FTC or 3TC when such individuals initiate HAART. Safety was high with no substantial differences in the rate of serious adverse events, including laboratory events, by arm. A modest but statistically significant 1% reduction in bone mineral density was observed in the FTC/TDF arm, compared with placebo [46]; this decline in bone mineral density is a known effect of tenofovir in studies of HIV-1 treatment – the effect appears to be early then stabilizes, and was not associated with increased risk of bone fracture. No risk compensation was observed based on self-reported condom use during anal sex. Thus, iPrEx indicates excellent safety and moderate efficacy in the context of moderate adherence. Rare resistance was detected -- among two subjects who initiated PrEP during acute seroconversion. Uptake, adherence, risk behaviors, and efficacy in the context of known efficacy of FTC/TDF PrEP is being studied among former iPrEx participants and a subset of new enrollees in a recently-initiated open-label extension study (iPrEx OLE).

The FEM-PrEP study enrolled 1951 high-risk HIV-1 uninfected women from Kenya, South Africa, and Tanzania into a placebo-controlled trial of daily oral FTC/TDF. The study was stopped by its Independent Data Monitoring Committee in April 2011 due to futility: an equal numbers of infections ($n=28$) were seen in each of the two study arms [47]. Adherence was low in FEM-PrEP; 26% of non-seroconverters had consistent tenofovir levels detected in plasma in a retrospective case-control analysis [48]. Intensive PK studies indicate that differential penetration and persistence of the metabolites of TDF and FTC in cervical, vaginal, and rectal tissues may make oral FTC/TDF PrEP more vulnerable to non-adherence in women than men [31, 49].

The TDF2 study enrolled 1200 heterosexual HIV-1 uninfected men and women between the ages of 18 and 40 in Botswana into a placebo-controlled trial of daily oral FTC/TDF; enrollment was terminated early due to larger-than-expected rates of loss-to-follow-up. The study reported results in July 2011: 63% efficacy (95% CI 22–83%, $p=0.01$) for HIV-1 protection due to PrEP [50]. The trial was underpowered to demonstrate sex-based differences with a total of 33 seroconversions, but the point estimates were protective for both men (80%, $p=0.03$) and women (49%, $p=0.1$). Safety and tolerability of FTC/TDF were high. There was no evidence of behavioral risk compensation.

The Partners PrEP Study (for which we are the lead investigators) is an ongoing 3 arm placebo-controlled trial of daily oral TDF and FTC/TDF among 4758 HIV-1 serodiscordant couples from Kenya and Africa among whom the HIV-infected partner is not eligible for ART according to national guidelines; the HIV-1 uninfected partners are randomized to receive PrEP or placebo [51]. On July 10, 2011, the study's Data Safety Monitoring Board recommended that the placebo arm be discontinued due to meeting pre-determined stopping guidelines for efficacy. The Partners PrEP Study demonstrated 67% efficacy of TDF (95% CI 44–81, $p<0.001$) and 75% efficacy of FTC/TDF (95% CI 55–87, $p<0.001$) compared to placebo; the difference between TDF and FTC/TDF was not statistically significant ($p=0.23$). Both TDF and FTC/TDF significantly reduced HIV-1 risk for both men and women [52]. Adherence to study drug was very high based on clinic-based pill counts of unreturned study medication, electronic monitoring and home visits for unannounced pill counts that were done at 3 of the 9 study sites [53], and tenofovir measurement in plasma which was detectable in 82% of randomly selected non-seroconverters [54]. Two of 8 individuals who were randomized to PrEP during seronegative acute HIV-1 infection developed resistance mutations (one K65R and one M184V mutation) but no participants who acquired HIV-1 after PrEP initiation acquired mutations conferring resistance to TDF or FTC [52]. Safety of both drugs was high and no evidence of risk compensation was observed.

The VOICE trial is an ongoing 5 arm study among 5021 HIV-1 uninfected women from South Africa, Uganda, and Zimbabwe in which daily 1% tenofovir gel, daily oral TDF, and daily oral FTC/TDF are being evaluated for safety and effectiveness compared to respective gel/oral placebos. The Data Safety Monitoring Committee for the VOICE trial recommended discontinuation of the oral TDF arm in September 2011 [55] and the daily vaginal 1% tenofovir gel arm in November 2011 due to inability to demonstrate efficacy [56]. The VOICE trial is continuing with the daily oral FTC/TDF arm, which will be completed by mid-2012 and will provide important safety and efficacy data for daily oral FTC/TDF among at-risk African women

Lastly, the Bangkok Tenofovir Study trial of daily oral TDF among HIV-1 uninfected injection drug users compared to placebo is fully enrolled with 2413 participants and is anticipated to have efficacy results in 2012. A majority of the participants are enrolled in methadone replacement programs, where they receive their study medication, essentially as directly-observed PrEP [57].

Current understanding and future directions for PrEP

Efficacy trials have demonstrated high safety and moderate to high efficacy of topical and oral PrEP among South African women (CAPRISA 004), MSM from diverse settings (iPrEx), young Botswana heterosexuals (TDF2), and East African heterosexuals with known HIV-1 infected partners (Partners PrEP). Unraveling potential behavioral and biologic explanations for lack of efficacy of oral PrEP in some populations of women –FTC/TDF in FEM-PrEP and TDF in VOICE -- is essential to understanding whether the lack of efficacy

is primarily due to adherence to study product, types of sexual exposure, and/or biologic co-factors. The futility of daily 1% tenofovir gel in the VOICE trial to prevent HIV-1 acquisition among African at-risk women from South Africa, Uganda and Zimbabwe was surprising, and will require analyses of adherence and biologic co-factors to explain the different outcome from CAPRISA 004 with peri-coital dosing. The recently-launched FACTS 001 study is a confirmatory study of the peri-coital BAT24 strategy among a diverse population of South African women [58] and will provide essential data on safety and effectiveness of peri-coital 1% tenofovir gel in order to hopefully move an effective microbicide forward to licensure and wide-scale use. Tenofovir gel has recently been reformulated in an isoosmolar formulation for rectal application, which will be evaluated for safety and pharmacokinetics among MSM in a Microbicide Trials Network study that will be initiated in 2012.

With respect to oral tenofovir-based PrEP in other populations, the iPrEx trial demonstrates that daily oral FTC/TDF has moderate efficacy (44%) among young, high-risk MSM who were recruited from diverse settings in the Americas, South Africa and Thailand. Importantly, efficacy appeared to be strongly associated with adherence. The iPrEx open label extension study will determine whether knowledge of efficacy of PrEP significantly increases adherence and alters risk behaviors among MSM and thus is a viable public health strategy for MSM. The potential for intermittent dosing of oral FTC/TDF is being evaluated in HPTN 067 with a focus on PK, adherence and risk behaviors [59] and in a recently-initiated trial among MSM in France and Canada (IPERGAY) [60].

The Partners PrEP and TDF2 trials provide strong evidence for PrEP efficacy among African heterosexual populations, who make up the largest portion of the global epidemic [61–65]. In Partners PrEP, both TDF and FTC/TDF were highly effective in both men and women. The study enrolled HIV-1 serodiscordant couples who recognize their risk of HIV-1 [66], and achieved high adherence to daily oral PrEP [53, 67]. Heterosexual HIV-1 serodiscordant couples in Africa account for a substantial proportion of new HIV-1 infections in Africa, and are an increasing focus of HIV-1 prevention efforts [24, 68]. ARV-based prevention for HIV-1 serodiscordant couples could involve a staged approach to ART for the HIV-1 infected partner and PrEP for the HIV-1 uninfected partner, in which PrEP is targeted to the highest-risk couples before the HIV-1 infected partner is eligible for ART by national guidelines and elects to initiate ART, and during the first 6 months after their partner initiates ART and achieves viral suppression [69]. This staged strategy will be evaluated in open label demonstration projects among Kenyan and Ugandan HIV-1 serodiscordant couples, an important next step after the HPTN 052 and Partners PrEP efficacy results, and which will provide data in the next few years for implementation of ARV-based HIV-1 prevention for HIV-1 serodiscordant couples.

Additional analyses – particularly examining adherence – in the FEM-PrEP and VOICE studies will be necessary to understand fully what those trials mean for PrEP use in high-risk women in some African settings. Partners PrEP and TDF2 have shown that PrEP is effective in women, but high-risk populations of women, for behavioral (e.g., adherence, sexual risk) or biologic (e.g., presence of risk factors that heighten HIV-1 risk) reasons, may not have received benefit from PrEP in FEM-PrEP and VOICE.

While awaiting findings from recently-completed and ongoing trials of oral and topical tenofovir-based PrEP in women and injection drug users, it is important to recognize the importance of these initial trials to demonstrating proof-of-concept of antiretroviral-based primary HIV-1 prevention. Much more needs to be understood about targeting of these strategies to those at highest risk in order to be a cost-effective intervention [70, 71], evaluating program delivery models to reduce the risk of PrEP use in acute HIV-1 infection

where the risk of resistance appears to be greatest [42, 72], motivating and monitoring adherence to PrEP, and messaging about PrEP as part of a combined risk reduction strategy given its partial efficacy for HIV-1 prevention (Table 2 and Figure 1). Longer-term safety, adherence, and efficacy needs to be assessed in the context of less frequent visits and briefer counseling than was provided in the intensive proof-of-concept trials.

Some persons at risk of HIV-1 would prefer to use PrEP intermittently, which requires greater understanding of the pharmacokinetics and adherence to intermittent dosing of tenofovir-based PrEP, as will be assessed in HPTN 067. While macaque studies demonstrate biologic feasibility of intermittent oral PrEP dosing, human studies are needed to determine the extent to which individuals at high risk of HIV-1 are able to anticipate sexual activity and achieve sufficient pre-exposure and post-exposure dosing to confer protection.

The PrEP field is also evaluating new candidate products, including non-nucleoside reverse transcriptase inhibitors (dapivirine and rilpivirine) and maraviroc, as well as sustained release delivery systems to provide more options with less dependence on coitally-dependent or daily dosing. Dapivirine has excellent safety and sustained dispersion in a vaginal ring formulation developed by the International Partnership for Microbicides [73], and will be studied for efficacy in collaboration with the US Microbicides Trials Network (MTN 020, ASPIRE trial), beginning in 2012. Oral maraviroc will be evaluated in a comparative safety and tolerability study in MSM (HPTN 069). Dapivirine and maraviroc are being co-formulated in a vaginal ring formulation. Rilpivirine, a NNRTI recently-licensed for treatment, has been formulated into a parenteral formulation for prolonged plasma exposure, and is being evaluated for pharmacokinetics and dose ranging in early clinical trials. Tenofovir is being developed for a vaginal ring formulation and is currently in pre-clinical evaluation. Other candidate compounds and delivery systems (e.g., a dissolvable film) are in pre-clinical stages of evaluation.

Summary

HPTN 052 provides definitive evidence that ART has substantial prevention benefits. The challenge now will be identification of resources to increase ART coverage for the approximate 54% of HIV-1 infected persons globally in 2010 who were eligible for ART under current WHO guidelines [74]. ART for HIV-1 infected partners with CD4 >350 cells/mm³ who are in a known serodiscordant partnership will be recommended in forthcoming WHO guidelines [24]. Further expansion of earlier ART to HIV-infected persons without regard to their partner's HIV-1 serostatus who are at risk of transmission based on other behavioral criteria is being considered and debated [75]. The challenges will not only be resources, but scale-up of HIV-1 testing to achieve high coverage of knowledge of HIV-1 serostatus, effectiveness of linkages to HIV-1 care after learning of HIV-1 infection, the willingness of HIV-1 infected persons to initiate ART at higher CD4 counts, and the proportion of those who initiate ART to achieve and sustain viral suppression.

Proof-of-concept has been demonstrated for tenofovir-based topical (peri-coital administration) and daily oral PrEP for primary prevention. Completion of the remaining trials and analyses of the VOICE and FEM-PrEP trials are needed to have a more complete understanding of effectiveness in different populations, particularly the relationship between adherence and efficacy. Demonstration projects in populations where PrEP has been found to be effective are needed to evaluate targeted implementation and cost-effectiveness of PrEP. Effective HIV-1 prevention requires choices of primary prevention strategies such as PrEP, and scale-up of secondary prevention including ART for HIV-infected persons. While the public health impact of PrEP is anticipated to be greater as products and delivery systems with sustained coverage are identified, those studies will take several years to

complete. In the meantime, targeted provision of daily oral tenofovir-based PrEP to populations at high-risk of HIV-1 acquisition should be evaluated for including as part of combination HIV-1 prevention programs, which should also include ART for HIV-1 infected persons as the cornerstone.

Implementation of both earlier ART for HIV-1 prevention as well as clinical benefits and PrEP will face significant logistical, cost, and commitment hurdles. However, there is now excellent scientific evidence that these strategies reduce HIV-1 risk to a significant degree and thus, could alter the course of the HIV-1 epidemic.

Acknowledgments

Gilead Sciences provided study drug for the Partners PrEP Study but did not provide funding to the authors or the institution.

Funding support: Bill and Melinda Gates Foundation 47674, US National Institutes of Health (R01 MH095507)

References

1. Quinn TC, Wawer MJ, Sewankambo N, et al. Viral load and heterosexual transmission of human immunodeficiency virus type 1. Rakai Project Study Group. *N Engl J Med*. 2000 Mar 30; 342(13): 921–9. [PubMed: 10738050]
2. Lingappa JR, Baeten JM, Wald A, et al. Daily aciclovir for HIV-1 disease progression in people dually infected with HIV-1 and herpes simplex virus type 2: a randomised placebo-controlled trial. *Lancet*. 2010 Mar 6; 375(9717):824–33. [PubMed: 20153888]
3. Baeten J, Kahle E, Lingappa JR, et al. Genital HIV-1 RNA levels predict risk of heterosexual HIV-1 transmission. *Sci Trans Med*. 2011; 3(77):77ra29.
4. Ellerbrock TV, Chiasson MA, Bush TJ, et al. Incidence of cervical squamous intraepithelial lesions in HIV-infected women. *J Am Med Assoc*. 2000 Feb 23; 283(8):1031–7.
5. Phillips AN, Staszewski S, Weber R, et al. HIV viral load response to antiretroviral therapy according to the baseline CD4 cell count and viral load. *J Am Med Assoc*. 2001 Nov 28; 286(20): 2560–7.
6. Graham SM, Holte SE, Peshu NM, et al. Initiation of antiretroviral therapy leads to a rapid decline in cervical and vaginal HIV-1 shedding. *AIDS*. 2007 Feb 19; 21(4):501–7. [PubMed: 17301569]
7. Gupta P, Mellors J, Kingsley L, et al. High viral load in semen of human immunodeficiency virus type 1-infected men at all stages of disease and its reduction by therapy with protease and nonnucleoside reverse transcriptase inhibitors. *J Virol*. 1997; 71(8):6271–5. [PubMed: 9223532]
8. Marcelin AG, Tubiana R, Lambert-Niclot S, et al. Detection of HIV-1 RNA in seminal plasma samples from treated patients with undetectable HIV-1 RNA in blood plasma. *AIDS*. 2008 Aug 20; 22(13):1677–9. [PubMed: 18670231]
9. Cu-Uvin S, Caliendo AM. Genital tract HIV-1 RNA shedding among women with below detectable plasma viral load. *AIDS*. 2011 Mar 27; 25(6):880–1. [PubMed: 21412068]
10. Vernazza PL. Genital shedding of HIV-1 despite successful antiretroviral therapy. *Lancet*. 2001 Nov 10.358(9293):1564. [PubMed: 11716875]
11. Mofenson LM. Can perinatal HIV infection be eliminated in the United States? *J Am Med Assoc*. 1999 Aug 11; 282(6):577–9.
12. Cohen MS, Gay C, Kashuba AD, Blower S, Paxton L. Narrative review: antiretroviral therapy to prevent the sexual transmission of HIV-1. *Ann Intern Med*. 2007 Apr 17; 146(8):591–601. [PubMed: 17438318]
13. Montaner JS, Hogg R, Wood E, et al. The case for expanding access to highly active antiretroviral therapy to curb the growth of the HIV epidemic. *Lancet*. 2006 Aug 5; 368(9534):531–6. [PubMed: 16890841]

14. Attia S, Egger M, Muller M, Zwahlen M, Low N. Sexual transmission of HIV according to viral load and antiretroviral therapy: systematic review and meta-analysis. *AIDS*. 2009 Jul 17; 23(11): 1397–404. [PubMed: 19381076]
15. Donnell D, Baeten JM, Kiari J, et al. Heterosexual HIV-1 transmission after initiation of antiretroviral therapy: a prospective cohort analysis. *Lancet*. 2010 Jun 12; 375(9731):2092–8. [PubMed: 20537376]
16. Cohen MS, Chen YQ, McCauley M, et al. Prevention of HIV-1 infection with early antiretroviral therapy. *N Engl J Med*. 2011 Aug 11; 365(6):493–505. [PubMed: 21767103]
17. Granich RM, Gilks CF, Dye C, De Cock KM, Williams BG. Universal voluntary HIV testing with immediate antiretroviral therapy as a strategy for elimination of HIV transmission: a mathematical model. *Lancet*. 2009 Jan 3; 373(9657):48–57. [PubMed: 19038438]
18. Dodd PJ, Garnett GP, Hallett TB. Examining the promise of HIV elimination by ‘test and treat’ in hyperendemic settings. *AIDS*. 2010; 24(5):729–35. [PubMed: 20154580]
19. Burns DN, Dieffenbach CW, Vermund SH. Rethinking prevention of HIV type 1 infection. *Clin Infect Dis*. 2010 Sep 15; 51(6):725–31. [PubMed: 20707698]
20. Rosen S, Fox MP. Retention in HIV care between testing and treatment in sub-Saharan Africa: a systematic review. *PLoS Med*. 2011 Jul.8(7):e1001056. [PubMed: 21811403]
21. HIV Prevention Trials Network. HPTN 065 - TLC-Plus: A study to evaluate the feasibility of an enhanced test, link to care, plus treat approach for HIV prevention in the United States. Washington, D.C: National Institute of Allergy and Infectious Diseases, National Institutes of Health; 2010. [cited 2012 10 Apr]; Available from: http://www.hptn.org/research_studies/hptn065.asp
22. El-Sadr, W., editor. ART for prevention: the science and the art. 19th Conference on Retroviruses and Opportunistic Infections (CROI); 2012 March 5–8; Seattle, WA.
23. El-Sadr WM, Coburn BJ, Blower S. Modeling the impact on the HIV epidemic of treating discordant couples with antiretrovirals to prevent transmission. *AIDS*. 2011 Nov 28; 25(18):2295–9. [PubMed: 21993304]
24. World Health Organization. Guidance on couples HIV testing and counselling including antiretroviral therapy for treatment and prevention in serodiscordant couples: Recommendations for a public health approach. Geneva, Switzerland: WHO Press; 2012. Available from: <http://www.who.int/hiv/pub/guidelines/9789241501972/en/index.html>
25. Connor EM, Sperling RS, Gelber R, et al. Reduction of maternal-infant transmission of human immunodeficiency virus type 1 with zidovudine treatment. Pediatric AIDS Clinical Trials Group Protocol 076 Study Group. *N Engl J Med*. 1994 Nov 3; 331(18):1173–80. [PubMed: 7935654]
26. Chasela CS, Hudgens MG, Jamieson DJ, et al. Maternal or infant antiretroviral drugs to reduce HIV-1 transmission. *N Engl J Med*. 2010 Jun 17; 362(24):2271–81. [PubMed: 20554982]
27. Mofenson LM. Protecting the next generation--eliminating perinatal HIV-1 infection. *N Engl J Med*. 2010 Jun 17; 362(24):2316–8. [PubMed: 20554987]
28. Garcia-Lerma JG, Otten RA, Qari SH, et al. Prevention of rectal SHIV transmission in macaques by daily or intermittent prophylaxis with emtricitabine and tenofovir. *PLoS Med*. 2008 Feb. 5(2):e28. [PubMed: 18254653]
29. Denton PW, Estes JD, Sun Z, et al. Antiretroviral pre-exposure prophylaxis prevents vaginal transmission of HIV-1 in humanized BLT mice. *PLoS Med*. 2008 Jan 15.5(1):e16. [PubMed: 18198941]
30. Dumond JB, Yeh RF, Patterson KB, et al. Antiretroviral drug exposure in the female genital tract: implications for oral pre- and post-exposure prophylaxis. *AIDS*. 2007 Sep 12; 21(14):1899–907. [PubMed: 17721097]
31. Patterson KB, Prince HA, Kraft E, et al. Penetration of tenofovir and emtricitabine in mucosal tissues: implications for prevention of HIV-1 transmission. *Sci Transl Med*. 2011 Dec 7.3(112): 112re4.
32. Barditch-Crovo P, Deeks SG, Collier A, et al. Phase i/ii trial of the pharmacokinetics, safety, and antiretroviral activity of tenofovir disoproxil fumarate in human immunodeficiency virus-infected adults. *Antimicrobial agents and chemotherapy*. 2001 Oct; 45(10):2733–9. [PubMed: 11557462]

33. Parikh UM, Dobard C, Sharma S, et al. Complete protection from repeated vaginal simian-human immunodeficiency virus exposures in macaques by a topical gel containing tenofovir alone or with emtricitabine. *J Virol*. 2009 Oct; 83(20):10358–65. [PubMed: 19656878]
34. Subbarao S, Otten RA, Ramos A, et al. Chemoprophylaxis with tenofovir disoproxil fumarate provided partial protection against infection with simian human immunodeficiency virus in macaques given multiple virus challenges. *J Infect Dis*. 2006 Oct 1; 194(7):904–11. [PubMed: 16960777]
35. Cong ME, Youngpairaj AS, Zheng Q, et al. Protection against rectal transmission of an emtricitabine-resistant simian/human immunodeficiency virus SHIV162p3M184V mutant by intermittent prophylaxis with Truvada. *J Virol*. 2011 Aug; 85(15):7933–6. [PubMed: 21632769]
36. Veazey, R.; Ketas, T.; Dufour, J.; Klasse, P.; Moore, J., editors. Protection of rhesus macaques from vaginal infection by maraviroc, an inhibitor of HIV-1 entry via the CCR5 co-receptor. 17th Conference on Retroviruses and Opportunistic Infections (CROI); 2010 16–19 Feb; San Francisco, CA.
37. Dobard, C.; Sharma, S.; Parikh, U., et al., editors. High protection against vaginal infection in macaques by PEP with gel containing RAL. 18th Conference on Retroviruses and Opportunistic Infections (CROI); 2011 27 Feb–2 Mar; Boston, MA.
38. Abdool Karim Q, Abdool Karim SS, Frohlich JA, et al. Effectiveness and safety of tenofovir gel, an antiretroviral microbicide, for the prevention of HIV infection in women. *Science*. 2010 Sep 3; 329(5996):1168–74. [PubMed: 20643915]
39. Abdool Karim SS, Kashuba AD, Werner L, Abdool Karim QA. Drug concentrations after topical and oral antiretroviral pre-exposure prophylaxis: implications for HIV prevention in women. *Lancet*. 2011 Jul 16; 378(9787):279–81. [PubMed: 21763939]
40. Andrei G, Lisco A, Vanpouille C, et al. Topical Tenofovir, a Microbicide Effective against HIV, Inhibits Herpes Simplex Virus-2 Replication. *Cell Host Microbe*. 2011 Oct 4; 10(4):379–89. [PubMed: 22018238]
41. Hendrix, C.; Minnis, A.; Guddera, V., et al., editors. MTN-001: A phase 2 cross-over study of daily oral and vaginal TFV in healthy, sexually active women results in significantly different product acceptability and vaginal tissue drug concentrations. 18th Conference on Retroviruses and Opportunistic Infections; 2011 February 27 – March 2; Boston, Massachusetts. p. Abstract 35LB
42. Grant RM, Lama JR, Anderson PL, et al. Preexposure chemoprophylaxis for HIV prevention in men who have sex with men. *N Engl J Med*. 2010 Dec 30; 363(27):2587–99. [PubMed: 21091279]
43. Anderson, P.; Lama, J.; Buchbinder, S., et al., editors. Interpreting detection rates of intracellular FTC-TP and TFV-DP: the iPrEx trial. 18th Conference on Retroviruses and Opportunistic Infections (CROI); 2011 February 27–March 2; Boston, MA.
44. Liegler, T.; Abdel-Mohsen, M.; Atchison, R., et al., editors. Drug resistance and minor drug resistant variants in iPrEx. 18th Conference on Retroviruses and Opportunistic Infections (CROI); 2011 27 Feb–2 Mar; Boston, MA.
45. Anderson, P.; Liu, A.; Buchbinder, S., et al., editors. Intracellular tenofovir-DP concentrations associated with PrEP efficacy in MSM from iPrEx. 19th Conference on Retroviruses and Opportunistic Infections (CROI); 2012 Mar 5–8; Seattle, WA.
46. Liu, A.; Vittinghoff, E.; Irby, R., et al., editors. BMD loss in HIV– men participating in a TDF PrEP clinical trial in San Francisco. 18th Conference on Retroviruses and Opportunistic Infections (CROI); 2011 27 Feb–2 Mar; Boston, MA.
47. Family Health International (FHI). VOICE HIV prevention trial continues, but researchers suspend oral tenofovir arm because of futility. Vol. 360. Research Triangle Park, NC: FHI; 2011. [updated 28 September 2011; cited 2011 28 September]; Available from: http://www.fhi360.org/en/AboutFHI/Media/Releases/res_VOICE.htm
48. Van Damme, L.; Cornelli, A.; Ahmed, K., et al., editors. The FEM-PrEP trial of emtricitabine/tenofovir disoproxil fumarate (Truvada) among African women. 19th Conference on Retroviruses and Opportunistic Infections (CROI); 2012 March 5–8; Seattle, WA.
49. Anderson, P.; Meditz, A.; Zheng, J-H., et al., editors. Cellular pharmacology of tenofovir and emtricitabine in blood, rectal, and cervical cells from HIV– volunteers. 19th Conference on Retroviruses and Opportunistic Infections (CROI); 2012 Mar 5–8; Seattle, WA.

50. Thigpen, M.; Kebaabetswe, P.; Smith, D., et al., editors. Daily oral antiretroviral use for the prevention of HIV infection in heterosexually active young adults in Botswana: results from the TDF2 study. 6th IAS Conference on HIV Pathogenesis, Treatment and Prevention; 2011 July 17–20; Rome, Italy.
51. Mujugira A, Baeten JM, Donnell D, et al. Characteristics of HIV-1 Serodiscordant Couples Enrolled in a Clinical Trial of Antiretroviral Pre-Exposure Prophylaxis for HIV-1 Prevention. PLoS ONE. 2011; 6(10):e25828. [PubMed: 21998703]
52. Baeten, J.; Donnell, D.; Ndase, P., et al., editors. ARV PrEP for HIV-1 prevention among heterosexual men and women. 19th Conference on Retroviruses and Opportunistic Infections (CROI); 2012 March 5–8; Seattle, WA.
53. Bangsberg, D.; Haberer, J.; Psaros, C., et al., editors. High adherence and high effectiveness observed in HIV discordant couples: Partners PrEP Study, adherence monitoring and counseling substudy. 19th Conference on Retroviruses and Opportunistic Infections (CROI); 2012 Mar 5–8; Seattle, WA.
54. Donnell, D.; Baeten, J.; Hendrix, C., et al., editors. Tenofovir disoproxil fumarate drug levels indicate PrEP use is strongly correlated with HIV-1 protective effects: Kenya and Uganda. 19th Conference on Retroviruses and Opportunistic Infections (CROI); 2012 March 5–8; Seattle, WA.
55. National Institute of Allergy and Infectious Diseases (NIAID). NIH modifies 'VOICE' HIV prevention study in women. National Institutes of Health. 2011. [updated 28 September 2011; cited 2011 28 September]; Available from: <http://www.nih.gov/news/health/sep2011/niaid-28.htm>
56. National Institute of Allergy and Infectious Diseases (NIAID). NIH discontinues tenofovir vaginal gel in 'Voice' HIV prevention study. 2011. [updated 25 November; cited 2011 27 January]; Available from: <http://www.nih.gov/news/health/nov2011/niaid-25.htm>
57. Martin M, Vanichseni S, Suntharasamai P, et al. Enrollment Characteristics and Risk Behaviors of Injection Drug Users Participating in the Bangkok Tenofovir Study, Thailand. PLoS ONE. 2011; 6(9):e25127. [PubMed: 21969870]
58. FACTS Consortium. FACTS 001 Study Design. 2012. [cited 2012 Apr 11]; Available from: http://www.facts-consortium.co.za/?page_id=227
59. HIV Prevention Trials Network. HPTN 067 - The ADAPT Study: A phase II, randomized, open-label, pharmacokinetic and behavioral study of the use of intermittent oral emtricitabine/tenofovir disoproxil fumarate pre-exposure prophylaxis (PrEP). Washington, D.C: National Institute of Allergy and Infectious Diseases, National Institutes of Health; 2011. [cited 2012 10 Apr]; Available from: http://www.hptn.org/research_studies/hptn067.asp
60. The IPERGAY Study. anRs. 2012. [cited 2012 Apr 11]; Available from: <http://www.ipergay.fr/>
61. Grabbe KL, Bunnell R. Reframing HIV prevention in sub-Saharan Africa using couple-centered approaches. J Am Med Assoc. 2010 Jul 21; 304(3):346–7.
62. Eyawo O, de Walque D, Ford N, et al. HIV status in discordant couples in sub-Saharan Africa: a systematic review and meta-analysis. Lancet Infect Dis. 2010 Nov; 10(11):770–7. [PubMed: 20926347]
63. Coburn BJ, Gerberry DJ, Blower S. Quantification of the role of discordant couples in driving incidence of HIV in sub-Saharan Africa. Lancet Infect Dis. 2011 Apr; 11(4):263–4. [PubMed: 21453868]
64. Chemaitelly H, Cremin I, Shelton J, Hallett TB, Abu-Raddad LJ. Distinct HIV discordancy patterns by epidemic size in stable sexual partnerships in sub-Saharan Africa. Sex Transm Infect. 2012 Feb; 88(1):51–7. [PubMed: 22250180]
65. Dunkle KL, Stephenson R, Karita E, et al. New heterosexually transmitted HIV infections in married or cohabiting couples in urban Zambia and Rwanda: an analysis of survey and clinical data. Lancet. 2008 Jun 28; 371(9631):2183–91. [PubMed: 18586173]
66. Ware N, Wyatt M, Haberer J, et al. What's love got to do with it? A theory of adherence to once-daily oral pre-exposure prophylaxis (PrEP) for HIV prevention derived from qualitative data among HIV serodiscordant couples. J Acquir Immune Defic Syndr. 2012; 59(5):463–8. [PubMed: 22267018]
67. Haberer, J.; Baeten, J.; Celum, C., et al., editors. Near perfect early adherence to antiretroviral PrEP against HIV infection among HIV serodiscordant couples as determined by multiple

- measures: preliminary data from the Partners PrEP Study. 18th Conference on Retroviruses and Opportunistic Infections (CROI); 2011 27 Feb-2 Mar; Boston, MA.
68. Curran K, Baeten J, Coates T, et al. HIV-1 Prevention for HIV-1 Serodiscordant Couples. *Curr HIV/AIDS Reports*. In Press.
69. Hallett T, Baeten J, Heffron R, et al. Optimal uses of antiretrovirals for prevention in HIV-1 serodiscordant heterosexual couples in South Africa: a modelling study. *PLoS ONE*. 2011; 8(11):e1001123.
70. Walensky, R., editor. Cost-effectiveness in HIV care: understanding value in a world of limited resources. 18th Conference on Retroviruses and Opportunistic Infections (CROI); 2011 27 Feb-2 Mar; Boston, MA.
71. Hallett TB, Baeten JM, Heffron R, et al. Optimal Uses of Antiretrovirals for Prevention in HIV-1 Serodiscordant Heterosexual Couples in South Africa: A Modelling Study. *PLoS Med*. 2011 Nov. 8(11):e1001123. [PubMed: 22110407]
72. Hurt CB, Eron JJ Jr, Cohen MS. Pre-exposure prophylaxis and antiretroviral resistance: HIV prevention at a cost? *Clin Infect Dis*. 2011 Dec; 53(12):1265–70. [PubMed: 21976467]
73. Romano J, Variano B, Coplan P, et al. Safety and availability of dapivirine (TMC120) delivered from an intravaginal ring. *AIDS Res Hum Retroviruses*. 2009 May; 25(5):483–8. [PubMed: 19388819]
74. UNAIDS. UNAIDS data tables: 2011. Geneva: Joint United Nations Programme on HIV/AIDS (UNAIDS); 2011. [cited 2012 10 Apr]; Available from: http://www.unaids.org/en/media/unaids/contentassets/documents/unaidspublication/2011/JC2225_UNAIDS_datatables_en.pdf
75. Cates W. HPTN 052 and the future of HIV treatment and prevention. *Lancet*. 2011 Jul 16; 378(9787):224–5. [PubMed: 21763928]
76. Baeten, J.; Celum, C., editors. Antiretroviral pre-exposure prophylaxis for HIV-1 prevention among heterosexual African men and women: the Partners PrEP Study. 6th IAS Conference on HIV Pathogenesis, Treatment and Prevention; 2011 July 17–20; Rome, Italy. p. Abstract MOAX0106

Figure 1. Treatment as Prevention (HPTN 052) and tenofovir-based topical and oral PrEP: HIV prevention efficacy from completed clinical trials

Figure does not include oral tenofovir data from VOICE, as data have not been released

Table 1

Key studies of HIV-1 prevention benefits of ART

Study	Population	N	HIV-1 transmission outcome associated with ART use	References
Meta-analysis of 5 observational studies	Heterosexual HIV-1 serodiscordant couples from 11 cohorts from 1996–2009	5021 couples (461 HIV-1 transmission events, not described whether virologically-linked)	92% decrease in HIV-1 transmission risk for couples in which the HIV-1 infected partner was receiving ART (5.64 compared to 0.46 transmissions per 100 person-years)	[14]
Partners in Prevention HSV/HIV Transmission Study	Observational analysis from a prospective study of HIV-1 serodiscordant couples (Botswana, Kenya, Rwanda, South Africa, Tanzania, Uganda, Zambia) from 2004–2008	3381 couples (103 virologically-linked HIV-1 transmissions)	92% decrease in HIV-1 transmission risk for couples in which the HIV-1 infected partner was receiving ART (2.24 compared to 0.37 transmissions per 100 person-years)	[15]
HPTN 052	Randomized trial of early (CD4 350–550 cells/mm ³) versus routine (CD4 250 cells/mm ³) initiation of ART in HIV-1 serodiscordant couples (Botswana, Brazil, India, Kenya, Malawi, South Africa, Thailand, U.S. Zimbabwe) from 2007–2010	1763 (28 virologically-linked HIV-1 transmissions)	96% decrease in HIV-1 transmission risk in couples assigned to receive early ART (versus routine ART initiation) (from 0.9 to 0.1 per 100 person-years)	[15]

Table 2

Efficacy trials of topical and oral PrEP

Study (location)	Population	N	PrEP Agent	Status	References
CAPRISA 004 (South Africa)	Women	889	Vaginal tenofovir gel (coitally-associated use)	39% reduction in HIV-1 incidence [38]	[38]
iPrEx (Brazil, Ecuador, Peru, South Africa, Thailand, US)	Men who have sex with men and transgender women	2499	FTC/TDF	44% reduction in HIV-1 incidence [42]	[42]
FEM-PrEP (Kenya, South Africa, Tanzania)	Higher-risk women	1950	FTC/TDF	Stopped for lack of efficacy April 2011 [47]	[47, 48]
Partners PrEP Study (Kenya, Uganda)	HIV-1 serodiscordant couples	4758	TDF, FTC/TDF	67% reduction in HIV-1 incidence for TDF 75% reduction in HIV-1 incidence for FTC/TDF [76]	[52]
TDF2 Study (Botswana)	Heterosexual men and women, ages 18–35	1200	FTC/TDF	62% reduction in HIV-1 incidence [50]	[50]
VOICE/MTN 003 (South Africa, Uganda, Zimbabwe)	Women	5021	TDF, FTC/TDF, Vaginal tenofovir gel (daily use)	Oral TDF stopped for lack of efficacy September 2011 [55] Vaginal tenofovir gel arm stopped November 2011 [56]	[55] [56]
Bangkok Tenofovir Study (Thailand)	Injection drug users	2400	TDF	Ongoing [57]	[57]
FACTS 001 (South Africa)	Women	2600	Vaginal tenofovir gel (coitally-associated use)	Initiated October 2011	[58]