Published in final edited form as:

Org Biomol Chem. 2012 October 21; 10(39): 7980–7985. doi:10.1039/c2ob26571k.

Versatile approach to α -alkoxy carbamate synthesis and stimulus-responsive alcohol release†

R. Adam Mosey and Paul E. Floreancig*

Department of Chemistry, University of Pittsburgh, Pittsburgh, Pennsylvania 15260, USA

Abstract

A series of α -alkoxy carbamates that cleave under mild conditions to release alcohols has been synthesized through a multicomponent process. The relationship between structural features in these compounds and the rate of alcohol release in the presence of basic hydrogen peroxide has been studied. The preparation of carbamates that cleave under other conditions has been demonstrated.

Introduction

Stimulus-responsive cleavage reactions have found broad use for applications in temporally-controlled drug release, chemical sensing, and material decomposition processes.¹ Carbamates have frequently been used to release amines in these processes due to the diverse array of stimuli that can be employed for the decomposition step. We have initiated a program by which α -alkoxy carbamate cleavage can serve to release alcohols in response to externally-applied stimuli, as shown in Scheme 1.

The capacity to direct drug release at a particular location would be a significant attribute for a drug-delivery agent. The benefit of employing α -alkoxy carbamates as delivery agents² is that the carbamate structure contains a branch that can be used to append cell-targeting agents. Thus a single molecule can contain directing, triggering, and cargo groups (Fig. 1). This design has been implemented in elegant studies from the Shabat,³ Shin,⁴ and Sinha⁵ groups. These successful constructs highlight the utility of the design yet reveal that enhanced synthetic accessibility would be desirable for rapid property optimization. In this manuscript we demonstrate that several α -alkoxy carbamates can be prepared rapidly through a multicomponent reaction and illustrate the manner in which this protocol can be used to access compounds that release alcohols at varying rates. Compounds with different triggering mechanisms have been prepared to demonstrate the merits of the multicomponent reaction for the preparation of these structures.

Results and discussion

Our approach to α -alkoxy carbamate synthesis employed a sequence of nitrile hydrozirconation, acylation, and alcohol addition.⁶ This protocol has successfully been applied to rapid syntheses of cyclic amides,⁷ the natural products pederin and psymberin, and many analogs.⁸ Conducting the multicomponent reaction at a late stage in the synthetic

†Electronic supplementary information (ESI) available. See DOI: 10.1039/c2ob26571k

© The Royal Society of Chemistry 2012

florean@pitt.edu; Fax: +1 412-624-8626; Tel: +1 412-624-8727.

sequence provides abundant opportunities for optimizing molecular properties through structural diversification.

The synthesis of a representative cleavable α -alkoxy carbamate is shown in Scheme 2. We chose to explore peroxide-triggered compounds in consideration of the potential for drug delivery to oxidant-rich medical conditions such as cancer⁹ and neurodegeneration.¹⁰ Our design was based on Chang's peroxide-released fluorophores¹¹ in which boronate oxidation forms a phenoxide that releases its cargo through *p*-quinone methide formation.¹² Thus, known benzylic alcohol **1** was treated with a solution of phosgene in toluene to form chloroformate **2**. Hydrozirconation of ^tBuCN (**3**) with Cp₂Zr(H)Cl followed by the addition of freshly-prepared **2** yielded intermediate acylimine **4**. The synthesis was completed through the addition of neopentyl alcohol to provide **5** in 64% yield.

The breakdown of compound **5** was effected by treatment with urea-H₂O₂ at pH = 8.0 (phosphate buffer) in a mixture of CD₃CN and D₂O (Scheme 3). This allowed for direct monitoring of starting material consumption and product formation by ¹H NMR. This experiment showed that the oxidation of the boronate to phenoxide **6** was rapid, as indicated by the subtle change in the chemical shifts of the diastereotopic hydrogens of the neopentyl ether group after 1 min (Fig. 2). Acyl amination proceeded through quinone methide loss to form **7**, decarboxylation to yield **8**, and amination breakdown to yield neopentyl alcohol and imine **9**. The imine was not observed in this reaction due to its rapid reaction with water to yield the corresponding aldehyde. Aldehydes oxidize rapidly with basic hydrogen peroxide, making the loss of this component difficult to monitor in this experiment. Under different conditions, however, this decomposition process could be used to release alcohol and aldehyde cargo through a single triggering event. Alcohol release was slower than phenoxide formation, indicating that oxidation is not the rate-determining step when high peroxide concentrations are employed for the breakdown.

Neopentyl alcohol was released at a similar rate to the decomposition of the carbamate (Fig. 2), as determined by integrating the growing methylene singlet from the alcohol and the AB system from the carbamate and comparing the intensities to THF, which was added as an internal standard. The yield of neopentyl alcohol was approximately 60% as determined by ¹H NMR, but subsequent experiments that released less volatile alcohols showed that isolated yields are much higher (see below), suggesting that yield determination can be influenced by relaxation rate differences in comparison to the internal standard. Therefore determinations of the half lives for product formation provides a valid comparison for determining the relative reactivities of different analogs. We used the conversion at 35 min as the end point for *t*_{1/2} determinations. Using conversion levels at later times does not significantly change the values. The half life for the release of neopentyl alcohol from **5** was 12 min.

This facile approach to α -alkoxy carbamate synthesis allowed us to prepare a number of analogous structures (Fig. 3)¹³ in accord with our objective of determining the influence of structural perturbations in the triggering group, the nitrile group, and the alcohol on the rate of peroxide-mediated alcohol release. Steric hindrance around the leaving group was altered by preparing **10** from acetonitrile, while the electronics of the nitrile component were altered by synthesizing **11** and **12** from the corresponding electron-poor and electron-rich aromatic nitriles. The carbamate group was altered through the incorporation of a fluorine, a methoxy group, and a benzylic methyl branch (**13–15**). Carbamate **16** was prepared to study the impact of increasing the leaving group ability of the cargo. The selection of alcohols in these analogs was based on the presence of methylene groups that appear as AB quartets in the ¹H NMR spectra of the carbamates then collapse to singlets upon release, thereby allowing decomposition to be monitored readily. Vinyl boronate **17** was prepared from the

corresponding allylic chloroformate to determine the relative merits of utilizing the β -elimination of an enolate intermediate¹⁴ to promote acyl aminal breakdown. Benzyl carbamate **18** was synthesized as a control compound to verify that alcohol release can be attributed to boronate oxidation.

The versatility of the synthetic protocol was further probed through the preparation of a number of additional structures that contain different triggering groups. Acetoxy carbamate **19** was prepared for cargo release in response to esterase-mediated cleavage. Azido carbamate **20** was prepared for cargo release in thioredoxin-rich environments. Nitro-substituted carbamate **21** was prepared for cargo release in the presence of nitrido reductase. The latter two compounds were designed for release in hypoxic environments that are associated with many forms of cancer.¹⁵

We exposed compounds **10**, **11**, and **12** to the oxidative fragmentation conditions to determine the influence of the group on the nitrile fragment on the rate of alcohol release. As shown in Fig. 4 very little difference was observed for these compounds, with **10** showing a $t_{1/2}$ of 9 min, **11** showing a $t_{1/2}$ of 12 min, and **12** showing a $t_{1/2}$ of 9 min. The similarities of these numbers contrasts with the significant steric and electronic differences between the structures and demonstrates that the rate determining step in this process is not the release of the neopentyl alcohol from a tetrahedral intermediate similar to **8**. The stability of control compound **18** toward oxidative fragmentation is also confirmed in Fig. 4. This demonstrates that the nitrile-derived subunit of these structures can be selected to facilitate the synthesis or to optimize a different property without consequence on the release rate.

The next phase of this project was to determine the impact of perturbations to the alkoxy group of the carbamate on the rate of alcohol release. The peroxide-mediated breakdowns of **13**, **14**, and **15** were compared to **5** (Fig. 5). While the incorporation of a fluorine atom to the benzene ring did not result in a significant change in the rate of alcohol release ($t_{1/2}$ for **13** = 11 min), the addition of a methoxy group to the arene provided a substantial rate increase, with a $t_{1/2}$ for **14** being 2 min. The addition of a methyl group at the benzylic site resulted in a noticeable rate increase, with a $t_{1/2}$ of 6 min for **15**. These results are consistent with benzyl cleavage being the rate determining step in this process.¹⁶ The addition of a branching alkyl group at the benzylic position and the incorporation of an electron-donating methoxy group to the arene facilitate the departure of the carboxyl group. These observations show that factors that stabilize a benzylic cation will promote breakdown. The negligible impact of incorporating a fluorine into the carbamate is therefore somewhat surprising since this substitution should inductively blunt the electron-donating capacity of the phenol that forms from boronate oxidation. We propose that this inductive stabilization can be balanced by the higher concentration of the more reactive phenoxide that results from the lower pK_a of the fluorinated analog. The decomposition rates for **5** and **14** did not vary significantly when the substrate concentration was 3 mM and $[H_2O_2]$ was 30 mM (see ESI[†]).

The influence of alcohol nucleofugacity on the cleavage rate was determined by comparing the rates of peroxide-mediated breakdowns of **5** and aryl ether **16**. These processes were shown to proceed with essentially identical rates ($t_{1/2}$ = 10 min for **16**), as shown in Fig. 6, despite the greater stability of the phenoxide leaving group in comparison to an alkoxide leaving group. This outcome is consistent with the hypothesis that the facility of benzylic carbon–oxygen bond cleavage, and not tetrahedral intermediate breakdown, determines the rate of alcohol release. The decomposition of **16** appears to proceed in higher yield than the

[†]Electronic supplementary information (ESI) available. See DOI: 10.1039/c2ob26571k

decomposition of **5**, though this can be attributed to a difference in ^1H relaxation rates in these NMR-derived plots.

Allylic carbamate **17** differs from the previous compounds in that its peroxide-mediated oxidation forms an enolate rather than a phenoxide. β -Elimination leads to cargo release and the formation of acrolein. The breakdown of this compound, in comparison to rapidly reacting substrate **14** and parent structure **5**, is shown in Fig. 7. This process is extremely fast, with a $t_{1/2}$ of <2 min, showing that this structural variant will be useful for applications where extremely rapid cargo release is desirable. We prepared carbamate **22** in an effort to determine the yield of the released alcohol.¹³ The alkoxy group in **22** was selected to be sufficiently non-volatile and UV-active to facilitate isolation. Exposing **22** to the standard cleavage conditions (Scheme 4) for 1 h provided alcohol **23** in an 82% isolated yield, demonstrating that cargo release proceeds more effectively than the ^1H NMR experiments would indicate.

This method was applied to the preparation of a compound that is designed to deliver an anti-oxidant to mitochondria¹⁷ in an effort to prepare agents that mitigate radiation damage by suppressing the formation of reactive oxygen species.¹⁸ The design of the compound and its synthesis are shown in Scheme 5. The boronate carbamate group was employed to trigger compound release in the presence of the high peroxide concentrations in mitochondria.¹⁹ Pentamethyl chromanol (PMC, **24**) was selected for release due to its ability to mitigate the effects of cytochrome *c*-mediated mitochondrial damage.²⁰ The triphenylphosphonium ion was included to promote passage through mitochondrial membranes.²¹ Hydrozirconation of nitrile **25** followed by acylation with **2** and nucleophilic addition of **24** provided carbamate **26** in 58% yield. Removal of the silyl group followed by acylation with commercially available acid **27** provided **28** in good yield. The rapid synthesis of this compound shows that a more stable linkage to the phosphonium ion could be incorporated if the ester linkage is susceptible to esterase cleavage in cellular studies. Benzyl carbamate **29** was prepared as a control compound for comparison in breakdown studies.

The peroxide-mediated breakdown of **28** to release PMC was studied in aqueous media (phosphate buffer containing 5% EtOH) at pH 7.2 and 8.0 (the pH of the mitochondrial matrix).²² These studies were performed with micromolar concentrations of the carbamate and H_2O_2 at 37 °C to mimic biological conditions. These concentrations required that the reactions be monitored by HPLC using an electrochemical detector. Treating **28** (40 μM) with varying concentrations of H_2O_2 (10, 20, and 40 μM) resulted in the liberation of PMC in a dose-dependant manner (Fig. 8). The release rate was notably accelerated when the reaction was performed at higher pH. A pH = 8.0 reaction mixture containing 40 μM of **28** and H_2O_2 generated approximately 20 μM of PMC within 30 min, whereas the reaction at pH = 7.2 required 90 min to reach a similar concentration. The necessity of the boronate ester for carbamate breakdown and alcohol release was also demonstrated as PMC accumulation was not observed following exposure of **29** to H_2O_2 and the small amounts of PMC that were detected were not distinguishable from noise. The release of PMC from **28** proceeded much more slowly than the release of neopentyl alcohol from **5**. This can be attributed to the rate reduction for boronate oxidation that results from the low concentrations. The rate of alcohol release is dependent on the rate of quinone methide formation, and the rate of quinone methide formation is dependent upon the concentration of the phenoxide intermediate that forms through a bimolecular reaction between the substrate and the hydroperoxide anion, thereby explaining the low release rates in these reactions.

Conclusions

We have shown that the multicomponent sequence of nitrile hydrozirconation, acylation, and alcohol addition can be used for the rapid preparation of structurally diverse α -alkoxy carbamates. These carbamates have been designed to release the alcohol component in response to a precise chemical signal as an entry into the development of drug delivery vehicles. The utility of this approach has been demonstrated in a quick study of the relationship between structural features and the rate of peroxide-mediated alcohol release from boronate-substituted benzylic carbamates. This work showed that the release rate is controlled by the rate of quinone methide formation rather than boronate oxidation or tetrahedral intermediate collapse. Carbamates that cleave in the presence of nucleophiles and reducing agents have also been prepared to highlight the generality of the pathway. A complex that contains a triphenyl phosphonium ion for mitochondrial transport, a boronate-substituted benzylic carbamate as a peroxide-mediated trigger, and a chromanol derivative has been prepared for the delivery of anti-oxidants for studies on the directed mitigation of damage from reactive oxygen species. The ability of **28** and related structures to act as radiation mitigators is currently under investigation and the results will be reported elsewhere. We anticipate that the facile preparation of structurally diverse agents that can release cargo under defined conditions will prove to be useful for applications in targeted drug delivery and for controlled tailoring of materials.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by generous funding from the National Science Foundation (CHE-0848299) and the National Institutes of Health (AI068021). We thank Ms Stephanie Garrell and Mr Akira Shimizu (recipient of a Brackenridge undergraduate research fellowship) for assistance in chloroformate synthesis. We thank Dr Detcho Stoyanovsky (Department of Occupational Health and Safety, University of Pittsburgh) for helpful discussions and technical assistance.

Notes and references

- (a) Blencowe CA, Russell AT, Greco F, Hayes W, Thornthwaite DW. *Polymer Chem.* 2011; 2:773.
(b) Shabat D, Amir RJ, Gopin A, Pessah N, Shamis M. *Chem.-Eur. J.* 2004; 10:2626. [PubMed: 15195294]
- For examples of alcohol and thiol release through *N*-acyl aminal cleavage, see: Böhm G, Dowden J, Rice DC, Burgess I, Pilard J-F, Guilbert B, Haxton A, Hunter RC, Turner NJ, Flitsch SL. *Tetrahedron Lett.* 1998; 39:3819. Meyer Y, Richard J-A, Delest B, Noack P, Renard P-Y, Romieu A. *Org. Biomol. Chem.* 2010; 8:1777. [PubMed: 20449478]
- Gopin A, Pessah N, Shamis M, Rader C, Shabat D. *Angew. Chem., Int. Ed.* 2003; 42:327.
- Lee M-R, Baek K-H, Jin HJ, Jung Y-G, Shin I. *Angew. Chem., Int. Ed.* 2004; 43:1675.
- Abraham S, Guo F, Li L-S, Rader C, Liu C, Barbas CF III, Lerner RA, Sinha SC. *Proc. Natl. Acad. Sci. U. S. A.* 2007; 104:5584. [PubMed: 17372220]
- (a) Wan S, Green ME, Park J-H, Floreancig PE. *Org. Lett.* 2007; 9:5385. [PubMed: 18020344] (b) DeBenedetto MV, Green ME, Wan S, Park J-H, Floreancig PE. *Org. Lett.* 2009; 11:835. [PubMed: 19152262]
- (a) Xiao Q, Floreancig PE. *Org. Lett.* 2008; 10:1139. [PubMed: 18278930] (b) Lu C, Xiao Q, Floreancig PE. *Org. Lett.* 2010; 12:5112. [PubMed: 20961073]
- (a) Wu F, Green ME, Floreancig PE. *Angew. Chem., Int. Ed.* 2011; 50:1131. (b) Wan S, Wu F, Rech JC, Green ME, Balachandran R, Horne WS, Day BW, Floreancig PE. *J. Am. Chem. Soc.* 2011; 133:16668. [PubMed: 21902245]

9. Fruehauf JP, Meysken FL Jr. Clin. Cancer Res. 2007; 13:789. [PubMed: 17289868]
10. DiMauro S, Schon EA. Annu. Rev. Neurosci. 2008; 31:91. [PubMed: 18333761]
11. (a) Lippert AR, Gschneidner T, Chang CJ. Chem. Commun. 2010; 46:7510. (b) Srikun D, Miller EW, Domaille DW, Chang CJ. J. Am. Chem. Soc. 2008; 130:4596. [PubMed: 18336027]
12. For other examples of peroxide-mediated fragmentation reactions of aryl boronates, see: Lippert AR, Van de Bittner GC, Chang CJ. Acc. Chem. Res. 2011; 44:793. [PubMed: 21834525] Major Jourden JL, Cohen SM. Angew. Chem., Int. Ed. 2010; 49:6795. Broaders KE, Grandhe S, Fréchet JMJ. J. Am. Chem. Soc. 2011; 133:756. [PubMed: 21171594] Sella E, Shabat D. Chem. Commun. 2008:5701. Nuñez SA, Yeung K, Fox NS, Phillips ST. J. Org. Chem. 2011; 76:10099. [PubMed: 22092043] Karton-Lifshin N, Segal E, Omer L, Portnoy M, Satchi-Fainaro R, Shabat D. J. Am. Chem. Soc. 2011; 133:10960. [PubMed: 21631116]
13. Please see the ESI[†] for details on the syntheses of these compounds.
14. Song F, Watanabe S, Floreancig PE, Koide K. J. Am. Chem. Soc. 2008; 130:16640.
15. Wilson WR, Hay MP. Nat. Rev. Cancer. 2011; 11:393. [PubMed: 21606941]
16. For similar fragmentation rate effects, see: Hay MP, Sykes BM, Denny WA, O'Connor CJ. J. Chem. Soc., Perkin Trans. 1. 1999:2759. Schmid KM, Jensen L, Phillips ST. J. Org. Chem. 2012; 77:4363. [PubMed: 22494313]
17. Hoye AT, Davoren JE, Wipf P, Fink MP, Kagan VE. Acc. Chem. Res. 2008; 41:87. [PubMed: 18193822]
18. (a) Zhao K, Zhao G-M, Wu D, Soong Y, Birk AV, Schiller PW, Szeto HH. J. Biol. Chem. 2004; 279:34682. [PubMed: 15178689] (b) Kanai A, Zabbarova I, Amoscato A, Epperly M, Xiao J, Wipf P. Org. Biomol. Chem. 2007; 5:307. [PubMed: 17205174]
19. Balaban RS, Nemeto S, Finkel T. Cell. 2005; 120:483. [PubMed: 15734681]
20. (a) Samhan-Arias AK, Tyurina YY, Kagan VE. J. Clin. Biochem. Nutr. 2011; 48:91. [PubMed: 21297919] (b) Staniek K, Rosenau T, Gregor W, Nohl H, Gille L. Biochem. Pharmacol. 2005; 70:1361. [PubMed: 16150421]
21. (a) Smith RAJ, Porteous CM, Coulter CV, Murphy MP. Eur. J. Biochem. 1999; 263:709. [PubMed: 10469134] (b) Muratovska A, Lightowlers RN, Taylor RW, Wilce JA, Murphy MP. Adv. Drug Delivery Rev. 2001; 49:189.
22. (a) Chacon E, Reece JM, Nieminen A-L, Zahrebelski G, Herman B, Lemasters JJ. Biophys. J. 1994; 66:942. [PubMed: 8038398] (b) Llopis J, McCaffery JM, Miyawaki A, Farquhar MG, Tsien RY. Proc. Natl. Acad. Sci. U. S. A. 1998; 95:6803. [PubMed: 9618493]

Fig. 1. Schematic of a cargo-releasing composite structure that contains localization and triggering groups.

Fig. 2. Time dependent release of neopentyl alcohol from **5**. The AB pattern arises from **5** and the growing singlet arises from neopentyl alcohol. See Scheme 3 for experimental conditions.

Fig. 3.
Structurally-varied carbamates.

Fig. 4. Comparison of the rates of alcohol release from structurally-varied nitrile precursors. See Scheme 3 for experimental conditions. Experiments were run in triplicate.

Fig. 5.
Impact of carbamate variation on alcohol release. See Scheme 3 for experimental conditions.
Experiments were run in triplicate.

Fig. 6.
Release as a function of alcohol structure. See Scheme 3 for experimental conditions.
Experiments were run in triplicate.

Fig. 7. Release from the allylic carbamate. See Scheme 3 for experimental conditions. Experiments were run in triplicate.

Fig. 8. PMC release from **28** in response to variations in pH and H₂O₂ concentration. Reactions were performed using 40 μM of **28** or **29** in 95% aqueous phosphate buffer/5% EtOH at 37 °C. Experiments were run in triplicate.

Scheme 1. Stimulus-promoted cleavage of α -alkoxy carbamates.

Scheme 2.
Multicomponent approach to α-alkoxy carbamate synthesis.

Scheme 3. Peroxide-mediated alcohol release. $[\mathbf{5}]_0 = 6 \text{ mM}$, $[\text{H}_2\text{O}_2]_0 = 90 \text{ mM}$, $T = 300 \text{ K}$.

Scheme 4.
Determination of an isolated yield.

Scheme 5.
Application to the synthesis of a potential mitigator of radiation damage.