Published in final edited form as:

Neurobiol Dis. 2012 December ; 48(3): 546–555. doi:10.1016/j.nbd.2012.08.002.

Thymosin β 4 Promotes the Recovery of Peripheral Neuropathy in Type II Diabetic Mice

Lei Wang¹, Michael Chopp^{1,2}, Alexandra Szalad¹, Zhongwu Liu¹, Mei Lu¹, Li Zhang¹, Jing Zhang¹, Rui Lan Zhang¹, Dan Morris¹, and Zheng Gang Zhang¹

¹Department of Neurology, Henry Ford Hospital, 2799 W. Grand Boulevard, Detroit, Michigan 48202

²Department of Physics, Oakland University, Rochester, Michigan 48309

Abstract

Peripheral neuropathy is one of the most common complications of diabetes mellitus. Using a mouse model of diabetic peripheral neuropathy, we tested the hypothesis that thymosin β 4 (T β 4) ameliorates diabetes-induced neurovascular dysfunction in the sciatic nerve and promotes recovery of neurological function from diabetic peripheral neuropathy. T β 4 treatment of diabetic mice increased functional vascular density and regional blood flow in the sciatic nerve, and improved nerve function. T β 4 upregulated angiopoietin-1 (Ang1) expression, but suppressed Ang2 expression in endothelial and Schwann cells in the diabetic sciatic nerve. In vitro, incubation of Human Umbilical Vein Endothelial Cells (HUVECs) with T β 4 under high glucose condition completely abolished high glucose-downregulated Ang1 expression and high glucose-reduced capillary-like tube formation. Moreover, incubation of HUVECs under high glucose with conditioned medium collected from Human Schwann cells (HSCs) treated with T β 4 significantly reversed high glucose-decreased capillary-like tube formation. PI3K/Akt signaling pathway is involved in T β 4-regulated Ang1 expression on endothelial and Schwann cells. These data indicate that T β 4 likely acts on endothelial cells and Schwann cells to preserve and/or restore vascular function in the sciatic nerve which facilitates improvement of peripheral nerve function under diabetic neuropathy. Thus, T β 4 has potential for the treatment of diabetic peripheral neuropathy.

Keywords

T β 4; peripheral neuropathy; diabetes; mice

INTRODUCTION

Peripheral neuropathy is one of the most common and disabling complications of diabetes mellitus. Studies of diabetic peripheral neuropathy from experimental animals and humans indicate that the development of diabetic neuropathy is closely associated with marked neurovascular dysfunction (Tefaye et al., 1993, Cameron et al., 2001, Ebenezer et al., 2011). Vascular dysfunction precedes the appearance of nerve conduction velocity deficits, leading to nerve damage (Cameron and Cotter, 1999, Cameron et al., 2005, Ebenezer et al., 2011). Therapies targeting neurovascular function have been shown to restore nerve

Please send all correspondence to: Dr. Zheng Gang Zhang Department of Neurology, Henry Ford Hospital, 2799 West Grand Boulevard, Detroit, MI 48202, Tel: 313-916-5456, Fax 313-916-1318, zhazh@neuro.hfh.edu.

Conflict of Interest: None

function in experimental diabetic peripheral neuropathy (Schratzberger et al., 2001, Kusano et al., 2004, Li et al., 2005).

Thymosin Beta4 (Tβ4), a small 4.9 kDa polypeptide of 43 amino acids, is a major intracellular G-actin-sequestering peptide (Goldstein et al., 1966) and is present in almost all cell types (Crockford et al., 2010). Tβ4 has multiple biological functions that include promoting diabetic wound healing by repairing and regenerating damaged tissues (Philp et al., 2003) and enhancing angiogenesis after myocardial infarction and vasculogenesis during development (Crockford, 2007, Smart et al., 2007a). Tβ4 is currently under a phase II clinic trial for the treatment of patients with acute myocardial infarction (ClinicalTrials.gov, <http://clinicaltrials.gov/ct2/show/NCT01311518>, Ruff et al., 2010). In the central nervous system, preclinical studies show that Tβ4 has neuroprotective and neurorestorative effects by reducing neuronal damage and enhancing oligodendrogenesis, which lead to improvement of neurological outcomes after stroke, traumatic brain injury and multiple sclerosis (Philp et al., 2003, Zhang et al., 2009, Morris et al., 2010, Xiong et al., 2011). However, the effect of Tβ4 on peripheral nerves, such as diabetic peripheral neuropathy, has not been investigated.

The angiopoietins (Ang1 and Ang2) and their receptor Tie-2 regulate vascular development and homeostasis (Suri et al., 1996, Teichert-Kuliszewska et al., 2001). Ang-1 promotes vascular stabilization and maturation whereas Ang2 acts as a partial agonist or antagonist of Ang1 signaling, depending on vascular endothelial growth factor (VEGF) bioavailability (Suri et al., 1996, Maisonpierre et al., 1997, Thebaud et al., 2005). The Ang/Tie2 signaling pathway plays an important role in mediating vascular function under diabetes (Chen and Stinnett, 2008b, Tuo et al., 2008). Hyperglycemia downregulates Ang1 and upregulates Ang2, which exacerbates myocardial infarction (Tuo et al., 2008, Morris et al., 2010). Increases in Ang1 levels normalize diabetes induced immature vasculature (Chen and Stinnett, 2008a). Patients with peripheral diabetic neuropathy have elevated levels of circulating Ang2 (Rasul et al., 2011). However, the effect of the Ang/Tie2 signaling pathway on peripheral diabetic neuropathy has not been extensively studied.

In the present study, using a mouse model of type II diabetes, we tested the hypothesis that treatment of diabetic peripheral neuropathy with Tβ4 ameliorates neurovascular dysfunction and improves peripheral nerve function. In addition, we investigated the effect of Tβ4 on the Ang/Tie2 signaling pathway under conditions of diabetic peripheral neuropathy.

MATERIAL AND METHODS

Animals

All experimental procedures were carried out in accordance with NIH Guide for the Care and Use of Laboratory Animals and approved by the institutional Animal Care and Use Committee of Henry Ford Hospital. Male BKS.Cg-*m*⁺/⁺*Lepr*^{db}/*J* (db/db) mice (Jackson Laboratories, Bar Harbor, Maine) aged 20 weeks were used. Age-matched heterozygotes mice (db/m), a non-penetrant genotype (Jackson Laboratories), were used as the control animals.

Tβ4 treatment

db/db mice at age 20 weeks were treated with Tβ4 at a dose of 6 mg/kg or 24 mg/kg (RegeneRx, Inc, intraperitoneal injection, i.p.), every 3 days for 4 weeks (n=10/group). db/db mice (n=10/group) at the same age treated with same volume of saline were used as a control group. Age-matched db/m mice treated with Tβ4 (6 mg/kg i.p. every 3 days, n=10/group) or saline (n=10/group) were used as additional control groups. All mice were sacrificed 8 weeks after the initial treatment. Doses of Tβ4 were selected based on published studies (Mora et al., 1997, Morris et al., 2010).

Blood glucose levels were measured from the mouse tail vein by using an instant check meter (Roche Diagnostics, Indianapolis, IN). Blood glucose levels, body weight and functional tests were measured at baseline (before treatment) and then every 2 weeks until sacrifice. Electrophysiological measurements were performed before treatment and then every 4 weeks until sacrifice. All procedures and analyses were performed by investigators who were blinded to the treatment administered.

Measurement of regional sciatic nerve blood flow by laser Doppler flowmetry

Regional sciatic nerve blood flow was measured at the end of the experiments (8 weeks after the initial treatment) using laser Doppler flowmetry (LDF PeriFlux PF4, Perimed AB, Järfälla, Sweden) (Zhang et al., 1997). Briefly, under anesthesia (ketamine/xylazine, i.p., 100/10 mg/kg, JHP Pharmaceuticals LLC, MI; LLOYD Inc. Iowa), the mouse was mounted on a Kopf stereotaxic apparatus. The left sciatic nerve was exposed in the mid-thigh region and animal rectal temperature was kept at $37 \pm 1.0^\circ\text{C}$ during the measurement period using a feedback controlled water bath. Using a micromanipulator, a LDF probe was placed at the surface of the sciatic nerve and relative flow values expressed as perfusion units were recorded every 5 minutes for total of 3 records. Regional sciatic nerve blood flow values from db/m mice were used as base line values and data are presented as a percentage of baseline values.

Neurophysiological Measurements

Sciatic nerve conduction velocity was assessed with orthodromic recording techniques, as previously described (Schratzberger et al., 2000, Li et al., 2005, Wang et al., 2011a). Briefly, Mice were anesthetized with ketamine/xylazine (i.p., 100/10 mg/kg). The stimulating electrodes were plated at the knee and sciatic notch. Trigger single square wave current pulses were delivered using an isolated pulse stimulator (Model 2100), A-M Systems, Everett, WA). The simultaneous electromyographies were recorded by two sterilized electrodes placed in the dorsum of the foot with a Grass Amplifier (Model P5, Grass Instruments, Quincy, MA). During the measurements, animal rectal temperature was kept at $37 \pm 1.0^\circ\text{C}$ using a feedback controlled water bath. Motor nerve conduction velocity (MCV) and sensory nerve conduction velocity (SCV) were calculated according to a published study (Li et al., 2005).

Tail-flick and hot plate tests

To examine thermal hyperalgesia, tail-flick test (the water immersion method) and hot plate test (the IITC Hot Plate Analgesia test) were employed according to published methods (Janssen et al., 1963, South and Smith, 1998, Vanderah et al., 2001, Wang et al., 2011a). Briefly, for tail-flick test, a mouse was restrained in a conical polypropylene tube with an opening through which its tail was exposed. Approximately 2 cm of the mouse's tail was immersed into a $52^\circ\text{C} \pm 0.2$ water bath and the time until the rodent flicks or removes its tail was recorded (Vanderah et al., 2001). For hot plate test, a mouse was placed within a plexiglass chamber on a transparent glass surface and allowed to acclimate for at least 20 min. A thermal stimulation meter (IITC Model 39 Hot Plate Analgesia Meter, IITC Life Science, CA) was used with floor temperature at 55°C (manufacturer's setup). The latency of paw withdrawal in response to the radiant heat was recorded (South and Smith, 1998). Cut-off periods of 10 and 15 seconds were employed to avoid tissue damage for the tail-flick and hot plate tests, respectively. In both tests, at least three readings per animal were taken at 15 min intervals, and the average was calculated.

Immunohistochemistry

The left and right side sciatic nerves were isolated at the mid-thigh level, fixed in 4% paraformaldehyde, and embedded in paraffin according to published protocol (Wang et al., 2011a). Three cross sections (6- μ m-thick) or three longitudinal sections (6- μ m-thick) in the one in ten series (60 μ m apart) for each animal were used for immunostaining according to our published protocols (Wang et al., 2011a). The following primary antibodies were used: polyclonal rabbit anti-Ang1 (1:2000; Abcam, Cambridge, MA), monoclonal mouse anti-CD31 antibody (1:500, BD Biosciences, San Jose, CA), polyclonal rabbit Anti-Von Willebrand Factor (vWF) (1:300, Dako, Carpinteria, CA), monoclonal mouse anti-occludin (1:200, Zymed, San Francisco, CA) and polyclonal rabbit anti-S100 (1:400, Abcam). Rabbit or goat IgG was used as a negative control. Sections were counterstained with 4',6-Diamidino-2-phenylindole (DAPI) (1:5000, Thermo Scientific, Rockford, IL).

Image Analysis and Quantification

To examine microvascular perfusion in the sciatic nerve, fluorescein isothiocyanate (FITC)-dextran (2×10^6 molecular weight, Sigma; 0.2 mL of 50 mg/mL) was administered intravenously to the mice 10 min before sacrifice (Zhang et al., 1999). The sciatic nerves were rapidly removed and placed in 2% of paraformaldehyde for 2 hours. The nerves were whole mounted and imaged under a 4x microscope objective (Zeiss Axiophot). Thereafter, the nerves were embedded in OCT compound and cross cryosections (20 μ m thick) prepared. Three sections at 60 μ m intervals from each mouse were used for further image analysis.

The cross sections were digitized under a 20x microscope objective (Zeiss Axiophot) via a Micro Computer Imaging Device (MCID) system (Imaging Research Inc, St. Catharines, ON, Canada) (Zhang et al., 1999). The total number of FITC-dextran perfused vessels was counted and divided by the total tissue-area to determine vascular density.

For analysis of vWF immunoreactive vascular morphology and density, three cross sections spaced at 60 μ m intervals from each mouse were used. Three fields of the view per section were randomly imaged under a 20x objective and vWF immunoreactive vascular perimeter and total number of vWF positive vessels were measured using MCID system.

All analysis was conducted with the examiner blinded to the identity of the samples being studied.

Cell culture

In the present study, we defined a normal glucose medium (NG) as a medium containing 5 mM glucose, while a high glucose medium (HG) is referred to a medium containing 30 mM glucose, which was chosen to match glucose levels prevalent in uncontrolled diabetic patients (Wu et al., 1999). These glucose concentrations for the in vitro hyperglycemia experiments have been used by others (Kim et al., 2008, Perrone et al., 2008).

Human Schwann cells (HSCs, ScienCell Research Laboratories, Carlsbad, CA) derived from primary culture and Human Umbilical Vein Endothelial Cells (HUVEC, American Type Culture Collection, ATCC, Manassas, VA) were cultured according to the manufactures' instructions (ScienCell Research Laboratories and ATCC). To examine the effect of T β 4 on HSCs, HSCs were cultured under the normal glucose or high glucose condition in the presence of different concentrations of T β 4 (0, 25, 50 and 100 ng/ml) for 24 or 72 hours. The cells were harvested for real-time RT-PCR and Western blot analysis. To collect conditioned medium from HSCs, 2.5×10^6 HSCs were plated onto a 35-mm-diameter dish in 1.2 ml of defined medium. The cells were cultured under the normal glucose or high

glucose conditions in the presence or absence of T β 4 (100 ng/ml) for 24 hours. HSCs were then washed three times with PBS and a fresh medium was added to avoid excessive T β 4 contamination. The cells were cultured for an additional 48 hours, and the supernatant (conditioned medium) was collected, centrifuged for 10 min at 1000 rpm, and stored at -80°C.

To assess the effect of T β 4 on in vitro angiogenesis, a capillary-like tube assay was used (Lee et al., 1999, Zhang et al., 2003, Zhang et al., 2004). Briefly, HUVECs (2×10^4 cells) were cultured on 96-well plate coated by Matrigel (BD Biosciences, Rockville, MD) in the conditioned medium or Dulbecco's Modified Eagle Medium (DMEM) in the presence or absence of T β 4 (0, 25, 50 and 100 ng/ml) for 5 hours. Total length of tubes was measured in 3 random fields from each well using MCID (Wang et al., 2004).

Real-time RT-PCR

Total RNA samples from cells were isolated using the Stratagene Absolutely RNA MicroRNA isolation kit (Stratagene, La Jolla, CA), according to the manufacturer's instructions. The complementary DNA (cDNA) was reversely transcribed from the same concentrations of total RNA products using random hexamers and M-MLV reverse-transcriptase (Invitrogen, Carlsbad, CA). Using the SYBR Green real-time PCR method (Wang et al., 2005), quantitative PCR was performed on an ABI 7000 PCR instrument (Applied Biosystems, Foster City, CA) by means of three-stage program parameters provided by the manufacturer, as follows; 2 min at 50 °C, 10 min at 95 °C, and then 40 cycles of 15 s at 95 °C and 1 min at 60 °C. Specificity of the produced amplification product was confirmed by examination of dissociation reaction plots. Each sample was tested in triplicate, and samples obtained from three independent experiments were used for analysis of relative gene expression using the $2^{-\Delta\Delta CT}$ method (Livak and Schmittgen, 2001). The following primers for real-time PCR were designed using Primer Express software (ABI): Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) (FWD, AGA ACA TCA TCC CTG CAT CC; REV, CAC ATT GGG GGT AGG AAC AC) and Ang1 (FWD GAA GGG AAC CGA GCC TAT TC; REV, GCT GAA ATC AGC ACC GTG TA), Ang2 (FWD, CAG ATC CGG GCT CTA GAC AG; REV, TCC GGA AAT CGT TCT TCA TC).

Western blot analysis

Western blot was performed according to published methods (Wang et al., 2005). Briefly, equal amounts of proteins were loaded on 10% SDS-polyacrylamide gel. After electrophoresis, the proteins were transferred to nitrocellulose membranes, and the blots were subsequently probed with the following antibodies: polyclonal rabbit anti-Ang1 (1:1000; Abcam, Cambridge, MA), polyclonal rabbit anti-Ang2 (1:1000; Abcam), Phospho-Akt (1:1000, Cell Signaling Technology, Inc. Danvers, MA) and Akt (1:1000, Cell Signaling Technology). For detection, horseradish peroxidase-conjugated secondary antibodies were used (1:2000) followed by enhanced chemiluminescence development (Pierce, Rockford, IL). Normalization of results was ensured by running parallel Western blot with β -actin antibody. The optical density was quantified using an image processing and analysis program (Scion Image, Ederick, MA).

Statistical analysis

For functional tests, data were evaluated for normality. Tail flick test data were not normal, and nonparametric Kruskal-Wallis Test was considered. The repeated measure analysis of variance (ANCOVA) was considered with dependent factor of time and independent factor of groups. The analysis started testing for group by time interaction, followed by the testing the main effect of group and subgroup analyses. Two-sample t-test or analysis of variance (ANOVA) was used to study the group difference on LDF, immunostaining, biochemistry,

Western blot, and tube formation analysis, respectively. Regression model was used to study the dose responses. The data are presented as mean $\pm$ SE. A value of $P < 0.05$ was taken as significant.

RESULTS

T β 4 improves diabetes-induced vascular dysfunction in the sciatic nerve

To examine whether db/db mice develop impairment of neurovasculature, we examined microvessels of the vasa nervora in the sciatic nerve of the mice at age of 28 weeks. Analysis of vWF immunoreactive vessels of the vasa nervora revealed that blood vessel perimeter in the db/db mice was significantly reduced compared to that in the age-matched db/m mice, although vascular density was not significantly different between these two groups (73.5 ± 10.5 vs. 56.9 ± 6.4 in db/m mice, $p > 0.05$) (Fig.1). In addition, the db/db mice exhibited substantial reduction of occludin, a tight junction protein, immunoreactive vessels (Fig.1). To examine whether the reduced perimeter in the db/db mice affects vascular function, we measured plasma-perfused microvessels, which represents functional vessels (Zhang et al., 1997, Ii et al., 2005) and regional blood flow. To measure plasma-perfused vessels, we intravenously injected FITC-dextran into the mice and then sacrificed them 10 minutes after injection, which provides sufficient time for the FITC-dextran to circulate throughout the entire vascular system under physiological conditions. Whole mount and cross sections showed that db/db mice exhibited overt reduction of FITC-dextran perfused vessels (Fig.1). Quantitative analysis of FITC-dextran perfused vessels on cross sections of the sciatic nerves revealed that the db/db mice had a significant reduction in microvascular densities perfused by FITC-dextran compared to the db/m mice (Fig.1). In parallel, sciatic nerve blood flow measured by LDF was significantly reduced in the db/db mice compared to that in the db/m mice (Fig.1). Together, these data indicate that diabetes induces vascular dysfunction in the sciatic nerve, which is consistent with published studies (Kusano et al., 2004, Ii et al., 2005, Jeong et al., 2009). However, the db/db mice treated with T β 4 at doses of 6 and 24 mg/kg for 4 weeks starting at animal age of 20 weeks exhibited significant increases in vascular perimeter, occludin immunoreactive vessels, the density of FITC-dextran perfused vessels and sciatic nerve blood flow at age of 28 weeks compared to the db/db mice treated with saline, which were close to levels measured in the db/m mice (Fig. 1). These data indicate that T β 4 ameliorates diabetes-induced vascular dysfunction in the sciatic nerve.

T β 4 improves neurological function in the diabetic mouse

Impairment of peripheral nerve conduction is a key indicator for diabetic patients having peripheral neuropathy (Said, 2007, Tesfaye et al., 2010) and vascular dysfunction affects nerve conduction (Cameron and Cotter, 1999, Cameron et al., 2005). We therefore investigated whether augmentation of regional blood flow by T β 4 affects motor and sensory conducting velocity (MCV and SCV) in the sciatic nerve. Electrophysiological recordings showed that MCV and SCV were significantly slowed in db/db mice compared to those of the age matched db/m mice (Fig.2), which are comparable to values reported by others (Ii et al., 2005, Pande et al., 2011). Treatment of the db/db mice with T β 4 at doses of 6 and 24 mg/kg for 4 weeks showed marked improvement in both MCV and SCV at the end of T β 4 treatment and at 4 weeks after termination of the treatment compared with saline-treated db/db mice (Fig.2). We then examined the effect of T β 4 treatment on sensory function by measuring the thermal latency with tail flick and hot plate tests. Treatment of the db/db mice with T β 4 markedly improved the thermal latency starting at the end of T β 4 treatment, which persisted for at least 4 weeks after termination of the treatment (the end of experimental period) (Fig.2). To examine the effect of T β 4 on non-diabetic mice, we treated db/m mice with T β 4 at a dose of 6 mg/kg and did not detect any functional changes as measured by the

methods listed above (data not shown). These data suggest that T β 4 improves peripheral nerve function in the diabetic mouse. Treatment of the db/db mouse with T β 4 did not significantly alter blood glucose levels and animal body weight (Table 1 and 2).

T β 4 regulates pro-angiogenic genes in the sciatic nerve

Alterations of Ang1 and Ang2 levels have been detected in diabetic patients and experimental diabetes (Tuo et al., 2008, Rasul et al., 2011). To examine the effect of T β 4 on these angiogenic genes, we measured protein levels of Ang1 and Ang2. Western blot analysis of the sciatic nerve showed substantial reduction of Ang1 levels and an increase in Ang2 levels in the db/db mouse (Fig.3), whereas treatment of the db/db mouse with T β 4 significantly increased Ang1 expression, but decreased Ang2 expression (Fig.3). Double immunostaining revealed that Ang1 immunoreactive cells were CD31 (a marker of endothelial cells) and S100 (a marker of Schwann cells) positive (Fig.3). Collectively, these data suggest that T β 4 upregulates Ang1 expression on endothelial cells and Schwann cells in the diabetic mouse.

The Ang/Tie2 signaling pathway mediates the effect of T β 4 on endothelial function

Aforementioned in vivo data suggest that Ang1 and Ang2 genes mediate T β 4-improved vascular function in the diabetic mouse. To examine a cause-effect of these genes on endothelial cells under hyperglycemia condition, we performed in vitro experiments using a capillary-like tube formation assay, which is widely used to examine in vitro endothelial function (Grant et al., 1995, Lee et al., 1999, Zhang et al., 2003, Wang et al., 2004). Incubation of HUVECs under high glucose conditions decreased capillary-like tube formation compared to HUVECs cultured under normal glucose conditions (Fig.4). However, T β 4 suppressed the effect of high glucose on reduction of capillary-like tube formation (Fig.4). Quantitative RT-PCR and Western blot analysis showed that incubation of HUVECs with high glucose substantially decreased and increased Ang1 and Ang2 expression, respectively (Fig.4), whereas T β 4 abolished high glucose-induced expression of Ang1 and Ang2 (Fig.4). Blockage of Tie2, a receptor of Ang1 and Ang2, with a neutralizing antibody against Tie2 inhibited T β 4-increased capillary-like tube formation (Fig.4). These data indicate that the Ang/Tie2 signaling pathway plays an important role in mediating T β 4-improved endothelial function under hyperglycemia.

Ang1 secreted by T β 4-treated Schwann cells improves endothelial function

In addition to endothelial cells, our in vivo double immunostaining data showed that Schwann cells expressed Ang1. We therefore, examined the effect of T β 4 on angiopoietin expression in Schwann cells. High glucose substantially downregulated Ang1 and upregulated Ang2, while T β 4 at a dose of 100 ng/ml suppressed the high glucose effect on the angiopoietin expression in HSCs (Fig.5), suggesting that T β 4 also regulates angiopoietin expression in Schwann cells. We then investigated whether Schwann cells secrete T β 4-upregulated Ang1 that consequently improves vascular function under high glucose condition. Using an ELISA specific to detect human Ang1, we measured Ang1 levels in supernatants harvested from HSCs cultured for 48 hours. ELISA showed that supernatants from HSCs cultured with high glucose had a significant reduction of Ang1 levels compared to levels in supernatants collected from normal glucose condition, while T β 4 reversed the effect of the high glucose on Ang1 levels (Fig.5).

Next, we examined the effect of the supernatants on capillary-like tube formation by culturing HUVECs with conditioned medium harvested from HSCs. Compared to the conditioned medium collected from HSCs cultured with normal glucose, the conditioned medium from HSCs cultured under high glucose condition resulted in a significant decrease of capillary-like tube formation (Fig.5). In contrast, the conditioned medium collected from

HSCs treated with T β 4 under high glucose condition significantly increased capillary-like tube formation. In the presence of the neutralizing antibody against Tie2, the effect of T β 4 conditioned medium on capillary-like tube formation was inhibited (Fig.5). Together, these data indicate that in addition to endothelial Ang1, soluble Ang1 secreted by T β 4-treated Schwann cells improves endothelial cell function.

The PI3K/Akt signaling pathway mediates the effect of T β 4 on Ang 1 expression

To further investigate whether intracellular signaling pathways are involved in T β 4-upregulated Ang1/Ang2 on endothelial cells and Schwann cells, we examined the PI3K/Akt signaling pathway that may mediate the effect of T β 4 on endothelial progenitor cell migration (Qiu et al., 2009). Western blot analysis of HUVECs and HSCs showed that the high glucose condition markedly decreased pAkt levels compared to the normal glucose (Fig.4 and 5). Treatment of HUVECs and HSCs with T β 4 under the high glucose condition significantly increased pAkt levels, which was fully blocked by a PI3K inhibitor LY294002 (Fig.4 and 5), indicating that T β 4 activates the PI3K/Akt signaling pathway. Moreover, LY294002 completely suppressed the effect of T β 4 on Ang1, but not Ang2 expression on HUVECs and HSCs under high glucose condition (Fig.4 and 5). These data suggest that the PI3K/Akt signaling pathway is involved in T β 4-regulated Ang1 expression on endothelial and Schwann cells. To verify these in vitro findings, we examined activation of Akt in the sciatic nerve of diabetic mice. Western blot analysis of the sciatic nerve tissue showed that diabetes substantially reduced pAkt compared to non-diabetic db/m mice, whereas treatment of db/db mice with T β 4 robustly elevated pAkt levels compared to db/db control mice (Fig. 4), suggesting that T β 4 also regulates the PI3K/Akt signaling pathway in vivo.

DISCUSSION

The present study for the first time demonstrates that T β 4 significantly improved sciatic nerve vascular function and peripheral nerve function in a mouse model of diabetic peripheral neuropathy. The Ang/Tie2 signaling pathway likely mediates the effect of T β 4 on improved vascular function.

T β 4, a naturally occurring peptide, enhances heart regeneration after myocardial infarction and promotes brain repair after stroke (Bock-Marquette et al., 2004, Morris et al., 2010). However, the effect of T β 4 on diabetic peripheral neuropathy has not been investigated. Using a well established mouse model of type II diabetes, the present study indicates that T β 4 ameliorates diabetic peripheral neuropathy, evidenced by reduction of sciatic nerve conduction velocity deficits, a key parameter for diabetic peripheral neuropathy, and improves responses to thermal stimuli. These data suggest that T β 4 has potential for the treatment of diabetic peripheral neuropathy.

Studies of diabetic peripheral neuropathy from experimental animals and human indicate that the development of diabetic neuropathy is closely associated with marked neurovascular dysfunction (Tesfaye et al., 1993, Cameron et al., 2001, Ebenezer et al., 2011). Vascular dysfunction precedes appearance of nerve conduction velocity deficits, leading to nerve damage (Cameron and Cotter, 1999, Cameron et al., 2005, Ebenezer et al., 2011). Our data demonstrated that T β 4 substantially increased plasma-perfused vessels and regional blood flow in the sciatic nerve, concomitantly with improvement of neurological function of diabetic neuropathy, suggesting that normalization of vascular function by T β 4 may contribute to observed reduction of nerve conduction velocity deficits. Our findings are consistent with prior reports that restoration of vascular function by either pro-angiogenic factors or cell therapies enhances neurological function in diabetic peripheral neuropathy (Kusano et al., 2004, Ii et al., 2005, Jeong et al., 2009). However, further studies on the direct effects of T β 4 on nerve fiber morphology and its relationship to improvement in

neurological function are warranted. Our data show that T β 4 at doses of 6 and 24 mg/kg significantly improved neurological outcome. Currently, we do not know why a dose of 24 mg/kg is not superior to a lower dose of 6 mg/kg to improve vascular function and neurological outcomes in diabetic mice with peripheral neuropathy. We previously demonstrated that T β 4 at a dose of 30 mg/kg significantly reduces brain injury and improves neurological outcome compared to a dose of 6 mg/kg (Xiong et al., 2012). A clinical phase 1A and 1B study demonstrates that T β 4 at a dose range of 42 to 1260 mg daily for 2 weeks did not have adverse effects, suggesting that T β 4 has a large dose range in the human (RegeneRx Biopharmaceuticals Inc). Further studies on the effects of doses at 30 mg/kg and higher on diabetic peripheral neuropathy are warranted.

The Ang/Tie2 signaling pathway regulates vascular homeostasis (Suri et al., 1996, Teichert-Kuliszewska et al., 2001). Ang1 promotes vascular maturation, while Ang2 acts as a competitive inhibitor of Ang1 for Tie2 binding and destabilizes blood vessels (Suri et al., 1996, Maisonpierre et al., 1997, Thebaud et al., 2005). Hyperglycemia downregulates Ang1 and upregulates Ang2 (Tuo et al., 2008). Increases in Ang1 levels normalize diabetes induced immature vasculature (Chen and Stinnett, 2008a). Ang1 by increasing angiogenesis reduces myocardial infarction, whereas an elevation of Ang2 levels exacerbates the infarction in diabetic rats (Tuo et al., 2008). Patients with diabetic peripheral neuropathy have significantly elevated levels of circulating Ang2 (Rasul et al., 2011). Our data show that hyperglycemia downregulated Ang1 and upregulated Ang2 on endothelial cells and Schwann cells, whereas T β 4 reversed expression of Ang1 and Ang2. T β 4 is a potent angiogenic factor and regulates angiogenesis and vasculogenesis during development by promoting progenitor cell differentiation and by directing endothelial cell migration (Bock-Marquette et al., 2004, Smart et al., 2007a, b). The effect of T β 4 on the Ang/Tie2 pathway has not been investigated. Our data that blockage of Tie2 with a neutralizing antibody suppressed the effect of T β 4 on in vitro angiogenesis implicates the Ang/Tie2 signaling pathway in mediating T β 4-improved vascular function observed in vivo. Our data further suggest that T β 4 regulates Ang1 through the activation of the PI3K/Akt pathway. A recent study shows that T β 4 interacts with ATP-responsive P2X4 receptor to regulate endothelial cell migration (Freeman et al., 2011). Angiogenesis involves endothelial cell proliferation and migration (Goukassian et al., 2001, Wang et al., 2011b). Therefore, further studies are warranted for investigating whether purinergic signaling is involved in the beneficial effects of T β 4 observed in the present study.

Schwann cells secrete numerous factors that regulate degeneration and regeneration peripheral nerves (Sobue, 1990, Frostick et al., 1998, Campana, 2007). The present study showed that Ang1 and Ang2 secreted by Schwann cells affected endothelial function under hyperglycemia condition, while T β 4 -elevated Ang1 levels on Schwann cells lead to enhancement of in vitro angiogenesis. We speculate that T β 4 acts on endothelial cells and Schwann cells to preserve and/or restore vascular function in the sciatic nerve, which facilitates improvement of peripheral nerve function under diabetic neuropathy.

Acknowledgments

This work was supported by the Dykstra Foundation, and NINDS grants PO1 NS23393, RO1 NS075084, RO1 AG037506, and RO1 NS075156. The authors thank Cynthia Roberts, and Qing-e Lu for their technical assistance and Deborah Jewell for secretarial support.

REFERENCES

- Bock-Marquette I, Saxena A, White MD, Dimaio JM, Srivastava D. Thymosin beta4 activates integrin-linked kinase and promotes cardiac cell migration, survival and cardiac repair. *Nature*. 2004; 432:466–472. [PubMed: 15565145]

- Cameron NE, Cotter MA. Effects of antioxidants on nerve and vascular dysfunction in experimental diabetes. *Diabetes Res Clin Pract.* 1999; 45:137–146. [PubMed: 10588366]
- Cameron NE, Eaton SE, Cotter MA, Tesfaye S. Vascular factors and metabolic interactions in the pathogenesis of diabetic neuropathy. *Diabetologia.* 2001; 44:1973–1988. [PubMed: 11719828]
- Cameron NE, Gibson TM, Nangle MR, Cotter MA. Inhibitors of advanced glycation end product formation and neurovascular dysfunction in experimental diabetes. *Annals of the New York Academy of Sciences.* 2005; 1043:784–792. [PubMed: 16037306]
- Campana WM. Schwann cells: activated peripheral glia and their role in neuropathic pain. *Brain Behav Immun.* 2007; 21:522–527. [PubMed: 17321718]
- Chen JX, Stinnett A. Ang-1 gene therapy inhibits hypoxia-inducible factor-1 α (HIF-1 α)-prolyl-4-hydroxylase-2, stabilizes HIF-1 α expression, and normalizes immature vasculature in db/db mice. *Diabetes.* 2008a; 57:3335–3343. [PubMed: 18835934]
- Chen JX, Stinnett A. Disruption of Ang-1/Tie-2 signaling contributes to the impaired myocardial vascular maturation and angiogenesis in type II diabetic mice. *Arterioscler Thromb Vasc Biol.* 2008b; 28:1606–1613. [PubMed: 18556567]
- ClinicalTrials.gov. (<http://clinicaltrials.gov/ct2/show/NCT01311518>)
- Crockford D. Development of thymosin beta4 for treatment of patients with ischemic heart disease. *Annals of the New York Academy of Sciences.* 2007; 1112:385–395. [PubMed: 17947592]
- Crockford D, Turjman N, Allan C, Angel J. Thymosin beta4: structure, function, and biological properties supporting current and future clinical applications. *Annals of the New York Academy of Sciences.* 2010; 1194:179–189. [PubMed: 20536467]
- Ebenezer GJ, O'Donnell R, Hauer P, Cimino NP, McArthur JC, Polydefkis M. Impaired neurovascular repair in subjects with diabetes following experimental intracutaneous axotomy. *Brain.* 2011; 134:1853–1863. [PubMed: 21616974]
- Freeman KW, Bowman BR, Zetter BR. Regenerative protein thymosin beta-4 is a novel regulator of purinergic signaling. *FASEB journal: official publication of the Federation of American Societies for Experimental Biology.* 2011; 25:907–915. [PubMed: 21106936]
- Frostick SP, Yin Q, Kemp GJ. Schwann cells, neurotrophic factors, and peripheral nerve regeneration. *Microsurgery.* 1998; 18:397–405. [PubMed: 9880154]
- Goldstein AL, Slater FD, White A. Preparation, assay, and partial purification of a thymic lymphocytopoietic factor (thymosin). *Proc Natl Acad Sci U S A.* 1996; 56:1010–1017. [PubMed: 5230175]
- Goukassian D, Diez-Juan A, Asahara T, Schratzberger P, Silver M, Murayama T, Isner JM, Andres V. Overexpression of p27(Kip1) by doxycycline-regulated adenoviral vectors inhibits endothelial cell proliferation and migration and impairs angiogenesis. *FASEB journal : official publication of the Federation of American Societies for Experimental Biology.* 2001; 15:1877–1885. [PubMed: 11532967]
- Grant DS, Kinsella JL, Kibbey MC, LaFlamme S, Burbelo PD, Goldstein AL, Kleinman HK. Matrigel induces thymosin beta 4 gene in differentiating endothelial cells. *J Cell Sci.* 1995; 108(Pt 12): 3685–3694. [PubMed: 8719875]
- Ii M, Nishimura H, Kusano KF, Qin G, Yoon YS, Wecker A, Asahara T, Losordo DW. Neuronal nitric oxide synthase mediates statin-induced restoration of vasa nervorum and reversal of diabetic neuropathy. *Circulation.* 2005; 112:93–102. [PubMed: 15983249]
- Janssen PA, Niemegeers CJ, Dony JG. The inhibitory effect of fentanyl and other morphine-like analgesics on the warm water induced tail withdrawal reflex in rats. *Arzneimittelforschung.* 1963; 13:502–507. [PubMed: 13957426]
- Jeong JO, Kim MO, Kim H, Lee MY, Kim SW, Ii M, Lee JU, Lee J, Choi YJ, Cho HJ, Lee N, Silver M, Wecker A, Kim DW, Yoon YS. Dual angiogenic and neurotrophic effects of bone marrow-derived endothelial progenitor cells on diabetic neuropathy. *Circulation.* 2009; 119:699–708. [PubMed: 19171856]
- Kim HK, Kim YJ, Kim JT, Kwon CH, Kim YK, Bae YC, Kim DH, Jung JS. Alterations in the proangiogenic functions of adipose tissue-derived stromal cells isolated from diabetic rats. *Stem Cells Dev.* 2008; 17:669–680. [PubMed: 18788931]

- Kusano KF, Allendoerfer KL, Munger W, Pola R, Bosch-Marce M, Kirchmair R, Yoon YS, Curry C, Silver M, Kearney M, Asahara T, Losordo DW. Sonic hedgehog induces arteriogenesis in diabetic vasa nervorum and restores function in diabetic neuropathy. *Arterioscler Thromb Vasc Biol.* 2004; 24:2102–2107. [PubMed: 15358602]
- Lee OH, Kim YM, Lee YM, Moon EJ, Lee DJ, Kim JH, Kim KW, Kwon YG. Sphingosine 1-phosphate induces angiogenesis: its angiogenic action and signaling mechanism in human umbilical vein endothelial cells. *Biochem Biophys Res Commun.* 1999; 264:743–750. [PubMed: 10544002]
- Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods.* 2001; 25:402–408. [PubMed: 11846609]
- Maisonpierre PC, Suri C, Jones PF, Bartunkova S, Wiegand SJ, Radziejewski C, Compton D, McClain J, Aldrich TH, Papadopoulos N, Daly TJ, Davis S, Sato TN, Yancopoulos GD. Angiopoietin-2, a natural antagonist for Tie2 that disrupts in vivo angiogenesis. *Science.* 1997; 277:55–60. [PubMed: 9204896]
- Mora CA, Baumann CA, Paino JE, Goldstein AL, Badamchian M. Biodistribution of synthetic thymosin beta 4 in the serum, urine, and major organs of mice. *International journal of immunopharmacology.* 1997; 19:1–8. [PubMed: 9226473]
- Morris DC, Chopp M, Zhang L, Lu M, Zhang ZG. Thymosin beta4 improves functional neurological outcome in a rat model of embolic stroke. *Neuroscience.* 2010; 169:674–682. [PubMed: 20627173]
- Pande M, Hur J, Hong Y, Backus C, Hayes JM, Oh SS, Kretzler M, Feldman EL. Transcriptional profiling of diabetic neuropathy in the BKS db/db mouse: a model of type 2 diabetes. *Diabetes.* 2011; 60:1981–1989. [PubMed: 21617178]
- Perrone L, Peluso G, Melone MA. RAGE recycles at the plasma membrane in S100B secretory vesicles and promotes Schwann cells morphological changes. *J Cell Physiol.* 2008; 217:60–71. [PubMed: 18452184]
- Philp D, Badamchian M, Scheremeta B, Nguyen M, Goldstein AL, Kleinman HK. Thymosin beta 4 and a synthetic peptide containing its actin-binding domain promote dermal wound repair in db/db diabetic mice and in aged mice. *Wound repair and regeneration : official publication of the Wound Healing Society [and] the European Tissue Repair Society.* 2003; 11:19–24.
- Qiu FY, Song XX, Zheng H, Zhao YB, Fu GS. Thymosin beta4 induces endothelial progenitor cell migration via PI3K/Akt/eNOS signal transduction pathway. *J Cardiovasc Pharmacol.* 2009; 53:209–214. [PubMed: 19247195]
- Rasul S, Reiter MH, Ilhan A, Lampichler K, Wagner L, Kautzky-Willer A. Circulating angiopoietin-2 and soluble Tie-2 in type 2 diabetes mellitus: a cross-sectional study. *Cardiovasc Diabetol.* 2011; 10:55. [PubMed: 21699724]
- Ruff D, Crockford D, Girardi G, Zhang Y. A randomized, placebo-controlled, single and multiple dose study of intravenous thymosin beta4 in healthy volunteers. *Annals of the New York Academy of Sciences.* 2010; 1194:223–229. [PubMed: 20536472]
- Said G. Diabetic neuropathy--a review. *Nat Clin Pract Neurol.* 2007; 3:331–340. [PubMed: 17549059]
- Schratzberger P, Schratzberger G, Silver M, Curry C, Kearney M, Magner M, Alroy J, Adelman LS, Weinberg DH, Ropper AH, Isner JM. Favorable effect of VEGF gene transfer on ischemic peripheral neuropathy. *Nature medicine.* 2000; 6:405–413.
- Schratzberger P, Walter DH, Rittig K, Bahlmann FH, Pola R, Curry C, Silver M, Krainin JG, Weinberg DH, Ropper AH, Isner JM. Reversal of experimental diabetic neuropathy by VEGF gene transfer. *J Clin Invest.* 2001; 107:1083–1092. [PubMed: 11342572]
- Smart N, Risebro CA, Melville AA, Moses K, Schwartz RJ, Chien KR, Riley PR. Thymosin beta4 induces adult epicardial progenitor mobilization and neovascularization. *Nature.* 2007a; 445:177–182. [PubMed: 17108969]
- Smart N, Risebro CA, Melville AA, Moses K, Schwartz RJ, Chien KR, Riley PR. Thymosin beta-4 is essential for coronary vessel development and promotes neovascularization via adult epicardium. *Annals of the New York Academy of Sciences.* 2007b; 1112:171–188. [PubMed: 17495252]
- Sobue G. The role of Schwann cells in peripheral nerve degeneration and regeneration--NGF-NGF receptor system. *Rinsho Shinkeigaku.* 1990; 30:1358–1360. [PubMed: 1966016]

- South SM, Smith MT. Apparent insensitivity of the hotplate latency test for detection of antinociception following intraperitoneal, intravenous or intracerebroventricular M6G administration to rats. *J Pharmacol Exp Ther*. 1998; 286:1326–1332. [PubMed: 9732396]
- Suri C, Jones PF, Patan S, Bartunkova S, Maisonpierre PC, Davis S, Sato TN, Yancopoulos GD. Requisite role of angiopoietin-1, a ligand for the TIE2 receptor, during embryonic angiogenesis. *Cell*. 1996; 87:1171–1180. [PubMed: 8980224]
- Teichert-Kuliszewska K, Maisonpierre PC, Jones N, Campbell AI, Master Z, Bendeck MP, Alitalo K, Dumont DJ, Yancopoulos GD, Stewart DJ. Biological action of angiopoietin-2 in a fibrin matrix model of angiogenesis is associated with activation of Tie2. *Cardiovasc Res*. 2001; 49:659–670. [PubMed: 11166279]
- Tesfaye S, Boulton AJ, Dyck PJ, Freeman R, Horowitz M, Kempner P, Lauria G, Malik RA, Spallone V, Vinik A, Bernardi L, Valensi P. Diabetic neuropathies: update on definitions, diagnostic criteria, estimation of severity, and treatments. *Diabetes Care*. 2010; 33:2285–2293. [PubMed: 20876709]
- Tesfaye S, Harris N, Jakubowski JJ, Mody C, Wilson RM, Rennie IG, Ward JD. Impaired blood flow and arterio-venous shunting in human diabetic neuropathy: a novel technique of nerve photography and fluorescein angiography. *Diabetologia*. 1993; 36:1266–1274. [PubMed: 8307254]
- Thebaud B, Ladha F, Michelakis ED, Sawicka M, Thurston G, Eaton F, Hashimoto K, Harry G, Haromy A, Korbitt G, Archer SL. Vascular endothelial growth factor gene therapy increases survival, promotes lung angiogenesis, and prevents alveolar damage in hyperoxia-induced lung injury: evidence that angiogenesis participates in alveolarization. *Circulation*. 2005; 112:2477–2486. [PubMed: 16230500]
- Tuo QH, Zeng H, Stinnett A, Yu H, Aschner JL, Liao DF, Chen JX. Critical role of angiopoietins/Tie-2 in hyperglycemic exacerbation of myocardial infarction and impaired angiogenesis. *Am J Physiol Heart Circ Physiol*. 2008; 294:H2547–H2557. [PubMed: 18408125]
- Vanderah TW, Suenaga NM, Ossipov MH, Malan TP Jr, Lai J, Porreca F. Tonic descending facilitation from the rostral ventromedial medulla mediates opioid-induced abnormal pain and antinociceptive tolerance. *J Neurosci*. 2001; 21:279–286. [PubMed: 11150345]
- Wang L, Chopp M, Szalad A, Liu Z, Bolz M, Alvarez FM, Lu M, Zhang L, Cui Y, Zhang RL, Zhang ZG. Phosphodiesterase-5 is a therapeutic target for peripheral neuropathy in diabetic mice. *Neuroscience*. 2011a; 193:399–410. [PubMed: 21820491]
- Wang L, Gang Zhang Z, Lan Zhang R, Chopp M. Activation of the PI3-K/Akt pathway mediates cGMP enhanced-neurogenesis in the adult progenitor cells derived from the subventricular zone. *J Cereb Blood Flow Metab*. 2005; 25:1150–1158. [PubMed: 15815584]
- Wang L, Zhang Z, Wang Y, Zhang R, Chopp M. Treatment of stroke with erythropoietin enhances neurogenesis and angiogenesis and improves neurological function in rats. *Stroke*. 2004; 35:1732–1737. [PubMed: 15178821]
- Wang Y, Li Z, Xu P, Huang L, Tong J, Huang H, Meng A. Angiomin-like2 gene (amotl2) is required for migration and proliferation of endothelial cells during angiogenesis. *The Journal of biological chemistry*. 2011b; 286:41095–41104. [PubMed: 21937427]
- Wu QD, Wang JH, Fennessy F, Redmond HP, Bouchier-Hayes D. Taurine prevents high-glucose-induced human vascular endothelial cell apoptosis. *Am J Physiol*. 1999; 277:C1229–C1238. [PubMed: 10600775]
- Xiong Y, Mahmood A, Meng Y, Zhang Y, Zhang ZG, Morris DC, Chopp M. Treatment of traumatic brain injury with thymosin beta in rats. *Journal of neurosurgery*. 2011; 114:102–115. [PubMed: 20486893]
- Xiong Y, Zhang Y, Mahmood A, Meng Y, Zhang ZG, Morris DC, Chopp M. Neuroprotective and neurorestorative effects of thymosin beta4 treatment initiated 6 hours after traumatic brain injury in rats. *Journal of neurosurgery*. 2012; 116:1081–1092. [PubMed: 22324420]
- Zhang J, Zhang ZG, Morris D, Li Y, Roberts C, Elias SB, Chopp M. Neurological functional recovery after thymosin beta4 treatment in mice with experimental auto encephalomyelitis. *Neuroscience*. 2009; 164:1887–1893. [PubMed: 19782721]

- Zhang R, Wang L, Zhang L, Chen J, Zhu Z, Zhang Z, Chopp M. Nitric oxide enhances angiogenesis via the synthesis of vascular endothelial growth factor and cGMP after stroke in the rat. *Circ Res.* 2003; 92:308–313. [PubMed: 12595343]
- Zhang R, Zhang Z, Wang L, Wang Y, Gousev A, Zhang L, Ho KL, Morshead C, Chopp M. Activated neural stem cells contribute to stroke-induced neurogenesis and neuroblast migration toward the infarct boundary in adult rats. *J Cereb Blood Flow Metab.* 2004; 24:441–448. [PubMed: 15087713]
- Zhang Z, Davies K, Probst J, Fenstermaker J, Chopp M. Quantitation of microvascular plasma perfusion and neuronal microtubule-associated protein in ischemic mouse brain by laser-scanning confocal microscopy. *J Cereb Blood Flow Metab.* 1999; 19:68–78. [PubMed: 9886357]
- Zhang Z, Zhang RL, Jiang Q, Raman SB, Cantwell L, Chopp M. A new rat model of thrombotic focal cerebral ischemia. *J Cereb Blood Flow Metab.* 1997; 17:123–135. [PubMed: 9040491]

Figure 1. Tβ4 improves vascular function in the sciatic nerve

Panels A to C show vWF immunoreactive blood vessels at the cross section of the sciatic nerve from a representative db/m mouse (A), db/db mouse treated with saline (B), and db/db mouse treated with Tβ4 (24 mg/kg, C). Whole mounted (D to F) and cross sections (G to I) of the sciatic nerve show FITC-dextran perfused vessels from a representative db/m mouse (D, G), db/db mouse treated with saline (E, H), and db/db mouse treated with Tβ4 (24 mg/kg, F, I). Panels J to M show quantitative data of vWF immunoreactive vascular perimeters (J, n=6/group), and density of FITC-dextran perfused vessels in cross section (K, n=4/group), percentage of occludin immunoreactive area (L, n=6/group) and percentage changes of sciatic nerve blood flow with a reference of db/m mice at 100% (M, n=4/group). *P<0.05

and #P<0.05 versus the db/m mouse and the saline treated db/db mouse, respectively.
Bar=100 μ m. dm=db/m mouse; db=db/db mouse.

Watermark-text

Watermark-text

Watermark-text

Figure 2. The effect of T β 4 on neurological function

Treatment of db/db mice with T β 4 improves neurological function measured by MCV (A), SCV (B), Tail flick test (C) and Hot plate test (D). * P <0.05 and # P <0.05 versus the db/m mouse and the db/db mouse treated with saline, respectively. n =10/group. dm=db/m mouse; db=db/db mouse.

Figure 3. The effect of T β 4 on Ang1 and Ang2 expression in endothelial cells and Schwann cells of the diabetic mouse

Western blot analysis (A and B) of Ang1 and Ang2 levels in sciatic nerve tissue and β -actin was used as an internal control. Representative images of double immunofluorescent staining show that Ang1 immunoreactivity (C, F, E, H, red, arrows) was colocalized to CD31 positive vessels (D, E, green, arrows) and S100 positive Schwann cells (G, H, green, arrows). * $P < 0.05$ and # $P < 0.05$ versus the db/m mouse and the saline treated db/db mouse, respectively. $n = 6$ /group. Bar = 100 μ m. dm=db/m mouse; db=db/db mouse.

Figure 4. The effect of Tβ4 on the Ang/Tie2 and PI3K/Akt signaling pathways

Representative microscopic images (A) and quantitative data (B) show capillary-like tube formation in HUVECs cultured in normal glucose (N), high glucose (H), high glucose with Tβ4 (H+TB4, 100 ng/ml), and high glucose with Tβ4 in the presence of a neutralizing antibody against Tie2 (+Tie2, 5 μg/ml). Real-time RT-PCR (C) and Western blot (D) data show mRNA and protein, respectively, levels of Ang1 and Ang2 in HUVECs cultured with normal glucose (N), high glucose (H), high glucose with Tβ4 (+TB4), and high glucose with Tβ4 in the presence of LY294003 (+LY, 10 μM). Panel E shows Western blot analysis of pAkt and total Akt in HUVECs cultured under different conditions listed above. GAPDH and β-actin were used as internal controls for mRNA and proteins, respectively. *P<0.05, #P<0.05 and \$P<0.05 versus the normal glucose (N), high glucose (H) and high glucose with Tβ4 (100 ng/ml) groups, respectively. n=6/group. Bar=50μm. Western blot analysis of sciatic nerve tissue (F) shows pAkt and total Akt levels in db/m (dm) and db/db (db) mice, db/db mice treated with Tβ4 at 6 mg/kg (db+6) or 24 mg/kg (db+24). *P<0.05 vs db/m mice and #P<0.05 vs db/db mice. n=6/group.

Figure 5. Ang1 secreted by Tβ4-treated Schwann cells improves endothelial function

Real-time RT-PCR (A) and Western blot (B) data show mRNA and protein levels of Ang1 and Ang2 in HSCs cultured with normal glucose (N), high glucose (H), high glucose with Tβ4 (+TB4), and high glucose with Tβ4 in the presence of LY294003 (+LY). Panel C shows Western blot analysis of pAkt and total Akt in HSCs cultured under different conditions listed above. GAPDH and β-actin were used as internal controls for mRNA and proteins, respectively. Panel D shows ELISA data of Ang1 levels in supernatants harvested from HSCs cultured with normal glucose (N), high glucose (H), high glucose and Tβ4 (+TB4, 100 ng/ml), and high glucose and Tβ4 in the presence of LY294003 (+LY). Representative microscopic images (E) and quantitative data (F) show capillary-like tube formation in HUVECs cultured with the conditioned medium collected from HSCs in normal glucose (N), high glucose (H), high glucose and Tβ4 (+TB4, 100 ng/ml), and high glucose and Tβ4 in the presence of the antibody against Tie2 (+Tie2). *P<0.05, #P<0.05 and \$P<0.05 versus the normal glucose (N), high glucose (H) and high glucose with Tβ4 (100 ng/ml) groups, respectively. n=6/group. Bar=50μm.

Watermark-text

Watermark-text

Watermark-text

Table 1

Effect of Tβ4 on bodyweight

Groups	Weight, g			
	0 w	2 w	4 w	8 w
dm-Saline	30.3±0.8 [*]	30.8±0.7 [*]	30.4±0.9 [*]	30.1±0.7 [*]
db-Saline	61.0±0.8	61.2±0.7	59.3±1.2	57.6±2.2
db-TB4(6mg/kg)	59.5±0.9	57.1±1.1	56.2±1.2	53.7±1.5
db-TB4(24mg/kg)	58.9±1.1	58.1±1.2	55.7±1.5	56.0±1.4

Values are mean±SE.

^{*} P<0.01 versus db+Saline group. n=10/group. W= week, 0 w represents before the treatment, while other numbers indicate after the treatment. dm=db/m mouse; db=db/db mouse.

Table 2Effect of T β 4 on blood glucose

Groups	Blood glucose(g/dl)				
	0 w	2 w	4 w	6 w	8 w
dm-Saline	141.6±14.5 [*]	138.6±6.4 [*]	141.4±7.7 [*]	140.1±8.0 [*]	141.3±4.6 [*]
db-Saline	513.8±14.0	516.6±18.6	517.6±10.7	516.4±32.3	524.2±23.3
db-TB4(6mg/kg)	503.4±24.0	513.4±17.5	512.8±24.8	517.6±21.4	503.3±18.0
db-TB4(24mg/kg)	517.6±15.0	515.9±13.1	504.7±16.5	521.6±16.0	515.8±19.3

Values are mean $\pm$ SE.

* P<0.01 versus db+Saline group. n=10/group. W= week, 0 w represents before the treatment, while other numbers indicate after the treatment. dm=db/m mouse; db=db/db mouse.