Published in final edited form as:

Obes Rev. 2011 October ; 12(10): 800–812. doi:10.1111/j.1467-789X.2011.00893.x.

The Neurocognitive Connection between Physical Activity and Eating Behavior

Richard J. Joseph, BS¹, Miguel Alonso-Alonso, MD, MPhil², Dale S. Bond, PhD³, Alvaro Pascual-Leone, MD, PhD^{2,4}, and George L. Blackburn, MD, PhD¹

¹Beth Israel Deaconess Medical Center, Center for the Study of Nutrition Medicine

²Beth Israel Deaconess Medical Center, Berenson Allen Center for Noninvasive Brain Stimulation

³The Miriam Hospital, Weight Control and Diabetes Research Center

⁴Institut Universitari de Neurorehabilitació Guttmann, Universidad Autónoma de Barcelona, Badalona, Spain

Abstract

As obesity rates increase worldwide, health care providers require methods to instill the lifestyle behaviors necessary for sustainable weight loss. Designing effective weight loss interventions requires an understanding of how these behaviors are elicited, how they relate to each other, and whether they are supported by common neurocognitive mechanisms. This may provide valuable insights to optimize existing interventions and develop novel approaches to weight control. Researchers have begun to investigate the neurocognitive underpinnings of eating behavior and the impact of physical activity on cognition and the brain. This review attempts to bring these somewhat disparate, yet interrelated lines of literature together in order to examine a hypothesis that eating behavior and physical activity share a common neurocognitive link. The link pertains to executive functions, which rely on brain circuits located in the prefrontal cortex. These advanced cognitive processes are of limited capacity and undergo relentless strain in the current obesogenic environment. The increased demand on these neurocognitive resources as well as their overuse and/or impairment may facilitate impulses to overeat, contributing to weight gain and obesity. This impulsive eating drive may be counteracted by physical activity due to its enhancement of neurocognitive resources for executive functions and goal-oriented behavior. By enhancing the resources that facilitate “top-down” inhibitory control, increased physical activity may help compensate and suppress the hedonic drive to overeat. Understanding how physical

Correspondence to: George L. Blackburn.

Author Contact Information

Richard Joseph, BS, Beth Israel Deaconess Medical Center, Department of Surgery, Center for the Study of Nutrition Medicine, 330 Brookline Ave, Feldberg 877A, Boston, MA 02215, Tel: 617-667-2654, Fax: 617-667-2685, rjoseph1@bidmc.harvard.edu

Miguel Alonso-Alonso, MD, MPhil, Beth Israel Deaconess Medical Center, Department of Neurology, Berenson Allen Center for Noninvasive Brain Stimulation, Kirshtein 158, East Campus, 330 Brookline Avenue, Boston, MA 02215, Tel: 617-667-0240, Fax: 617-975-5322, malonso@bidmc.harvard.edu

Dale S. Bond, PhD, Warren Alpert Medical School of Brown University, Department of Psychiatry and Human Behavior, The Miriam Hospital, Weight Control and Diabetes Research Center, 196 Richmond Street, Providence, RI 02903, Tel 401-793-8970, Fax 401-793-8944, dbond@lifespan.org

Alvaro Pascual-Leone, MD, PhD, Beth Israel Deaconess Medical Center, Department of Neurology, Berenson Allen Center for Noninvasive Brain Stimulation, Kirshtein 158, East Campus, 330 Brookline Avenue, Boston, MA 02215, Tel: 617-667-0203, Fax: 617-975-5322, apalone@bidmc.harvard.edu

George L. Blackburn, MD, PhD, Beth Israel Deaconess Medical Center, Department of Surgery, Center for the Study of Nutrition Medicine, Feldberg 880, East Campus, 330 Brookline Avenue, Boston, MA 02215, Tel: 617-667-2603, Fax: 617-667-2608, gblackbu@bidmc.harvard.edu

Conflict of Interest Statement

No conflict of interest was declared.

activity and eating behaviors interact on a neurocognitive level may help to maintain a healthy lifestyle in an obesogenic environment.

Keywords

obesity; neurocognition; eating behavior; physical activity

Introduction

Despite increased awareness, public health campaigns, and novel therapies, obesity rates remain at alarming levels in the United States (1) and are increasing worldwide (2). Lifestyle-based weight loss interventions consistently produce a 7–10% decline in initial body weight (3), but the condition is refractory, and poor retention rates (4,5) and relapse of weight gain (6) are common. The limited long-term success of such interventions may be attributable in part to a failure to fully understand and exploit the connection between the two therapeutic cornerstones, physical activity and eating behaviors. Indeed, while much emphasis is placed on modifying both behaviors to promote an immediate energy deficit, there has been less emphasis on understanding how a change in one of these behaviors may support the longer-term maintenance of change in the other. A clearer understanding of the interaction between physical activity and eating behaviors based on underlying neurocognitive processes and adaptations may help to optimize the implementation, adoption, and combined effect of each for healthy, sustainable weight loss (7,8).

The analysis of behavior change via neurocognitive modeling holds promise as one of this decade's most beneficial developments in the treatment of obesity. Behavior change implies a critical switch whereby an externally imposed intervention transitions into self-regulated actions that facilitate a desired outcome. Adapted cognitive processes and resultant behavior patterns underlie these so-called "goal-oriented" actions, such as resisting high-calorie foods and increasing physical activity to align with a weight loss goal. On a neurocognitive level, these behavior changes involve the alteration of the brain's neurobiological substrates and plastic processing pathways that are cemented over time (9). Identifying these pathways and determining how to modify the neurocognitive processes involved in adopting healthy lifestyle behaviors is essential to achieving sustained weight loss. By linking two emerging lines of neuroscience literature, we posit that increased physical activity and consumption of a healthy diet share a common neurocognitive underpinning. This may help to explain the observational association and behavioral interplay between these lifestyle factors and may be useful in developing more effective weight loss interventions.

The associative link between physical activity and diet: observations and theoretical foundations

Whereas the effect of physical activity and dietary habits on disease has been established (10,11), the ability of these two behaviors to indirectly modify each other has been sparsely documented. The few studies that examined the interaction between habitual physical activity and healthy diet suggest an interactive and reinforcing relationship (12,13). A study by Brodney et al. found that higher cardiorespiratory fitness levels were associated with lower percent of energy from fat, higher dietary fiber intake, reduced cholesterol intake, and closer adherence to dietary recommendations (14). Findings from a study of 2,004 New England households suggested that moderately to very active individuals tend to consume more fiber, less total fat, and less saturated fat than sedentary participants (15). In one of the few prospective studies in this area, men and women who reported increases in their activity levels from age 33 to 42 also reported improvement in the overall quality of their diet (16).

Taken together, these studies suggest a link between physical activity and diet, opening the possibility that engagement in higher levels of physical activity may influence the amount and quality of dietary intake or vice versa. However, crude and imprecise measures of physical activity, small sample sizes, reliance on dietary self-report, and mixed samples of normal and overweight/obese individuals may undermine the reliability of the data. Such epidemiologic studies can, at best, identify potential associations and help formulate hypothesis that warrant follow-up testing. The extent and direction of causality and underlying mechanisms of this behavioral covariance remains largely unexplored given the observational nature of these studies.

Studies also indicate that physical activity and dietary behaviors comprise a constellation of similar lifestyle behaviors that cluster together (17,18). A review article by Blair et al. indicates that physical activity may indirectly influence health behaviors like smoking, substance abuse, risk taking, overeating, and stress management (19). Conversely, sedentary behavior and unhealthy eating habits fit into the cluster of behaviors that are adverse to health. One study, which documented the prevalence and cluster patterns of lifestyle factors among a random sample of adolescents, adults, and seniors from a large Midwestern health plan, found that overall, 14.5% of the health plan members met recommended guidelines for four healthy lifestyle factors—physical activity, nonsmoking, high-quality diet, and healthy weight (20). An investigation into the prevalence and clustering of lifestyle risk factors such as sedentary behavior, smoking, low fruit and vegetable consumption, and excessive alcohol consumption among Dutch men and women found significant clustering, with about 20% of the participants having three risk factors simultaneously (21). Analysis of data from the Cardiovascular Risk in Young Finns Study reported similar grouping (22). Specifically, physical inactivity has been associated with smoking (23) and increased dietary fat consumption (24), and those who smoke and drink regularly tend to consume less fruits and vegetables (25). In one study that specifically examined the patterns of adolescent physical activity and dietary behaviors, the most prevalent cluster of related health behaviors was formed by not meeting the physical activity and fruit and vegetable recommendations (26). These studies imply that physical activity and dietary behaviors do not occur in isolation. The synchronized tracking of lifestyle factors suggests that these are complex, multidimensional behaviors that coexist, occur collectively in distinct patterns, and perhaps reinforce each other.

For decades, health psychologists, who examine how biological, behavioral, and psychological factors influence health and illness, have laid the groundwork to explain the connection between lifestyle behaviors through observation, theories, and tested psychological constructs. A variety of social cognitive models (27), each of which provides a slightly different view or perspective, have been developed and adapted to describe the cognitive determinants of health-related behavior change. Intra-personal models that have built a reputable evidence base to explain and guide lifestyle changes include the Health Belief Model (28), Social Cognitive Theory (29), the Theory of Reasoned Action/Theory of Planned Behavior (30), Self-Determination Theory (31), and the Transtheoretical Model of Change (32). Each model includes a set of health cognitions, thoughts and feelings that the individual associates with health-related behaviors. In combining physical activity, eating behavior, smoking, alcohol consumption, etc. into an indistinguishable group of health behaviors, these models inherently support observational evidence of clustering.

Models from social cognitive theory have been adapted to behavioral weight loss interventions, many of which report a positive interaction between reduced caloric intake and increased physical activity. In an examination of the comparative and cumulative effects of diet and physical activity behavior changes during a 2-year weight loss program, investigators noted synergistic improvements in both factors (33). A study by Jakicic et al.

showed that after 18 months of participation in a behavioral weight loss program, subjects reported increased leisure-time physical activity and adherence to eating behavior associated with weight loss (34). In turn, new behavior modification strategies are being developed (35) and integrated into the lifestyle interventions of large-scale clinical trials such as Look AHEAD (36). Although multiple lines of research imply a symbiotic interaction, it remains to be determined whether or not the concurrent adoption of elevated physical activity and dietary choices are independent processes or if these lifestyle behaviors somehow track together and might even be mediated by shared neurobiological substrates, which if identified, might lead to more effective interventions.

Physical Activity and Homeostatic vs. Hedonic Eating

Different lines of research suggest a positive influence of physical activity on the self-regulation of eating behavior by enhancing the sensitivity of the physiological satiety signaling system, adjusting macronutrient food choices or requirements, and/or changing the hedonic response to food stimuli. One area of research, which investigates the effect of physical activity on the homeostatic mechanisms that control appetite, indicates that active people have sharper hunger-satiety mechanisms, and hence, improved control of appetite (37). A review by Blundell et al. (38) showed that contrary to the popular belief that physical activity increases appetite and calorie intake, men and women can tolerate exercise-induced acute energy deficits and do not compensate by eating more. Data indicates that this holds true for both lean (39) and obese (40) cohorts. Although compensation may increase in the long term (16+ days), this compensation is partial or incomplete. Conversely, it has been shown that reducing physical activity from 1.8 to 1.4 $\times$ RMR (resting metabolic rate) does not induce a compensatory reduction in energy intake, leading instead to a positive energy balance (41). The above findings suggest that increased physical activity may have significant effects on metabolic processes, and hence, energy balance in the short term. However, these studies are limited by short duration (days or weeks) and use of predominantly normal weight subjects. Although some prospective studies that relate physical activity levels to longer-term changes in energy intake (42) and macronutrient preference (43) do exist, assessment via dietary recall has shortcomings in accurately reflecting nutrient intake and eating patterns. Measures that capture changes in the behavioral components of physical activity and food intake will help to distinguish between physiologic and behavioral changes over yearlong periods.

This discussion introduces the important distinction between the impact of physical activity on short-term metabolic adaptations versus long-term behavioral adaptations, both of which influence energy intake, but through separate mechanisms. The metabolically regulated, more automatic appetite is controlled by homeostatic mechanisms. The gut influences feeding behavior by generating hunger and satiety signals, which are communicated to the brain via peptides from the gastrointestinal tract (e.g. ghrelin) and adipose tissue (e.g. leptin) (44). The homeostatic system is comprised of the hypothalamus and the brainstem, which drive food intake based on caloric needs and energy balance (45,46). The characterization of the biochemical signals and neurobiological processes involved in hunger, meal initiation, and satiety has been the subject of extensive research since the early 20th century (47). Numerous theories—lipostatic (48), aminostatic (49), thermostatic (50), glucostatic (51), and others (52,53)—have explored the metabolic basis of food intake. The glucostatic theory, for example, incorporates neurophysiologic considerations and builds upon the shared premise that glucose uptake and utilization is central to the control of hunger, satiety, and control of energy balance. Experimental evidence from prospective studies links hypoglycemia to weight gain (54), which may be exacerbated by modern realities including sleep-deprivation or increased use of computers for knowledge-based work, both of which have been shown to induce spontaneous energy intake related to changes in glycemic control

(55,56). Although this review limits its focus to the neurocognitive dimension of hedonic eating behavior, it is important to recognize that such hypotheses have generated complementary, rather than alternative evidence of the obesogenic consequences of modern lifestyle.

Regardless of whether physical activity transiently impacts energy intake (57) or whether perturbations in glycemia increase caloric consumption in the short term (54), it is evident that sustainable weight loss requires enduring changes in the motivation to eat (58). In contrast to metabolic mechanisms that regulate satiety, this dimension of food intake is cognitive-behavioral in nature. Within the last decade, it has been recognized that an increasing proportion of human food consumption is driven by the pleasure, known as “hedonic hunger” (59), rather than the homeostatic principles of energy balance. In turn, increased attention has been focused on the influence of reward sensitivity (60), the brain reward system (61), and the mechanisms underlying liking and wanting (62). This reward-related appetite is centered in the brain’s mesolimbic dopamine system, including the ventral tegmental area of the midbrain and extending to the nucleus accumbens in the striatum (63). In an obesity-promoting environment, the hedonic response to food stimuli can disrupt and override the homeostatic mechanisms that regulate satiety (64).

Presently, the distinction between the homeostatic and hedonic appetite is blurred by evidence that these influences may change over time (65). In future studies, it will be important to delineate the separation between these forces. For the purposes of this review, which focuses on long-term weight control for 2+ years, the discussion concentrates on the behavioral self-regulation of the hedonic response to food. Changes in behavior patterns from those that promote hedonic overeating to those that support healthy weight maintenance result from changes in neurobiological substrates and the formation of new connections between neural circuits. In order to better identify, effectively target, and ultimately adapt the impulsive, reward-seeking component of eating behavior, a deeper understanding of the underlying neurocognitive mechanisms is essential.

Transitioning from behavior to neurocognition

Health psychology and social cognitive theory built the framework upon which cognitive neuroscience is now investigating neural systems and neurophysiologic mechanisms that underlie behaviors. Theoretical concepts from behavioral psychology such as reasoning, self-control, and decision-making are now being identified through quantitative tests that target cognitive processes and identify underlying brain regions and circuits (66). With regard to physical activity and eating behaviors, in order to build upon the behavioral association and to propose a neurocognitive explanation, a brief overview of the neurocognitive basis of human behavior is instructive.

Behaviors often involve decision making, irrespective of whether these choices are evaluated rationally (67) or influenced by emotions (68). This decision-making process is mediated by our neural systems (69). Individuals constantly monitor behavior through a process called self-regulation, which involves re-evaluating beliefs, setting goals, and keeping thoughts and behaviors in line with such goals (70). Self-regulation falls under the loosely defined umbrella of executive functions, a group of advanced cognitive processes that break the normal flow from impulse to action. By halting automatic actions, executive processes facilitate the execution of goal-directed behavior, intentioned steps to complete a task. Goal-oriented processes include the ability to initiate and stop actions, to monitor and change behavior as needed, and to plan future behavior when presented with novel situations and environments. Experimentally, executive functions manifest in various cognitive components such as working memory, mental set shifting and inhibitory control (71), as

well as more complex functions such as decision-making, delayed-gratification or self-regulation. Neuroimaging studies have mapped these executive functions (also called cognitive control) to the brain's prefrontal cortex, a region that coordinates thought and action in accordance with internal goals (72). A growing body of literature indicates that in addition to the homeostasis and reward centers of the brain, the prefrontal cortex is integrally involved in the regulation of the lifestyle behaviors that are contributing to the obesity epidemic (73).

Until recently, neuroscientific obesity-related research has focused primarily on the connection between the hypothalamus and the gastrointestinal system in the regulation of appetite (74). However, as evidenced by the prevalence of obesity, the metabolic circuits that regulate energy homeostasis can be impaired and desensitized, leading to overeating; the human perception of and behavior towards food can override neuroendocrine satiety signals. For our ancestors, food intake was a means of subsistence and vigorous physical activity was a daily requirement. Modern realities have now shifted these necessities into the category of "lifestyle behaviors" (75). The "toxic environment" (76) in which we live promotes sedentary behavior and overconsumption of calories. From ubiquitous junk food advertisements to the diminished imperative for physical activity, our neural systems are repeatedly bombarded with health-detering stimuli. In essence, healthy behaviors are no longer supported externally; we must cultivate and fortify them internally. Aside from considerations of socioeconomic status, cultural norms, and religious beliefs, each individual now bears the onus of implementing a set of health behaviors that collectively comprise lifestyle.

The prefrontal cortex "top-down" inhibitory control, and eating behavior

Our cognitive processes underlie our behavior, which in turn, determines aspects of lifestyle, such as dietary pattern, and hence, nutritional intake. For the purposes of this discussion, our eating behavior is best conceptualized as an extension of Satpute and Lieberman's neurocognitive approaches (77) of the reflexive and reflective dual process models (73, 78) in the field of social cognition. The reflexive system is the "default responder," meaning that it is responsible for our automatic, unconscious reactions. We share the reflexive system with other primates, suggesting that this system is evolutionarily old. The reflexive system favors reward and has been traced to dopamine-responsive neural circuits in the limbic and paralimbic structures. Hence, this system drives hedonic overeating. In contrast, the reflective system is the more recent human adaptation that enables us to make rational, typically conscious decisions. This system demands greater cognitive resources in order to override the reflexive system and execute goal-oriented actions. The reflective system controls our executive processes and has been traced to the prefrontal cortex and associated cortices, anatomical brain regions that have undergone a remarkable expansion in the human species. Importantly, the reflective system is also more plastic (79), meaning that it is adaptable and capable of integrating new habits (80). Thus, despite predetermined genetic differences in anatomy and connections, such initial substrates might be modified and ultimately lead to new and improved lifestyle behaviors.

With regard to nutrient intake, executive functions are necessary to regulate and inhibit the impulsive, hedonic response to food (81). Impulsivity can be generally defined as the tendency to think, control, and plan insufficiently, which is therefore inaccurate or maladaptive (82). This behavioral trait suggests a shortfall or repression of our reflective system, the system that includes our executive functions. When defined as an insufficient response inhibition, impulsivity can be seen as the inability to stop a predominant action, e.g. a movement that has been overlearned. Inhibitory control exerts the stop signal necessary to override these automatic responses, thereby enabling goal-oriented actions. The

prefrontal cortex guides this “top-down” processing in order to align behavior with internal states or intentions (83). For example, if a cookie lover is presented with a warm chocolate chip cookie, the automatic response would be to eat the cookie; however, if this impulse to eat conflicts with that person’s dietary goals, executive functions must suppress the impulse to maintain dietary restriction.

A useful cognitive test to assess inhibitory control is the stop-signal task (84), during which participants are directed to respond to a visual cue by pressing a button, but to stop pushing when they hear a stop tone. Neederkorn and colleagues have employed the stop-signal task to show that in general, obese women (85) and children (86) tend to be more impulsive. From this follows a series of studies showing that a less efficient response inhibition (i.e. greater impulsivity) is related to increased food intake and overeating (87), increased weight and obesity (88), and less weight loss during weight reduction treatment (89). This group has also demonstrated a causal connection between overeating and impulsivity by inducing impulsivity in high- and low-restrained eaters (90). Data indicates that overeating follows from an interaction between restraint and impulsivity. High-restraint, high-impulsive subjects tend to overeat after being primed with a tasty preload or the aroma of food; in contrast, those who exhibit high-restraint but are lower on the impulsivity scale do not overeat to the same extent (91). Thus, behavioral disinhibition, a deficit in or repression of this inhibitory check, may play a prominent role in overeating, and hence facilitate the development of obesity over time.

Insight into the neurocognitive link between disinhibition and overeating comes from conditions explained by defective inhibitory control, most notably attention deficit/hyperactivity disorder (ADHD). In his Unifying Theory of ADHD, Russell A. Barkley argues that inhibitory control sets the occasion for the performance of self-directed actions by providing the delay necessary for them to occur (92). The prefrontal cortex provides the brake necessary for the execution of goal-oriented behavior, and this brake is deficient in ADHD. Overeating may be partially explained by this model as studies confirm strong associations between overweight/obesity and symptoms of ADHD in children, adolescents, and adults (93,94). Research has also shown that higher than expected rates of binge eating, which is characterized by bouts of unrestrained eating, occur in individuals with ADHD (95).

A second way to gauge impulsivity is to think of it as sensitivity to reward, which can be assessed by the ability to delay gratification, or to wait in order to attain something one wants (96). Walter Mischel’s famous “marshmallow experiments” (97), which examined the cognitive mechanisms that enable a young child to forego immediate satisfaction and to wait for a larger but delayed reward, understood the inherent link between delay of gratification and eating behavior. Continuing research with the original participants suggests that the ability to delay gratification during the preschool years predicts future outcomes, including body mass index and drug use (98). Recently, a prospective study showed that students who failed the delayed gratification task at 4 years of age were more likely to be overweight at 11 years old (99). Similarly, increased adiposity has been linked to poorer performance on the Iowa Gambling Task (100), a psychological test that assesses reward sensitivity, among other cognitive components, by choosing between immediate versus eventual monetary gains. Delay of gratification and delay discounting in the Iowa Gambling Task are two constructs that have been useful for examining inhibition of impulsive overeating (101,102). Of course, the intuitive link between overeating and impulsivity is that successful weight loss, a delayed reward, requires avoiding triggers and sacrificing the immediate gratification of overeating.

Neuroimaging studies in both the right (103) and the left (104) prefrontal cortex bolster Mischel's theory by suggesting this region's involvement in delaying gratification. Furthermore, neuroimaging studies confirm a role for the prefrontal cortex in hedonic feeding inhibition by linking activation patterns with motivation to consume palatable foods. A study published in 2001 (105) examined the activation patterns in response to continuous ingestion of chocolate by self-identified chocolate fanatics. Decreased motivation to consume more chocolate over time was associated with increased activation of the lateral prefrontal cortex, suggesting an inhibitory signal. In contrast, exposure to appetizing food cues results in preferential activation of hedonic regions of the brain, with obese individuals showing greater activation in these areas (106,107). Transcranial magnetic stimulation, a technique that noninvasively modulates cortical activity, is now being tested as a therapeutic intervention for such impulsive eating habits (supported by the Klarman Family Foundation). Preliminary trials indicate that enhancing activity in both the left (108) and the right (109) dorsolateral prefrontal cortices with noninvasive brain stimulation can suppress food cravings (110). These studies suggest that the amount of activity in the prefrontal cortex may determine the degree of inhibition over downstream reward circuits that encourage what was once an evolutionarily advantageous impulse to overeat.

Conversely, the ability to adopt and maintain healthy eating habits has been linked to greater inhibitory control. Studies show that increases in cognitive restraint and self-efficacy (111,112) and decreases in emotional, disinhibited eating (113) are predictive of successful weight maintenance. As shown in analysis of members of the National Weight Control Registry (114), an ongoing study of more than 7000 weight loss maintainers, one of the critical predictors of successful weight loss maintenance is a lower level of dietary disinhibition (115,116). This is supported by neuroimaging studies of successful weight loss participants, which report greater activation in the prefrontal regions in response to food cues (117). After ingestion of a meal, successful dieters show increased neural activity in the dorsolateral prefrontal cortex, which is involved in behavioral control and self-restraint (118). In contrast, obese men (119) and women (120) demonstrate less activation of the dorsolateral prefrontal cortex in response to a meal than do lean or formerly obese men and women.

This evidence lends strong support to the notion that eating behavior is not regulated solely by homeostatic mechanisms centered in the hypothalamus; executive functions, particularly inhibitory control, and the prefrontal lobes also appear to play an important role. While this concept of impulsivity may connote a lack of self-control or willpower, we maintain that impulsivity is becoming a more compulsory response in our obesogenic environment. The reflective brain network, which encompasses the prefrontal circuits that guide executive functions and goal-oriented behaviors, is of a limited capacity (77). Controlled processes, such as abiding by dietary goals and resisting temptations to indulge, tax and eventually exhaust the cognitive resources that regulate more emotional, automatic responses. This strain is exacerbated by stimuli in the obesogenic environment, which suggests that maintaining a healthy weight in our modern lifestyle demands an elevated and durable cognitive reserve. Regardless of whether or not the individual has a predisposed deficit in executive function or a dysfunction of prefrontal circuits, the absence of external support to enable healthy eating habits places the burden of behavior change on the individual. The aforementioned association between cognitive disinhibition and excessive caloric intake begs the question of whether or not the cognitive resources necessary to counteract obesogenic stimuli and fortify inhibitory control can somehow be "exercised" and enhanced internally. If so, it logically follows that goal-oriented behavior, as epitomized by healthy dietary choices, can be improved, refined, and perhaps maintained. In determining how to boost executive functions, it is now instructive to consider the burgeoning literature on the cognitive benefits associated with the companion health behavior: physical activity.

Physical activity can improve executive functions, self-efficacy, and mood

A surge of research over the past decade provides supportive evidence that physical activity is positively correlated with neurogenerative, neuroadaptive, and neuroprotective processes. Data from cross-sectional studies and randomized clinical trials suggest positive influences of fitness training on human brain structure and cognitive function (121). After assessing 18 studies published from 1966 to 2001 in a meta-analysis, it was reported that exercise broadly improves cognitive function across a number of domains, particularly executive functions, which appear to be more sensitive to enhancement as compared to other aspects of cognition (122). Kramer et al. showed that over a 6-month period, a group that engaged in moderate walking not only became more aerobically fit, but also showed enhanced executive processing, as indexed by better performance in task switching, stopping, and selective attention tasks, compared with the stretching and toning control group (123). In agreement with these findings, functional MRI data shows that modulation of physical activity disproportionately influences tasks that necessitate greater amounts of executive control (124). This coincides with evidence that brain regions related to executive control, including prefrontal circuits, are more plastic (125), structurally sound (126), and even prone to growth (127) with elevated levels of aerobic fitness.

Given the evidence that increasing physical activity improves executive function, it seems that decreased physical activity would contribute to or at least be associated with deficits in similar cognitive processes. Indeed, when researchers at Berlin's Center for Space Medicine put 24 healthy males into the bed rest model of prolonged weightlessness, scores on the Iowa Gambling Task, which has been associated with regulation of eating behavior, were significantly lower during bed rest than before or after the intervention (128). This finding is supported by studies which report that a combination of high disinhibition, a disposition for opportunistic eating, and low self-restraint is associated with higher sedentary behavior (129). Exercise has been shown to exert a positive influence on those with a high disinhibition score by decreasing motivation to eat (130) and increasing preference for low-fat foods (131). These cognitive measures are supported anatomically by the intertwined development between the adjacent prefrontal and motor cortices (132). Since exercise has been shown to enhance connectivity between distinct functional networks in separate brain regions (124), increased use of the motor cortex during physical activity may enhance executive processes and vice versa. Although this remains speculative, neurocognitive underpinnings may help to explain the reported correlation between sedentary activities, such as time spent watching television, and energy intake (133).

Both cognitive testing and neuroimaging studies show that regular physical activity enhances executive functions, and these same cognitive processes are implicated in the regulation of impulsive eating behavior. This cognitive convergence is further supported by research on depression and mood, which are integrally linked to cognition and for which both physical activity and eating behavior appear to be key mediators (134,135). Depression and obesity are positively related in a bidirectional relationship (136). Emotional eating, which refers to the tendency to eat in response to negative emotions, has been associated with elevated levels of depressive symptoms (137), the consumption of high calorie foods (138), and higher body mass index (139). In line with this, a lack of physical activity self-efficacy, a person's confidence in his or her ability to be physically active on a regular basis, has also been linked to higher depressive symptoms (140). A recent study by Kontinen et al. (141) used mediation analysis to confirm both of these associations within the same cohort. Although this study examined emotional eating and physical activity self-efficacy as independent pathways between depressive symptoms and higher adiposity, the negative correlation between these factors suggests an interaction.

In contrast, successful adoption of a physically activity lifestyle may positively influence eating behavior through its effect on variables such as motivation, commitment, and self-efficacy (31,142). Evidence shows that increases in exercise motivation and self-efficacy predict positive changes in the self-regulation of eating behavior, and the resulting confidence could “spill-over” into other behaviors important for weight management (143). Both structured exercise (e.g. running, cycling, fitness classes) and lifestyle activities (e.g. walking) have been positively associated with weight control (144) and eating self-regulation. Andrade et al. recently showed that both vigorous exercise and lifestyle activities improve several eating behavior variables that describe dietary self-regulation (145). These variables mediated the relationship between physical activity and 1-year change in body weight. This indicates that adoption of physical activity may promote a maintained decrease in body weight not only through an immediate increase in energy expenditure, but also by changing psychological markers of eating behavior, and may also explain why physical activity has been shown to play the paramount role in long-term weight management (7).

Depression provides insight into possible mechanisms by which increased physical activity can improve weight control via strengthening the cognitive resources needed for hedonic restraint. Negative emotional states promote overeating by depleting the cognitive resources necessary to resist the temptation to overeat (146). Physical activity, perhaps by improving mood (147) and/or reducing anxiety and stress (148) counters the reflexive impulse to overeat. In the largest randomized control trial to date, with a cohort of 200 clinically depressed adults, Blumenthal et al. showed that four months of aerobic exercise achieved reductions in depression comparable to standard antidepressant medication and greater than placebo (149). Neurocognitive research shows that depression is characterized by hypoactivity of the prefrontal cortex (150). In the presented framework of prefrontal inhibitory control, this would be predicted to be associated with a dampened or suppressed reflective system. Hence, given physical activity’s therapeutic effect on depression and depression’s link to unrestrained eating, it seems plausible that physical activity could enhance those same cognitive processes that regulate hedonic eating behavior.

Physical activity modifies eating behavior through a neurocognitive connection

Connecting the pieces between physical activity and executive function with eating behavior and executive function suggests that rather than a coincidental association, physical activity and eating behavior may share a common neurocognitive platform. As shown diagrammatically in Figure 1, physical activity builds the cognitive resources, namely inhibitory control, necessary to block impulsive actions that may have adverse health consequences, specifically overeating. In this sense, physical activity is a “gateway behavior” that may induce people to improve their diets over time (151, 152). Since the mid 1980s, health psychologists have posited that physical activity can be an inherently rewarding activity that contributes to both health and subjective vitality. By filling a psychological need of competence and enjoyment, adherence to physical activity may affirm other healthy behaviors (153). In today’s world, in an environment that fosters sedentary behavior and overeating, practicing regular physical activity and adhering to healthy diet are necessarily self-regulatory, goal-oriented behaviors. From a neurocognitive perspective, top-down inhibitory control enables us to repress health-averse stimuli and execute health-promoting behaviors.

In addition to this neurocognitive basis, the choice of physical activity as an effective first target for weight loss also garners critical support from the physiologic/homeostatic angle. Regular physical activity improves insulin sensitivity (154), increases fat oxidation, and enhances the body’s response to leptin (155). Starting weight loss with physical activity may

represent an investment in physiological functionality to counteract the hypoglycemia and weight regain that often follows weight loss (156). This point reinforces the idea that the more immediate, homeostatic regulation of appetite and energy intake complements and perhaps interacts with and/or stimulates some of the neurocognitive adaptations and subsequent behavior changes over time. Physical activity has been shown to be one of, if not the most important factor in long-term weight maintenance (8), but whether this is due to metabolic changes, neurocognitive adaptations, or more likely a combination of both remains to be determined. Collaborative projects that integrate multiple aspects of eating behavior and physical activity are a priority for future research.

This hypothesis is grounded in the link between executive function and eating behavior, which suggests that suboptimal prefrontal cortex function, or more aptly, cognitive resources under constant strain in an obesogenic environment, may contribute to the etiology of obesity. The mounting evidence that positively correlates increased physical activity with enhanced executive function may represent a viable therapeutic approach to controlling and modifying eating behavior. The first step in building an evidence base for this hypothesis would be to establish the causal relationship between physical activity and enhanced cognition. Then, it would be important to investigate whether or not exercise causes an enhancement in inhibitory control, or other key components of executive functions. It would also be instructive to concurrently encourage and monitor new studies investigating the behavioral impact of physical activity interventions and pharmacotherapies on ADHD patients as a positive impact would presumably act through a similar neurocognitive mechanism. Whereas this review covers physical activity in the broad sense, including both exercise and lifestyle activities, it will be equally critical to determine intensity thresholds, durations, or specific activities that best strengthen and refine executive functions. Can different modes of exercise, such as aerobic and resistance training, produce the same outcome on these cognitive tests? We would need to develop an evidence-based program to adopt physical activity in order to induce and sustain the desired neurobiological effects and consequential eating behavior modifications. Connecting these spheres of research opens expansive investigative opportunities with important translational implications for the treatment of obesity and adoption of a healthy lifestyle.

Conclusion

To the best of our knowledge, this is the first review to combine the robust literature on physical activity and eating behavior in order to hypothesize a neurocognitive link. This proposal is meant to spark interaction among neuroscientists, psychologists, endocrinologists, exercise physiologists, dieticians, and clinicians to promote integrative, translational research. Our hypothetical model has widespread implications for both clinical care and public policy. When designing the most effective and efficient approach to weight loss and subsequent weight loss maintenance, it is critical to understand which to target first — physical activity or diet—based not only on its effect on energy balance, but on its ability to positively impact the other behavior. Perhaps elevated physical activity can facilitate changes in related healthy lifestyle behaviors by “training” the cognitive resources necessary for their execution. Considering the fallibility of fad diets and the underperformance of clinical treatment strategies, it seems plausible to approach from the physical activity angle. In this way, we may facilitate healthy and steady weight loss by dimming the often overbearing spotlight on weight loss. With regard to public policy, we take particular issue with the fact that financially-strained school systems are cutting physical education classes and unstructured recess time. By reducing physical activity in schools, we may be propagating the childhood obesity crisis not only by reducing energy expenditure, but also by worsening eating habits. Ironically, too much work with no play may be weakening those cognitive abilities that we are so eager to test (157).

As embodied by our developmentally advanced prefrontal cortex, humans display complex yet interrelated behaviors, the underlying neurocognitive components of which must be elucidated to help treat lifestyle-related diseases. Investigating physical activity and other possible modes of enhancing executive function may increase our understanding of how to control appetite and enhance inhibitory mechanisms that regulate eating behavior. This could improve adherence to diet and exercise therapy, ultimately facilitating adoption of these behaviors.

Acknowledgments

MA-A and AP-L were partly supported by a Davis Foundation Award of the Klarman Family Foundation Grants Program in Eating Disorders, and by Grant Number UL1 RR025758 – Harvard Clinical and Translational Science Center, National Center for Research Resources. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Center for Research Resources or the National Institutes of Health.

References

1. Flegal KM, Carroll MD, Ogden CL, Curtin LR. Prevalence and trends in obesity among US adults, 1999–2008. *JAMA*. 2010; 303:235–241. [PubMed: 20071471]
2. World Health Organization. [accessed 6/1/2011] Controlling the global obesity epidemic [WWW document]. URL <http://www.who.int/nutrition/topics/obesity/en/index.html>
3. Curioni CC, Lourenco PM. Long-term weight loss after diet and exercise: a systematic review. *Int J Obes (Lond)*. 2005; 29:1168–1174. [PubMed: 15925949]
4. Davis MJ, Addis ME. Predictors of attrition from behavioral medicine treatments. *Ann Behav Med*. 1999; 21:339–349. [PubMed: 10721442]
5. Brownell KD, Wadden TA. Etiology and treatment of obesity: understanding a serious, prevalent and refractory disorder. *J Consult Clin Psychol*. 1992; 60:505–517. [PubMed: 1506499]
6. Jeffery RW, Drewnowski A, Epstein LH, Stunkard AJ, Wilson GT, Wing RR, Hill DR. Long-term maintenance of weight loss: current status. *Health Psychol*. 2000; 19:5–16. [PubMed: 10709944]
7. Pronk NP, Wing RR. Physical activity and long-term maintenance of weight loss. *Obes Res*. 1994; 2:587–599. [PubMed: 16358405]
8. Hankinson AL, Daviglus ML, Bouchard C, Carnethon M, Lewis CE, Schreiner PJ, Liu K, Sidney S. Maintaining a high physical activity level over 20 years and weight gain. *JAMA*. 2010; 304:3603–3610.
9. James, W. *The Principles of Psychology*. New York: Henry Holt and Company; 1890.
10. The Diabetes Prevention Program Research Group. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *N Engl J Med*. 2002; 346:393–403. [PubMed: 11832527]
11. Look AHEAD Research Group, Wing RR. Long-term effects of a lifestyle intervention on weight and cardiovascular risk factors in individuals with type 2 diabetes mellitus: four-year results of the Look AHEAD trial. *Arch Internal Med*. 2010; 170:1566–1575. [PubMed: 20876408]
12. Johnson NA, Boyle CA, Heller RF. Leisure-time physical activity and other health behaviors: are they related? *Aust J Public Health*. 1995; 19:69–75. [PubMed: 7734598]
13. Matthews CE, Herbert JR, Ockene IS, Saperia G, Merriam PA. Relationship between leisure-time physical activity and selected dietary variables in the Worcester Area Trial for Counseling in Hyperlipidemia. *Med Sci Sports Exerc*. 1997; 29:1199–1207. [PubMed: 9309632]
14. Brodney S, Mcpherson RS, Carpenter RS, Welten D, Blair SN. Nutrient intake of fit and unfit men and women. *Med Sci Sports Exerc*. 2001; 33:459–467. [PubMed: 11252075]
15. Eaton CB, McPhillips JB, Gans KM, Garber CE, Assaf AR, Lasater TM, Carleton RA. Cross sectional relationship between diet and physical activity in two southeastern New England communities. *Am J Prev Med*. 1995; 11:238–244. [PubMed: 7495600]
16. Parsons TJ, Power C, Manor O. Longitudinal physical activity and diet patterns in the 1958 British birth cohort. *Med Sci Sports Exerc*. 2006; 38:547–554.

17. Berrigan D, Dodd K, Troiano RP, Krebs-Smith SM, Barbash RB. Patterns of health behavior in U.S. adults. *Prev Med.* 2003; 36:615–623. [PubMed: 12689807]
18. Patterson RE, Haines PS, Popkin BM. Health lifestyle patterns of US adults. *Prev Med.* 1994; 23:453–460. [PubMed: 7971872]
19. Blair SN, Jacobs DR, Powell KE. Relationships between exercise or physical activity and other health behaviors. *Public Health Rep.* 1985; 100:172–180. [PubMed: 3920715]
20. Pronk NP, Anderson LH, Crain AL, Martinson BC, O'Connor PJ, Sherwood NE, Whitebird RR. Meeting recommendations for multiple healthy lifestyle factors. Prevalence, clustering, and predictors among adolescent, adult, and senior health plan members. *Am J Prev Med.* 2004; 27:25–33. [PubMed: 15275671]
21. Schuit AJ, van Loon AJ, Tijhuis M, Ocké M. Clustering of lifestyle risk factors in a general adult population. *Prev Med.* 2002; 35:219–224. [PubMed: 12202063]
22. Raitakari OT, Leino M, Rääkkönen K, Porkka KV, Taimela S, Räsänen L, Viikari JS. Clustering of risk habits in young adults. The Cardiovascular Risk in Young Finns Study. *Am J Epidemiol.* 1995; 142:36–44. [PubMed: 7785672]
23. Emmons KM, Marcus BH, Linnan L, Rossi JS, Abrams DB. Mechanisms in multiple risk factor interventions: smoking, physical activity, and dietary fat intake among manufacturing workers. *Prev Med.* 1994; 23:481–489. [PubMed: 7971876]
24. Simoes EJ, Byers T, Coates RJ, Serdual MK, Mokdad AH, Heath GW. The association between leisure-time physical activity and dietary fat in American adults. *Am J Public Health.* 1995; 85:240–244. [PubMed: 7856785]
25. Subar AF, Harlan LC, Mattson ME. Food and nutrient intake differences between smokers and nonsmokers in the US. *Am J Public Health.* 1990; 80:1323–1329. [PubMed: 2240298]
26. Pearson N, Atkin AJ, Biddle SJ, Gorely T, Edwardson C. Patterns of adolescent physical activity and dietary behaviours. *Int J Behav Nutr Phys Act.* 2009; 6:45. URL <http://www.ijbnpa.org/content/6/1/45>. [PubMed: 19624822]
27. Conner, M. Cognitive determinants of health behavior. In: Steptoe, A., editor. *Handbook of Behavioral Medicine: Methods and Applications*. Springer; New York: 2010. p. 19-30.
28. Abraham, C.; Sheeran, P. The health belief model. In: Connor, M.; Norman, P., editors. *Predicting Health Behavior: Research and Practice with Social Cognition Models*. 2nd Ed.. Maidenhead: Open University Press; 2005. p. 28-80.
29. Bandura A. Self-efficacy: Toward a unifying theory of behavioral change. *Psychol Rev.* 1977; 84:191–215. [PubMed: 847061]
30. Ajzen I. The theory of planned behavior. *Organiz Behav Hum Dec Proc.* 1991; 50:179–211.
31. Deci, EL.; Ryan, RM. *Intrinsic motivation and self-determination in human behavior*. New York, NY: Plenum Press; 1985.
32. Prochaska, JO.; DiClemente, CC. *The Transtheoretical Approach: Crossing Traditional Boundaries of Therapy*. Homewood, IL: Dow Jones Irwin; 1984.
33. Dunn CL, Hannan PJ, Jeffery RW, Sherwood NE, Pronk NP, Boyle R. The comparative and cumulative effects of a dietary restriction and exercise on weight loss. *Int J Obes.* 2006; 30:112–121.
34. Jakicic JM, Wing RR, Winters-Heart C. Relationship of physical activity to eating behaviors and weight loss in women. *Med Sci Sports Exer.* 2002; 34:1653–1659.
35. West DS, Gorin AA, Subak LL, Foster G, Bragg C, Hecht J, Schembri M, Wing RR. A motivation-focused weight loss maintenance program is an effective alternative to a skill-based approach. *Int J Obes (Lond).* 2011; 35:259–269. [PubMed: 20680012]
36. Ryan DH, Espeland MA, Foster GD, Haffner SM, Hubbard VS, Johnson KC, Kahn SE, Knowler WC, Yanovski SZ. Look AHEAD Research Group. Look AHEAD (Action for Health in Diabetes): design and methods for a clinical trial of weight loss for the prevention of cardiovascular disease in type 2 diabetes. *Control Clin Trials.* 2003; 24:610–628. [PubMed: 14500058]
37. Martins C, Morgan L, Truby H. A review of the effects of exercise on appetite regulation: an obesity perspective. *Int J Obes.* 2008; 32:1337–1347.

38. Blundell JE, Stubbs RJ, Hughes DA, Whybrow S, King NA. Cross-talk between physical activity and appetite control: does PA stimulate appetite? *Proc Nutr Soc.* 2003; 62:651–661. [PubMed: 14692601]
39. Whybrow S, Hughes DA, Ritz P, Johnstone AM, Horgan GW, King N, Blundell JE, Stubbs RJ. The effect of an incremental increase in exercise on appetite, eating behavior and energy balance in lean men and women feeding ad libitum. *Br J Nutr.* 2008; 100:1109–1115. [PubMed: 18377694]
40. Unick JL, Otto AD, Goodpaster BH, Helsel DL, Pellegrini CA, Jakicic JM. The acute effect of walking on energy intake in overweight/obese women. *Appetite.* 2010; 55:413–419. [PubMed: 20674640]
41. Stubbs RJ, Hughes DA, Johnstone AM, Horgan GW, King N, Blundell JE. A decrease in physical activity affects appetite, energy, and nutrient balance in lean men feeding ad libitum. *Am J Clin Nutr.* 2004; 79:62–69. [PubMed: 14684398]
42. Waling M, Lind T, Hernell O, Larsson C. A one-year intervention has modest effects on energy and macronutrient intakes of overweight and obese Swedish children. *J Nutr.* 2010; 140:1793–1798. [PubMed: 20739446]
43. Dutton GR, Napolitano MA, Whiteley JA, Marcus BH. Is physical activity a gateway behavior for diet? Findings from a physical activity trial. *Prev Med.* 2008; 46:216–221. [PubMed: 18234327]
44. Wren AM, Bloom SR. Got hormones and appetite control. *Gastroenterology.* 2007; 132:2116–2130. [PubMed: 17498507]
45. Arora S, Chitkara A. Role of neuropeptides in appetite regulation and obesity—a review. *Neuropeptides.* 2006; 40:375–401. [PubMed: 16935329]
46. Berthoud HR, Morrison C. The brain, appetite, and obesity. *Annu Rev Psychol.* 2008; 59:55–92. [PubMed: 18154499]
47. Carlson, AJ., editor. *The Control of Hunger in Health and Disease.* Chicago, IL: University of Chicago Press; 1916.
48. Kennedy GC. The hypothalamic control of food intake in rats. *Proc R Soc Lond B Biol Sci.* 1950; 889:535–549. [PubMed: 14797669]
49. Mellinkoff SM, Frankland M, Boyle D, Greipel M. Relationship between serum amino acid concentration and fluctuations in appetite. *J Appl Physiol.* 1956; 8:535–538. [PubMed: 13295170]
50. Brobeck JR. Food intake as a mechanism of temperature regulation. *Yale J Biol Med.* 1948; 20:545–552. [PubMed: 18872321]
51. Mayer J. Glucostatic mechanism of regulation of food intake. *N Engl J Med.* 1953; 249:13–16. [PubMed: 13063674]
52. Schwartz MW, Woods SC, Porte D Jr, Seely RJ, Baskin DG. Central nervous system control of food intake. *Nature.* 2000; 404:661–671. [PubMed: 10766253]
53. Bray GA, Campfield LA. Metabolic factors in the control of energy stores. *Metabolism.* 1975; 24:99–117. [PubMed: 1089189]
54. Boulé NG, Chaput JP, Doucet E, Richard D, Després JP, Bouchard C, et al. Glucose homeostasis predicts weight gain: prospective and clinical evidence. *Diabetes Metab Res Rev.* 2008; 24:123–129. [PubMed: 17879957]
55. Chaput JP, Despre's JP, Bouchard C, Tremblay A. Association of sleep duration with type 2 diabetes and impaired glucose tolerance. *Diabetologia.* 2007; 50:2298–2304. [PubMed: 17717644]
56. Chaput JP, Drapeau V, Poirier P, Teasdale N, Tremblay A. Glycemic instability and spontaneous energy intake: association with knowledge-based work. *Psychosom Med.* 2008; 70:797–804. [PubMed: 18725427]
57. King NA, Hopkins M, Caudwell P, Stubbs RJ, Blundell JE. Individual variability following 12 weeks of supervised exercise: identification and characterization of compensation for exercise-induced weight loss. *Int J Obes.* 2008; 32:177–184.
58. Wing RR, Papandonatos G, Fava JL, Gorin AA, Phelan S, McCaffery J, Tate DF. Maintaining large weight losses: the role of behavioral and psychological factors. *J Consult Clin Psychol.* 2008; 76:1015–1021. [PubMed: 19045969]
59. Lowe MR, Butryn ML. Hedonic hunger: a new dimension of appetite? *Physiol Behav.* 2007; 91:432–439. [PubMed: 17531274]

60. Davis C, Levitan R, Kaplan A, Carter J, Reid C, Curtis C, et al. Reward sensitivity and the D2 dopamine receptor gene: a case-control study of binge eating disorder. *Prog Neuropsychopharmacol Biol Psychiatry*. 2008; 32:620–628. [PubMed: 18262320]
61. Berridge K, Kringelbach M. Affective neuroscience of pleasure: reward in humans and animals. *Psychopharmacology*. 2008; 199:457–480. [PubMed: 18311558]
62. Finlayson G, King N, Blundell J. Liking vs. wanting food: importance for human appetite control and weight regulation. *Neurosci Biobehav Rev*. 2007; 31:987–1002. [PubMed: 17559933]
63. Haber SN, Knutson B. The reward circuit: linking primary anatomy and human imaging. *Neuropsychopharmacology*. 2010; 35:4–26. [PubMed: 19812543]
64. Hofmann W, van Koningsbruggen GM, Stroebe W, Ramanathan S, Aarts H. As pleasure unfolds: hedonic responses to tempting food. *Psychol Sci*. 2010; 21:1863–1870. [PubMed: 21106885]
65. King NA, Caudwell P, Hopkins M, Byrne NM, Colley R, Hills AP, Stubbs JR, Blundell JE. Metabolic and behavioral compensatory responses to exercise interventions: barriers to weight loss. *Obesity*. 2007; 15:1373–1383. [PubMed: 17557973]
66. Carlson, NR. *Physiology of Behavior*. 10th edn.. Boston, MA: Allyn & Bacon; 2009.
67. Hsu, M.; Preuschoff, K. The Neurobiological Foundations of Valuation in Human Decision Making under Uncertainty. In: Glimcher, PW.; Camerer, CF.; Fehr, E.; Poldrack, RA., editors. *Neuroeconomics: Decision Making and the Brain*. New York, NY: Academic Press; 2008. p. 351–363.
68. Loewenstein GF, Weber EU, Hsee CK, Welch N. Risk as feelings. *Psychol Bull*. 2001; 127:267–286. [PubMed: 11316014]
69. Heekeren HR, Marrett S, Ungerleider LG. The neural systems that mediate human perceptual decision-making. *Nat Rev Neurosci*. 2008; 9:467–479. [PubMed: 18464792]
70. Austin JT, Vancouver JB. Goal constructs in psychology: Structure, process, and content. *Psychol Bull*. 1996; 3:338–375.
71. Miyake A, Friedman NP, Emerson MJ, Witzki AH, Howerter A, Wager TD. The unity and diversity of executive functions and their contributions to complex “Frontal Lobe” tasks: a latent variable analysis. *Cogn Psychol*. 2000; 41:49–100. [PubMed: 10945922]
72. Miller EK, Cohen JD. An integrative theory of prefrontal cortex function. *Annu Rev Neurosci*. 2001; 24:167–202. [PubMed: 11283309]
73. Alonso-Alonso M, Pascual-Leone A. The right brain hypothesis for obesity. *JAMA*. 2007; 297:1819–1822. [PubMed: 17456824]
74. Neary NM, Goldstone AP, Bloom SR. Appetite regulation: from the gut to the hypothalamus. *Clin Endocrinol (Oxford)*. 2004; 60:153–160.
75. Jones LR, Wilson CI, Wadden TA. Lifestyle modification in the treatment of obesity: an educational challenge and opportunity. *Clin Pharmacol Ther*. 2007; 81:776–779. [PubMed: 17361122]
76. Wadden TA, Brownell KD, Foster GD. Obesity: Responding to the global epidemic. *J Consult Clin Psych*. 2002; 70:510–25.
77. Satpute AB, Leiberhan MD. Integrating automatic and controlled processes into neurocognitive models of social cognition. *Brain Res*. 2006; 1079:86–97. [PubMed: 16490183]
78. Evans JS. Dual Processing accounts of reasoning, judgement, and social cognition. *Annu Rev Psychol*. 2008; 59:255–278. [PubMed: 18154502]
79. Pascual-Leone A, Amedi A, Fregni F, Merabet LB. The plastic human brain cortex. *Annu Rev Neurosci*. 2005; 28:377–401. [PubMed: 16022601]
80. Graybiel AM. Habits, rituals, and the evaluative brain. *Annu Rev Neurosci*. 2008; 31:359–87. [PubMed: 18558860]
81. Guerrieri R, Nederkoorn C, Jansen A. The effect of an impulsive personality on overeating and obesity: current state of affairs. *Psychological Topics*. 2008; 17:265–286.
82. Solanto MV, Abikoff H, Sonuga-Barke E, Schachar R, Logan GD, Wigal T, Hechtman T, Hinshaw S, Turkel E. The ecological validity of delay aversion and response inhibition as measures of impulsivity in AD/HD: A supplement to the NIMH multimodal treatment study of AD/HD. *J Abnorm Child Psychol*. 2001; 29:215–228. [PubMed: 11411784]

83. Goghari VM, MacDonald AW. The neural basis of inhibition in cognitive control: response selection and inhibition. *Brain Cogn.* 2009; 71:72–83. [PubMed: 19427089]
84. Logan GD, Schachar RJ, Tannock R. Impulsivity and inhibitory control. *Psychol Sci.* 1997; 8:60–64.
85. Nederkoorn C, Smulders FTY, Havermans RC, Roefs A, Jansen A. Impulsivity in obese women. *Appetite.* 2006; 47:253–256. [PubMed: 16782231]
86. Nederkoorn C, Braet C, Van Eijs Y, Tanghe A, Jansen A. Why obese children cannot resist food: The role of impulsivity. *Eat Behav.* 2006; 7:315–322. [PubMed: 17056407]
87. Guerrieri R, Nederkoorn C, Jansen A. How impulsiveness and variety influence food intake in a sample of healthy women. *Appetite.* 1997; 49:119–122.
88. Guerrieri R, Nederkoorn C, Jansen A. The interaction between impulsivity and a varied food environment: its influence on food intake and overweight. *Int J Obes.* 2008; 32:708–714.
89. Nederkoorn C, Jansen E, Mulkens S, Jansen A. Impulsivity predicts treatment outcome in obese children. *Behav Res Ther.* 2007; 45:1071–1075. [PubMed: 16828053]
90. Guerrieri R, Nederkoorn C, Schrooten M, Martijn C, Jansen A. Inducing impulsivity leads high and low restrained eaters into overeating, whereas current dieters stick to their diet. *Appetite.* 2009; 53:93–100. [PubMed: 19467278]
91. Jansen A, Nederkoorn C, Van Baak L, Keirse C, Guerrieri R, Havermans R. High-restrained eaters only overeat when they are also impulsive. *Behav Res Ther.* 2009; 47:105–110. [PubMed: 19038379]
92. Barkley RA. Behavioral inhibition, sustained attention, and executive functions: constructing a unifying theory of ADHD. *Psychol Bull.* 1997; 121:65–94. [PubMed: 9000892]
93. David C. Attention-deficit/hyperactivity disorder: associations with overeating and obesity. *Curr Psychiatry Rep.* 2010; 12:389–395. [PubMed: 20632134]
94. Agranat-Meged AN, Deitcher C, Goldzweig G, Leibenson L, Stein M, Galili-Weisstub E. Childhood obesity and attention deficit/hyperactivity disorder: a newly described comorbidity in obese hospitalized children. *Int J Eat Disord.* 2005; 37:357–359. [PubMed: 15856493]
95. Cortese S, Bernardina BD, Mouren MC. Attention deficit/hyperactivity disorder (ADHD) and binge eating. *Nutr Rev.* 2007; 65:404–411. [PubMed: 17958207]
96. Mischel W, Ebbeson EB, Zeiss AR. Cognitive and attentional mechanisms in delay of gratification. *J Pers Soc Psychol.* 1972; 21:204–218. [PubMed: 5010404]
97. Mischel W, Shoda Y, Rodriguez ML. Delay of Gratification in children. *Science.* 1989; 244:933–938. [PubMed: 2658056]
98. Mischel W, Shoda Y, Peake PK. The nature of adolescent competencies predicted by preschool delay of gratification. *J Pers Soc Psychol.* 1988; 54:687–696. [PubMed: 3367285]
99. Seeyave DM, Coleman S, Appugliese D, Corwyn RF, Bradley RH, Davidson NS, Kaciroti N, Lumeng JC. Ability to delay gratification at age 4 years and risk of overweight at age 11 years. *Arch Pediatr Adolesc Med.* 2009; 163:303–308. [PubMed: 19349558]
100. Pignatti R, Bertella L, Albani G, Mauro A, Molinari E, Semenza C. Decision-making in obesity: a study using the gambling task. *Eat Weight Discord.* 2006; 11:126–132.
101. Appelhans BM. Neurobehavioral inhibition of reward-driven feeding: implications for dieting and obesity. *Obesity (Silver Springs).* 2009; 17:640–647.
102. Davis C, Patte K, Levitan R, Reid C, Tweed S, Curtis C. From motivation to behavior: a model for reward sensitivity, overeating, and food preferences in the risk profile for obesity. *Appetite.* 2007; 48:12–19. [PubMed: 16875757]
103. McClure SM, Laibson DI, Loewenstein G, Cohen JD. Separate neural systems value immediate and delayed monetary rewards. *Science.* 2004; 306:503–507. [PubMed: 15486304]
104. Figner B, Knoch D, Johnson EJ, Krosch AR, Lisanby SH, Fehr E, Weber EU. Lateral prefrontal cortex and self control in intertemporal choice. *Nat Neurosci.* 2010; 13:538–539. [PubMed: 20348919]
105. Small DM, Zatorre RJ, Dagher A, Evans AC, Jones-Gotman M. Changes in brain activity related to eating chocolate: from pleasure to aversion. *Brain.* 2001; 124:1720–1733. [PubMed: 11522575]

106. Rothemund Y, Preuschhof C, Böhner G, Bauknecht HC, Klingebiel R, Flor H, Klapp BF. Differential activation of the dorsal striatum by high calorie visual food stimuli in obese individuals. *Neuroimage*. 2007; 37:410–421. [PubMed: 17566768]
107. Geliebter A, Ladell T, Logan M, Schneider T, Sharafi M, Hirsch J. Responsivity to food stimuli in obese and lean binge eaters using functional MRI. *Appetite*. 2006; 46:31–35. [PubMed: 16364498]
108. Uher R, Yoganathan D, Mogg A, Eranti SV, Treasure J, Campbell IC, McLoughlin DM, Schmidt U. Effect of left prefrontal repetitive transcranial magnetic stimulation on food craving. *Biol Psychiatry*. 2005; 58:840–842. [PubMed: 16084855]
109. Fregni F, Orsati F, Pedrosa W, Fecteau S, Tome FA, Nitsche MA, Mecca T, Macedo EC, Pascual-Leone A, Boggio PS. Transcranial direct current stimulation of the prefrontal cortex modulates the desire for specific foods. *Appetite*. 2008; 51:34–41. [PubMed: 18243412]
110. Goldman RL, Borckardt JJ, Frohman HA, O'Neil PM, Madan A, Campbell LK, Budak A, George MS. Prefrontal cortex transcranial direct current stimulation (tDCS) temporarily reduces food cravings and increases the self-reported ability to resist food in adults with frequent food craving. *Appetite*. 2011; 56:741–746. [PubMed: 21352881]
111. Linde JA, Rothman AJ, Baldwin AS, Jeffery RW. The impact of self-efficacy on behavior change and weight change among overweight participants in a weight loss trial. *Health Psychol*. 2006; 25:282–291. [PubMed: 16719599]
112. Teixeira PJ, Going SB, Houtkooper LB, Cussler EC, Metcalfe LL, Blew RM, Sardinha LB, Lohman TG. Exercise motivation, eating, and body image variables as predictors of weight control. *Med Sci Sports Exerc*. 2006; 38:179–188. [PubMed: 16394972]
113. Bryant EJ, King NA, Blundell JE. Disinhibition: its effect on appetite and weight regulation. *Obes Rev*. 2008; 9:409–419. [PubMed: 18179615]
114. Klem ML, Wing RR, McGuire MT, Seagle HM, Hill OJ. A descriptive study of individuals successful at long-term weight maintenance of substantial weight loss. *Am J Clin Nutr*. 1997; 66:239–246. [PubMed: 9250100]
115. Neimeier HM, Phelan S, Fava JL, Wing RR. Internal disinhibition predicts weight regain following weight loss and weight loss maintenance. *Obesity*. 2007; 15:2485–2494. [PubMed: 17925475]
116. Bond DS, Phelan S, Leahey TM, Hill JO, Wing RR. Weight loss maintenance in successful weight losers: surgical vs. non-surgical methods. *Int J Obes*. 2009; 33:173–180.
117. McCaffery JM, Haley AP, Sweet LH, Phelan S, Raynor HA, Del Parigi A, Cohen R, Wing RR. Differential frontal magnetic resonance imaging response to food pictures in successful weight loss maintainers relative to normal weight and obese controls. *Am J Clin Nutr*. 2009; 90:928–934. [PubMed: 19675107]
118. DelParigi A, Chen K, Salbe AD, Hill JO, Wing RR, Reiman EM, Tataranni PA. Successful dieters have increased neural activity in cortical areas involved in the control of behavior. *Int J Obes*. 2007; 31:440–448.
119. Le DS, Pannacciulli N, Chen K, Del Parigi A, Salbe AD, Reiman EM, Krakoff J. Less activation of the left dorsolateral prefrontal cortex in response to a meal: a feature of obesity. *Am J Clin Nutr*. 2006; 84:725–731. [PubMed: 17023697]
120. Le DS, Pannacciulli N, Chen K, Salbe AD, Del Parigi A, Hill JO, Wing RR, Reiman EM, Krakoff J. Less activation in the left dorsolateral prefrontal cortex in the reanalysis of the response to a meal in obese than in lean women and its association with successful weight loss. *Am J Clin Nutr*. 2007; 86:573–579. [PubMed: 17823419]
121. Kramer AF, Erickson KI. Capitalizing on cortical plasticity: Influence of physical activity on cognition and brain function. *Trends Cogn Sci*. 2007; 11:342–348. [PubMed: 17629545]
122. Colcombe SJ, Kramer AF. Fitness effects on the cognitive function of older adults: a meta-analytic study. *Psychol Sci*. 2003; 14:125–130. [PubMed: 12661673]
123. Kramer AF, Hahn S, Cohen NJ, Banich MT, McAuley E, Harrison CR, Chason J, Vakil E, Bardell L, Boileau RA, Colcombe A. Ageing, fitness, and neurocognitive function. *Nature*. 1999; 400:418–419. [PubMed: 10440369]

124. Colcombe SJ, Kramer AF, Erickson KI, Scalf P, McAuley E, Cohen NJ, Webb A, Jerome GJ, Marquez DX, Elavsky S. Cardiovascular fitness, cortical plasticity and aging. *Proc Nat Acad Sci USA*. 2004; 101:3316–3321. [PubMed: 14978288]
125. Voss MW, Prakash RS, Erickson KI, Basak C, Chaddock L, Kim JS, Alves H, Heo S, Szabo AN, White SM, Wójcicki TR, Mailey EL, Gothe N, Olson EA, McAuley E, Kramer AF. Plasticity of brain networks in a randomized intervention trial of exercise training in older adults. *Front Aging Neurosci*. 2010; 26:2, pii: 32.
126. Colcombe SJ, Erickson KI, Raz N, Webb AG, Cohen NJ, McAuley E, Kramer AF. Aerobic fitness reduced brain tissue loss in aging humans. *J Gerontol A Biol Sci Med Sci*. 2003; 58:176–180.
127. Colcombe SJ, Erickson KI, Scalf PE, Kim JS, Prakash R, McAuley E, Elavsky S, Marquez DX, Hu L, Kramer AF. Aerobic exercise training increases brain volume in aging humans. *J Gerontol A Biol Sci Med Sci*. 2006; 61:1166–1170. [PubMed: 17167157]
128. Lipnicki DM, Gunga HC, Belavy DL, Felsenberg D. Bed rest and cognition: effects on executive functioning and reaction time. *Aviat Space Environ Med*. 2009; 80:1018–1024.
129. Bryant EJ, Keizebrink K, King NA, Blundell JE. Interaction between disinhibition and restraint: Implications for body weight and eating disturbance. *Eat Weight Discord*. 2010; 15:e43–e45.
130. Bryant E, King N, Blundell J. Effect of exercise on appetite control in women with high trait disinhibition. *Appetite*. 2005; 45:363.
131. Bryant E, Finlayson G, King N, Blundell J. The influence of acute exercise on linking and preferences for food on high trait disinhibition women. *Obes Rev*. 2006; 17:343.
132. Diamond A. Close interrelation of motor development and cognitive development and of the cerebellum and prefrontal cortex. *Child Dev*. 2000; 71:44–56. [PubMed: 10836557]
133. Temple JL, Giacomelli AM, Kent KM, Roemmich KN, Epstein LH. Television watching increases motivated responding for food and energy intake in children. *Am J Clin Nutr*. 2007; 85:355–361. [PubMed: 17284729]
134. Stunkard AJ, Faith MS, Allison KC. Depression and obesity. *Biol Psychiatry*. 2003; 54:330–337. [PubMed: 12893108]
135. Markowitz S, Friedman MA, Arent SM. Understanding the relation between obesity and depression: causal mechanisms and implications for treatment. *Clin Psychol Sci Pract*. 2008; 15:1–20.
136. Rooke SE, Thorsteinsson EB. Examining the temporal relationship between depression and obesity: a meta-analyses of prospective research. *Health Psych Rev*. 2008; 2:94–109.
137. Ouwens MA, van Strien T, van Leeuwe JF. Possible pathways between depression, emotional and external eating. A structural equation model. *Appetite*. 2009; 53:245–248. [PubMed: 19505515]
138. Kontinen H, Mäanistö S, Sarlio-Lähteenkorva S, Silventoinen K, Haukkala A. Emotional eating, depressive symptoms and self-reported food consumption. A population based study. *Appetite*. 2010; 54:473–479. [PubMed: 20138944]
139. Keskitalo K, Tuorila H, Spector TD, Cherkas LF, Knaapila A, Kaprio J, Silventoinen K, Perola M. The Three-Factor Eating Questionnaire, body mass index, and responses to sweet and salty fatty foods: a twin study of genetic and environmental associations. *Am J Clin Nutr*. 2008; 88:263–267. [PubMed: 18689360]
140. Craft LL, Perna FA, Freund KM, Culpepper L. Psychosocial correlates of exercise in women with self-reported depressive symptoms. *J Phys Act Health*. 2008; 5:469–480. [PubMed: 18579923]
141. Kontinen H, Silventoinen K, Sarlio-Lähteenkorva S, Mäanistö S, Haukkala A. Emotional eating and physical activity self-efficacy as pathways in the association between depressive symptoms and adiposity indicators. *Am J Clin Nutr*. 2010; 92:1031–1039. [PubMed: 20861176]
142. Ryan R, Deci E. Self-determination theory and the facilitation of intrinsic motivation, social development, and well-being. *Am Psychol*. 2000; 55:68–78. [PubMed: 11392867]
143. Mata J, Silva MN, Vieira PN, Carraca EV, Andrade AM, Coutinho SR, Sardinha LB, Teixeira PJ. Motivational “spill-over” during weight control: increased self-determination and exercise intrinsic motivation predict eating self-regulation. *Health Psychol*. 2009; 28:709–716. [PubMed: 19916639]

144. Donnelly JE, Blair SN, Jakicic JM, Manore MM, Rankin JW, Smith BK. Appropriate physical activity intervention strategies for weight loss and prevention of weight regain for adults. *Med Sci Sports Exer.* 2009; 41:459–471.
145. Andrade AM, Coutinho SR, Silva MN, Matta J, Vieira PN, Minderico CS, Melanson KJ, Baptista FB, Sardinha LB, Teixeira PJ. The effect of physical activity on weight loss is mediated by eating selfregulation. *Patient Educ Couns.* 2010; 79:320–326. [PubMed: 20149955]
146. Macht M. How emotions affect eating: a five-way model. *Appetite.* 2008; 50:1–11. [PubMed: 17707947]
147. Melanson KJ, Dell'Olio J, Carpenter MR, Angelopoulos TJ. Changes in multiple health outcomes at 12 and 24 weeks resulting from 12 weeks of exercise counseling with or without dietary counseling in obese adults. *Nutrition.* 2004; 20:849–856. [PubMed: 15474871]
148. Salmon P. Effects of physical exercise on anxiety, depression, and sensitivity to stress: a unifying theory. *Clin Psychol Rev.* 2001; 21:33–61. [PubMed: 11148895]
149. Blumenthal JA, Babyak MA, Doraiswamy PM, Watkins L, Hoffman BM, Barbour KA, Herman S, Craighead WE, Brosse AL, Waugh R, Hinderliter A, Sherwood A. Exercise and pharmacotherapy in the treatment of major depressive disorder. *Psychosom Med.* 2007; 69:587–596. [PubMed: 17846259]
150. George MS, Ketter TA, Post RM. Prefrontal cortex dysfunction in clinical depression. *Depression.* 1994; 2:59–72.
151. Tucker M, Reicks M. Exercise as a gateway behavior for healthful eating among older adults: an exploratory study. *J Nutr Edu Behav.* 2002; 34(Suppl 1):S14–S19.
152. Costakis CE, Dunnagan T, Haynes G. The relationship between the stages of exercise adoption and other health behaviors. *Am J Health Promot.* 1999; 14:22–30. [PubMed: 10621521]
153. Ryan RM, Frederick CM, Lepes D, Rubio N, Sheldon KM. Intrinsic motivation and exercise adherence. *Int J Sport Psychol.* 1997; 28:335–354.
154. Boulé NG, Weisnagel SJ, Lakka TA, Tremblay A, Bergman RN, Rankinen T, Leon AS, Skinner JS, Wilmore JH, Rao DC, Bouchard C. HERITAGE Family Study. Effects of exercise training on glucose homeostasis: the HERITAGE Family Study. *Diabetes Care.* 2005; 28:108–114. [PubMed: 15616242]
155. Tremblay A, Therrien F. Physical activity and body functionality: implications for obesity prevention and treatment. *Can J Physiol Pharmacol.* 2006; 84:149–156. [PubMed: 16900940]
156. Chaput JP, Tremblay A. The glucostatic theory of appetite control and the risk of obesity and diabetes. *Int J Obes (Lond).* 2009; 33:46–53. [PubMed: 19002144]
157. Trost SG, van der Mars H. Why we should not cut P.E. *Health and Learning.* 2010; 67:60–65.

Figure 1.

A diagrammatic representation of the hypothesis linking physical activity to eating behavior via executive function. In the proposed model, physical activity can enhance executive function by generating, strengthening, and refining neural circuits in the prefrontal cortex. These executive functions are requisite for inhibitory control, the ability to suppress impulsive behavior and responses to external stimuli. Due to the abundance of negative food stimuli in the environment that prompt unhealthy dietary habits, practicing healthy dietary choices demands a high degree of inhibitory control. Inhibitory control acts as the brake, thereby enabling the execution of goalsetting behavior and self-control. Thus, physical activity indirectly modifies eating behavior by strengthening executive function. Over time, physical activity and healthy dietary habits become behavioral adaptations connected by their goal-oriented nature.