Published in final edited form as:

Mol Microbiol. 2013 January ; 87(1): 211–229. doi:10.1111/mmi.12103.

The cell biology of peritrichous flagella in *Bacillus subtilis*

Sarah B. Guttenplan, Sidney Shaw, and Daniel B. Kearns*

Indiana University, Department of Biology, Bloomington, IN 47408

SUMMARY

Bacterial flagella are highly conserved molecular machines that have been extensively studied for assembly, function, and gene regulation. Less studied is how and why bacteria differ based on the number and arrangement of the flagella they synthesize. Here we explore the cell biology of peritrichous flagella in the model bacterium *Bacillus subtilis* by fluorescently labeling flagellar basal bodies, hooks, and filaments. We find that the average *B. subtilis* cell assembles approximately 26 flagellar basal bodies and we show that basal body number is controlled by the protein of unknown function SwrA. Basal bodies are assembled rapidly (< 5 minutes) but the assembly of flagella capable of supporting motility (> 40 minutes) is rate limited by filament polymerization. We find that basal bodies are not positioned randomly on the cell surface. Rather, basal bodies occupy a grid-like pattern organized symmetrically around the midcell and that flagella are discouraged at the poles. Basal body position is genetically determined by FlhF and FlhG homologs to control spatial patterning differently from what is seen in bacteria with polar flagella. Finally, spatial control of flagella in *B. subtilis* seems more relevant to the inheritance of flagella in individual cells than the motile behavior of populations.

Keywords

Motility; flagella; SwrA; FlhF; FlhG

INTRODUCTION

Many species of bacteria assemble rotary flagella to swim in liquid and swarm over solid surfaces (Fig 1A). Each flagellum is a machine constructed from over 30 different proteins, regulated to ensure proper stoichiometry and timing, and integrated into the fundamental architecture of the cell envelope (Kalir *et al.*, 2001; Macnab, 2003; Brown *et al.*, 2008; Chevance and Hughes, 2008). First to be assembled is the flagellar basal body that anchors the flagellum to the cell membrane, powers rotation, and houses the type III secretion apparatus that exports the more distal components like the hook, a flexible universal joint, and finally the long helical filament (Fig 1B). Complex regulatory feedback mechanisms coordinate flagellar assembly. Some systems globally regulate all flagella in the cell by controlling common pools of structural components like flagellin used to build the filament (Hughes *et al.*, 1993; Kutsukake, 1994; Mukherjee *et al.*, 2011). By contrast, some systems locally regulate individual flagella such as the mechanism that controls secretion apparatus specificity in response to the length of the associated hook (Journet *et al.*, 2003; Shibata *et al.*, 2007; Erhardt *et al.*, 2011). The function of global and local regulators, however, likely depends on the relative number of flagella being controlled.

Flagellar number varies dramatically according to the bacterial species (Fig 1A). One flagellum is sufficient to swim in liquid media and the reason why some bacteria make more

*Corresponding author, dbkearns@indiana.edu, Phone: 1-812-856-2523, Fax: 1-812-855-6705.

than one flagellum isn't precisely known. Past studies have indicated that cells with a single polar flagellum perform better in environments with low viscosity but worse in environments with high viscosity when compared to bacteria with multiple flagella (Atsumi *et al.*, 1996). Motility in viscous environments has been correlated with the ability to swarm atop solid surfaces and an increase in flagellar number has been reported for a variety of bacteria during swarming (Atsumi *et al.*, 1996; for review see Kearns, 2010). In an extreme case, some bacteria make one polar flagellum while swimming but induce multiple flagella, expressed from an entirely separate set of genes, to swarm (for review see McCarter, 2004). The mechanisms that control flagellar number likely operate at the level of expression of basal body genes but when, how, and why flagellar number changes within a species is poorly understood (for review see Patrick and Kearns, 2012).

Flagellar research has focused primarily on flagellar structure and how regulation ensures the sequential addition of the subunits. Flagellar assembly, however, takes place in the context of the whole cell and when a flagellum is built, it must be inserted through, and immobilized in, the cell wall. Although the proteins that assemble flagella are highly conserved, bacterial species differ in the placement of flagella on their cell surface (Fig 1A) (Boltjes, 1948; Leifson, 1951; Pallen *et al.*, 2005; Liu and Ochman, 2007). Some bacteria restrict flagella to the cell poles (called a polar or lophotrichous arrangement) whereas other bacteria insert flagella along the length of the cell (called a medial or peritrichous arrangement). The pattern of flagellar arrangement is sufficiently specific and constant that it was used as a defining characteristic of a bacterial species in numerical taxonomy (Stanier and Van Niel, 1941; Boltjes, 1948). Thus, flagella were recognized as having specific cellular addresses long before the first evidence of subcellular localization of individual proteins in bacteria (Bi and Lutkenhaus, 1991; Alley *et al.*, 1992; Maddock and Shapiro, 1993; Ma *et al.*, 1996). The biological relevance of any particular flagellar arrangement is poorly understood.

How flagellar position is controlled is only beginning to be understood and work has primarily been conducted in bacteria with the most conspicuous bias of polar flagella. Two proteins have been found to be important to the positioning of polar flagella, FlhF and FlhG. FlhF is a signal-recognition particle-like GTPase, which when mutated mislocalizes flagella from the pole to the sides of the cell (Pandza *et al.*, 2000; Correa *et al.*, 2005; Murray and Kazmierczak, 2006). FlhF binds GTP in vitro and GTP binding, but perhaps not hydrolysis, is required to localize the basal body protein FliF in the membrane (Correa *et al.*, 2005; Bange *et al.*, 2007; Green *et al.*, 2009; Kusumoto *et al.*, 2009; Balaban and Hendrixson, 2011). FlhG is a MinD-like ATPase which when mutated often results in an increase in flagellar number (Dasgupta *et al.*, 2000; Correa *et al.*, 2005; Balaban and Hendrixson, 2011). Biochemically, FlhG binds to FlhF and enhances FlhF GTP hydrolysis (Kusumoto *et al.*, 2008; Bange *et al.*, 2011). Precisely how the two proteins dictate polar basal body positioning and number is unknown but both FlhF and FlhG colocalize with polar flagella (Murray and Kazmierczak, 2006; Kusumoto *et al.*, 2008; Green *et al.*, 2009; Kusumoto *et al.*, 2009; Balaban and Hendrixson, 2011).

Here we ask and answer fundamental questions about flagellar placement in a bacterium with a peritrichous flagellar arrangement, *Bacillus subtilis*. How many flagella does *B. subtilis* make? How long does it take to synthesize a flagellum? Where are the flagella located on the cell surface? In *B. subtilis* cells, the genes that synthesize flagellar basal body proteins are encoded within the large 31 gene, *fla/che* operon activated by the master regulator SwrA (Fig 2A; Albertini *et al.*, 1991; Márquez-Magaña and Chamberlin, 1994; Kearns and Losick, 2005). Here we show that SwrA increases basal body number and we show that homologs of FlhF and FlhG encoded within the *fla/che* operon cooperate to control the distance between basal bodies on the cell surface. In sum, we find that flagellar

placement is non-random and genetically determined. Finally, our work suggests that basal body patterning is important for fidelity of flagella segregation during binary fission and thus the motile behavior of individual cells.

RESULTS

FliM-GFP puncta indicate the location of flagellar basal bodies

B. subtilis encodes many proteins homologous to those required for flagellar biosynthesis in *E. coli* and *S. enterica*, but the structure and positioning of Gram positive flagella is poorly understood (Aizawa et al., 2002). To localize flagella basal bodies, we constructed a translational fusion between FliM and GFP expressed under the control of the native promoter of the *fla/che* operon, *P_{fla/che}* (Fig 1B, Fig 2A, *amyE::P_{fla/che}-fliM-GFP*). FliM is part of the basal body C-ring that controls the direction of flagellum rotation and has been used to localize basal bodies in other model systems (Toker and Macnab, 1997; Sourjik and Berg, 2000; Macnab, 2003; Aldridge et al., 2006). The FliM-GFP fusion was partly functional for complementing a Δ *fliM* deletion mutation in the chromosome. The swarming motility of a Δ *fliM* mutant complemented with FliM-GFP was slightly worse than a strain complemented with FliM alone (Fig 2B), but the swimming motility was slightly better when complemented with FliM-GFP (Fig 2C). Although the reason for the differential efficacy in complementing different motile behaviors is unknown, Western blot analysis indicated that the FliM-GFP fusion was relatively stable and primarily resolved as a 65 kDa protein, the expected size of the translational fusion (Fig S1A). We conclude that the *B. subtilis* FliM-GFP translational fusion is partly functional and largely intact *in vivo*.

FliM-GFP localized as puncta when introduced as an extra copy to the wild type but the puncta were obscured by a high background of cytoplasmic GFP presumably due either to competition with the unlabeled native FliM or due to limited proteolysis of GFP from the FliM fusion protein (Fig. 3A; Fig S1A). FliM-GFP puncta were more intense and the background fluorescence was diminished when the native copy of *fliM* was deleted from the chromosome (Fig 3B). We infer that competition with the native FliM protein was significant and for the clearest labeling, all subsequent FliM-GFP experiments were conducted in a Δ *fliM* background. Next we determined whether the FliM-GFP punctate localization pattern was dependent upon other flagella proteins. The FliM-GFP translational fusion showed diffuse mislocalization to the cytoplasm in cells lacking FliF, the basal body membrane anchor required for flagellar assembly (Fig 1B; Fig 3C; Ueno et al., 1992). In contrast, punctate localization was not altered in cells mutated for MotB, part of the proton channel that powers flagellar rotation (Fig 1B; Fig 3D), or in cells mutated for σ^D , the alternative sigma factor that expresses flagellar biosynthesis genes after the basal body and hook are completed (Fig 3E) (Zhou et al., 1998; Serizawa et al., 2004; Courtney et al., 2012). We conclude that FliM-GFP forms puncta that require the flagellar basal body but do not require flagellar rotation or flagellar completion.

To further characterize the FliM-fluorescent protein fusion, we determined whether it co-localized with other flagella-associated proteins: the flagellar hook and the flagellar clutch. The hook is the external, flexible joint that connects the filament to the basal body-rod and is polymerized from the protein FlgE (Fig 1B; Komeda et al., 1978; Samatey et al., 2004; Courtney et al., 2012). The clutch, EpsE, is normally made during biofilm formation and inhibits flagellar rotation by separating the cytoplasmic FliG rotor from the MotA/MotB stator (Fig 1B; Blair et al., 2008; Guttenplan et al., 2010). Cells co-expressing an allele of FlgE that could be fluorescently labeled with a cysteine-reactive maleimide stain (FlgE^{T123C}) and FliM-GFP showed puncta of hooks and puncta of FliM that co-localized (Fig 4A; Fig S2) (Courtney et al., 2012). Similarly, cells co-expressing EpsE-YFP and FliM-CFP showed puncta of clutch and puncta of FliM that co-localized (Fig 4B). Thus, FliM

puncta not only required genes involved in basal body synthesis but also co-localized with intracellular and extracellular markers associated with the flagellum. We conclude that puncta of FliM indicate the location of flagellar basal bodies.

The timing of flagellar assembly

Over 30 structural proteins contribute to flagellar biosynthesis, and the flagellum is assembled in stages: the basal body is completed first, followed by the hook, and finally the filament (Fig 1B) (Macnab, 2003). We wanted to determine how long it took to complete each stage of flagellum assembly using the FliM-GFP fusion to fluorescently label basal bodies and previously established constructs to fluorescently label the hook and the filament (Blair *et al.*, 2008; Courtney *et al.*, 2012). To measure the timing of flagellar biosynthesis we first generated a strain in which the expression of flagella genes in the *fla/che* operon was under artificial control by integrating an IPTG-inducible $P_{hysp\alpha nk}$ promoter between the native $P_{fla/che}$ promoter and the *fla/che* operon ($P_{fla/che} \Omega P_{hysp\alpha nk} - fla/che$). In the absence of IPTG, the cells were non-motile and grew in long chains due to the failure to express the *fla/che*-operon-encoded alternative sigma factor σ^D which in turn directs the expression of the vegetative cell separating enzymes LytC, LytD, and LytF (Blackman *et al.*, 1998; Margot *et al.*, 1999; Chen *et al.*, 2009). Motility and cell separation was incrementally restored as the amount of IPTG was increased in the medium (Fig S3). The addition of 0.3 mM IPTG was sufficient to restore both swarming motility and the expression of the basal body protein FliG (encoded by *fliG* within the *fla/che* operon) to near wild type levels (Fig S1B, Fig S3).

To determine how long it took to synthesize flagellar basal bodies, the FliM-GFP construct was introduced to the strain containing the IPTG-inducible *fla/che* operon ($P_{fla/che} \Omega P_{hysp\alpha nk} - fla/che$). FliM-GFP was ectopically expressed from the native $P_{fla/che}$ promoter, and FliM-GFP localized as a diffuse haze in the cytoplasm prior to the induction of the $P_{hysp\alpha nk}$ promoter controlling the rest of the *fla/che* operon (Fig 5A). Five minutes after IPTG induction, membrane associated puncta of FliM-GFP were visible amidst the background haze, and by 10 minutes puncta of FliM-GFP resembled the wild type (Fig 5A). We infer that the pre-existing pool of FliM-GFP manifests as a diffuse haze but nucleates into puncta once other basal body proteins are expressed and assembled. We conclude that basal bodies assemble rapidly in less than five minutes post initiation (Fig 5B).

To determine how long it took to assemble flagellar hooks, the FlgE^{T123C} allele of the hook protein that can be fluorescently labeled with a cysteine-reactive maleimide stain was introduced to the strain containing the IPTG-inducible *fla/che* operon (Courtney *et al.*, 2012). Puncta of fluorescently labeled hooks were visible on the cell membrane after 10 minutes and resembled the wild type after 20 minutes of IPTG induction (Fig 5C). Hooks must be assembled prior to the filament and we conclude that it takes 10–20 minutes post initiation to assemble flagellar hooks (Fig 5B). To determine how long it took to synthesize flagellar filaments, the Hag^{T209C} allele of flagellin that can be fluorescently labeled with a cysteine-reactive maleimide stain was introduced to the strain containing the IPTG-inducible *fla/che* operon (Blair *et al.*, 2008). Short fluorescent stubs were visible as early as 15 minutes after IPTG induction and after 60 minutes cells had long filaments on the cell surface that resembled the wild type (Fig 5D). Cells were found to be motile between 40–60 minutes after IPTG induction when observed microscopically. We conclude that flagellar filament synthesis is initiated within 15 minutes after induction and completed, with respect to permitting motility, after approximately another 25–45 minutes of assembly (Fig 5B). We infer that a threshold filament length is required to generate propulsion. It has been estimated that the filament is comprised of ~20,000 flagellin subunits, and if the filament takes ~30 minutes to assemble, we estimate that the *B. subtilis* basal body exports approximately ~11 subunits per second, a rate comparable to that estimated for *Salmonella* (Iino, 1969; Macnab, 1992; Chevance and Hughes, 2008).

Basal body number is determined by SwrA

The number of flagella resident on a *B. subtilis* cell is unknown. We attempted to directly count flagellar filaments by fluorescently labeling the flagellin protein (Hag^{T209C}) and observing with a super-resolution OMX 3D structured illumination microscope (SIM). Accurate filament counts, however, proved difficult because filaments were too numerous, had the tendency to form bundles, and appeared to shear from some cells (Fig 6A). Counting flagella by EM requires a large number of high quality images and is hampered not only by the same problems as fluorescent microscopy mentioned above but also the inability to distinguish which flagella originate from which cell when cells are conjoined (Aizawa *et al.*, 2002). To count flagella indirectly, we counted basal body number by super-resolution images of FliM-GFP (Fig 6B). Wild type cells had an average of 26 ± 6 basal bodies ($N = 42$ cells, std given for all measures) and the number of basal bodies generally increased with increasing cell length (Fig 6B, C). If one makes the likely imprecise but practical approximation that each basal body produces a flagellum, *B. subtilis* synthesizes 3–8 times more flagella per cell than *E. coli* (Iino, 1969; Sourjik and Berg, 2000; Wozniak *et al.*, 2010; Erhardt and Hughes, 2010).

Flagellar number in *B. subtilis* is thought to be controlled by SwrA, a protein of unknown molecular function that enhances the expression of the *fla/che* operon (Kearns and Losick, 2005; Calvio *et al.*, 2005; Cozy and Kearns, 2010; Patrick and Kearns, 2012). When SwrA was either mutated ($\Delta swrA$) or overexpressed from an artificial *P_{hyspank}* promoter ($swrA^{++}$) in a strain encoding the Hag^{T209C} filament protein, the number of flagella per cell appeared to decrease and increase, respectively, relative to wild type (Fig 6A; Kearns and Losick, 2005). To determine whether the qualitative changes in flagellar filament number corresponded with the number of basal bodies, SwrA was either mutated or overexpressed in a background expressing FliM-GFP (Fig 6B). As reported previously, strains lacking SwrA showed an increase in the number of chaining cells in the population (Kearns and Losick, 2005). Cells in chains were excluded from further analysis because they lacked both flagellar filaments and basal bodies (FliM-GFP puncta) (Fig. 3F; Kearns and Losick, 2005). Single cells of the *swrA* mutant synthesized flagellar filaments and an average of 9 ± 2 basal bodies per cell ($N = 43$ cells) but unlike the wild type, the number of basal bodies did not appear to increase proportionally with cell length (Fig 6C). Conversely, cells overexpressing SwrA ($swrA^{++}$) synthesized an average of 41 ± 6 basal bodies ($N = 42$ cells) and the basal body number appeared to be more tightly correlated with cell length than the wild type (Fig 3G, 6C). We note that the changes in basal body number were commensurate with the changes in *P_{fla/che}* expression observed in backgrounds either mutated for or overexpressing *swrA* (Fig S4; Kearns and Losick, 2005). We conclude that SwrA controls basal body number over a 4-fold range.

Basal body distribution is not random

Reviewing 3-D rendered data sets of FliM-GFP expressing cells suggested that basal bodies might be arranged in a non-random pattern that was perhaps helical (Supplemental Movies S1–4). To analyze basal body positions, the outer surface of the cell was labeled with the membrane staining dye, FM4-64, and each basal body position in the super-resolved SIM data was assigned an X-Y-Z coordinate using a center of mass calculation with Imaris software. The nearest neighbor distances for all basal bodies were determined using scripts developed in MATLAB and the minimum separation was found to be, on average, $0.39 \pm 0.14 \mu\text{m}$ ($N = 42$ cells) in the wild type (Fig 7A). When SwrA was mutated, the mean nearest neighbor distances increased nearly two-fold, likely because cells lacking SwrA synthesized half the number of basal bodies per cell as compared to the wild type (Fig 6C, Fig 7A). The two-fold over-production of basal bodies in cells overexpressing SwrA ($swrA^{++}$), however, reduced the mean nearest neighbor distance by only ~20% (Fig 7A). To

evaluate how the distribution of nearest neighbor distances in each case might vary from a uniform random distribution, the basal bodies in each cell were mathematically reassigned to random surface positions within the confines of each cell's dimensions, and the nearest neighbor positions were re-measured for all cells (Fig. 7A grey bars). For each cell type, the random distribution was skewed toward a shorter average nearest neighbor distance (Fig 7A). We conclude that there is a minimum distance between basal bodies that is greater than expected for a random distribution.

To quantitatively assess how basal bodies were dispersed across the cell surface, independent of the number of basal bodies per cell, we calculated a Clark-Evans ratio (Clark and Evans, 1954). The Clark-Evans ratio sets an expectation that a random 2-dimensional distribution would produce a mean nearest neighbor distance that is half the square root of the area density. Therefore, a value close to 1 indicates a random dispersion while values < 1 indicate clustering and values between 1 and 2 suggest a more uniform gridded distribution (Fig 7B). The nearest neighbor distances were taken for each cell and the calculated Clark-Evans ratios ($N > 40$ cells) were plotted as a histogram and compared to a randomized data set. The distribution of Clark-Evans ratios from 1000 randomly generated cells produced the expected mean of ~ 1 for each cell type (Fig 7B grey lines). For all cell types, the actual mean Clark-Evans ratios were significantly greater than 1, when compared to the randomized control data ($p < 0.01$), indicating a non-random distribution of basal bodies tending toward a higher degree of dispersion (Fig 7B). We found that cells either lacking SwrA or overexpressing SwrA did not significantly reduce the Clark-Evans ratio relative to wild type. We conclude that the distribution of basal bodies on the surface of *B. subtilis* is more dispersed and grid-like than would be expected for a random distribution. We further conclude that although SwrA controls the basal body density, SwrA does not control basal body distribution.

To further characterize the general pattern of basal body placement on the cell surface, we examined the distribution of basal bodies along the cell length. To account for differences in cell length and the arbitrary orientation of the rod shaped cells, we measured the distance of each basal body from the midcell and normalized the distance from values of 0 (midcell) to 1 (cell pole). The accumulated values from all cells of a specific class were grouped and presented as a histogram (Fig 7C). To determine if the distribution of basal bodies along the cell length departs from a random distribution, the basal bodies of each cell were randomly redistributed as previously described (Fig 7C grey bars). As expected, the randomized basal body positions showed a uniform distribution with a standard deviation related to the average number of basal bodies per cell. In general, the spatial distribution of basal bodies of wild type, SwrA mutant ($\Delta swrA$), and SwrA overexpressing cells ($swrA^{++}$) were relatively even along the length of the cell until the cell poles (Fig 7C). At the cell poles, there was a significant reduction in basal bodies relative to the remainder of the cell and a significant reduction relative to the randomized control set. We conclude that basal bodies are underrepresented at the cell poles in wild type cells and in cells for which basal body number has been increased or decreased by the presence or absence of SwrA respectively.

To quantitatively assess the asymmetry of basal body distribution, the number of puncta on each side of a cell's midpoint was counted and the higher number was divided by the lower number such that perfect symmetry would give a value of 1. Mean asymmetry ratios and standard deviations were calculated for each cell type and were then recalculated in parallel for a randomized data set (Table 1). The mean basal body asymmetry of the wild type was close to one and it was significantly lower (P value = 0.0005) than the asymmetry predicted for the randomized data set (Table 1). The mean basal body asymmetry of the SwrA overexpressed strain ($swrA^{++}$) was close to one, was significantly lower than the randomized data set (P value = 0.0455), and was not significantly different than the wild

type. For the SwrA mutated strain ($\Delta swrA$), statistical analysis indicated that the basal bodies appeared highly asymmetric but the value was uninformative due to the reduced number of basal bodies overall that caused smaller values to divide into larger ratios (Fig 6D). We infer that the wild type patterning of basal bodies leads to a distribution that is more symmetrical than one would expect from a random process (Table 1; Fig 6D).

The well-dispersed basal bodies often appeared to create spiral or helical patterns on the cell surface. To test whether the distribution of basal bodies could be organized on a helical principle, an incremental series of helices was produced with increasing pitch from 0.1 to 5 rotations around the bacterial cell, starting from positions 0–360° around the cell, and in both left- and right-handed orientations. The cumulative distance for 5 basal bodies nearest to each helix was used to find the best fitting helices and the algorithm was run against wild type FliM-GFP expressing cells and control cells with randomized basal body positions. Randomly generated basal body distributions conformed to the regular helices better than wild type cell data and no helix of constant pitch was found to best fit the wild type basal body positions (data not shown). We conclude from modeling analysis that the apparent helical nature of the basal body patterning either arises from our perception of the non-random spacing on the cell surface or that the grid-like pattern of basal bodies is more complicated than a regular spiral.

Basal body distribution is genetically determined

Patterns may arise either by spontaneous stochastic fluctuations or patterns may be dictated by cellular localization systems (Thiem and Sourjik, 2008; Wang et al., 2008). To identify potential regulators of basal body patterning in *B. subtilis*, we took a reverse genetic candidate gene approach. One candidate gene is *flhF* encoding FlhF a putative GTPase homolog which has been shown to regulate either basal body number and/or basal body position in bacteria with polar flagella (Pandza et al., 2000; Correa et al., 2005; Murray and Lazmierczak, 2006; Salvetti et al., 2007; Green et al., 2009). In other model organisms, mutation of *flhF* confers a motility defect but an in-frame markerless deletion of *flhF* in *B. subtilis* exhibited wild type swimming and swarming motility (Fig. 2 C, D). Furthermore, the *flhF* mutant produced an average number of flagellar basal bodies per cell with a nearest neighbor distance comparable to the wild type (Fig 6C; Fig 7A). The Clark-Evans ratio of the *flhF* mutant was closer to one, however, suggesting that basal body distribution was more random than the wild type (Fig. 7B). In addition, the basal bodies of the *flhF* mutant were less discouraged at the cell poles (Fig 7C). We conclude that FlhF in *B. subtilis* does not control the number of flagella per cell but instead has an effect on flagellar positioning.

With respect to basal body positioning, the most noticeable phenotype of the *flhF* mutant relative to the wild type was the frequent accumulation of basal bodies at one pole of the cell (Fig 8 A,B). At low resolution, FliM-GFP appeared as a polar fluorescent cap, but with 3D SIM, the cap resolved as an accumulation of individual polar puncta (Fig 6B, Fig 8B). Despite having an equivalent number of basal bodies, basal bodies of the *flhF* mutant were significantly more asymmetrical than the wild type ($p = 0.00039$) (Table 1). To further explore basal body mislocalization, a strain expressing FliM-GFP was generated in which the *lytC*, *lytD*, and *lytF* genes encoding the vegetative autolysins LytC, LytD, and LytF respectively were disrupted causing the population to grow exclusively in long chains (Chen et al., 2009; Cozy and Kearns, 2010). Whereas wild type cells showed an even distribution of basal bodies throughout the chains, the *flhF* mutant had a non-homogeneous distribution of basal bodies that clumped near cell poles (Fig S5). To confirm that disruption of *flhF* was responsible for FliM-GFP asymmetric distribution, a complementation construct was generated in which *flhF* was cloned downstream of the $P_{fla/che}$ promoter and inserted at the ectopic *thrC* site of a *flhF* mutant (*thrC::P_{fla/che}-flhF*). Complementation of *flhF* restored an

even basal body distribution pattern (Fig 8C). We conclude that FlhF contributes to the symmetrical distribution of basal bodies.

Another candidate gene for controlling the distribution of basal bodies is *flhG* (formerly *ylxH*), that encodes FlhG, a MinD-like ATPase which has been shown to inhibit basal body number in bacteria with polar flagella (Dasgupta *et al.*, 2000; Correa *et al.*, 2005; Kusumoto *et al.*, 2008; Balaban and Hendrixson, 2011). In other organisms, mutation of *flhG* confers a motility defect but an in-frame markerless deletion of *flhG* in *B. subtilis* exhibited wild type swimming and swarming motility (Fig 2 C,D). When *flhG* was mutated in the context of the FliM-GFP fusion, there appeared to be a dramatic reduction in the number of puncta, and therefore basal bodies per cell (Fig 8D). We noted, however, that the FliM-GFP puncta in the *flhG* mutant appeared to be considerably larger than in the wild type using SIM (Fig 6B). Staining for flagella filaments revealed that cells mutated for *flhG* synthesized multiple filaments that originated as tufts from constrained loci on the cell (Fig 6A). We infer that the larger FliM-GFP puncta observed in the *flhG* mutant actually represent aggregates of basal bodies with a nearest neighbor distance below the limit of resolution of 3D SIM. Thus, although the distance between flagellar basal bodies in the *flhG* mutant is below the limit of resolution for this approach, we infer that the Clark-Evans ratio would be closer to zero than one, representing a clumped distribution.

To confirm that disruption of *flhG* was responsible for the clumped FliM-GFP localization phenotype, a complementation construct was generated in which *flhG* was cloned downstream of the *P_{fla/che}* promoter and inserted at the ectopic *thrC* site of a *flhG* mutant (*thrC::P_{fla/che}-flhG*). Complementation with *flhG* alone produced cells with aggregated basal body foci and failed to restore a wild type distribution of basal bodies to the *flhG* mutant (Fig 8E). The *flhG* gene is encoded immediately downstream of *flhF* in the *fla/che* operon and the start and stop codons of the two genes overlap, perhaps suggesting translational coupling (Fig 2A). In a second attempt to complement the *flhG* phenotype, a complementation construct was generated in which both *flhF* and *flhG* were simultaneously cloned downstream of the *P_{fla/che}* promoter and inserted at the ectopic *lacA* site of a *flhG* mutant (*lacA::P_{fla/che}-flhFflhG*). Co-expressing both *flhF* and *flhG* restored a wild type distribution of basal bodies to the *flhG* mutant (Fig 8F). We conclude that the absence of FlhG causes basal body aggregation and thus FlhG somehow governs the nearest neighbor distance of flagella.

One way in which FlhG might control the nearest neighbor distance between basal bodies, is by regulating FlhF. FlhF and FlhG have been shown to directly interact and the possible translational coupling of the *flhF* and *flhG* genes in *B. subtilis* suggests a stoichiometric relationship between the two gene products (Kusumoto *et al.*, 2008; Bange *et al.*, 2011). To determine the epistatic relationship of mutations in *flhF* and *flhG*, an *flhF flhG* double mutant was generated in a strain expressing FliM-GFP. The *flhF flhG* double mutant did not show large foci of basal body aggregates like the *flhG* mutant and instead resembled the *flhF* mutant with frequent caps of fluorescence at one pole (Fig 8G). Furthermore, when *flhF* was overexpressed from an artificial *P_{hyspank}* promoter, large puncta of FliM-GFP were found reminiscent of the *flhG* mutant (Fig 8H). When both *flhF* and *flhG* were simultaneously overexpressed from a *P_{hyspank}* promoter, FliM-GFP localized as the wild type (Fig. 8I). We conclude that FlhG controls basal body patterning by antagonizing FlhF. We infer that the stoichiometric ratio between FlhF and FlhG is important for the correct spacing of basal bodies.

FlhF was required for proper basal body positioning but it was not required for motility in either the swimming or swarming assays. We therefore infer that the significance of basal body positioning cannot be detected in behavior at the level of whole populations. Instead,

perhaps, an even distribution of basal bodies may be important for the behavior of individual cells. We infer that each daughter cell must inherit functional flagella to be motile as the duration of flagellar assembly completion exceeded the doubling time. To measure the inheritance of flagella, exponentially growing cells expressing the Hag^{T209C} allele were stained for their flagella with a green fluorescent dye, and then diluted two-fold, four-fold, or eight-fold in fresh media and allowed to grow for one, two, or three generations respectively. After the set period of growth, cells were re-stained for newly synthesized flagella with a red fluorescent dye and visualized microscopically (Fig 9A). After three generations, 77% of wild type cells had both green and red flagella but only 43% of *flhF* mutant cells had flagella with both labels (Fig 9B). Whereas third generation cells of the wild type often appeared to have single well-separated green flagella, cells of the *flhF* mutant tended to show multiple green flagella in groups. We conclude that non-random patterning in a peritrichous arrangement ensures the more faithful segregation of flagella during cell division than a random distribution.

DISCUSSION

Once stained and visualized by light or electron microscopy, flagella are the most conspicuous landmark on a bacterial cell's surface. Different species of bacteria have different numbers of flagella and different patterns of flagellar localization. Both the mechanism that controls, and the relevance of, flagellar patterning is poorly understood. Here we explore flagellar patterning in a peritrichously flagellated bacterium, *B. subtilis*. The term peritrichous, often considered to mean the random insertion of flagella (Macnab, 1996), actually refers to the synthesis of multiple flagella arranged along the length of a rod-shaped cell (Leifson, 1951). We used super resolution microscopy and statistical analyses to show that the peritrichous patterning is non-random. Mutants that disrupted the pattern of basal bodies indicated that the patterning was genetically determined. We infer that the rules that govern positioning must accommodate rapid basal body assembly in the context of a pre-existing pattern of large number of flagella. Despite the fact that classical models for peritrichous arrangement like *E. coli* and *S. enterica* have been better studied for motility, *B. subtilis* was especially amenable to the cell biology of flagella due to the larger cell size and the greater number of flagella per cell.

Each *B. subtilis* cell synthesizes a large number of basal bodies and basal body number is governed by a protein of poorly understood function, SwrA (Kearns *et al.*, 2004; Calvio *et al.*, 2005). SwrA activates the expression the *fla/che* operon that encodes flagellar basal body genes over a four-fold range, and here we demonstrate that SwrA commensurately controls basal body number in motile cells (Kearns and Losick, 2005). The sub-log variation in basal body number, while narrow, appears to be biologically relevant as cells mutated for *swrA* fail to swarm over solid surfaces and cells that overexpress SwrA swarm without lag (Kearns *et al.*, 2004; Kearns and Losick, 2005). Thus, there may be a threshold number of flagella required to traverse solid surfaces (Patrick and Kearns, 2012). SwrA also counteracts the SlrA/SinR/SlrR system that reduces *fla/che* operon transcript levels and increases the frequency of long non-motile chains of cells in the population (Kobayashi, 2008; Chai *et al.*, 2009; Cozy and Kearns, 2010; Chai *et al.*, 2010; Cozy *et al.*, 2012). Consistent with SwrA's dual role in regulating the *fla/che* operon, we now show that the chaining subpopulation of a *swrA* mutant fails to synthesize flagellar basal bodies (Fig 3F). While the mechanism(s) by which SwrA activates the P_{*fla/che*} promoter and antagonizes SlrA/SinR/SlrR is(are) unknown, we infer that basal body number contributes to both swarming motility and the development of heterogeneous cell types in a population.

The *B. subtilis* flagellum assembles rapidly. Basal body formation began within 5 minutes, hook formation began within 10 minutes, and the first filaments were observed as early as

15 minutes after the induction of the *fla/che* operon. The flagellar assembly process appears to be 2 to 3-fold faster in *B. subtilis* than in *S. enterica* (Karlinsey *et al.*, 2000). Flagella assembly time may appear shorter in *B. subtilis* for technical reasons because the basal body structural genes were directly induced whereas in *S. enterica* basal body genes were indirectly induced by expression of the master regulator FlhDC (Karlinsey *et al.*, 2000; Brown *et al.*, 2008; Wozniak *et al.*, 2010). Flagella assembly time may also be shorter in *B. subtilis* for biological reasons. Although the mechanism by which the *B. subtilis* flagellar rod penetrates the Gram positive peptidoglycan is currently unknown, basal body completion may be faster than the Gram negative *S. enterica* due to the absence of the outer membrane. Although basal body assembly occurs quickly, we note that the duration of de novo flagellar assembly required to permit motility still exceeds two doubling times under the conditions we tested.

Flagella were found to be inserted non-randomly along the body of the *B. subtilis* cell, and five aspects of flagellar patterning were observed in the wild type. First, basal bodies were discouraged at the poles. Second, basal bodies appeared absent from the division septum. Third, individual basal bodies were separated from one another by a minimum distance greater than predicted by chance. Fourth, basal bodies were distributed in a grid-like pattern. Fifth, basal bodies were symmetrically distributed on either side of the midcell. The pattern of *B. subtilis* basal body formation was controlled by FlhF and FlhG, the effects of which differed from what has been observed in bacteria with polar flagella. In bacteria with polar flagella, the lack of FlhG results in an increase in flagellar number, whereas in *B. subtilis*, the lack of FlhG resulted in aggregation of flagellar basal bodies (Dasgupta *et al.*, 2000; Correa *et al.*, 2005; Balaban and Hendrixson, 2011). In bacteria with polar flagella, the lack of FlhF misplaces flagella along the sides of the cell, whereas in *B. subtilis*, the lack of FlhF resulted in the accumulation of flagella at one pole (Pandza *et al.*, 2000; Correa *et al.*, 2005; Murray and Lazmierczak, 2006; Green *et al.*, 2009). Even among closely-related peritrichously-flagellated bacteria, FlhF was shown to control flagella number in *Bacillus cereus* but number control was not observed in *B. subtilis* (Salvetti *et al.*, 2007). We infer that the mechanism by which FlhF and FlhG control flagellar number and position is conserved but sufficiently plastic to govern species-specific patterning.

The symmetrical distribution of flagella in *B. subtilis* appears to be related to growth and cell division. Flagella are large, complex, stable structures that are embedded within the fundamental architecture of the cell and when patterning is defective, flagellar segregation may fail such that flagellar inheritance in daughters is stochastic rather than determinative. Inheritance of unequal numbers of flagella may result in perhaps disadvantageous variable motile behavior in individual cells, the consequences of which are lost in the average of the entire population. Furthermore, flagella are imbued with pleiotropic regulatory modules that sense and respond to flagellar assembly state. Thus, unequal inheritance of flagella may also have regulatory consequences as flagellar assembly regulators may become imbalanced and have cascading effects on the expression of genes co-regulated with motility. Thus, optimum spacing and symmetrical distribution of flagella may be a strategy to ensure equal inheritance of a powerful motor that carries epigenetic information. We note that mutants that disrupt either flagellar patterning and/or number in bacteria that normally produce a single flagellum exacerbate inheritance defects perhaps explaining the more severe phenotypes seen in these organisms. Perhaps the high probability of inheriting more than one flagellum is as disabling as inheriting no flagella in patterning deficient mutants of otherwise monotrichous bacteria.

How FlhG antagonism of FlhF controls flagellar patterning with respect to cell division is unknown, but it is intriguing that FlhF is a GTPase, like the cell division protein FtsZ, and that FlhG is an ATPase, like the FtsZ-antagonizing protein MinD (Meinhardt and de Boer,

2001). MinD homologs have been implicated not only in cell division, but are also involved in partitioning plasmids and flagellar-regulating chemotaxis clusters (Thompson *et al.*, 2006; Ringgaard *et al.*, 2009; Ringgaard *et al.*, 2011; Roberts *et al.*, 2012; Vecchiarelli *et al.*, 2012). In each case, the ATPase governs the spacing between cargo. For *B. subtilis* motility, FlhG may set the distance between molecules of FlhF which in turn instructs the cell where to insert new basal bodies relative to pre-existing flagella. Regular spacing may be sufficient to account for the symmetrical distribution of flagella in a grid-like pattern. Finally, we note that flagella do not appear to insert into the division septum even in the absence of FlhF, despite the fact that basal body assembly duration is on the same order as synthesis of the division plane (<10 minutes). How basal bodies would discriminate septal peptidoglycan is unclear but an as-yet-undiscovered active exclusion mechanism may discourage synthesis of intervening flagella that could impair cell separation of incipient daughter cells.

EXPERIMENTAL PROCEDURES

Strains and growth conditions

B. subtilis strains were grown in Luria-Bertani (LB) (10 g tryptone, 5 g yeast extract, 5 g NaCl per L) broth or on LB plates fortified with 1.5% Bacto agar at 37°C. When appropriate, antibiotics were included at the following concentrations: 10 µg/ml tetracycline, 100 µg/ml spectinomycin, 5 µg/ml chloramphenicol, 5 µg/ml kanamycin, and 1 µg/ml erythromycin plus 25 µg/ml lincomycin (*m/s*).

For the swim motility assay, swim agar plates containing 25 ml LB fortified with 0.3% Bacto agar were prepared fresh and each plate was toothpick inoculated from an overnight colony and scored for motility after 18 hours incubation at 37°C.

Swarm expansion assay

Cells were grown to mid-log phase at 37°C in LB broth and resuspended to 10 OD₆₀₀ in pH 8.0 PBS buffer (137 mM NaCl, 2.7 mM KCl, 10 mM Na₂HPO₄, and 2 mM KH₂PO₄) containing 0.5% India ink (Higgins). Freshly prepared LB containing 0.7% Bacto agar (25 ml/plate) was dried for 20 minutes in a laminar flow hood, centrally inoculated with 10 µl of the cell suspension, dried for another 10 minutes, and incubated at 37°C. The India ink demarks the origin of the colony and the swarm radius was measured relative to the origin. For consistency, an axis was drawn on the back of the plate and swarm radii measurements were taken along this transect. For experiments including IPTG, cells were propagated in broth in the presence of IPTG, and IPTG was included in the swarm agar plates.

Microscopy

Fluorescence microscopy was performed with a Nikon 80i microscope with a phase contrast objective Nikon Plan Apo 100X and an Excite 120 metal halide lamp. FM4-64 and Alexa Fluor 594 C₅ maleimide fluorescent signals were visualized with a C-FL HYQ Texas Red Filter Cube (excitation filter 532–587 nm, barrier filter >590 nm). YFP was visualized using a C-FL HYQ YFP Filter Cube (excitation filter 490–510 nm, barrier filter 520–550 nm). GFP and Alexa Fluor 488 C₅ maleimide fluorescent signals were visualized using a C-FL HYQ FITC Filter Cube (FITC, excitation filter 460–500 nm, barrier filter 515–550 nm). CFP fluorescent signals were viewed using a C-FL HYQ CFP Filter Cube (excitation filter 426–446 nm, barrier filter 460–500 nm). TMA-DPH fluorescent signals was visualized using a UV-2E/C DAPI Filter Cube (excitation filter 340–380 nm, barrier filter 435–485 nm). Images were captured with a Photometrics Coolsnap HQ2 camera in black and white, false colored, and superimposed using Metamorph image software.

For FliM-GFP microscopy, cells were grown at 37°C in LB broth until they reached an OD₆₀₀ of 0.5–0.8. 1.0 ml was pelleted and resuspended in 50 µl of PBS containing 5 µg/ml FM4-64 (Molecular Probes) to stain the membrane. The membrane stain was incubated for 5 min at room temperature. Cells were pelleted, washed in 0.2 ml of PBS, and pelleted again. Samples were resuspended to 10OD. Samples were observed by spotting 3 µl of the suspension on a glass microscope slide and were immobilized with a poly-L-lysine-treated glass coverslip.

For co-localization of FliM and EpsE microscopy, cells were grown overnight at 22°C in LB broth. 250µl was pelleted and resuspended in 50µl of PBS containing 5 µg/ml FM4-64 to stain the membrane. The membrane stain was incubated for 5 min at room temperature. Cells were pelleted, washed in 0.2 ml of PBS, and pelleted again. Samples were resuspended in 40µl. Samples were observed by spotting 3 µl of the suspension on a glass microscope slide and were immobilized with a poly-L-lysine-treated glass coverslip.

For co-localization of FliM and FlgE microscopy, cells were grown at 37°C in LB broth until they reached an OD₆₀₀ of 0.5–0.8. 1.0 ml was pelleted and resuspended in 50 µl of PBS containing 5µg/ml Alexa Fluor 594 C₅ maleimide (Molecular Probes), and incubated for 5 min at room temperature. Cells were then washed with 1ml of PBS. Membranes were stained by resuspension in 50 µl of PBS containing 100 µg/ml of TMA-DPH (Molecular Probes) and incubated for 5 min at room temperature. Cells were pelleted, washed in 0.2 ml of PBS, and pelleted again. Samples were resuspended to 10OD. Samples were observed by spotting 3 µl of the suspension on a glass microscope slide and were immobilized with a poly-L-lysine-treated glass coverslip.

For fluorescent microscopy of flagella filaments or hooks, cells were grown at 37°C in LB broth until they reached an OD₆₀₀ of 0.5–0.8. 1.0 ml of broth culture was harvested and resuspended in 50 µl of PBS containing 5µg/ml Alexa Fluor 488 C₅ maleimide (Molecular Probes), and incubated for 5 min at room temperature. Cells were then washed with 1ml of PBS. Membranes were stained by resuspension in 50 µl of PBS containing 5 µg/ml FM4-64 and incubated for 5 min at room temperature. Cells were pelleted, washed in 0.2 ml of PBS, and pelleted again. Samples were resuspended to 10OD. Samples were observed by spotting 3 µl of the suspension on a glass microscope slide and were immobilized with a poly-L-lysine-treated glass coverslip.

For fluorescent microscopy of flagella filaments over three generations, 1.5 ml of broth culture was harvested at approximately OD₆₀₀=0.6, resuspended in 75 µl of PBS containing 5µg/ml Alexa Fluor 488, and incubated for 5 min at room temperature. Cells were then washed twice with 1.5 ml of PBS and resuspended in 1.5ml of LB broth. The stained cells were diluted two, four, and eight-fold in fresh LB and grown back to an approximate OD₆₀₀ of 0.6. 1.0 ml of broth culture was harvested, resuspended in 50 µl of PBS containing 5µg/ml Alexa Fluor 594 C₅ maleimide, and incubated for 5 min at room temperature. Cells were then washed twice with 1ml of PBS. Membranes were stained by resuspension in 50 µl of PBS containing 50 µl of PBS containing 100 µg/ml of TMA-DPH and incubated for 5 min at room temperature. Samples were observed by spotting 3 µl of the suspension on a glass microscope slide and were immobilized with a poly-L-lysine-treated glass coverslip.

For the timing of FliM, FlgE, and Hag assembly, the indicated strains were grown at 37°C in LB kan⁵ to an OD₆₀₀ of 0.4–0.5. A final concentration of 0.3mM IPTG was then added to the cells and incubated further for the times indicated. Cells were harvested at the indicated time points and cells were prepared and imaged as described above.

OMX Microscopy

A Deltavision OMX 3D-SIM Super Resolution System V3.0 (Applied Precision) was used for Super Resolution and 3D microscopy. Images were acquired with the E Cascade II 512 EM-CCD camera (Photometrics, Tuscon AZ) and a 1.4NA Olympus 100X oil objective sequentially using the 561 (emission filter 609–654) and 488 (emission filter 500–550) nm lasers. Exposure times were between 50–100 ms and immersion oil with a 1.514 refractive index was used. Image acquisition was directed by OMX controller Software (Applied Precision Inc.). Image reconstructions were made with the OMX specific SoftWoRx software package (Applied Precision Inc).

OMX Image Analysis

Images that were reconstructed with softWoRx were used in Imaris (Bitplane Inc.) to determine the number and location of FliM-GFP in the cell. The spots feature labeled each FliM-GFP foci by the search parameter of identifying spots of $.1\mu\text{M}$ in the 488 wavelength. We verified by eye that the spots labeling identified all the foci. Cell pole positions were determined by using the 561 nm wavelength to create a surface. Cells were rotated to put a spot on the surface to label each cell pole. The x, y, and z coordinates of each foci in each cell was exported from Imaris. Scripts were developed in MATLAB (The Mathworks) for importing 3-dimensional Cartesian coordinates for basal body positions and cell end positions from Imaris. Scripts were developed to center and orient cells using an Eigenvector transform approach and the cell endwall positions. The cell length and midpoint on the long axis were determined using the manually selected or verified endwall positions. The 3 dimensional positions of the basal bodies were transformed on to a 2 dimensional area by computing the distance and angle from a midline through the oriented cell. Nearest neighbor distances were determined by calculating all distances from an axially centered basal body on the 2 dimensional transform and finding the shortest distance. All fits to the data are simple linear regression functions.

The cell length, circumferential distance, and number of basal bodies was recorded for each wild type and mutant cell and used as input for creating each simulated cell with a uniform random distribution of basal bodies. For each trial, the data was input for a cell and positions on the long axis and around the circumferential distance of the cell were drawn from independent random uniform distributions. To create a population of cell estimates, cell data was drawn from the population of wild type or mutant cells at random with replacement.

Clark-Evans ratios were calculated per cell as the mean nearest neighbor distance divided by half the square root of the basal body area density on the cell. Area density was calculated as the surface area of the cylindrical cell (long axis length multiplied by the cylindrical distance excluding the end wall area) divided by the number of basal bodies.

The normalized distribution of positions on the long axis of the cell was performed by setting the midpoint of the cell equal to zero and the end wall positions as 1 and -1 . Since there is no orientation marker for direction, the position of each spot was normalized to a position between the midpoint and end wall and the absolute value was used to construct histograms. In all cases, p-values were determined using a 2-tailed Student's t-test.

Strain construction

All constructs were first introduced into the domesticated strain PY79 by natural competence and then transferred to the 3610 background using SPP1-mediated generalized phage transduction (Yasbin and Young, 1974). All strains used in this study are listed in Table 2. All plasmids used in this study are listed in Supplemental Table S1. All primers used in this study are listed in Supplemental Table S2.

Complementation constructs

To generate the *amyE::P_{fla/che}-fliM* complementation construct (pSG29), *P_{fla/che}* was amplified from 3610 with primer pair 321/1406 and digested with *EcoRI* and *Sall* and *fliM* was amplified from 3610 with primer pair 1707/1708 and digested with *Sall* and *NheI*. Both digested PCR fragments were simultaneously ligated into the *EcoRI/NheI* sites of pAH25 containing a spectinomycin resistance gene between the arms of the *amyE* gene (Blair *et al.*, 2008).

To generate the *thrC::P_{fla/che}-flhF* complementation construct (pSG59), *P_{fla/che}* was amplified from 3610 with primer pair 321/1406 and digested with *EcoRI* and *Sall* and *flhF* was amplified from 3610 with primer pair 2800/2801 and digested with *Sall* and *BamHI*. Both digested PCR fragments were simultaneously ligated into the *EcoRI/BamHI* sites of pDG1664 containing a erythromycin (*mls*) resistance cassette and a polylinker downstream between the arms of the *thrC* gene (Guérout-Fleury *et al.*, 1996).

To generate the *thrC::P_{fla/che}-flhG* complementation construct (pSG62), *P_{fla/che}* was amplified from 3610 with primer pair 321/1406 and digested with *EcoRI* and *Sall* and *flhG* was amplified from 3610 with primer pair 2804/2805 and digested with *Sall* and *BamHI*. Both digested PCR fragments were simultaneously ligated into the *EcoRI/BamHI* sites of pDG1664 containing a erythromycin (*mls*) resistance cassette and a polylinker downstream between the arms of the *thrC* gene.

To generate the *lacA::P_{fla/che}-flhFflhG* complementation construct (pSG71), *P_{fla/che}* was amplified from 3610 with primer pair 321/1406 and digested with *EcoRI* and *Sall* and *flhFflhG* were amplified from 3610 with primer pair 2800/2805 and digested with *Sall* and *BamHI*. Both digested PCR fragments were simultaneously ligated into the *EcoRI/BamHI* sites of pNCO18 containing a tet resistance gene between the arms of the *lacA* gene.

To generate *thrC::P_{fla/che}-flgE^{T123C}* complementation construct (pSG68), the *P_{fla/che}-flgE^{T123C}* region was amplified with primer pair 321/1854 from pCC16 as a template and digested with *EcoRI* and *BamHI*. The PCR fragment was ligated into the *EcoRI/BamHI* sites of pDG1664.

To generate *lacA::P_{hag}-hag^{T209C}* (pKB142), a fragment containing *P_{hag}-hag^{T209C}* was digested from pND4 using *BamHI* and *Sall* (Blair *et al.*, 2008). This fragment was ligated into the *BamHI/Sall* sites of pDR183 which contains a *mls* resistance gene between the arms of the *lacA* gene.

Translational Fusions

To generate the translational fusion of FliM to CFP (pSG30), a fragment containing *fliM* was amplified using 3610 as template and primer pair 1707/1709 and was digested with *Sall* and *NheI*. The *cfp* gene was amplified using pDR200 as a template and primer pair 1706/1173 and was digested with *NheI* and *BamHI*. Both digested fragments were simultaneously ligated into the *Sall/BamHI* sites of pSG29 which contains the *P_{fla/che}* promoter and spectinomycin resistance between the arms of the *amyE* gene to generate pSG30. Primer 1709 (reverse primer for *fliM* amplification) incorporated the sequence of a flexible linker domain (Waldo *et al.*, 1999).

To generate the translational fusion of FliM to GFP, the same strategy was employed as described for FliM-CFP except the *gfp* gene was amplified using pKB59 as template and primer pair 2301/996. This generated pSG49.

To generate the translational fusion of EpsE to YFP, a fragment containing *P_{eps}-epsE* was amplified using pKB59 as a template and primer pair 709/975 and was digested with *EcoRI* and *XhoI*. The *yfp* gene was amplified from pKL183 with primer pair 995/996 and was digested with *XhoI* and *BamHI*. Both digested fragments were simultaneously ligated into the *EcoRI/BamHI* sites of pDG1664 to generate pSG23. A flexible linker domain was incorporated in the reverse primer for *epsE* (Primer 975).

In-frame deletions

To generate the $\Delta fliM$ in frame marker-less deletion construct, the region upstream of *fliM* was PCR amplified using the primer pair 1569/1570 and digested with *EcoRI* and *BamHI*, and the region downstream of *fliM* was PCR amplified using the primer pair 1813/1572 and digested with *BamHI* and *Sall*. The two fragments were then simultaneously ligated into the *EcoRI/Sall* sites of pMiniMAD which carries a temperature sensitive origin of replication and an erythromycin resistance cassette to generate pSG32 (Patrick and Kearns, 2008). The plasmid pSG32 was introduced to PY79 by single cross-over integration by transformation at the restrictive temperature for plasmid replication (37°C) using mls resistance as a selection. The integrated plasmid was then transduced into 3610. To evict the plasmid, the strain was incubated in 3ml LB broth at a permissive temperature for plasmid replication (22°C) for 14 hours, diluted 30-fold in fresh LB broth, and incubated at 22°C for another 8 hours. Dilution and outgrowth was repeated 2 more times. Cells were then serially diluted and plated on LB agar at 37°C. Individual colonies were patched on LB plates and LB plates containing mls to identify mls sensitive colonies that had evicted the plasmid. Colony PCR was used on mls sensitive colonies using primer pair 1569/1572 to determine which colony retained the $\Delta fliM$ allele.

To generate pSG70 ($\Omega \Delta flhF flhG$) the region upstream of *flhF* was PCR amplified using the primer pair 2103/2104 and digested with *EcoRI* and *XhoI*, and the region downstream of *flhG* was PCR amplified using the primer pair 2894/2895 and digested with *XhoI* and *BamHI*. The two fragments were then simultaneously ligated into the *EcoRI/BamHI* sites of pMiniMAD. pSG70 was integrated into the *B. subtilis* PY79 genome, transduced to strain 3610, and evicted as described above. Colonies were screened by PCR using primers 2103/2895 to determine which isolate had retained the $\Delta flhF flhG$ allele.

To generate pLC22 ($\Omega \Delta flhG$) the region upstream of *flhG* was PCR amplified using the primer pair 826/827 and digested with *EcoRI* and *BamHI*, and the region downstream of *flhG* was PCR amplified using the primer pair 828/829 and digested with *BamHI* and *Sall*. The two fragments were then simultaneously ligated into the *EcoRI/Sall* sites of pMiniMAD. pLC22 was integrated into the *B. subtilis* PY79 genome, transduced to strain 3610, and evicted as described above. Colonies were screened by PCR using primers 826/829 to determine which isolate had retained the $\Delta flhG$ allele.

To generate pEC2 ($\Omega \Delta motB$) the region upstream of *motB* was PCR amplified using the primer pair 2453/2454 and digested with *EcoRI* and *XhoI*, and the region downstream of *motB* was PCR amplified using the primer pair 2455/2456 and digested with *XhoI* and *BamHI*. The two fragments were then simultaneously ligated into the *EcoRI/BamHI* sites of pMiniMAD. pEC2 was integrated into the *B. subtilis* PY79 genome, transduced to strain 3610, and evicted as described above. Colonies were screened by PCR using primers 2453/2456 to determine which isolate had retained the $\Delta motB$ allele.

To generate pDP333 ($\Omega \Delta flhF$) the region upstream of *flhF* was PCR amplified using the primer pair 2103/2104 and digested with *EcoRI* and *XhoI*, and the region downstream of *flhF* was PCR amplified using the primer pair 2105/2106 and digested with *XhoI* and *BamHI*. The two fragments were then simultaneously ligated into the *EcoRI/BamHI* sites of

pMiniMAD. pDP333 was integrated into the *B. subtilis* PY79 genome, transduced to strain 3610, and evicted as described above. Colonies were screened by PCR using primers 2103/2106 to determine which isolate had retained the $\Delta flhF$ allele.

IPTG inducible constructs

To generate the inducible *P_{fla/che}OP_{hyspank}-fla/che* operon construct pSG79, a fragment containing the *P_{hyspank}* promoter, *flgB* gene, and gene for the LacI repressor was digested from pDP161 using *EcoRI* and *BamHI* (Kearns and Losick, 2005). This fragment was ligated into the *EcoRI/BamHI* sites of pDG780 which contains a kanamycin resistance gene to generate pSG79 (Guérout-Fleury *et al.*, 1996).

To generate the inducible *thrC::P_{hyspank}-flhF* construct pSG60, a fragment containing *flhF* was amplified from 3610 using primer pair 2802/2803 and digested with *NheI* and *SphI*. The fragment was ligated into the *NheI/SphI* sites of pDP150.

To generate the inducible *thrC::P_{hyspank}-flhFflhG* construct pSG82, a fragment containing *flhFflhG* was amplified from 3610 using primer pair 2802/2807 and digested with *NheI* and *SphI*. The fragment was ligated into the *NheI/SphI* sites of pDP150.

Allelic Replacement

The *sinR::kan* mutation was constructed as previously described (Blair *et al.*, 2008). The *epsH::tet* mutation was constructed as previously described (Branda *et al.*, 2001). The *sigD::tet* mutation was constructed as previously described (Kearns and Losick, 2005). The *srfAA::mls* mutation was constructed as previously described (Kearns and Losick, 2003).

SPP1 phage transduction

To 0.2 ml of dense culture grown in TY broth (LB broth supplemented after autoclaving with 10 mM MgSO₄ and 100 μ M MnSO₄), serial dilutions of SPP1 phage stock were added and statically incubated for 15 minutes at 37°C. To each mixture, 3 ml TYSA (molten TY supplemented with 0.5% agar) was added, poured atop fresh TY plates, and incubated at 30°C overnight. Top agar from the plate containing near confluent plaques was harvested by scraping into a 15 ml conical tube, treated with 25 μ g/ml DNase final concentration, vortexed and centrifuged at 5,000 $\times$ g for 10 minutes. The supernatant was passed through a 0.45 μ m syringe filter and stored at 4°C.

Recipient cells were grown to stationary phase in 3 ml TY broth at 37°C. 1.0 ml cells were mixed with 25 μ l of SPP1 donor phage stock. 9 ml of TY broth was added to the mixture and incubated statically at 37°C for 30 minutes. The transduction mixture was then centrifuged at 5,000 $\times$ g for 10 minutes, the supernatant was discarded and the pellet was resuspended in the remaining volume. 100 μ l of cell suspension was then plated on TY fortified with 1.5% agar, the appropriate antibiotic, and 10 mM sodium citrate.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Jim Powers for 3D OMX SIM microscope training. We thank Sampri Mukherjee and Anna Bree for technical assistance and Kris Blair, Loralyn Cozy, Joyce Patrick, and Colleen Courtney for strains and reagents. We thank Masaya Fujita for the gift of the anti-SigA antibody and David Rudner for the gift of the anti-GFP antibody. Some of this work was supported by the Indiana METACyt Initiative of Indiana University, funded in part through a major grant from the Lilly Endowment. The 3D OMX SIM microscope was funded by NIH grant

S10RR028697-01 and the work was supported by NSF Grant MCB1157982 to SLS and NIH Grant GM093030 to DBK.

REFERENCES

- Aizawa, S-I.; Zhulin, IB.; Márquez-Magaña, L.; Ordal, GW. Chemotaxis and motility. In: Sonenshein, AL.; Hock, JA.; Losick, R., editors. *Bacillus subtilis* and Its Closest Relatives: From Genes to Cells. Washington, DC: American Society of Microbiology; 2002. p. 437-451.
- Albertini AM, Caramori T, Crabb WD, Scoffone F, Galizzi A. The *flaA* locus of *Bacillus subtilis* is part of a large operon coding for flagellar structures, motility functions, and an ATPase-like polypeptide. *J Bacteriol.* 1991; 173:3573–3579. [PubMed: 1828465]
- Aldridge P, Karlinsey JE, Becker E, Chevance FFV, Hughes KT. Flk prevents premature secretion of the anti-sigma factor FlgM into the periplasm. *Mol Microbiol.* 2006; 60:630–642. [PubMed: 16629666]
- Alley MR, Maddock JR, Shapiro L. Polar localization of a bacterial chemoreceptor. *Genes Dev.* 1992; 6:825–836. [PubMed: 1577276]
- Atsumi T, Maekawa Y, Yamada T, Kawagishi I, Imae Y, Homma M. Effect of viscosity on swimming by the lateral and polar flagella of *Vibrio alginolyticus*. *J Bacteriol.* 1996; 178:5024–5026. [PubMed: 8759871]
- Balaban M, Hendrixson DR. Polar flagellar biosynthesis and a regulator of flagellar number influence spatial parameters of cell division in *Campylobacter jejuni*. *PLoS Path.* 2011; 7 e1002420.
- Bange G, Petzold G, Wild K, Parlitz RO, Sinning I. The crystal structure of the third signal-recognition particle GTPase FlhF reveals a homodimer with bound GTP. *Proc Natl Acad Sci USA.* 2007; 104:13621–13625. [PubMed: 17699634]
- Bange G, Kümmerer N, Grudnik P, Linder R, Petzold G, Kressler D, Hurt E, Wild K, Sinning I. Structural basis for the molecular evolution of SRP-GTPase activation by protein. *Nat Struct Mol Biol.* 2011; 18:1376–1380. [PubMed: 22056770]
- Bi E, Lutkenhaus J. FtsZ ring structure associated with division in *Escherichia coli*. *Nature.* 1991; 354:161–164. [PubMed: 1944597]
- Blackman SA, Smith TJ, Foster SJ. The role of autolysins during vegetative growth of *Bacillus subtilis* 168. *Microbiol.* 1998; 144:73–82.
- Blair KM, Turner L, Winkelman JT, Berg HC, Kearns DB. A molecular clutch disables flagella in the *Bacillus subtilis* biofilm. *Science.* 2008; 320:1636–1638. [PubMed: 18566286]
- Boltjes TYK. Function and arrangement of bacterial flagella. *J Pathol Bacteriol.* 1948; 60:275–287. [PubMed: 18099664]
- Branda SS, Gonzalez-Pastor JE, Ben-Yehuda S, Losick R, Kolter R. Fruiting body formation by *Bacillus subtilis*. *Proc Natl Acad Sci USA.* 2001; 98:11621–11626. [PubMed: 11572999]
- Brown JD, Saini S, Aldridge C, Herbert J, Rao CV, Aldridge PD. The rate of protein secretion dictates the temporal dynamics of flagellar gene expression. *Mol Microbiol.* 2008; 70:924–937. [PubMed: 18811728]
- Calvio C, Celandroni F, Ghelardi E, Amati G, Salvetti S, Cecilian F, et al. Swarming differentiation and swimming motility in *Bacillus subtilis* are controlled by *swrA*, a newly identified dicistronic operon. *J Bacteriol.* 2005; 187:5356–5366. [PubMed: 16030230]
- Chai Y, Kolter R, Losick R. Paralogous antirepressors acting on the master regulator for biofilm formation in *Bacillus subtilis*. *Mol Microbiol.* 2009; 74:876–887. [PubMed: 19788541]
- Chai Y, Norman T, Kolter R, Losick R. An epigenetic switch governing daughter cell separation in *Bacillus subtilis*. *Genes Dev.* 2010; 24:754–765. [PubMed: 20351052]
- Chen R, Guttenplan SB, Blair KM, Kearns DB. Role of the σ^D -dependent autolysins in *Bacillus subtilis* population heterogeneity. *J Bacteriol.* 2009; 191:5775–5784. [PubMed: 19542270]
- Chevance FFV, Hughes KT. Coordinating assembly of a bacterial macromolecular machine. *Nat Rev Microbiol.* 2008; 6:445–465.
- Clark PJ, Evans FC. Distance to nearest neighbor as a measure of spatial relationships in populations. *Ecology.* 1954; 35:445–453.

- Correa NE, Peng F, Klose KE. Roles of the regulatory proteins FlhF and FlhG in the *Vibrio cholerae* flagellar transcription hierarchy. *J Bacteriol.* 2005; 187:6324–6332. [PubMed: 16159765]
- Courtney CR, Cozy LM, Kearns DB. Molecular characterization of the flagellar hook in *Bacillus subtilis*. *J Bacteriol.* 2012; 194:4619–4629. [PubMed: 22730131]
- Cozy LM, Kearns DB. Gene position in a long operon governs motility development in *Bacillus subtilis*. *Mol Microbiol.* 2010; 76:273–285. [PubMed: 20233303]
- Cozy LM, Phillips AM, Calvo RA, Bate AR, Hsueh Y-H, Bonneau R, et al. SlrA/SinR/SlrR inhibits motility gene expression upstream of a hypersensitive and hysteretic switch at the level of σ^D in *Bacillus subtilis*. *Mol Microbiol.* 2012; 83:1210–1228. [PubMed: 22329926]
- Dasgupta N, Arora SK, Ramphal R. *flaN*, a gene that regulates flagellar number in *Pseudomonas aeruginosa*. *J Bacteriol.* 2000; 182:357–364. [PubMed: 10629180]
- Erhardt M, Hughes KT. C-ring requirement in flagellar type III secretion is bypassed by FlhDC upregulation. *Mol Microbiol.* 2010; 75:376–393. [PubMed: 19919668]
- Erhardt M, Singer HM, Wee DH, Kenner JP, Hughes KT. An infrequent molecular ruler controls flagellar hook length in *Salmonella enterica*. *EMBO J.* 2011; 30:2948–2961. [PubMed: 21654632]
- Green JCD, Kahramanoglou C, Rahman A, Pender AMC, Charbonnel N, Fraser GM. Recruitment of the earliest component of the bacterial flagellum to the old cell division pole by a membrane-associated signal recognition particle family GTP-binding protein. *J Mol Biol.* 2009; 391:679–690. [PubMed: 19497327]
- Guérout-Fleury AM, Frandsen N, Stragier P. Plasmids for ectopic integration in *Bacillus subtilis*. *Gene.* 1996; 180:57–61. [PubMed: 8973347]
- Guttenplan SB, Blair KM, Kearns DB. The EpsE flagellar clutch is bifunctional and synergizes with EPS biosynthesis to promote *Bacillus subtilis* biofilm formation. *PLoS Genet.* 2010; 6:e1001243.
- Hughes KT, Gillen KL, Semon MJ, Karlinsey JE. Sensing structural intermediates in bacterial flagellar assembly by export of a negative regulator. *Science.* 1993; 262:1277–1280. [PubMed: 8235660]
- Iino T. Polarity of flagellar growth in *Salmonella*. *J Gen Microbiol.* 1969; 56:227–239. [PubMed: 4893344]
- Journet L, Agrain C, Broz P, Cornelis GR. The needle length of bacterial injectisomes is determined by a molecular ruler. *Science.* 2003; 302:1757–1760. [PubMed: 14657497]
- Kalir S, McClure J, Pabbaraju K, Southward C, Ronen M, Leibler S, et al. Ordering genes in a flagella pathway by analysis of expression kinetics from living bacteria. *Science.* 2001; 292:2080–2083. [PubMed: 11408658]
- Karlinsey JE, Tanaka S, Bettenworth V, Yamaguchi S, Boos W, Aizawa S-I, Hughes KT. Completion of the hook-basal body complex of the *Salmonella typhimurium* flagellum is coupled to FlgM secretion and *fliC* transcription. *Mol Microbiol.* 2000; 37:1220–1231. [PubMed: 10972838]
- Kearns DB, Losick R. Swarming motility in undomesticated *Bacillus subtilis*. *Mol Microbiol.* 2003; 49:581–590. [PubMed: 12864845]
- Kearns DB, Chu F, Rudner R, Losick R. Genes governing swarming in *Bacillus subtilis* and evidence for a phase variation mechanism controlling surface motility. *Mol Microbiol.* 2004; 52:357–369. [PubMed: 15066026]
- Kearns DB, Losick R. Cell population heterogeneity during growth of *Bacillus subtilis*. *Genes Dev.* 2005; 19:3083–3094. [PubMed: 16357223]
- Kearns DB. A field guide to bacterial swarming motility. *Nat Rev Microbiol.* 2010; 8:634–644. [PubMed: 20694026]
- Kobayashi K. SlrR/SlrA controls the initiation of biofilm formation in *Bacillus subtilis*. *Mol Microbiol.* 2008; 69:1399–1410. [PubMed: 18647168]
- Komeda Y, Silverman M, Simon M. Identification of the structural gene for the hook subunit protein of *Escherichia coli* flagella. *J Bacteriol.* 1978; 133:364–371. [PubMed: 338588]
- Kusumoto A, Shinohara A, Terashima H, Kojima S, Yakushi T, Homma M. Collaboration of FlhF and FlhG to regulate polar-flagella number and localization in *Vibrio alginolyticus*. *Microbiol.* 2008; 154:1390–1399.

- Kusumoto A, Nishioka N, Kojima S, Homma M. Mutational analysis of the GTP-binding motif of FlhF which regulates the number and placement of the polar flagellum in *Vibrio alginolyticus*. *J Biochem*. 2009; 146:643–650. [PubMed: 19605463]
- Kutsukake K. Excretion of the anti-sigma factor through a flagellar substructure couples flagellar gene expression with flagellar assembly in *Salmonella typhimurium*. *Mol Gen Genet*. 1994; 243:605–612. [PubMed: 8028576]
- Leifson E. Staining, shape, and arrangement of bacterial flagella. *J Bacteriol*. 1951; 62:377–389. [PubMed: 14897809]
- Lemon KP, Grossman AD. Movement of replicating DNA through a stationary replisome. *Mol Cell*. 2000; 6:1321–1330. [PubMed: 11163206]
- Liu R, Ochman H. Stepwise formation of the bacterial flagellar system. *Proc Natl Acad Sci USA*. 2007; 104:7116–7121. [PubMed: 17438286]
- Ma X, Ehrhardt DW, Margolin W. Colocalization of cell division proteins FtsZ and FtsA to cytoskeletal structures in living *Escherichia coli* cells by using green fluorescent protein. *Proc Natl Acad Sci USA*. 1996; 93:12998–13003. [PubMed: 8917533]
- Macnab RM. Genetics and biogenesis of bacterial flagella. *Annu Rev Genet*. 1992; 26:131–158. [PubMed: 1482109]
- Macnab, RM. Flagella and motility. In: Neidhart, FC., editor. *Escherichia coli and Salmonella typhimurium: Cellular and Molecular Biology*. 2nd edition. Washington, DC: American Society of Microbiology; 1996. p. 123-145.
- Macnab RM. How bacteria assemble flagella. *Annu Rev Microbiol*. 2003; 57:77–100. [PubMed: 12730325]
- Maddock JR, Shapiro L. Polar localization of the chemoreceptor complex in the *Escherichia coli* cell. *Science*. 1993; 259:1717–1723. [PubMed: 8456299]
- Margot P, Pagni M, Karamata D. *Bacillus subtilis* 168 gene *lytF* encodes a γ -D-glutamate-meso-diaminopimelate murepeptidase expressed by the alternative sigma factor, σ^D . *Microbiol*. 1999; 145:57–65.
- Márquez-Magaña LM, Chamberlin MJ. Characterization of the *sigD* transcription unit of *Bacillus subtilis*. *J Bacteriol*. 1994; 176:24247–22434.
- McCarter LL. Dual flagellar systems enable motility under different circumstances. *J Mol Microbiol Biotechnol*. 2004; 7:18–29. [PubMed: 15170400]
- Meinhardt H, de Boer PAJ. Pattern formation in *Escherichia coli*: a model for the pole-to-pole oscillation of Min proteins and the localization of the division site. *Proc Natl Acad Sci USA*. 2001; 98:14202–14207. [PubMed: 11734639]
- Mukherjee S, Yakhnin H, Kysela D, Sokoloski J, Babitzke P, Kearns DB. CsrA-FliW interaction governs flagellin homeostasis and a checkpoint on flagellar morphogenesis in *Bacillus subtilis*. *Mol Microbiol*. 2011; 82:447–461. [PubMed: 21895793]
- Murray TS, Kazmierczak BI. FlhF is required for swimming and swarming of *Pseudomonas aeruginosa*. *J Bacteriol*. 2006; 188:6995–7004. [PubMed: 16980502]
- Pallen MJ, Penn CW, Chaudhuri RR. Bacterial flagellar diversity in the post-genomic era. *Trends Microbiol*. 2005; 13:143–149. [PubMed: 15817382]
- Pandza S, Baetens M, Park CH, Au T, Keyhan M, Matin A. The G-protein FlhF has a role in polar flagellar placement and general stress response induction in *Pseudomonas putida*. *Mol Microbiol*. 2000; 36:414–423. [PubMed: 10792727]
- Patrick JE, Kearns DB. MinJ (YvjD) is a topological determinant of cell division in *Bacillus subtilis*. *Mol Microbiol*. 2008; 70:1166–1179. [PubMed: 18976281]
- Patrick JE, Kearns DB. Laboratory strains of *Bacillus subtilis* do not exhibit swarming motility. *J Bacteriol*. 2009; 191:7129–7133. [PubMed: 19749039]
- Patrick JE, Kearns DB. Swarming motility and the control of master regulators of flagellar biosynthesis. *Mol Microbiol*. 2012; 83:14–23. [PubMed: 22092493]
- Ringgaard S, van Zon J, Howard M, Gerdes K. Movement and equipositioning of plasmids by ParA filament assembly. *Proc Natl Acad Sci USA*. 2009; 46:19369–19374. [PubMed: 19906997]

- Ringgaard S, Schirner K, Davis BM, Waldor MK. A family of ParA-like ATPases promotes cell pole maturation by facilitating polar localization of chemotaxis proteins. *Genes Dev.* 2011; 25:1544–1555. [PubMed: 21764856]
- Roberts MAJ, Wadhams GH, Hadfield KA, Tickner S, Armitage JP. ParA-like protein uses nonspecific chromosomal DNA binding to partition protein complexes. *Proc Natl Acad Sci USA.* 2012; 109:6698–6703. [PubMed: 22496588]
- Salveti S, Ghelardi E, Celandroni F, Ceragioli M, Giannessi F, Senesi S. FlhF, a signal recognition particle-like GTPase, is involved in the regulation of flagellar arrangement, motility behavior and protein secretion in *Bacillus cereus*. *Microbiol.* 2007; 153:2541–2552.
- Samatey FA, Matusnami H, Imada K, Nagashima S, Shaikh TR, Thomas DR, et al. Structure of the bacterial flagellar hook and implication for the molecular universal joint mechanism. *Nature.* 2004; 431:1062–1068. [PubMed: 15510139]
- Serizawa M, Yamamoto H, Yamaguchi H, Fujita Y, Kobayashi K, Ogasawara N, Sekiguchi J. Systematic analysis of SigD-regulated genes in *Bacillus subtilis* by DNA microarray and Northern blotting analyses. *Gene.* 2004; 329:125–136. [PubMed: 15033535]
- Shibata S, Takahashi N, Chevance FFV, Karlinsey JE, Hughes KT, Aizawa S-I. FliK regulates flagellar hook length as an internal ruler. *Mol Microbiol.* 2007; 64:1404–1415. [PubMed: 17542929]
- Sourjik V, Berg HC. Localization of components of the chemotaxis machinery of *Escherichia coli* using fluorescent protein fusions. *Mol Microbiol.* 2000; 37:740–751. [PubMed: 10972797]
- Stanier RY, Van Niel CB. The main outlines of bacterial classification. *J Bacteriol.* 1941; 42:437–466. [PubMed: 16560462]
- Thiem S, Sourjik V. Stochastic assembly of chemoreceptor clusters in *Escherichia coli*. *Mol Microbiol.* 2008; 68:1228–1236. [PubMed: 18476921]
- Thompson SR, Wadhams GH, Armitage JP. The positioning of cytoplasmic protein clusters in bacteria. *Proc Natl Acad Sci USA.* 2006; 103:8209–8214. [PubMed: 16702547]
- Toker AS, Macnab RM. Distinct regions of bacterial flagellar switch protein FliM interact with FliG, FliN, and CheY. *J Mol Biol.* 1997; 273:623–634. [PubMed: 9356251]
- Ueno T, Oosawa K, Aizawa S-I. M ring, S ring, and proximal rod of the flagellar basal body of *Salmonella typhimurium* are composed of a subunits of a single protein, FliF. *J Mol Biol.* 1992; 227:672–677. [PubMed: 1404383]
- Vecchiarelli AG, Mizuuchi K, Funnell BE. Surfing biological surfaces: exploiting the nucleoid for partition and transport in bacteria. *Mol Microbiol.* 2012 (in press).
- Waldo GS, Standish BM, Berendzen J, Terwilliger TC. Rapid protein-folding assay using green fluorescent protein. *Nat Biotechnol.* 1999; 17:691–695. [PubMed: 10404163]
- Wang H, Wingreen NS, Mukhopadhyay R. Self-organized periodicity of protein clusters in growing bacteria. *Phys Rev Lett.* 2008; 101 218101.
- Wozniak CE, Chevance FFV, Hughes KT. Multiple promoters contribute to swarming and the coordination of transcription with flagellar assembly in *Salmonella*. *J Bacteriol.* 2010; 192:4752–4762. [PubMed: 20639318]
- Yasbin RE, Young FE. Transduction in *Bacillus subtilis* by Bacteriophage SPP1. *J Virol.* 1974; 14:1343–1348. [PubMed: 4214946]
- Zhou J, Sharp LL, Tang HL, Lloyd SA, Billings S, Braun TF, Blair DF. Function of protonatable residues in the flagellar motor of *Escherichia coli*: a critical role for Asp32 of MotB. *J Bacteriol.* 1998; 180:2729–2735. [PubMed: 9573160]

Figure 1. Flagellar arrangement and architecture

A) Various spatial arrangements of bacterial flagella. Polar indicates a single flagellum at one or both poles. Lophotrichous indicates multiple flagella at one or both poles. Medial indicates a single flagellum arising from along the cell body. Peritrichous indicates multiple flagella arising from along the cell body. Flagellar filaments indicated in green, flagellar hooks indicated in blue, flagellar basal bodies indicated in red. B) The putative architecture of the *B. subtilis* Gram positive flagellum. The flagellar filament indicated in green, the flagellar hook is indicated in blue, and flagellar basal body is indicated in pink. The locations of specific proteins are indicated with a color matching their position. The membrane is colored light gray and peptidoglycan is colored dark gray.

Figure 3. FliM-GFP forms puncta that depend on the flagellar basal body FliF

Fluorescence microscopy of the indicated strains stained for membranes with FM4-64 (false colored red) and FliM-GFP (false colored green). All strains expressed FliM-GFP inserted at an ectopic location (*amyE*). Scale bar is 1 micron. The following strains were used to generate this figure: A) wild type (DK15), B) *fliM* (DS8521), C) *fliF* (DS8598), D) *motB* (DS8979), E) *sigD* (DS9105), F) *swrA* (DS8600), and G) *swrA*⁺⁺ (DS8608) where *swrA* was expressed from an artificial *P_{hyspank}* promoter and induced with 1 mM IPTG.

Figure 4. FliM-GFP co-localizes with the flagella clutch and hook proteins

A) Fluorescence microscopy of strain DS8996 stained for membranes with TMA-DPH (false colored blue), FlgE^{T123C} (false colored red) and FliM-GFP (false colored green). Scale bar is 0.5 micron. White carets indicate points of basal body/hook co-localization. B) Fluorescence microscopy of strain DK16 stained for membranes with FM4-64 (false colored red), EpsE-YFP (false colored green) and FliM-CFP (false colored blue). Scale bar is 1 micron. White carets indicate points of basal body/clutch co-localization.

Figure 5. Timing of filament, hook, and basal body assembly

A) Fluorescence microscopy overlay of strain DK31 ($\Delta fliM$ $amyE::P_{fla/che}$ - $fliM$ -GFP spec $P_{fla/che}\Omega P_{hy-spank}$ - fla/che operon kan) stained with membrane dye FM 4-64 (false colored red) and FliM-GFP (false colored green) at the indicated time points following induction with 0.3 mM IPTG. Carets indicate the first appearance of puncta. B) Timeline of flagella assembly in minutes following induction. Red bar indicates period during which FliM-GFP puncta (basal bodies) first appear. Blue bar indicates period during which FlgE^{T123C} puncta (hooks) first appear. Green bar indicates period during which Hag^{T209C} filaments (filaments) first appear. Gray bar indicates period when motile cells first appear as determined by wet mount microscopy. Dotted lines indicate that assembly likely occurred prior to the timepoint of first observation. C) Fluorescence microscopy overlay of strain DK30 ($\Delta flgE$ $amyE::P_{fla/che}$ - $flgE^{T123C}$ cat $P_{fla/che}\Omega P_{hy-spank}$ - fla/che operon kan) stained with membrane dye FM 4-64 (false colored red) and FlgE stained with a maleimide reactive dye (false colored green) at the indicated time points following induction with 0.3 mM IPTG. Carets indicate the first appearance of puncta. D) Fluorescence microscopy overlay of strain DK62 (Δhag $amyE::P_{hag}$ - hag^{T209C} spec $P_{fla/che}\Omega P_{hy-spank}$ - fla/che operon kan) stained with membrane dye FM 4-64 (false colored red) and Hag stained with a maleimide reactive dye (false colored green) at the indicated time points following induction with 0.3 mM IPTG. Carets indicate the first appearance of filaments. Scale bar is 1 micron.

Figure 6. Super-resolution microscopy of flagellar filaments and basal bodies

A) Super resolution fluorescence microscopy of strains of the indicated genotypes stained for membranes with FM4-64 (false colored red) and flagellar filaments (false colored green). The following strains were used to generate this panel: WT (DS1916), *swrA* (DS8022), *swrA*⁺⁺ (DS6543), *flhF* (DS9235), and *flhG* (DS9236). Scale bar is 1 micron. B) Super resolution fluorescence microscopy of strains of the indicated genotypes stained for membranes with FM4-64 (false colored red) and FliM-GFP (false colored green). The following strains were used to generate this panel: WT (DS8521), *swrA* (DS8600), *swrA*⁺⁺ (DS8608) where *swrA* was expressed from an artificial *P_{hyspank}* promoter and induced with 1 mM IPTG, *flhF* (DS8599), and *flhG* (DS8614). Scale bar is 0.5 micron. C) Graphical representation of flagellar number as determined by counting the number of FliM-GFP puncta per cell per cell length using Imaris image analysis software. Individual cell data presented as a scatter plot. Average of length and spot number indicated as a larger solid dot with standard deviations that emerge from the solid dot as horizontal and vertical lines. Best fit trend line indicated as a diagonal line. Black symbols indicate wild type (DS8521), red symbols indicate *swrA* (DS8600), blue symbols indicate *swrA*⁺⁺ (DS8608) where *swrA* was expressed from an artificial *P_{hyspank}* promoter and induced with 1 mM IPTG, and green symbols indicate *flhF* (DS8599). D) Graphical representation of flagellar asymmetry as determined by counting the number of FliM-GFP puncta on either side of the cells midpoint and dividing the larger number by the smaller number. Individual cell data presented as a scatter plot. Average of ratio of asymmetry presented as a larger solid dot with standard deviation that emerges from the solid dot as a horizontal line. A randomized data set was generated for each cell type and presented as an average ratio of asymmetry in a pale

larger dot of the same color with standard deviation that emerges from the pale dot as a horizontal line.

\$watermark-text

\$watermark-text

\$watermark-text

Figure 7. Statistical analysis of basal body puncta distribution

All data gathered by measuring the distance between FliM-GFP puncta and cell length using 3D SIM and Imaris image analysis software. A) Nearest neighbor distance between FliM-GFP puncta. Top, a cartoon depicting the interpretation of the data below. Bottom, the distances between all FliM-GFP puncta were calculated and a histogram of the minimum distances is presented as solid bars. Gray bars represent the remeasured data set for mathematically randomized FliM-GFP puncta. B) The Clark-Evens distribution of FliM-GFP puncta. Top, a cartoon depicting the interpretation of the data below. Bottom, solid bars represent histograms of the Clark-Evens distribution value for individual cells of the indicated genotype. Gray line indicates the expected histogram of Clark-Evens ratios for 1000 cells with FliM-GFP at randomized positions. C) The distance of FliM-GFP puncta from the midcell. Top, a cartoon depicting the interpretation of the data below. Bottom, solid bars represent histograms of the number of FliM-GFP puncta found at the indicated distance from the midcell. Gray dots and standard deviations indicate the histogram of FliM-GFP distribution of cells with FliM-GFP puncta at randomized positions.

Figure 8. FlhF and FlhG control flagellar distribution

Fluorescence microscopy of the indicated strains stained for membranes with FM4-64 (false colored red) and FliM-GFP (false colored green). Carets indicate a polar enrichment of FliM-GFP puncta. All strains were mutated for *fliM* at the native locus and expressed FliM-GFP inserted at an ectopic location (*amyE*). Scale bar is 1 micron. The following strains were used to generate this figure: A) wild type (DS8521), B) *flhF* (DS8599), C) *flhF* (*flhF*⁺) (DS8813), D) *flhG* (DS8614), E) *flhG* (*flhG*⁺) (DS8810), F) *flhG* (*flhFflhG*⁺) (DS9253), G) *flhFflhG* (DS9296), H) *flhF* (*P_{hyspank}-flhF*) (DS8811) and I) *flhFflhG* (*P_{hyspank}-flhFflhG*) (DK219).

Figure 9. The *flhF* mutant is defective in flagellar segregation

A) Fluorescence microscopy of wild type and *flhF* mutants grown to OD₆₀₀ 0.6, stained for flagella with Alexafluor 488 (false colored green), diluted 8-fold in growth media, regrown to OD₆₀₀ 0.6, and re-stained for flagella in Alexafluor 594 (false colored red) and stained for membranes with TMA DPH (false colored blue). Panels shown are the representative images of the three generation experiment but cells grown for one and two generations respectively were treated similarly. Scale bar is 10 μ m. B) For each generation and cell type, 600 cells with red flagella filaments were counted and then scored for the frequency of green flagella filaments. Note, cell chains of 4 cells or more often lacked flagella and were excluded from the analysis.

Table 1

Observed and randomized asymmetry ratios of basal body distribution

Genotype	Observed ¹		Randomized ²		P value ³	Conclusion ⁴
	Mean	Std dev	Mean	Std dev		
wild type	1.1910	0.1880	1.4450	0.4716	0.0005	Accept
(<i>swrA</i> ⁺⁺)	1.2321	0.1948	1.3241	0.3016	0.0455	Accept
$\Delta swrA$	1.7872	0.8979	2.0466	1.3399	0.000001	Accept
$\Delta flhF$	1.4110	0.3376	1.4446	0.4692	0.6390	Reject

¹ Asymmetry ratio means and standard deviations from observed data.² Asymmetry ratio means and standard deviations from randomized data.³ Student's T-test P value comparison between the randomized and observed data.⁴ Either accept or reject whether the randomized data set and the actual data are statistically significance based on a 0.05 P value cutoff.

Table 2

Strains

Strain	Genotype
3610	Wild type
DK15	<i>amyE::P_{fla/che}-fliM-GFP spec</i>
DK16	<i>ΔfliM ΔepsE epsH::tet amyE::P_{fla/che}-fliM-CFP spec thrC::P_{eps}-epsE-YFP mls sinR::kan</i>
DK29	<i>P_{fla/che}ΩP_{hy-spank}-fla/che operon kan</i>
DK30	<i>ΔflgE amyE::P_{fla/che}-flgE^{T123C} cat P_{fla/che}ΩP_{hy-spank}-fla/che operon kan</i>
DK31	<i>ΔfliM amyE::P_{fla/che}-fliM-GFP spec P_{fla/che}ΩP_{hy-spank}-fla/che operon kan</i>
DK62	<i>Δhag amyE::P_{hag}-hag^{T209C} spec P_{fla/che}ΩP_{hy-spank}-fla/che operon kan</i>
DK219	<i>ΔflhFflhG ΔfliM amyE::P_{fla/che}-fliM-GFP spec thrC::P_{hyspank}-flhFflhG mls</i>
DK415	<i>srfAA::mls epsA-O::tet ΔflhF</i>
DK416	<i>srfAA::mls epsA-O::tet ΔfliM amyE::P_{fla/che}-fliM spec</i>
DK417	<i>srfAA::mls epsA-O::tet</i>
DK429	<i>srfAA::mls epsA-O::tet ΔfliM amyE::P_{fla/che}-fliM-GFP spec</i>
DK430	<i>srfAA::mls epsA-O::tet ΔfliM</i>
DK431	<i>srfAA::mls epsA-O::tet ΔflhG</i>
DS1916	<i>amyE::P_{hag}-hag^{T209C} spec</i> (Blair et al., 2008)
DS6543	<i>thrC::P_{hyspank}-swrA mls amyE::P_{hag}-hag^{T209C} spec</i>
DS6658	<i>ΔflhF</i>
DS6775	<i>ΔfliM</i>
DS6799	<i>ΔfliM amyE::P_{fla/che}-fliM spec</i>
DS7358	<i>ΔflhG</i>
DS8022	<i>ΔswrA lacA::P_{hag}-hag^{T209C} mls</i>
DS8521	<i>ΔfliM amyE::P_{fla/che}-fliM-GFP spec</i>
DS8598	<i>ΔfliF ΔfliM amyE::P_{fla/che}-fliM-GFP spec</i>
DS8599	<i>ΔflhF ΔfliM amyE::P_{fla/che}-fliM-GFP spec</i>
DS8600	<i>ΔswrA ΔfliM amyE::P_{fla/che}-fliM-GFP spec</i>
DS8608	<i>ΔfliM thrC::P_{hyspank}-swrA mls amyE::P_{fla/che}-fliM-GFP spec</i>
DS8614	<i>ΔflhG ΔfliM amyE::P_{fla/che}-fliM-GFP spec</i>
DS8810	<i>ΔfliM ΔflhG amyE::P_{fla/che}-fliM-GFP spec thrC::P_{fla/che}-flhG mls</i>
DS8811	<i>ΔflhF ΔfliM amyE::P_{fla/che}-fliM-GFP spec thrC::P_{hyspank}-flhF mls</i>
DS8813	<i>ΔfliM ΔflhF amyE::P_{fla/che}-fliM-GFP spec thrC::P_{fla/che}-flhF mls</i>
DS8978	<i>ΔlytC ΔlytD ΔlytF ΔflhF ΔfliM amyE::P_{fla/che}-fliM-GFP spec</i>
DS8979	<i>ΔmotB ΔfliM amyE::P_{fla/che}-fliM-GFP spec</i>
DS8980	<i>ΔlytC ΔlytD ΔlytF ΔfliM amyE::P_{fla/che}-fliM-GFP spec</i>
DS8996	<i>ΔfliM ΔflgE amyE::P_{fla/che}-fliM-GFP spec thrC::P_{fla/che}-flgE^{T123C} mls</i>
DS9105	<i>sigD::tet ΔfliM amyE::P_{fla/che}-fliM-GFP spec</i>
DS9235	<i>ΔflhF lacA::P_{hag}-hag^{T209C} mls</i>

Strain	Genotype
DS9236	$\Delta flhG$ $lacA::P_{hag}-hag^{T209C}$ mls
DS9253	$\Delta fliM$ $\Delta flhG$ $amyE::P_{fla/che}-fliM-GFP$ $spec$ $lacA::P_{fla/che}-flhFflhG$ tet
DS9296	$\Delta fliM$ $\Delta flhFflhG$ $amyE::P_{fla/che}-fliM-GFP$ $spec$