Aflibercept in wet age-related macular degeneration: a perspective review

Matthew Ohr and Peter K. Kaiser

Ther Adv Chronic Dis

[2012] 3(4) 153–161

DOI: 10.1177/

2040622312446007

© The Author(s), 2012.

Reprints and permissions:
[http://www.sagepub.co.uk/
journalsPermissions.nav](http://www.sagepub.co.uk/journalsPermissions.nav)

Abstract: In the treatment of neovascular age-related macular degeneration (AMD), vascular endothelial growth factor (VEGF) has emerged as a key target of therapy. Currently, patients with neovascular AMD are treated with monthly intravitreal injections of anti-VEGF medications. Aflibercept is a novel recombinant fusion protein engineered to bind all isoforms of VEGF-A, VEGF-B, and placental growth factor. It is the latest medication to receive US Federal Drug Administration (FDA) approval for the treatment of neovascular AMD. Theoretical models suggest this molecule may have a longer duration of action compared with current treatments. The results of the VEGF Trap-Eye: Investigation of Efficacy and Safety in wet Age-related Macular Degeneration studies (VIEW 1 and VIEW 2) support this by demonstrating that aflibercept, dosed every 2 months after a monthly loading dose for 3 months, was noninferior in the proportion of patients who maintained or improved vision at 52 weeks compared with monthly injections of ranibizumab. These results were maintained over the 2 years of the studies. Aflibercept (Eylea; Regeneron Pharmaceuticals, Inc., Tarrytown, NY, USA and Bayer, Basel, Switzerland) was approved by the FDA for the treatment of neovascular AMD on 18 November 2011.

Keywords: aflibercept, neovascular age-related macular degeneration, ranibizumab, vascular endothelial growth factor, wet age-related macular degeneration

Introduction

In the USA, age-related macular degeneration (AMD) is the leading cause of vision loss in older patients. It is estimated that the prevalence of AMD is 6.5% among people aged 40 years and older [Klein *et al.* 2011]. AMD also remains a leading cause of vision loss among older adults in other Western countries. Most of this vision loss stems from advanced AMD. Advanced AMD can be classified into two major forms: the non-neovascular, atrophic (dry) form or the neovascular (wet) form. The majority of people with severe vision loss (20/200 or worse) from AMD have the neovascular form, which is estimated to occur in 10–20% of patients [Ferris *et al.* 1984]. Currently, there is no effective treatment for advanced, dry AMD [Meleth *et al.* 2011]. However, neovascular AMD has been successfully targeted by a number of treatment strategies.

Overview of current therapy

The hallmark of wet AMD is the formation of new, anomalous blood vessels that typically arise

from the choroidal vasculature and can grow into the subretinal pigment epithelial or subretinal space. Rarely, this process may originate from the retina and extend posteriorly into the subretinal space, a form of neovascular AMD termed retinal angiomatous proliferation. These neovascular vessels commonly hemorrhage and leak and can compromise vision by distorting the retinal and subretinal architecture with fluid, blood, or fibrovascular tissue [Spilisbury *et al.* 2000]. Untreated, choroidal neovascularization (CNV) usually leads to permanent loss of central vision.

The pathogenesis of CNV is not completely understood. However, the overexpression of vascular endothelial growth factor (VEGF), a pro-angiogenic cytokine, has been shown to play a crucial role [Spilisbury *et al.* 2000]. Previous studies have demonstrated increased levels of VEGF in the presence of inflammatory cytokines, suggesting that inflammation is a key component of AMD [Nagineni *et al.* 2012]. Others have suggested that ischemia, also associated with increased VEGF [Witmer *et al.* 2003], may play a role in AMD

Correspondence to:
Peter K. Kaiser, MD
Cleveland Clinic – Cole
Eye Institute, 9500
Euclid Avenue, Desk i3,
Cleveland, OH 44122, USA
pkkaiser@mac.com

Matthew Ohr, MD
Cleveland Clinic – Cole
Eye Institute, Cleveland,
OH, USA

[Feigl, 2009]. All of these reports clearly indicate that VEGF is vital to the pathogenesis of CNV in AMD.

Early treatment strategies focused on destruction of choroidal neovascular membranes using laser photocoagulation. The Macular Photocoagulation Study (MPS) established guidelines for treatment of these lesions [Macular Photocoagulation Study Group, 1982, 1986, 1991]. Although this treatment reduced the likelihood of severe vision loss compared with the natural course of the disease, there were many limitations, especially when treating lesions in the fovea. The primary downsides were related to the fact that the laser induced a permanent scotoma, and recurrence of the CNV occurred in over 50% of treated eyes [Macular Photocoagulation Study Group, 1991].

Until 1999, laser photocoagulation was the only treatment for neovascular AMD that had been shown to reduce the risk of vision loss. At that time, the Treatment of Age-Related Macular Degeneration with Photodynamic Therapy (TAP) Study reported that photodynamic therapy (PDT) with verteporfin (Visudyne; Novartis Pharma AG, Basel, Switzerland) reduced the risk of moderate to severe vision loss for at least 5 years in patients who presented with subfoveal lesions classified as predominantly classic [TAP Study Group, 1999; Azab *et al.* 2004; Blumenkranz *et al.* 2002; Bressler and TAP Study Group, 2001; Bressler *et al.* 2002; Kaiser *et al.* 2006]. PDT is a two-step process that involves the intravenous injection of verteporfin, a photosensitizing molecule, which is taken up by dividing cells within CNV. The drug is activated by local application of energy from a diode laser source at a wavelength that corresponds to an absorption peak of the molecule. A photochemical reaction occurs and activated free radicals are generated that can lead to capillary endothelial cell damage and vessel thrombosis. At 2 years, 59% of verteporfin treated eyes *versus* 31% of placebo eyes avoided at least moderate vision loss [Bressler and TAP Study Group, 2001]. While PDT improved the results seen with laser photocoagulation, there remained a pressing need for better treatment modalities.

The first VEGF inhibitor to obtain US Federal Drug Administration (FDA) approval for CNV in AMD was pegaptanib (Macugen; OSI/Eyetech Pharmaceuticals, New York, NY, USA) in 2004. Pegaptanib is an RNA aptamer that binds human VEGF₁₆₅ with high affinity and

specificity [Gragoudas *et al.* 2004]. The drug, however, did not bind other active VEGF isoforms such as VEGF₁₂₁. Pegaptanib is administered as an intravitreal injection every 6 weeks. The VEGF Inhibition Study in Ocular Neovascularization (VISION) trial was a prospective, randomized, double-masked, controlled, dose-ranging phase III clinical trial in which 1186 patients with AMD and subfoveal CNV received one of three doses of pegaptanib or sham injections every 6 weeks for 48 weeks [Gragoudas *et al.* 2004]. The results of this study were promising, with 70% of patients losing less than three lines of vision compared with 55% of controls ($p < 0.001$). Unfortunately, similar to the results with PDT, a minority of patients gained vision with this therapy.

One of the most exciting advances in the treatment of CNV in AMD came with the introduction of ranibizumab (Lucentis; Genentech, South San Francisco, CA, USA) in 2006. Ranibizumab is a recombinantly produced, humanized, antibody (Fab) fragment that binds VEGF [Rosenfeld *et al.* 2006]. Unlike pegaptanib, ranibizumab binds to and inhibits the biological activity of all active forms of VEGF-A. The Minimally Classic/Occult Trial of the Anti-VEGF Antibody Ranibizumab in the Treatment of Neovascular AMD (MARINA) study was a randomized, double-masked, sham-controlled clinical trial of 716 patients with minimally classic or occult CNV secondary to AMD treated with one of two different doses of intravitreal ranibizumab or sham injections given every 4 weeks for 2 years [Rosenfeld *et al.* 2006]. The results of this study were revolutionary with 94.5% of patients treated with ranibizumab 0.3 mg and 94.6% of patients treated with ranibizumab 0.5 mg experiencing vision stabilization or improvement compared with 62.2% of patients receiving sham injections ($p < 0.001$). In fact, visual acuity improved by 15 letters or more in 24.8% of patients receiving 0.3 mg and 33.8% of patients receiving 0.5 mg ranibizumab compared with 5.0% of the sham injection group ($p < 0.001$). These results were further supported by the Anti-VEGF Antibody for the Treatment of Predominantly Classic Choroidal Neovascularization in AMD (ANCHOR) study, which was a randomized, controlled, double-masked phase III clinical trial of 423 patients that compared patients treated with ranibizumab with patients treated with PDT with verteporfin treatment [Brown *et al.* 2006, 2009]. These results showed that 94.3% of patients treated with

0.3 mg ranibizumab and 96.4% of patients treated with 0.5 mg ranibizumab lost less than 15 letters of vision compared with 64.3% of patients treated with PDT at 1 year ($p < 0.001$). Patients receiving ranibizumab again showed increased vision in 35.7% of patients treated with 0.3 mg ranibizumab and 40.3% of patients treated with 0.5 mg ranibizumab compared with only 5.6% of patients treated with PDT ($p < 0.001$). The results of these trials resulted in anti-VEGF therapies largely replacing previous treatment modalities.

Bevacizumab (Avastin; Genentech) is a full-length monoclonal antibody that binds all isoforms of VEGF-A. The FDA originally approved it in February 2004 for the treatment of metastatic colorectal cancer. Soon thereafter, physicians started to use it off label as an intravenous or intravitreal treatment for neovascular AMD. Despite the lack of clinical research to support its safety or efficacy, anecdotal evidence led to its widespread popularity prior to the approval of ranibizumab in 2006. To deliver an intravitreal injection, a physician or pharmacy takes a vial of bevacizumab and makes numerous unit doses. This dramatically lowers the cost of the drug. The approximate cost differential between ranibizumab (US\$2000) and bevacizumab (US\$50) was prohibitive for some patients, and bevacizumab continued to be utilized, despite not being FDA approved for intravitreal use. In fact, in a review of Medicare claims for neovascular AMD in 2008, it was noted that 58% of all intravitreal injections given were bevacizumab and 41% were ranibizumab [Brechtner *et al.* 2011].

To address the safety and therapeutic concerns of the widespread, off-label use of bevacizumab in the treatment of wet AMD, the National Eye Institute commissioned the Comparison of Age-Related Macular Degeneration Treatment Trial (CATT) [CATT Research Group *et al.* 2011]. In this multicenter, single-blind, noninferiority trial, 1208 patients with neovascular AMD were randomized into four groups. After the first mandatory intravitreal injection, patients received ranibizumab every 28 days (ranibizumab monthly), bevacizumab every 28 days (bevacizumab monthly), ranibizumab only when signs of active neovascularization were present (ranibizumab as needed), and bevacizumab only when signs of active neovascularization were present (bevacizumab as needed). The 1-year results of this study demonstrated that monthly bevacizumab was equivalent to monthly ranibizumab with 8.0 and 8.5 letters

gained, respectively. Bevacizumab as needed was found to be equivalent to ranibizumab as needed with 5.9 and 6.8 letters gained, respectively. While ranibizumab as needed was found to be equivalent to ranibizumab monthly, the equivalence of bevacizumab as needed compared with bevacizumab monthly was found to be inconclusive.

Aflibercept

Aflibercept (Eylea; Regeneron, Tarrytown, NY, USA and Bayer, Basel, Switzerland) is a fully human, recombinant fusion protein composed of the second immunoglobulin (Ig) binding domain of VEGF receptor 1 and the third Ig binding domain of VEGF receptor 2, fused to the Fc region of human IgG1. It binds to all VEGF-A isoforms, VEGF-B, and placental growth factor (PlGF) [Papadopoulos *et al.* 2012]. Aflibercept is a member of Regeneron's proprietary family of 'Trap' products that catch, hold, and block (i.e. trap) certain cytokines [Adis R&D Profile, 2008]. Aflibercept is being developed for the treatment of cancer (Zaltrap; Regeneron and Sanofi, Bridgewater, NJ, USA) and eye disorders. The eye formulation, also referred to in the literature as VEGF Trap-Eye, is identical in structure to the intravenous cancer treatment, with further purification steps and buffer modification to allow for comfortable, nonirritating intravitreal injection [Dixon *et al.* 2009].

Unlike currently available anti-VEGF therapies, aflibercept binds PlGF in addition to all isoforms of VEGF-A and VEGF-B. Like VEGF, PlGF is present in human CNV membranes, and animal studies have shown that PlGF contributes to the development of experimental CNV [Rakic *et al.* 2003]. Another differentiating feature of aflibercept is that the binding affinity for VEGF is 0.5 pM K_d, which is considerably stronger than ranibizumab, bevacizumab, or native VEGF receptors. This allows for effective blocking of VEGF, even at low concentrations, which may translate into a longer duration of action and extended dosing intervals [Stewart and Rosenfeld, 2008].

The results of preclinical studies were promising. In Matrigel-induced models of CNV in rats, aflibercept was shown to arrest the growth of CNV and led to the regression of recently established lesions [Cao *et al.* 2010]. Primate studies of laser-induced CNV also showed promise for the drug. When aflibercept was given prior to and following

attempted laser induction of CNV, minimal neovascularization was noted compared with placebo. In drug-naïve eyes with previously established CNV, aflibercept was successful in causing regression of the CNV and resolving vascular leakage [Nork *et al.* 2011]. These encouraging results coupled with the apparent safety of the drug, fueled the demand for human clinical trials.

Phase I

A phase I, randomized, multicenter, masked, placebo-controlled clinical trial of *intravenous* aflibercept in patients with subfoveal CNV from AMD showed a dose-dependent decrease in retinal thickness [Nguyen *et al.* 2006]. However, at systemic doses of 3 mg/kg, hypertension and proteinuria were observed, and the study was halted for safety concerns. This led to investigation of alternative delivery methods.

The safety, tolerability, maximum tolerated dose, and bioactivity of intravitreal injection of aflibercept were evaluated in a phase I, multicenter, dose-escalation study [Nguyen *et al.* 2009]. In the study, 21 patients received a single dose of aflibercept. Patients were monitored for 12 weeks after injection. There were no serious ocular or systemic events noted. With any dose of aflibercept, stable or improved vision was seen in 95% of patients at 6 weeks. The mean decrease in foveal thickness was $-105.5\ \mu\text{m}$, and the mean increase in visual acuity was $+4.43$ letters. Half of the patients receiving 2 or 4 mg doses showed no retinal leakage and maintained vision gains at 12 weeks after a single injection. These positive results paved the way for further development of an intravitreal formulation of aflibercept.

Phase II

The clinical evaluation of anti-angiogenesis in the retina study (CLEAR-IT) 2 trial was a phase II multicenter, prospective, randomized, double-masked clinical trial designed to study the effect of intravitreal aflibercept in patients with neovascular AMD [Brown *et al.* 2011; Heier *et al.* 2011]. This trial was divided into two parts. In the first part, patients were treated with a fixed dosing interval up to 12 weeks. The second part of the study was designed to be as needed (PRN) dosing and took place from week 16 to 52. The primary endpoint of the study was the change in central retinal thickness. The mean change in best corrected visual acuity (BCVA) was evaluated as a

secondary outcome. The study included 159 patients who were randomized into five treatment groups. The first two groups received treatment every 4 weeks and were dosed at 0.5 mg (group 1) or 2 mg (group 2). The last three groups were treated every 12 weeks and were dosed at 0.5 mg (group 3), 2 mg (group 4), or 4 mg (group 5). The primary outcome was at 12 weeks, following the fixed dosing period. The mean decrease in central retinal thickness from baseline to 12 weeks in all groups was $-119\ \mu\text{m}$. Monthly dosing with either 0.5 or 2 mg (groups 1 and 2) provided a more profound and consistent effect than any of the groups treated every 12 weeks. Overall, there was a mean increase in BCVA of $+5.7$ Early Treatment Diabetic Retinopathy Study (ETDRS) letters in all groups. The greatest mean increase in BCVA, more than $+8$ letters, was seen in the monthly dosing groups compared with the patients receiving only one injection [Brown *et al.* 2011].

For the PRN dosing arm of the study, patients were evaluated every 4 weeks to determine the need for continued treatment. Patients received an injection of the baseline dose at week 12. At week 16 and thereafter, eyes were reinjected with aflibercept if any of the following conditions were noted: increase in central retinal thickness of at least $100\ \mu\text{m}$ by optical coherence tomography (OCT); loss of at least 5 ETDRS letters with recurrent fluid on OCT; persistent fluid on OCT; new-onset of classic CNV; new or persistent leak on fluorescein angiography; or new macular hemorrhage on clinical examination. Using these criteria, the mean decrease in central retinal thickness in all groups from baseline to 52 weeks was $-130\ \mu\text{m}$. The mean increase in BCVA was $+5.3$ ETDRS letters in all groups. The greatest increase in BCVA occurred in the group initially treated with 2 mg every 4 weeks for 12 weeks before PRN dosing with a mean increase of $+9.0$ letters at 1 year. To achieve these excellent visual gains, an average of two additional injections was administered after the 12-week fixed-dosing phase across all groups. The mean time to the first reinjection was 129 days, with 19% of patients receiving no injections and 45% receiving one or two additional injections [Heier *et al.* 2011].

Phase III

Two parallel, phase III, double-masked, randomized studies were initiated in August 2007. The VEGF Trap-Eye: Investigation of Efficacy and Safety in wet Age-Related Macular Degeneration

Table 1. Summary of the 1-year results of VIEW 1 and VIEW 2 studies compared with MARINA and CATT trials.

	MARINA		CATT		VIEW 1 (12 months)				VIEW 2 (12 months)			
	Ran 0.5 mg	Sham	Ran 0.5 mg	Bev 0.5 mg	Afl 0.5 mg	Afl 2 mg	Afl 2 mg Q*	Ran 0.5 mg	Afl 0.5 mg	Afl 2 mg	Afl 2 mg Q*	Ran 0.5 mg
≥15 letter gain (%)	33	4	34	31								
Stable vision (%)	90	53	94	94	96	95	95	94	96	96	96	94
Mean gain in VA from baseline at 12 months	+7.2	-10.4	+8.5	+8.0	+8.1	+10.9	+7.9	+8.1	+9.7	+7.6	+8.9	+9.4
Number of injections	12		12	12								

*Dosed every 8 weeks after treatment initiation with 3 monthly doses.

Afl, aflibercept; Bev, bevacizumab; CATT, Comparison of Age-Related Macular Degeneration Treatment Trial; MARINA, Minimally Classic/Occult Trial of the Anti-VEGF Antibody Ranibizumab in the Treatment of Neovascular AMD; Ran, ranibizumab; VA, visual acuity; VIEW, VEGF Trap-Eye: Investigation of Efficacy and Safety in Wet Age-related Macular Degeneration.

(VIEW 1) study was performed in North America. The VIEW 2 study was an international study including Europe, Asia Pacific, Japan, and Latin America. The studies were designed as noninferiority studies comparing intravitreal aflibercept with ranibizumab. Patients with subfoveal CNV due to AMD were randomized into four groups. The first two groups received intravitreal injections of aflibercept at doses of 0.5 and 2 mg administered at 4-week intervals. The third group received 2 mg of aflibercept at 8-week intervals following three loading doses given every 4 weeks. These were compared with the fourth group, the control, receiving 0.5 mg of ranibizumab administered every 4 weeks. The primary endpoint was statistical noninferiority in the proportion of patients who maintained or improved vision over 52 weeks compared with ranibizumab.

The 1-year results of the VIEW 1 study showed that vision was maintained, defined as losing fewer than 15 ETDRS letters, in 96% of patients receiving aflibercept 0.5 mg monthly, 95% of patients receiving 2 mg monthly, and 95% of patients receiving 2 mg every 2 months. These results compared favorably with the 94% of patients maintaining vision in the group receiving ranibizumab 0.5 mg monthly [Regeneron, 2010; Heier, 2011]. The patients receiving aflibercept 2 mg monthly on average gained 10.9 letters compared with a mean 8.1 letter gain with ranibizumab 0.5 mg dosed every month ($p < 0.01$). The VIEW 2 study showed similar results, with maintenance of vision in 96% of patients receiving 0.5 mg monthly, 96% of patients receiving 2 mg monthly, and 96% of patients receiving 2 mg every 2 months. These results also compared

favorably with the 94% of patients maintaining vision in the group treated with ranibizumab 0.5 mg monthly [Schmidt-Erfurth, 2011]. They are similar to results found in the MARINA and CATT trial (Table 1). The safety of both VIEW 1 and VIEW 2 studies was excellent with no difference seen between any aflibercept group and the ranibizumab group. The fact that 2 mg aflibercept dosed every 8 weeks after three loading doses was noninferior to ranibizumab dosed every 4 weeks in terms of safety and efficacy is exciting, as it offers the hope of similar visual gains with less treatment burden.

The 2-year results of the VIEW 1 and VIEW 2 studies were recently released [Regeneron, 2011]. The integrated analysis of these two studies (Table 2) shows that patients receiving aflibercept 2 mg every 8 weeks gained +7.6 letters from baseline at week 96 compared with +8.4 letters at week 52. The visual acuity gain in from baseline in patients receiving monthly ranibizumab was +7.9 letters at week 96 compared with +8.7 letters at week 52. Patients receiving aflibercept 2 mg every 8 weeks received an average of 11.2 injections over 2 years while patients treated with ranibizumab had an average of 16.5 injections over 2 years. Aflibercept (Eylea) was approved by the FDA for the treatment of wet AMD on 18 November 2011.

Conclusions

The evolution of treatment strategies for neovascular AMD has resulted in a paradigm shift in terms of expectations among patients and physicians. Prior to these recent advances, patients

Table 2. Summary of combined 2-year results of VIEW 1 and VIEW 2 studies.

	Aflibercept	Ranibizumab
Study arm	2 mg Q*	0.5 mg
Stabilization of VA (%)	95	94
Mean gain in VA (ETDRS letters)		
Year 1	+8.4	+8.7
Year 2	+7.6	+7.9
Average number of injections over 24 months	11.2	16.5
*Dosed every 8 weeks after treatment initiation with 3 monthly doses. ETDRS, Early Treatment Diabetic Retinopathy Study; VA, visual acuity; VIEW, VEGF Trap-Eye: Investigation of Efficacy and Safety in Wet Age-related Macular Degeneration.		

who developed neovascular changes could anticipate a progressive and permanent decrease in vision. While destruction of the CNV lesion with laser photocoagulation was possible, in particular when the lesion was located outside of the visual axis, it offered only modest hope of maintaining vision compared with the natural history of the disease, and results were less than satisfactory. The introduction of PDT with verteporfin offered some improvement, especially to patients with subfoveal disease. However, many patients continued to lose vision, and only minimal visual gains were achieved.

The advent of anti-VEGF therapy marked a turning point in the treatment of neovascular AMD. The first FDA-approved anti-VEGF therapy for neovascular AMD was pegaptanib. Unfortunately, the specific targeting of VEGF₁₆₅ seemed to limit its effectiveness, and as with PDT, patients generally continued to slowly lose vision. It was the phase III results of ranibizumab and the off-label intravitreal use of the cancer drug, bevacizumab, that began to change expectations. Anecdotal evidence of improvements in vision and retinal thickness after a single treatment were reported in patients whose condition had failed to respond to pegaptanib therapy [Rosenfeld *et al.* 2005]. With the publication of the ANCHOR and MARINA trials, and the subsequent FDA approval of ranibizumab, vision could be expected to stabilize in close to 95% of cases, with improvement of BCVA by three or more lines in approximately 40% of patients.

While anti-VEGF therapy has changed the efficacy of treatment, it is not without drawbacks. Patients are subjected to intravitreal injections as often as every 4 weeks to maintain vision. Frequent office visits, testing, and medication costs represent a burden to patients, physicians, and society. Although there have been no proven, adverse systemic effects from intravitreal injections, every injection puts patients at risk for endophthalmitis, intraocular inflammation, vitreous hemorrhage, retinal tear, retinal detachment, and iatrogenic cataract. Recent studies have also suggested a sustained rise in intraocular pressure may occur with serial injections of anti-VEGF agents [Tseng *et al.* 2011]. Aside from serious complications, patients are commonly subject to anxiety, discomfort, and the undesirable aesthetics of conjunctival hyperemia or hemorrhage.

Current efforts have focused on extending the benefits of anti-VEGF treatment with less frequent dosing. In the phase IIb, multi-center, randomized, double-masked, sham injection-controlled study of efficacy and safety of ranibizumab in subjects with subfoveal CNV with or without Classic CNV secondary to AMD (PIER) study, patients received monthly injections of ranibizumab for 3 months followed by quarterly dosing. Gains in visual acuity were noted at 3 months, only to be lost during the quarterly dosing phase of the trial [Regillo *et al.* 2008]. In the prospective optical coherence tomography OCT imaging of patients with neovascular age-related macular degeneration treated with intra-ocular lucentis (PrONTO) study patients also received monthly injections of ranibizumab for 3 months. Following the initial dosing, additional treatments were given on a PRN basis. After 2 years, 78% of patients maintained vision and 43% of patients showed improvement in more than three lines of vision. An average of 9.9 injections was given over the 2-year period [Lalwani *et al.* 2009].

One of the criticisms of PRN dosing is the fact that monthly visits are still required. One approach to treatment that aims to reduce the number of injections and visits is the 'inject and extend' method [Spaide, 2007]. This strategy involves treatment with 3-monthly injections followed by a follow up appointment extended to 6 weeks. At the follow-up visit OCT and biomicroscopy are performed. If edema or any other sign of exudation is present, the patient is given an injection and told to follow up in 4 weeks. Patients without any evidence of exudation are injected and have

their follow-up visit extended to 8 weeks. The same evaluation occurs at the 8-week visit. However, patients with signs of exudation are injected and scheduled to follow up at 6 weeks. Patients without evidence of exudation are further extended to follow up at 10 weeks. In this way, an optimal, individualized treatment strategy can be obtained. While a few small studies have reported positive results with this method, there have been no large, prospective studies to support its effectiveness [Engelbert *et al.* 2009; Oubraham *et al.* 2011].

Twelve-month results from the VIEW 1 and VIEW 2 studies appear promising for aflibercept. Although the data will have to be analyzed further, the apparent noninferiority of the various aflibercept dosing regimens compared with ranibizumab represents a new milestone in the treatment of CNV due to AMD. Of particular interest is the 2 mg every 8 weeks dosing interval studied in the trials. The recent results of the 2-year data for the VIEW 1 and VIEW 2 studies have further demonstrated the ability of aflibercept to maintain the visual gains attained in the first year of the study with a significantly smaller number of injections compared with ranibizumab. Aflibercept was recently approved by the FDA for the treatment of CNV in AMD. The wholesale price of aflibercept (US\$1850) is slightly below that of ranibizumab. However, the reduced treatment requirements of every 8-week dosing *versus* monthly dosing of ranibizumab may represent a substantial savings in cost and treatment burden to patients.

Funding

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Conflict of interest statement

Dr Kaiser is a consultant for Regeneron and Bayer that has been disclosed and approved by the Cleveland Clinic Conflict of Interest Committee. In addition, the Cole Eye Institute, the employer of Drs Ohr and Kaiser has received research grant support from Regeneron.

References

Adis R&D Profile (2008) Aflibercept: AVE 0005, AVE 005, AVE0005, VEGF Trap – Regeneron, VEGF Trap (R1R2), VEGF Trap-Eye. *Drugs R D* 9: 261–269.

Azab, M., Benchaboune, M., Blinder, K.J., Bressler, N.M., Bressler, S.B., Gragoudas, E.S. *et al.* (2004) Verteporfin therapy of subfoveal choroidal neovascularization in age-related macular degeneration: meta-analysis of 2-year safety results in three randomized clinical trials: Treatment of Age-Related Macular Degeneration With Photodynamic Therapy and Verteporfin in Photodynamic Therapy Study report no. 4. *Retina* 24: 1–12.

Blumenkranz, M.S., Bressler, N.M., Bressler, S.B., Donati, G., Fish, G.E., Haynes, L.A. *et al.* (2002) Verteporfin therapy for subfoveal choroidal neovascularization in age-related macular degeneration: three-year results of an open-label extension of 2 randomized clinical trials – TAP report no. 5. *Arch Ophthalmol* 120: 1307–1314.

Brechner, R.J., Rosenfeld, P.J., Babish, J.D. and Caplan, S. (2011) Pharmacotherapy for neovascular age-related macular degeneration: an analysis of the 100% 2008 Medicare fee-for-service part B claims file. *Am J Ophthalmol* 151: 887–895.e1.

Bressler, N.M., Arnold, J., Benchaboune, M., Blumenkranz, M.S., Fish, G.E., Gragoudas, E.S. *et al.* (2002) Verteporfin therapy of subfoveal choroidal neovascularization in patients with age-related macular degeneration: additional information regarding baseline lesion composition's impact on vision outcomes – TAP report no. 3. *Arch Ophthalmol* 120: 1443–1454.

Bressler, N.M. and Treatment of Age-Related Macular Degeneration with Photodynamic Therapy (TAP) Study Group (2001) Photodynamic therapy of subfoveal choroidal neovascularization in age-related macular degeneration with verteporfin: two-year results of 2 randomized clinical trials – TAP report 2. *Arch Ophthalmol* 119: 198–207.

Brown, D.M., Heier, J.S., Ciulla, T., Benz, M., Abraham, P., Yancopoulos, G. *et al.* (2011) Primary endpoint results of a phase II study of vascular endothelial growth factor trap-eye in wet age-related macular degeneration. *Ophthalmology* 118: 1089–1097.

Brown, D.M., Kaiser, P.K., Michels, M., Soubrane, G., Heier, J.S., Kim, R.Y. *et al.* (2006) Ranibizumab versus verteporfin for neovascular age-related macular degeneration. *N Engl J Med* 355: 1432–1444.

Brown, D.M., Michels, M., Kaiser, P.K., Heier, J.S., Sy, J.P., Ianchulev, T. *et al.* (2009) Ranibizumab versus verteporfin photodynamic therapy for neovascular age-related macular degeneration: two-year results of the ANCHOR study. *Ophthalmology* 116: 57–65.e5.

Cao, J., Zhao, L., Li, Y., Liu, Y., Xiao, W., Song, Y. *et al.* (2010) A subretinal matrigel rat choroidal neovascularization (CNV) model and inhibition of CNV and associated inflammation

- and fibrosis by VEGF trap. *Investigative Ophthalmol Vis Sci* 51: 6009–6017.
- CATT Research Group, Martin, D.F., Maguire, M.G., Ying, G.S., Grunwald, J.E., Fine, S.L., Jaffe, G.J. *et al.* (2011) Ranibizumab and bevacizumab for neovascular age-related macular degeneration. *N Engl J Med* 364: 1897–1908.
- Dixon, J.A., Oliver, S.C., Olson, J.L. and Mandava, N. (2009) VEGF Trap-Eye for the treatment of neovascular age-related macular degeneration. *Exp Opin Invest Drugs* 18: 1573–1580.
- Engelbert, M., Zweifel, S.A. and Freund, K.B. (2009) ‘Treat and extend’ dosing of intravitreal anti-vascular endothelial growth factor therapy for type 3 neovascularization/retinal angiomatous proliferation. *Retina* 29: 1424–1431.
- Feigl, B. (2009) Age-related maculopathy – linking aetiology and pathophysiological changes to the ischaemia hypothesis. *Prog Retin Eye Res* 28: 63–86.
- Ferris, F.L., 3rd, Fine, S.L. and Hyman, L. (1984) Age-related macular degeneration and blindness due to neovascular maculopathy. *Arch Ophthalmol* 102: 1640–1642.
- Gragoudas, E.S., Adamis, A.P., Cunningham, E.T., Jr, Feinsod, M., Guyer, D.R. and VEGF Inhibition Study in Ocular Neovascularization Clinical Trial Group (2004) Pegaptanib for neovascular age-related macular degeneration. *N Engl J Med* 351: 2805–2816.
- Heier, J.S. (2011). *VEGF Trap-Eye Phase III Trial Results. VIEW 1 results*. Presented at: Angiogenesis, Exudation, and Degeneration, February 12, 2011; Miami FL.
- Heier, J.S., Boyer, D., Nguyen, Q.D., Marcus, D., Roth, D.B., Yancopoulos, G. *et al.* (2011) The 1-year results of CLEAR-IT 2, a phase 2 study of vascular endothelial growth factor trap-eye dosed as-needed after 12-week fixed dosing. *Ophthalmology* 118: 1098–1106.
- Kaiser, P.K. and Treatment of Age-Related Macular Degeneration with Photodynamic Therapy (TAP) Study Group (2006) Verteporfin therapy of subfoveal choroidal neovascularization in age-related macular degeneration: 5-year results of two randomized clinical trials with an open-label extension: TAP report no. 8. *Graefes Arch Clin Exp Ophthalmol* 244: 1132–1142.
- Klein, R., Chou, C.F., Klein, B.E., Zhang, X., Meuer, S.M. and Saaddine, J.B. (2011) Prevalence of age-related macular degeneration in the US population. *Arch Ophthalmol* 129: 75–80.
- Lalwani, G.A., Rosenfeld, P.J., Fung, A.E., Dubovy, S.R., Michels, S., Feuer, W. *et al.* (2009) A variable-dosing regimen with intravitreal ranibizumab for neovascular age-related macular degeneration: year 2 of the PrONTO Study. *Am J Ophthalmol* 148: 43–58.e1.
- Macular Photocoagulation Study Group (1982) Argon laser photocoagulation for senile macular degeneration. Results of a randomized clinical trial. *Arch Ophthalmol* 100: 912–918.
- Macular Photocoagulation Study Group (1986) Argon laser photocoagulation for neovascular maculopathy. Three-year results from randomized clinical trials. *Arch Ophthalmol* 104: 694–701.
- Macular Photocoagulation Study Group (1991) Argon laser photocoagulation for neovascular maculopathy. Five-year results from randomized clinical trials. *Arch Ophthalmol* 109: 1109–1114.
- Meleth, A.D., Wong, W.T. and Chew, E.Y. (2011) Treatment for atrophic macular degeneration. *Curr Opin Ophthalmol* 22(3): 190–193.
- Nagineni, C.N., Kommineni, V.K., William, A., Detrick, B. and Hooks, J.J. (2012) Regulation of VEGF expression in human retinal cells by cytokines: implications for the role of inflammation in age-related macular degeneration. *J Cell Physiol* 227: 116–126.
- Nguyen, Q.D., Shah, S.M., Browning, D.J., Hudson, H., Sonkin, P., Hariprasad, S.M. *et al.* (2009) A phase I study of intravitreal vascular endothelial growth factor trap-eye in patients with neovascular age-related macular degeneration. *Ophthalmology* 116: 2141–2148.e1.
- Nguyen, Q.D., Shah, S.M., Hafiz, G., Quinlan, E., Sung, J., Chu, K. *et al.* (2006) A phase I trial of an IV-administered vascular endothelial growth factor trap for treatment in patients with choroidal neovascularization due to age-related macular degeneration. *Ophthalmology* 113: 1522.e1–1522.e14.
- Nork, T.M., Dubielzig, R.R., Christian, B.J., Miller, P.E., Miller, J.M., Cao, J. *et al.* (2011) Prevention of experimental choroidal neovascularization and resolution of active lesions by VEGF trap in nonhuman primates. *Arch Ophthalmol* 129: 1042–1052.
- Oubraham, H., Cohen, S.Y., Samimi, S., Marotte, D., Bouzaher, I., Bonicel, P. *et al.* (2011) Inject and extend dosing versus dosing as needed: a comparative retrospective study of ranibizumab in exudative age-related macular degeneration. *Retina* 31: 26–30.
- Papadopoulos, N., Martin, J., Ruan, Q., Rafique, A., Rosconi, M.P., Shi, E. *et al.* (2012) Binding and neutralization of vascular endothelial growth factor (VEGF) and related ligands by VEGF Trap, ranibizumab and bevacizumab. *Angiogenesis* 3 February [Epub ahead of print].
- Rakic, J.M., Lambert, V., Devy, L., Luttun, A., Carmeliet, P., Claes, C. *et al.* (2003) Placental growth factor, a member of the VEGF family, contributes

to the development of choroidal neovascularization. *Invest Ophthalmol Vis Sci* 44: 3186–3193.

Regeneron (2010) Bayer and Regeneron report positive top-line results of two phase 3 studies with VEGF Trap-Eye in wet age-related macular degeneration (press release). Available at: <http://investor.regeneron.com/releasedetail.cfm?ReleaseID=532099> (accessed February 2012).

Regeneron (2011) Two year results of phase 3 studies with EYLEA™ (aflibercept) injection in wet AMD show sustained improvement in visual acuity (press release). Available at: <http://investor.regeneron.com/releasedetail.cfm?ReleaseID=629800> (accessed February 2012).

Regillo, C.D., Brown, D.M., Abraham, P., Yue, H., Ianchulev, T., Schneider, S. *et al.* (2008) Randomized, double-masked, sham-controlled trial of ranibizumab for neovascular age-related macular degeneration: PIER Study year 1. *Am J Ophthalmol* 145: 239–248.

Rosenfeld, P.J., Brown, D.M., Heier, J.S., Boyer, D.S., Kaiser, P.K., Chung, C.Y. *et al.* (2006) Ranibizumab for neovascular age-related macular degeneration. *N Engl J Med* 355: 1419–1431.

Rosenfeld, P.J., Moshfeghi, A.A. and Puliafito, C.A. (2005) Optical coherence tomography findings after an intravitreal injection of bevacizumab (avastin) for neovascular age-related macular degeneration. *Ophthalm Surg Lasers Imaging* 36: 331–335.

Schmidt-Erfurth, U. (2011). *VEGF Trap-Eye Phase III Trial Results. VIEW 2 results*. Presented at:

Angiogenesis, Exudation, and Degeneration, February 12, 2011; Miami FL.

Spaide, R. (2007) Ranibizumab according to need: a treatment for age-related macular degeneration. *Am J Ophthalmol* 143: 679–680.

Spilsbury, K., Garrett, K.L., Shen, W.Y., Constable, I.J. and Rakoczy, P.E. (2000) Overexpression of vascular endothelial growth factor (VEGF) in the retinal pigment epithelium leads to the development of choroidal neovascularization. *Am J Pathol* 157: 135–144.

Stewart, M.W. and Rosenfeld, P.J. (2008) Predicted biological activity of intravitreal VEGF Trap. *Br J Ophthalmol* 92: 667–668.

Treatment of Age-related Macular Degeneration with Photodynamic Therapy (TAP) Study Group (1999) Photodynamic therapy of subfoveal choroidal neovascularization in age-related macular degeneration with verteporfin: one-year results of 2 randomized clinical trials–TAP report. *Arch Ophthalmol* 117: 1329–1345.

Tseng, J.J., Vance, S.K., Della Torre, K.E., Mendonca, L.S., Cooney, M.J., Klancnik, J.M. *et al.* (2011) Sustained increased intraocular pressure related to intravitreal antivascular endothelial growth factor therapy for neovascular age-related macular degeneration. *J Glaucoma* 21: 241–247.

Witmer, A.N., Vrensen, G.F., Van Noorden, C.J. and Schlingemann, R.O. (2003) Vascular endothelial growth factors and angiogenesis in eye disease. *Prog Retin Eye Res* 22: 1–29.

Visit SAGE journals online
<http://taj.sagepub.com>

SAGE journals