Published in final edited form as:

Ann Biomed Eng. 2013 February ; 41(2): 327–337. doi:10.1007/s10439-012-0653-x.

Analysis of Kinetics of Dihydroethidium Fluorescence with Superoxide Using Xanthine Oxidase and Hypoxanthine Assay

Juan Chen¹, Steven C. Rogers², and Mahendra Kavdia^{1,*}

¹Department of Biomedical Engineering, Wayne State University, Detroit, MI 48202

²Cell and Molecular Biology Program, University of Arkansas, Fayetteville, AR72701

Abstract

Superoxide (O_2^-) is an important reactive oxygen species (ROS), and has an essential role in physiology and pathophysiology. An accurate detection of O_2^- is needed to better understand numerous vascular pathologies. In this study, we performed a mechanistic study by using the xanthine oxidase (XOD)/hypoxanthine (HX) assay for O_2^- generation and a O_2^- sensitive fluorescent dye dihydroethidium (DHE) for O_2^- measurement. To quantify O_2^- and DHE interactions, we measured fluorescence using a microplate reader. We conducted a detailed reaction kinetic analysis for DHE- O_2^- interaction to understand the effect of O_2^- self-dismutation and to quantify DHE- O_2^- reaction rate. Fluorescence of DHE and 2-hydroethidium (EOH), a product of DHE and O_2^- interaction, were dependent on reaction conditions. Kinetic analysis resulted in a reaction rate constant of $2.169 \pm 0.059 \times 10^3 \text{ M}^{-1}\text{s}^{-1}$ for DHE- O_2^- reaction that is $\sim 100\times$ slower than the reported value of $2.6 \pm 0.6 \times 10^5 \text{ M}^{-1}\text{s}^{-1}$. In addition, the O_2^- self-dismutation has significant effect on DHE- O_2^- interaction. A slower reaction rate of DHE with O_2^- is more reasonable for O_2^- measurements. In this manner, the DHE is not competing with superoxide dismutase and NO for O_2^- . Results suggest that an accurate measurement of O_2^- production rate may be difficult due to competitive interference for many factors; however O_2^- concentration may be quantified.

Keywords

DHE; Reaction rate constant; Kinetic analysis; Oxidative stress; cytochrome c

Introduction

Superoxide (O_2^-), which is one of the major reactive oxygen species (ROS), is one-electron reduction product of molecular oxygen. O_2^- participates in signaling pathways^{4, 9, 12} and is a common marker for oxidative stress. O_2^- leads to the formation of other ROS including hydrogen peroxide (H_2O_2), and hydroxyl radical ($\bullet OH$), and reactive nitrogen species (RNS)^{7, 22, 25}. The accurate detection of O_2^- levels is critical for understanding numerous physiological phenomenon and pathologies such as cell proliferation, apoptosis, cardiovascular disease, diabetes and cancer^{8, 15, 26, 30}.

In the previous literature, several methods have been described to detect O_2^- . These methods include lucigenin and luminol assays, electron paramagnetic resonance, high performance liquid chromatography (HPLC) coupled with mass spectrometry (MS), and fluorescence^{7, 8, 30, 32}. Most of these methods are not specific for O_2^- detection. For

*Corresponding Author: Mahendra Kavdia, Ph.D., Department of Biomedical Engineering, Wayne State University, 2152 Engineering Bldg, 5050 Anthony Wayne Dr., Detroit, MI 48202., Telephone: 313-577-3803, Fax: 313-577-8333, kavdia@wayne.edu.

example, lucigenin and dichlorodihydrofluorescein-diacetate (H₂DCF-DA) can generate O₂⁻ and react with other oxidative stress. In addition, some methods can be very harmful to the human biological system. For example, lucigenin can stimulate oxidative stress formation to inhibit endothelium-dependent relaxation²⁹, and hydroethidine (45 μM) has cytotoxic effect and there is an inverse correlation between hydroethidine uptake and cell survival³⁵.

Since the early 1990's, fluorescence resulting from the oxidation of dihydroethidium (DHE) has been used as an O₂⁻ probe with success²⁷. DHE has been established to detect O₂⁻ with end-product fluorescence measurement by many researchers^{5, 6, 24, 33, 34, 37}. DHE fluorescence is inhibited by O₂⁻ scavengers such as superoxide dismutase (SOD) and nitric oxide (NO)²⁰. Products for the oxidation of DHE by O₂⁻ are well established. In the presence of O₂⁻, DHE is oxidized to EOH and intermediate products. These intermediate products can react with •OH or H₂O₂ to form ethidium (E⁺) to a much lesser extent. Whereas E⁺ fluorescence is measured at excitation of 500–530nm and emission of 590–620nm, the EOH fluorescence is measured at an excitation and emission wavelength of 480nm and 567nm, respectively^{8, 33}. EOH is stable within the cell, allowing for precise measurement of DHE fluorescence without risk of intra-conversion variability.

Fluorescence measurement of DHE yields O₂⁻ levels specific for either intracellular or extracellular locations²². However, it is difficult to detect and quantify O₂⁻ because of its short life-time (a few seconds), low flux rate, fast self-dismutation rate, rapid reaction with multiple intra- and extracellular scavengers such as SOD, vitamin C and E, glutathione (GSH) and NO^{3, 8}, and fluorescent dye reacting with various species to form other products or intermediate products^{10, 21, 35}. For the detection of O₂⁻ using DHE fluorescence, HPLC, microplate reader analysis, and fluorometry have been proven effective for qualitative and quantitative measurement of O₂⁻ concentrations¹⁶. However, the quantitative understanding of O₂⁻ interaction with DHE in experiments is needed. Without this, only qualitative measurement of O₂⁻ is possible. A quantitative measurement of O₂⁻ interaction will help us to further study the impact and mechanism of O₂⁻ resulting damage.

Several studies have focused on quantitative interaction of DHE with O₂⁻^{2, 6, 13, 18, 33, 36}. Laurindo et al.¹³ described the possible way to separate using HPLC and to quantify EOH, which is the main product of DHE and O₂⁻ interaction, and E⁺, which is the main product of DHE and H₂O₂ interaction. Fink Bruno et al.⁶ reported that EOH formation is inhibited in the presence of SOD, which catalyzes O₂⁻ dismutation. After the addition of H₂O₂, peroxynitrite (ONOO⁻) or hypochlorous acid to O₂⁻ generation system of potassium superoxide and XOD, they reported no additional change in EOH formation. Benov et al.² showed that the ratio of E⁺ formation/O₂⁻ decreased as the flux of O₂⁻ increased. In addition, they showed that the DHE can catalyze the dismutation of O₂⁻. Fernandes et al.⁵ showed that EOH increased after treatment of smooth muscle cells with AngII by detecting with HPLC and fluorescence micro-plate reader.

Zhao et al.³³ reported that the addition of DEA/NO (an NO donor) inhibits the EOH fluorescence in xanthine/XOD system. In addition, using competition kinetic analysis with SOD, they estimated a reaction rate constant for DHE and O₂⁻ interaction of $2.6 \pm 0.6 \times 10^5 \text{ M}^{-1}\text{s}^{-1}$. Later, Zielonka et al.³⁶ used pulse radiolysis technique to calculate the rate constant between O₂⁻ and DHE is $2.0 \times 10^6 \text{ M}^{-1}\text{s}^{-1}$. They also reported that DHE interaction with only O₂⁻ results in EOH formation³⁴. Previous measurements of DHE interactions with O₂⁻ ignored the self-dismutation of O₂⁻, which can lead to the overestimation of this reaction rate constant.

In this study, we systematically investigated interactions of O₂⁻ with DHE by measuring DHE and oxidized product EOH using XOD catalysis of HX oxidation to form O₂⁻ using a

microplate reader. We varied enzyme (XOD), substrate (HX), and reactant (DHE) concentration over a wide range to understand how these can affect the DHE and O_2^- interaction. We also developed a biochemical kinetic model to quantify DHE and O_2^- interactions and to understand the effect of self-dismutation of O_2^- . As a result of this study, we provide a new reaction rate constant for the reaction between DHE and O_2^- .

Materials and Methods

Materials

XOD, HX, catalase, ferricytochrome-c and phosphate buffer solution (PBS) were obtained from Sigma Chemical Co. (St. Louis, MO, USA). DHE was purchased from Invitrogen Corp. (Carlsbad, CA, USA).

O_2^- and DHE reaction

Reactions were conducted in a 96-well Microtest™ Optilux™ Plate with a transparent bottom holding a total reagent volume of 250 μ L in PBS (10 mM, pH 7.4). O_2^- was produced using HX and XOD enzyme system¹¹. Catalase was used to remove excess H_2O_2 , which is generated from the dismutation of O_2^- and the XOD/HX system. The reaction volume of 250 μ L contained DHE, HX, catalase (100 U/ml) with or without XOD.

To understand the kinetics of the DHE and O_2^- reaction, we modified the kinetic analysis from an earlier study by Zhao et al.³³ that used the competition kinetic analysis to provide a rate constant of $2.6 \pm 0.6 \times 10^5 \text{ M}^{-1}\text{s}^{-1}$ for the O_2^- and DHE reaction. Three sets of experiments were conducted. First, we varied XOD concentration 1, 1.5, 2.5, 5 mU/ml, and kept HX and DHE concentration constant at 0.25 mM and 5 μ M, respectively. Second, we varied DHE concentration 2, 5, 10, 20, and 50 μ M and kept HX and XOD concentration constant at 0.25 mM and 1.5 mU/ml, respectively. Finally, the HX concentration was varied to 0.0625, 0.125, and 0.25 mM and XOD and DHE concentration was kept at 1.5 mU/ml and 5 μ M, respectively.

Fluorescence measurement

In this study, EOH fluorescence was monitored by a Synergy 2 multi-detection microplate reader using a Gen5 microplate data collection & analysis software (BioTek Instruments Inc., VT, USA). As described by Zhao³³, Robinson²⁴ and Laurindo¹³, the excitation/emission wavelength was chosen as 420/590 nm with a 50nm or 35nm slit width, respectively. The DHE fluorescence was measured at excitation/emission wavelength of 360/460 nm with a 40nm slit width. The plate was read from 50% top, which was suitable for all these filters, and the sensitivity was set to be 70. The kinetic measurements were performed for 120 min and fluorescence was recorded every 3 min. The fluorescence values were obtained by subtracting the background fluorescence values obtained in PBS with HX, catalase and DHE in the absence of XOD.

To obtain the DHE concentration, we generated a standard curve of DHE (concentration vs. fluorescence) by measuring DHE fluorescence at DHE concentration of 1, 2, 5, 10, 20, and 50 μ M in the same solution as experiment group without XOD. We observed that the DHE fluorescence without XOD in the solution (i.e. without superoxide) does not change significantly over 120 min.

O_2^- production

O_2^- production was measured using a widely used rapid reduction of ferri-cytochrome c reduction assay^{14, 16}. Absorbance readings at 550 nm were collected every min over 30 min. A molar extinction coefficient $21,000 \text{ M}^{-1}\text{cm}^{-1}$ ³³ and a path length 0.975 cm were

used to calculate O_2^- production of different HX and XOD concentration experiments. The absorbance assay volume was 250 μ L in PBS with reagents including XOD (1, 1.5, 2.5, and 5 mU/ml), HX (0.0625, 0.125, and 0.25 mM), catalase (100 U/mL), and ferri-cytochrome c (81 μ M). We determined the rate of O_2^- generation by multiplying the slope of ferri-cytochrome c generation with the reciprocal of extinction coefficient of ferri-cytochrome c and path length.

DHE and O_2^- reaction kinetic analysis

We consider the following reactions occurring in the system to quantify the DHE and O_2^- interactions.

- i. Reaction between DHE and O_2^- yields a fluorescent product at a rate constant of k_{DHE} as follows:

- ii. We included the self-dismutation of O_2^- in the analysis. Earlier studies neglected this reaction^{31, 33}. We also analyzed the effect of this reaction on the results of our study. In addition, we added large amount of catalase to scavenge H_2O_2 . The rate constant for self-dismutation of O_2^- (k_D)^{1, 23} is $8 \times 10^7 \text{ M}^{-1}\text{s}^{-1}$ and the reaction occurs as follows:

Thus, the mass balance equation in the reaction system for the species of interests, O_2^- , and DHE, can be written as:

$$\frac{dC_{O_2^-}}{dt} = S_{O_2^-} - k_{DHE} C_{DHE} C_{O_2^-} - k_D C_{HO_2} C_{O_2^-}, \quad (3)$$

$$\frac{dC_{DHE}}{dt} = -k_{DHE} C_{DHE} C_{O_2^-} \quad (4)$$

Note that i) $S_{O_2^-}$ is the rate of O_2^- production in the system and was obtained from ferri-cytochrome c reduction assay as described in the previous subsection, and ii) the rate of O_2^- dismutation can be simplified assuming a rapid equilibrium of HO_2/O_2^- with $C_{HO_2}/C_{O_2^-} = 0.0025$, which is based on $pK_a = 4.8$ and the pH value of 7.4 and the relationship

for the mole fraction of HO_2 ($f = \frac{1}{1 + 10^{(pK_{per} - pH)}}$). We assumed initial concentration of O_2^- is zero.

For a given experimental condition, Equations 3 and 4 were solved using the MatLab[®] with initial concentrations of DHE (varied) and O_2^- . We fitted the experimental data of DHE concentration (measured at ex/em of 360/460 nm) with the model predictions by varying k_{DHE} . As shown in Equation 5, we used least square estimation and minimized the sum of squares of error to obtain the rate constant, k_{DHE} .

$$Q = \sum_{j=1}^k \sum_{i=1}^n (y_{exp} - y_{model})^2 \quad (5)$$

Where y_{exp} and y_{model} are the average measured and model predicted DHE concentration values, respectively for one experiment condition. For each of the three sets of experiments mentioned earlier, k represents the number of varying conditions in one set of experiments and n represents the number of time data points in one experimental conditions. We fit the data only for first 18 min of experiments, therefore n is equal to 7. For each set of experiments, a new value of Q was calculated. The reported rate constant in this manuscript corresponds to the minimum Q .

Statistical analysis

Three runs were performed for each experimental condition and all values are reported as mean $\pm$ st. err. We used the Matlab function `ode15s` to solve the simultaneous differential equations and used absolute error and relative error of 1×10^{-13} . For optimization of k_{DHE} , we used `fminsearch` function with a toleration error for Q of 1×10^{-13} .

Results

DHE concentration related change in DHE and EOH fluorescence

Figure 1A and B show the reduction of DHE fluorescence and the accumulation of EOH fluorescence for DHE concentration of 2, 5, 10, 20, and 50 μ M over 120 minutes. The initial XOD and HX concentration was 1.5 mU/ml and 0.25 mM, respectively. Both DHE and EOH fluorescence had two phases of fluorescence change. The fluorescence changed at an initial high slope followed by a small slope. The DHE fluorescence decreased linearly for the first 20 min for all DHE concentrations. In addition, an initial faster decrease in DHE fluorescence and increase in EOH fluorescence was observed at higher DHE concentrations ($> 10 \mu$ M). After 40 min, DHE reacted completely and the formation of EOH reached a plateau except for the DHE concentration of 20 and 50 μ M.

Enzyme XOD concentration related change in DHE and EOH fluorescence

Figure 2A and B show DHE and EOH fluorescence for XOD concentration of 1, 1.5, 2.5, and 5 mU/ml over 120 min. The initial DHE and HX concentration was 5 μ M and 0.25 mM, respectively. The DHE and EOH fluorescence again show the two phases for changes of fluorescence density; an initial steep change for the first 20 min followed by a slow change. In addition, Figure 2A and B show that the higher the XOD concentration, the faster the initial decrease in DHE fluorescence and the increase in EOH fluorescence. At XOD concentration of 1 mU/ml, Figure 2B shows that the DHE did not completely transform into EOH as demonstrated by non-convergence of RFU at 120 min with other XOD concentrations. This may be possible because of auto-oxidation of DHE resulting in other products and not EOH or sufficient superoxide formation to react with DHE.

Substrate HX concentration related change in DHE and EOH fluorescence

Figure 3A and B show the effect of HX concentration of 0.0625, 0.125, and 0.25 mM on DHE and EOH fluorescence, respectively. The initial DHE concentration was 5 μ M and 1.5 mU/ml for XOD. The same two phases in DHE and EOH fluorescence were also present in this set of experiment. However, the initial DHE and EOH fluorescence changes were higher at increased HX concentrations. Figure 3B also shows that DHE did not completely convert into EOH at HX concentration of 0.0625 and 0.125 mM as compared with HX concentration of 0.25 mM.

Absorbance measurements of O_2^- generation (product of XOD and HX)

Figure 4A and B describe ferrocyanochrome c absorbance for the measurement of O_2^- formation rate as a function of XOD concentration and HX concentration, respectively. These data is required for the kinetic analysis. As shown, the ferrocyanochrome c absorbance increased linearly with varying slope for the first 20 min for all conditions. It implies that O_2^- formation rate is constant for a given condition over the first 20 min. For a given HX and XOD concentration, the O_2^- formation rate was calculated by multiplying the slope of increase in ferrocyanochrome c absorbance with the reciprocal of EC value ($21 \text{ mM}^{-1} \text{ cm}^{-1}$) and path length (0.975 cm). At 1, 1.5, 2.5, and 5 mU/ml XOD concentration, the O_2^- formation rate was 0.943, 1.465, 1.636, and 2.623 $\mu\text{M}/\text{min}$, respectively. The O_2^- formation rate was 0.866, 1.189, and 1.473 $\mu\text{M}/\text{min}$ at 0.0625, 0.125, and 0.25 mM HX concentration. This data indicated the higher substrate and enzyme concentration leads to the higher O_2^- formation rate.

Standard curve of DHE fluorescence vs. concentration

As shown in Figure 5, we generated the standard curve of DHE fluorescence vs. DHE concentration by fitting DHE concentrations (1, 2, 5, 10, 20, and 50 μM) with their corresponding initial fluorescence. As we used a very wide range of DHE concentration, the optimal fit was not linear. However, the nonlinear relationship ($R^2 = 0.9965$) between DHE concentration and RFU is $[\text{DHE}], M = 1 \times 10^{-7} \times (\text{DHE RFU})^2 + 0.0014 \times (\text{DHE RFU})$.

Kinetic analysis of DHE and O_2^- reaction

In order to obtain the kinetic rate constant, we fitted measured experimental data with model data by varying k_{DHE} . We used non-linear relationships from previous section to convert DHE RFU into DHE concentration.

First, we fitted experimental data from Figure 1A with varying DHE concentration and the constant O_2^- formation rate of 1.469 $\mu\text{M}/\text{min}$, which is the average value of 1.465 and 1.473 $\mu\text{M}/\text{min}$ based on the ferrocyanochrome c absorbance data at XOD=1.5 mU/ml and HX=0.25 mM. The minimum sum of squares of error between experimental data and predicted model data (Q) was $4.045 \times 10^3 \text{ M}^{-1}\text{s}^{-1}$. The minimum Q was obtained with a reaction rate constant of DHE and O_2^- of $2.116 \times 10^3 \text{ M}^{-1}\text{s}^{-1}$. The resulting fit with experimental data is shown in Figure 6A. As seen, the model data closely predicted each point of experimental data for 18 min for all DHE concentrations.

Next, we fitted experimental data from Figure 2A and 3A for varying XOD concentration and HX concentration, respectively. These conditions resulted in varying O_2^- formation rates as described in previous sections for ferrocyanochrome c absorbance data. Using different O_2^- formation rates, we obtained k_{DHE} for each of the XOD and HX concentration variation. The resulting fit with experimental data is shown in Figure 6B and 6C. At 1, 1.5, 2.5, and 5 mU/ml XOD concentration, the k_{DHE} is 1.933×10^3 , 1.827×10^3 , 2.626×10^3 , and $2.372 \times 10^3 \text{ M}^{-1}\text{s}^{-1}$, and at the HX concentration of 0.0625, 0.125, and 0.25 mM, the k_{DHE} reaction is 2.191×10^3 , 2.208×10^3 , and $2.295 \times 10^3 \text{ M}^{-1}\text{s}^{-1}$ resulted in the best fit between experimental and predicted data. The respective Q value was 5.436×10^{-15} , 2.415×10^{-14} , 1.447×10^{-13} , 7.072×10^{-15} , 7.327×10^{-15} , 4.122×10^{-15} , and 2.628×10^{-15} . Based on all these reaction rate constant of DHE and O_2^- , the mean rate of reaction of DHE with O_2^- , k_{DHE} , is $2.169 \pm 0.059 \times 10^3 \text{ M}^{-1}\text{s}^{-1}$.

Discussion

O_2^- is the precursor of many derivative ROS, and can rapidly react with NO to inactivate NO bioavailability and augment production of RNS. Therefore, the quantitative

measurement of O_2^- measurement is critical. Traditionally, the fluorescence product of DHE was measured at the excitation/emission wavelength of 510/595 nm, which is the wavelength for detecting E^+ . The reaction of O_2^- and DHE also generates EOH that is confirmed to be the specific products formed. Researchers have started using excitation/emission wavelength to detect EOH fluorescence at either sensitive or specific wavelengths of 480/580 nm and 396/579 nm, respectively^{5, 13, 24, 33}. In this study, we have confirmed that DHE fluorescence can be used to quantify O_2^- by HX-XOD system using microplate readers like other methods of O_2^- measurement^{8, 13, 30, 32, 33}. The development of mathematical model for DHE interaction with O_2^- is novel and can be extended further for understanding many aspects of oxidative stress interactions including NO at molecular level. In addition, we provide a calibration curve for DHE concentration using RFU for the first time. We propose a DHE and O_2^- reaction rate constant of $2.169 \pm 0.059 \times 10^3 \text{ M}^{-1}\text{s}^{-1}$.

O_2^- measurements

A predictable cell-free O_2^- generating system was utilized to achieve measurements of O_2^- concentration using a common fluorescence detection method and microplate reader analysis. The formation of EOH and depletion of DHE was specific to O_2^- as seen in Figure 1 with varying DHE concentration. The change in the rate of DHE fluorescence was dependent on O_2^- formation from the HX-XOD system as seen in Figures 2 and 3. However, in the absence of O_2^- , DHE depletion was also influenced by autooxidation of DHE at later time points (>40 min). Both the increase in EOH and the decrease in DHE occurred in two linear phases: a steep initial change followed by a slower, steadier change. As the DHE concentration or the rate of O_2^- formation increased through the increase in the concentration of HX and XOD, both the decrease in DHE fluorescence and increase in EOH fluorescence would be accelerated. From these profiles, we showed that the higher the concentration of reactant, the faster the reaction between DHE and O_2^- . The biphasic curve in Figures 1–3 was because of the change in the superoxide production rate at later time points that were seen after 30 min in most cases (data not shown) during absorbance assay (Figure 4). We showed that using a wide ranging superoxide production rate (varying HX and XOD concentrations for the first 18 min), we were able to fit the data with the k_{DHE} .

Currently, much research begins to pay attention to the accuracy of fluorescence detection of DHE- O_2^- reaction^{13, 35}. These studies suggest that HPLC analysis is a reliable method for measurement of DHE detection of O_2^- . Due to the extremely short O_2^- half-life and rapid dismutation reaction rate, DHE fluorescence detection is prone to underestimation of actual O_2^- production^{34, 37}. Additionally, studies have shown that DHE can react with various oxidants found within the cell, including mitochondrial cytochrome hemes, hypochlorous acid, myeloperoxidase, horseradish peroxidase $ONOO^-$, *OH , H_2O_2 , compounds I and II, Fenton reagent, and so on^{10, 19, 21, 35}. Shao et al.²⁸ also demonstrated that certain potent diet antioxidants can greatly decrease O_2^- concentration even in extreme acute oxidative stress. The total O_2^- production measurements are impossible as reported⁸. At best, it will be an estimate due to the varying amount of O_2^- consumption among various interactions with other species.

However, Munzel¹⁷ has suggested that it is wise to use two or more techniques to detect O_2^- . Reliable conclusions could be drawn from similar results by using different measurements. In this study, we used cytochrome c absorbance and DHE fluorescence to detect O_2^- generation. Our results show that the formed products between superoxide and DHE or Cyt c respectively increase linearly. Moreover, by fitting our experimental data with computational model, we confirmed that DHE fluorescence measurement and microplate reader analysis can describe as a quantitative measurement of localized O_2^- concentration in vivo, which will provide us an effective method to detect O_2^- .

DHE-O₂⁻ reaction rate constant of $2.169 \pm 0.059 \times 10^3 \text{ M}^{-1}\text{s}^{-1}$

Based on experimental measurements and computational modeling in this study, we report a reaction rate constant of $\sim 2.169 \pm 0.059 \times 10^3 \text{ M}^{-1}\text{s}^{-1}$, which is $\sim 100 \times$ lower than the previously reported value of $2.6 \pm 0.6 \times 10^5 \text{ M}^{-1}\text{s}^{-1}$ ³³. There are two possible reasons for this discrepancy. Firstly, the kinetic analysis in our study is more detailed with respect to the reactions occurring in the system. Secondly, the self-dismutation of O₂⁻ was considered in present study that increased O₂⁻ consumption and was not considered in previous study. The rate constant of O₂⁻ self-dismutation ($8.0 \times 10^7 \text{ M}^{-1}\text{s}^{-1}$) is lower than that of O₂⁻ and SOD reaction ($1.6 \times 10^9 \text{ M}^{-1}\text{s}^{-1}$)²³, but higher than that of DHE and O₂⁻ reaction ($2.169 \pm 0.059 \times 10^3 \text{ M}^{-1}\text{s}^{-1}$). Therefore, it is necessary to include O₂⁻ self-dismutation in DHE and O₂⁻ reaction system as done in this study.

We analyzed reasons behind a lower reaction rate constant for DHE reaction with O₂⁻. We simulated the experiment performed by Zhao et al.³³ for k_{DHE} of 2.17×10^3 and $2.69 \times 10^5 \text{ M}^{-1}\text{s}^{-1}$ ³³ with and without the inclusion of self-dismutation of O₂⁻. We used a 50 μM initial DHE and 8 $\mu\text{M}/\text{min}$ O₂⁻ generation rate³³, and the results are plotted in Figure 7. Figures 7A and 7B show results for k_{DHE} of $2.69 \times 10^5 \text{ M}^{-1}\text{s}^{-1}$ without and with the inclusion of self-dismutation of O₂⁻, respectively, while Figure 7C and 7D show results for k_{DHE} of $2.17 \times 10^3 \text{ M}^{-1}\text{s}^{-1}$ without and with the inclusion of self-dismutation of O₂⁻, respectively. Figures 7A–C are similar to each other and show that DHE is completely consumed and transformed to EOH at 6, 7, and 8 min respectively. However, as Zhao et al.³³ described that the intensity of product fluorescence linear increased (Figure 10 in reference³³) and kept the same trend for first 20 minutes until they added DEA/NO into this reaction buffer. The model simulation results are not in agreement with experimental results. Therefore, we can conclude that DHE would fast consume for both k_{DHE} when self-dismutation of O₂⁻ is neglected.

However, when we include self-dismutation with the lower k_{DHE} , Figure 7D shows that there is an initial linear increase in EOH and decrease in DHE for the first 20 min followed by a slow change in EOH and DHE. In addition, we also simulated the experimental results of the current study without (Figure 7E) or with (Figure 7F) self-dismutation reaction at 5 μM DHE and 1.465 $\mu\text{M}/\text{min}$ superoxide production rate (HX (0.25 mM) and XOD (1.5 mU/ml)). As shown in Figure 7E, the depletion of DHE occurred in the first 10 minutes without self-dismutation reaction. Figure 7F shows that the model predictions with self-dismutation reaction are similar to the experiment result i.e. DHE can last much longer. This indicated that O₂⁻ self-dismutation cannot be neglected and model results are in agreement with experimental results.

Implications of slower reaction rate of DHE with O₂⁻

A slower reaction rate of DHE with O₂⁻ is more reasonable for O₂⁻ measurements. In addition to superoxide production rate, the most important factors for the determination of O₂⁻ concentration in a given system are SOD and NO. SOD can fast dismutate O₂⁻ into H₂O₂ and NO can quickly react with O₂⁻ to generate RNS. The slower reaction rate of DHE-O₂⁻ implies that the presence of DHE does not affect O₂⁻ concentration, whereas the faster reaction rate implies that DHE competes with SOD and NO for O₂⁻ and reduces O₂⁻ concentration. Therefore, the respective changes in DHE fluorescence indicate change in O₂⁻ concentration in the system for the slower reaction rate. Many other chemical reactions simultaneously occur in the physiological system, the results from this study can be extrapolated to explain the effect of these reactions on oxidative stress. The change in DHE fluorescence will be in the same trend as these other reactions affect the O₂⁻ concentration. Thus, the fluorescence probes designed to exhibit a relatively low rate constant with the analyte detected will be ideal in vivo detection of O₂⁻. In ROS and RNS mathematical

modeling, SOD can fast dismutate O_2^- into H_2O_2 and NO can quickly react with O_2^- to generate RNS. We conclude that DHE can be used to detect the O_2^- concentration without the interference of pathways in the cell system. It will be useful to determine the accumulated O_2^- concentrations and to predict damages in cell systems.

In conclusion, an accurate method for O_2^- measurement would be highly beneficial to the potential diagnosis and treatment of a number of different biological issues. This current study describes some necessary considerations for experimental measurement and analysis of O_2^- concentrations. DHE measurement of O_2^- may provide a more accurate description of oxidative stress, especially for O_2^- . This detailed kinetic analysis study provided a DHE and O_2^- reaction rate constant of $\sim 2.169 \pm 0.059 \times 10^3 \text{ M}^{-1}\text{s}^{-1}$ that is $\sim 100 \times$ lower than the previously reported value of $2.6 \pm 0.6 \times 10^5 \text{ M}^{-1}\text{s}^{-1}$, and is more appropriate for DHE O_2^- measurement.

Acknowledgments

This study was supported by NIH grant # R01 HL084337.

Abbreviations

DHE	Dihydroethidium
E⁺	Ethidium
EOH	2-Hydroethidium
GSH	Glutathione
H₂DCF-DA	Dichlorodihydrofluorescein-diacetate
H₂O₂	Hydrogen peroxide
HPLC	High performance liquid chromatography
HX	Hypoxanthine
MS	Mass spectrometry
NO	Nitric oxide
O₂⁻	Superoxide
•OH	Hydroxyl radical
ONOO⁻	Peroxynitrite
PBS	Phosphate buffer solution
ROS	Reactive oxygen species
RNS	Reactive nitrogen species
SOD	Superoxide dismutase
XOD	Xanthine oxidase

References

- Behar D, Czapski G, Rabani J, Dorfman LM, Schwarz HA. Acid Dissociation Constant and Decay Kinetics of Perhydroxyl Radical. *J Phys Chem-Us*. 1970; 74:3209.
- Benov L, Szejnberg L, Fridovich I. Critical evaluation of the use of hydroethidine as a measure of superoxide anion radical. *Free Radical Biology and Medicine*. 1998; 25:826–831. [PubMed: 9823548]

3. Deng T, Xu K, Zhang L, Zheng X. Dynamic determination of Ox-LDL-induced oxidative/nitrosative stress in single macrophage by using fluorescent probes. *Cell Biol Int*. 2008; 32:1425–32. [PubMed: 18782627]
4. Droge W. Free radicals in the physiological control of cell function. *Physiol Rev*. 2002; 82:47–95. [PubMed: 11773609]
5. Fernandes DC, Wosniak J Jr, Pescatore LA, Bertoline MA, Liberman M, Laurindo FR, Santos CX. Analysis of DHE-derived oxidation products by HPLC in the assessment of superoxide production and NADPH oxidase activity in vascular systems. *Am J Physiol Cell Physiol*. 2007; 292:C413–22. [PubMed: 16971501]
6. Fink B, Laude K, McCann L, Doughan A, Harrison DG, Dikalov S. Detection of intracellular superoxide formation in endothelial cells and intact tissues using dihydroethidium and an HPLC-based assay. *Am J Physiol Cell Physiol*. 2004; 287:C895–902. [PubMed: 15306539]
7. Fujita M, Tsuruta R, Kasaoka S, Fujimoto K, Tanaka R, Oda Y, Nanba M, Igarashi M, Yuasa M, Yoshikawa T, Maekawa T. In vivo real-time measurement of superoxide anion radical with a novel electrochemical sensor. *Free Radic Biol Med*. 2009; 47:1039–48. [PubMed: 19616620]
8. Georgiou CD, Papapostolou I, Patsoukis N, Tsegenidis T, Sideris T. An ultrasensitive fluorescent assay for the in vivo quantification of superoxide radical in organisms. *Analytical Biochemistry*. 2005; 347:144–51. [PubMed: 16246291]
9. Hernandez MS, Britto LR, Real CC, Martins DO, Lopes LR. Reactive oxygen species and the structural remodeling of the visual system after ocular enucleation. *Neuroscience*. 2010; 170:1249–60. [PubMed: 20728508]
10. Kalyanaraman B, Darley-Usmar V, Davies KJA, Dennery PA, Forman HJ, Grisham MB, Mann GE, Moore K, Roberts LJ, Ischiropoulos H. Measuring reactive oxygen and nitrogen species with fluorescent probes: challenges and limitations. *Free Radical Bio Med*. 2012; 52:1–6. [PubMed: 22027063]
11. Kavdia M, Stanfield JL, Lewis RS. Nitric oxide, superoxide, and peroxynitrite effects on the insulin secretion and viability of betaTC3 cells. *Ann Biomed Eng*. 2000; 28:102–9. [PubMed: 10645793]
12. Kishida KT, Klann E. Sources and targets of reactive oxygen species in synaptic plasticity and memory. *Antioxid Redox Signal*. 2007; 9:233–44. [PubMed: 17115936]
13. Laurindo FR, Fernandes DC, Santos CX. Assessment of superoxide production and NADPH oxidase activity by HPLC analysis of dihydroethidium oxidation products. *Methods Enzymol*. 2008; 441:237–60. [PubMed: 18554538]
14. Massey V. The microestimation of succinate and the extinction coefficient of cytochrome c. *Biochim Biophys Acta*. 1959; 34:255–6. [PubMed: 14422133]
15. McFarland R, Blokhin A, Sydnor J, Mariani J, Vogel MW. Oxidative stress, nitric oxide, and the mechanisms of cell death in Lurcher Purkinje cells. *Dev Neurobiol*. 2007; 67:1032–46. [PubMed: 17565706]
16. Messner KR, Imlay JA. In vitro quantitation of biological superoxide and hydrogen peroxide generation. *Methods Enzymol*. 2002; 349:354–61. [PubMed: 11912927]
17. Munzel T I, Afanas'ev B, Kleschyov AL, Harrison DG. Detection of superoxide in vascular tissue. *Arterioscler Thromb Vasc Biol*. 2002; 22:1761–8. [PubMed: 12426202]
18. Papaharalambus CA, Griendling KK. Basic mechanisms of oxidative stress and reactive oxygen species in cardiovascular injury. *Trends in Cardiovascular Medicine*. 2007; 17:48–54. [PubMed: 17292046]
19. Papapostolou I, Patsoukis N, Georgiou CD. The fluorescence detection of superoxide radical using hydroethidine could be complicated by the presence of heme proteins. *Anal Biochem*. 2004; 332:290–8. [PubMed: 15325298]
20. Paravicini TM, Touyz RM. NADPH oxidases, reactive oxygen species, and hypertension: clinical implications and therapeutic possibilities. *Diabetes Care*. 2008; 31(Suppl 2):S170–80. [PubMed: 18227481]
21. Patsoukis N, Papapostolou I, Georgiou CD. Interference of non-specific peroxidases in the fluorescence detection of superoxide radical by hydroethidine oxidation: a new assay for H₂O₂. *Anal Bioanal Chem*. 2005; 381:1065–1072. [PubMed: 15690180]

22. Peshavariya HM, Dusting GJ, Selemidis S. Analysis of dihydroethidium fluorescence for the detection of intracellular and extracellular superoxide produced by NADPH oxidase. *Free Radic Res.* 2007; 41:699–712. [PubMed: 17516243]
23. Potdar S, Kavdia M. NO/peroxynitrite dynamics of high glucose-exposed HUVECs: chemiluminescent measurement and computational model. *Microvascular research.* 2009; 78:191–8. [PubMed: 19362569]
24. Robinson KM, Janes MS, Pehar M, Monette JS, Ross MF, Hagen TM, Murphy MP, Beckman JS. Selective fluorescent imaging of superoxide in vivo using ethidium-based probes. *Proc Natl Acad Sci U S A.* 2006; 103:15038–43. [PubMed: 17015830]
25. Rojas A, Figueroa H, Re L, Morales MA. Oxidative stress at the vascular wall. Mechanistic and pharmacological aspects. *Arch Med Res.* 2006; 37:436–48. [PubMed: 16624640]
26. Ryter SW, Kim HP, Hoetzel A, Park JW, Nakahira K, Wang X, Choi AM. Mechanisms of cell death in oxidative stress. *Antioxid Redox Signal.* 2007; 9:49–89. [PubMed: 17115887]
27. Selemidis S, Dusting GJ, Peshavariya H, Kemp-Harper BK, Drummond GR. Nitric oxide suppresses NADPH oxidase-dependent superoxide production by S-nitrosylation in human endothelial cells. *Cardiovasc Res.* 2007; 75:349–58. [PubMed: 17568572]
28. Shao ZH, Xie JT, Vanden Hoek TL, Mehendale S, Aung H, Li CQ, Qin Y, Schumacker PT, Becker LB, Yuan CS. Antioxidant effects of American ginseng berry extract in cardiomyocytes exposed to acute oxidant stress. *Biochim Biophys Acta.* 2004; 1670:165–71. [PubMed: 14980443]
29. Tarpey MM, White CR, Suarez E, Richardson G, Radi R, Freeman BA. Chemiluminescent detection of oxidants in vascular tissue. Lucigenin but not coelenterazine enhances superoxide formation. *Circ Res.* 1999; 84:1203–11. [PubMed: 10347095]
30. Tarpey MM, Wink DA, Grisham MB. Methods for detection of reactive metabolites of oxygen and nitrogen: in vitro and in vivo considerations. *Am J Physiol Regul Integr Comp Physiol.* 2004; 286:R431–44. [PubMed: 14761864]
31. Vasquez-Vivar J, Whitsett J, Martasek P, Hogg N, Kalyanaraman B. Reaction of tetrahydrobiopterin with superoxide: EPR-kinetic analysis and characterization of the pteridine radical. *Free Radic Biol Med.* 2001; 31:975–85. [PubMed: 11595382]
32. Wardman P. Fluorescent and luminescent probes for measurement of oxidative and nitrosative species in cells and tissues: progress, pitfalls, and prospects. *Free Radic Biol Med.* 2007; 43:995–1022. [PubMed: 17761297]
33. Zhao H, Kalivendi S, Zhang H, Joseph J, Nithipatikom K, Vasquez-Vivar J, Kalyanaraman B. Superoxide reacts with hydroethidine but forms a fluorescent product that is distinctly different from ethidium: potential implications in intracellular fluorescence detection of superoxide. *Free Radic Biol Med.* 2003; 34:1359–68. [PubMed: 12757846]
34. Zielonka J, Hardy M, Kalyanaraman B. HPLC study of oxidation products of hydroethidine in chemical and biological systems: ramifications in superoxide measurements. *Free Radic Biol Med.* 2009; 46:329–38. [PubMed: 19026738]
35. Zielonka J, Kalyanaraman B. Hydroethidine- and MitoSOX-derived red fluorescence is not a reliable indicator of intracellular superoxide formation: another inconvenient truth. *Free Radic Biol Med.* 2010; 48:983–1001. [PubMed: 20116425]
36. Zielonka J, Sarna T, Roberts JE, Wishart JF, Kalyanaraman B. Pulse radiolysis and steady-state analyses of the reaction between hydroethidine and superoxide and other oxidants. *Archives of Biochemistry and Biophysics.* 2006; 456:39–47. [PubMed: 17081495]
37. Zielonka J, Vasquez-Vivar J, Kalyanaraman B. Detection of 2-hydroxyethidium in cellular systems: a unique marker product of superoxide and hydroethidine. *Nat Protoc.* 2008; 3:8–21. [PubMed: 18193017]

Figure 1. Measured DHE and EOH fluorescence as a function of DHE concentration

Figure 1A and B shows the DHE and EOH fluorescence with respect to time for DHE concentration of 2, 5, 10, 20, and 50 M with XOD concentration of 1.5 mU/ml and HX concentration of 0.25 mM for 120 min.

Figure 2. Measured DHE and EOH fluorescence as a function of XOD concentration

Figure 2A and B shows the DHE and EOH fluorescence for XOD concentration of 1.0, 1.5, 2.5 and 5 mU/ml for HX (=0.25 mM) and DHE (=5 mM) for 120 min.

Figure 3. Measured DHE and EOH fluorescence as a function of HX concentration
 Figure 3A and B shows the DHE and EOH fluorescence for HX concentration of 0.0625, 0.125 and 0.25 mM with XOD (=1.5 mU/ml) and DHE (=5 M) for 120 min.

Figure 4. O_2^- generation rate by HX-XOD system determined by measuring ferrocytochrome c at 550 nm ($EC=21 \text{ mM}^{-1}\text{cm}^{-1}$)

Figure 4A shows the absorbance of ferrocytochrome c for XOD concentration of 1.0, 1.5, 2.5 and 5 mU/ml for HX (=0.25 mM). Figure 4B shows the absorbance of ferrocytochrome c for HX concentration of 0.0625, 0.125 and 0.25 mM with XOD (=1.5 mU/ml).

Figure 5. Standard curve of DHE concentration vs. DHE fluorescence

Figure 5 shows DHE relative fluorescence unit (RFU) vs. DHE concentration for standard curve.

Figure 6. Measured and Predicted DHE concentration profiles using kinetic analysis

Figure 6A-C shows the kinetic model results (solid lines) for experimental data (discrete symbols) of varying DHE concentration, XOD concentration and HX concentrations, respectively.

Figure 7. Predicted O_2^- , EOH, and DHE concentration profiles using kinetic analysis

Figure 7 shows the predicted concentration profiles from the kinetic model for DHE and O_2^- reaction for 120 min. DHE and EOH concentration are shown on left axis. O_2^- concentration is shown on right axis. Figure A–D shows DHE and O_2^- reaction under O_2^- formation of $8 \mu\text{M}/\text{min}$ and DHE of $50 \mu\text{M}$, and Figure E–F shows DHE and O_2^- reaction under O_2^- formation of $1.465 \mu\text{M}/\text{min}$ and DHE of $5 \mu\text{M}$. Figure A and B shows the profile for without and with O_2^- self-dismutation, respectively for $2.6 \times 10^5 \text{ M}^{-1}\text{s}^{-1}$ reaction rate constant of DHE and O_2^- reaction. Figure C and E shows the profile for without O_2^- self-dismutation for $2.17 \times 10^3 \text{ M}^{-1}\text{s}^{-1}$ reaction rate constant of DHE and O_2^- reaction. Figure D

and F shows the profile for with O_2^- self-dismutation for $2.17 \times 10^3 \text{ M}^{-1}\text{s}^{-1}$ reaction rate constant of DHE and O_2^- reaction

\$watermark-text

\$watermark-text

\$watermark-text