Published in final edited form as:

Mol Microbiol. 2012 November ; 86(4): 1007–1022. doi:10.1111/mmi.12039.

A Role for Topoisomerase III in *Escherichia coli* Chromosome Segregation

Brenda A. Perez-Cheeks¹, Chong Lee¹, Ryo Hayama², and Kenneth J. Mariani^{1,*}

¹Molecular Biology Program, Memorial Sloan-Kettering Cancer Center, New York, NY 10065, USA

²Physiology, Biophysics, and Systems Biology Graduate Program, Weill-Cornell Graduate School of Medical Sciences, New York, NY 10065, USA

Summary

The cellular function of *Escherichia coli* topoisomerase III remains elusive. We show that rescue of temperature-sensitive mutants in *parE* and *parC* (encoding the subunits of the chromosomal decatenase topoisomerase IV) at restrictive temperatures by high-copy suppressors is strictly dependent on *topB* (encoding topoisomerase III). Double mutants of *parEΔtopB* and *parCΔtopB* were barely viable, grew slowly, and were defective in chromosome segregation at permissive temperatures. The *topB* mutant phenotype did not result from accumulation of toxic recombination intermediates, because it was not relieved by mutations in either *recQ* or *recA*. In addition, in an otherwise wild-type genetic background, *ΔtopB* cells treated with the type II topoisomerase inhibitor novobiocin displayed aberrant chromosome segregation. This novobiocin sensitivity was attributable to an increased demand for topoisomerase IV and is unlikely to define a new role for topoisomerase III; therefore, these results suggest that topoisomerase III participates in orderly and efficient chromosome segregation in *E. coli*.

Introduction

Topoisomerases are essential enzymes that regulate and maintain the topological state of chromosomal DNA by making a transient DNA break, passing a DNA segment through that break, and subsequently resealing the break (Wigley, 1995, Champoux, 2001, Bates & Maxwell, 2005). Type I topoisomerases cleave and pass one strand of DNA, whereas type II enzymes cleave and pass both strands of DNA (Wigley, 1995, Champoux, 2001, Bates & Maxwell, 2005).

Topoisomerases are essential for managing DNA topology during all macromolecular transactions on chromosomal DNA. In particular, their activities ensure the removal of excess positive topological linkages that accumulate during DNA replication because of the unwinding of the parental DNA strands. Either a type I or type II enzyme can be effective at this task. Once replication is completed, any remaining linkages between the two parental strands are manifested as catenanes between the two daughter chromosomes. If the replicated strands are completely sealed, then only a type II enzyme can unlink the daughter molecules (Wang, 1991, Sundin & Varshavsky, 1980, Sundin & Varshavsky, 1981). Champoux and Been (Champoux & Been, 1980) noted that the excess positive linkages that accumulate ahead of the replication fork can diffuse behind the replisome to form windings of the partially replicated daughter duplexes about each other that have come to be called

*Correspondence: Memorial Sloan-Kettering Cancer Center 1275 York Avenue New York, NY 10065 Tel. 212-639-5890 Fax 212-717-3627 kmarians@sloankettering.edu.

precatenanes (because once replication is completed they convert to catenanes). We demonstrated that precatenanes do form during replication of a circular DNA in vitro (Peter *et al.*, 1998). Precatenanes can clearly be resolved by type II topoisomerases, however, we have suggested that a type I topoisomerase operating at or near the replication fork where single-stranded gaps are available could also unlink them (Nurse *et al.*, 2003).

E. coli harbors four DNA topoisomerases: two type I enzymes [topoisomerase (Topo) I and III] and two type II enzymes (DNA gyrase and Topo IV) (Bates & Maxwell, 2005). Extensive studies and the properties of these enzymes have implicated some of them in clearly defined roles in the cell. DNA gyrase (encoded by *gyrA* and *gyrB*) (Gellert *et al.*, 1976a, Mizuuchi *et al.*, 1978) is required for preserving the global negative supercoiled state of the chromosome (Champoux, 2001), which it does primarily via the conversion of positive supercoils generated by DNA replication and transcription directly to negative supercoils (Levine *et al.*, 1998, Wang, 2002). Topo I (encoded by *topA*) is important for preventing the accumulation of excess negative supercoils by specifically relaxing negatively-supercoiled DNA (Wang, 1991), thus joining with DNA gyrase to homeostatically regulate the global level of chromosomal supercoiling (Menzel & Gellert, 1983).

Topo IV is required for the complete topological separation of the linked daughter chromosomes generated by DNA replication and is essential for viability (Kato *et al.*, 1990, Adams *et al.*, 1992, Peng & Marians, 1993). Topo IV consists of a heterotetramer of ParE and ParC dimers (Kato *et al.*, 1992, Peng & Marians, 1993). Temperature-sensitive mutants of *parE* and *parC* manifest a partitioning defect (*par*) (Kato *et al.*, 1989, Kato *et al.*, 1990) where cell division is arrested, leading to extreme filamentation coincident with the accumulation of one or two large masses of unsegregated, replicating DNA at the cell center.

More is known about the activities of Topo III in vitro than its role in vivo. This enzyme binds tightly to single-stranded DNA and does not appear to have the ability to locally unwind duplex DNA to create a binding site for catalysis. Thus, it is a potent decatenase on DNA rings that have small gaps, but is very poor at relaxing supercoils (Srivenugopal *et al.*, 1984, DiGate & Marians, 1988), suggesting that it is unlikely to contribute to the maintenance of global supercoiling. On the other hand, Topo III can support replication fork progression in the absence of either DNA gyrase or Topo IV, and generate completely unlinked daughter chromosomes (Hiasa & Marians, 1994). The ability of the enzyme to remove precatenanes is thought to contribute to these activities (Nurse *et al.*, 2003).

Mutants of *topB* (encoding Topo III) are viable and do not display obvious chromosome abnormalities or growth defects, thus it is not clear if Topo III is capable of resolving catenanes in vivo, but these mutants do show an increase in deletions between short homologous regions and an increase in frameshift mutations, suggesting that cells lacking *topB* have either an increased incidence of recombination at short DNA repeats or DNA replication mispairing slippage (Yi *et al.*, 1988, Schofield *et al.*, 1992, Uematsu *et al.*, 1997). In addition, *topB* mutations are synthetically lethal with *topA* (Zhu *et al.*, 2001) and *parE* mutations (Lopez *et al.*, 2005) and it has been reported that deletion of *recA* rescues the synthetic lethality in both cases, suggesting a role for *topB* in the resolution of recombination intermediates (Zhu *et al.*, 2001, Lopez *et al.*, 2005). This observation is consistent with the known role of Topo III in eukaryotes, where it has been shown to associate with various RecQ family helicases in resolving toxic, unscheduled recombination intermediates that can occur, for example, at stalled replication forks (Oakley & Hickson, 2002). Failure to resolve such intermediates can lead to chromosome breakage, chromosome rearrangements, and faulty chromosome segregation.

Functional cooperation between the *E. coli* enzymes has been demonstrated in vitro as well: Topo III and RecQ can catenate covalently-closed, double-stranded DNA rings (Harmon *et al.*, 1999) and catalyze the resolution of stalled, converging replication forks (Suski & Mariani, 2008). Collectively, these data point to a role for Topo III in vivo of either clearing toxic recombination and replication intermediates or unlinking sister chromosomes during replication, possibly with the assistance of its cognate helicase RecQ.

To understand how Topo IV could be regulated and to identify other proteins involved in resolving topologically-linked sister chromosomes, we identified high-copy suppressors of the temperature-sensitive (ts) alleles *parE10* and *parC1215*. Overproduction of several replisome components: the γ and τ subunits of the DNA polymerase III holoenzyme (both encoded by *dnaX*) and the β processivity clamp (encoded by *dnaN*), as well as an integral membrane protein encoded by *setB*, and Topo III could suppress the Topo IV mutant *par* and growth defects at nonpermissive temperatures (Levine & Mariani, 1998, Espeli *et al.*, 2003b, Nurse *et al.*, 2003).

As a potent decatenase (DiGate & Mariani, 1988), it was reasonable to speculate that Topo III, when overproduced, could suppress *parE* and *parC* ts mutations at the restrictive temperature by decatenation of linked sister chromosomes. However, the question remained: Does Topo III participate in the removal of catenanes and/or precatenanes under physiological conditions?

Suppressors of the *parE10* and *parC1215* mutations are likely to either play a role in chromosome segregation or influence Topo IV activity. Here we have investigated the role of these suppressors. We find that *topB* is required for the suppressors to promote growth of *parE* and *parC* ts mutant strains at restrictive temperatures, suggesting that the mechanism of suppression could be potentiation of Topo III activity. Combinations of $\Delta topB$ with the Topo IV ts mutations were synthetically lethal at permissive temperatures and these double mutant strains exhibited chromosome segregation defects. Contrary to previously reported findings (Lopez *et al.*, 2005), neither *recQ* nor *recA* were required for the *parE10* $\Delta topB$ synthetic lethality, because mutations in *recQ* and *recA* did not rescue the *parE10* $\Delta topB$ double mutant strain. Furthermore, in the absence of these recombination proteins, the high-copy suppressor γ [*dnaX*(γ)] is still unable to rescue the *parE10* $\Delta topB$ double mutant strain at 30 °C, arguing that the observed phenotype was not the result of the accumulation of toxic recombination intermediates. However, we discovered that mutants of *topB* are hypersensitive to the type II topoisomerase inhibitor novobiocin: when exposed to novobiocin these cells are less viable and have severe chromosome segregation defects. These data suggest that Topo III plays a role in orderly and efficient chromosome segregation.

Results

Rederivation of the parE10 and parC1215 mutations in W3110

As will be obvious below, some of our results deviate significantly from those reported by Lopez *et al.* (Lopez *et al.*, 2005). To eliminate the possibility that these differences could be the result of different suppressors acquired in the W3110*parE10* and C600*parC1215* (EJ182) strains originally isolated by Kato (Kato *et al.*, 1990, Kato *et al.*, 1988) over the years in different laboratories, we engineered both the *parE10* mutation (C316Y) and the *parC1215* mutation (G725D) into a fresh W3110 background. The resultant W3110*parC1215* (CL31) strain was barely viable even at 25 °C and could not be used for any analyses. The W3110*parE10* (CL001) strain used in this report is the newly rederived strain.

Topo III is required for γ -mediated rescue of *parE10* at the restrictive temperature

The isolation of high-copy suppressors of the *par* and growth defects of ts Topo IV mutations suggested that these suppressor genes either stabilized the ts Topo IV, formed a new topoisomerase, or either enhanced activity or activated one of the other topoisomerases in the cell. The first two possibilities have been ruled out in previous studies (Levine and Mariani, 1998; Nurse et al., 2003; Hassing and Mariani, unpublished data; Levine, 2000). Because Topo III was one of the suppressors isolated, we considered the possibility that it played a role in the suppression by the other suppressor genes identified. To test this possibility, we constructed (Table S1) the W3110*parE10* Δ *topB::kan* (CL002) and C600*parC1215* Δ *topB::kan* (BP079) strains and their respective isogenic parents, W3110*topB::kan* (BP093) and C600*topB::kan* (BP105) and transformed W3110*parE10* Δ *topB::kan* and W3110*topB::kan* with the IPTG-regulated overexpression plasmids pLEX5BA-*dnaX*(γ), which produces only the truncated γ gene product, and pLEX5BA-*topB* (pBP42).

Overexpression of γ in W3110*parE10* Δ *topB::kan* failed to bypass the growth impediment at 37 °C and 42 °C under any growth condition tested (even slow-growth conditions on nutrient-limited media) (Fig. 1A). Overexpression of SetB and τ also failed to suppress the growth defect displayed by W3110*parE10* Δ *topB::kan* at 37 °C and 42 °C (data not shown). As expected, overexpression of *topB* suppresses lethality of the W3110*parE10* Δ *topB::kan* mutant at 42 °C (Fig. 2). Because of the severe growth defect of C600*parC1215* Δ *topB::kan*, discussed below, we were unable to perform similar suppression tests on this mutant. These results imply that the mechanism of suppression of the Topo IV ts mutations by *dnaX*(γ) and possibly the other suppressors is either through direct stimulation of Topo III or that Topo III provides either a supporting or backup function to Topo IV during chromosome decatenation. As to the former possibility, however, we have been unable to detect any stimulation of Topo III activity by either DnaX or DnaN in our assays *in vitro* [(Levine, 2000) and data not shown].

The catalytic activity of Topo III is required for suppression

To determine if the suppression of the *parE10* mutation at non-permissive temperatures is specific to Topo III, we tested whether the other type I topoisomerase present in *E. coli*, Topo I, could also rescue W3110*parE10* at 42 °C when overproduced. This proved not to be the case. Whereas overexpression of Topo I [from pLEX5BA-*topA* (pBP59)] was not toxic to either parental strain (Fig. S2), it could not rescue growth of either W3110*parE10* or C600*parC1215* at 42 °C (Fig. 3). Therefore, Topo III is the only topoisomerase capable of suppressing a Topo IV deficiency efficiently. [Overexpression of DNA gyrase results in incomplete suppression (Kato et al., 1992).]

If Topo III was stimulating Topo IV directly under suppression conditions, one would expect that a catalytically-inactive *topB* allele would work as a suppressor. Using targeted mutagenesis, we replaced the catalytic Tyr with Phe (*topBY328F*) (Zhang et al., 1995, Harmon et al., 1999) in the *topB* ORF in pLEX5BA-*topB*, generating pLEX5BA-*topBY328F* (pBP47). Unlike wild-type Topo III, overexpression of this inactive mutant of Topo III (TopBY328F) did not enable either W3110*parE10* or C600*parC1215* to grow at 42 °C (Fig. 3). [The level of expression of both the wild-type and mutant Topo III was identical (Fig. S3).] These data suggest that the enzymatic activity of Topo III, likely the decatenase function, is important for chromosome segregation under suppression conditions.

Topo III and γ can suppress the *parE10* mutation in the absence of either *RecQ* or *RecA*

Because of the functional relationship between Topo III and *RecQ*, and a possible role for Topo III in recombination, we explored a potential function for *recQ* and *recA* in the Topo

III-dependent rescue of W3110*parE10* growth at 42 °C. We showed previously that *recQ* was not required for rescue of W3110*parE10* by overexpression of *topB* [Fig. 4A and (Nurse et al., 2003)]. However, it was possible that the high levels of Topo III present under those conditions could have obscured a *recQ* requirement. Therefore, because *topB* is essential for rescue of W3110*parE10* by the other suppressors, we asked if either *recQ* or *recA* were necessary for *topB*-dependent suppression of W3110*parE10* temperature sensitivity by overexpression of *dnaX*(γ). Interestingly, we found that overexpression of *dnaX*(γ) could bypass the Topo IV defect in cells lacking either *recQ* (Fig. 1A), suggesting that concerted action of RecQ and Topo III is not required for suppression, or *recA* (Fig. 1B), suggesting that recombination in general was not required for suppression. Similarly, a disruption in *recA* did not affect the ability of overproduced Topo III to suppress the growth defect of W3110*parE10* at 42 °C (Fig. 4), suggesting that RecA-dependent recombination is not required for rescue of W3110*parE10* at restrictive temperatures by Topo III. Interestingly, removal of RecQ from W3110*parE10* made rescue by γ overexpression more efficient at 42 °C (Fig. 1A). The reason for this effect is unclear.

Deletion of *topB* in Topo IV mutants leads to pronounced chromosome segregation defects at permissive temperatures

The inability of the suppressors (excluding *topB*) to rescue the growth defect of W3110*parE10* Δ *topB::kan* and the failure to detect direct stimulation of Topo III activity by any of the suppressor proteins *in vitro*, strongly suggested that Topo III was directly involved in chromosome segregation by resolving topological linkages between daughter molecules. We therefore investigated whether Topo III was required for growth of the Topo IV ts mutants at permissive temperatures. In a previous report (Lopez et al., 2005), a synthetic lethal effect between the *parE10* and Δ *topB* mutations was observed as a failure to obtain at 37 °C, bacteriophage P1-mediated co-transduction into W3110*parE10* of Δ *topB* when a drug marker 50 kbp away from *topB* was selected, whereas at 30 °C the double mutant strain could be recovered. These authors also reported that even at 30 °C, the C600*parC1215* Δ *topB* double mutant strain could not be constructed by similar means (Lopez et al., 2005). However, we were able to construct both the C600*parC1215* Δ *topB::kan* and W3110*parE10* Δ *topB::kan* double mutant strains by growing all P1 recombinants at 25 °C. In agreement with the previous report (Lopez et al., 2005), the *parE10* and Δ *topB::kan* mutations are synthetically lethal at 37 °C (Figs. 5 and 6), and the double mutant strain displayed a significant growth impairment at 30 °C compared to W3110*parE10* (Fig. 5N). In addition, the C600*parC1215* Δ *topB::kan* double mutant strain was severely compromised, failing to grow at 30 °C (Fig. 5M) and higher temperatures (data not shown). Ectopic expression of either Topo III or Topo IV from pLEX5BA permits growth of W3110*parE10* Δ *topB::kan* at restrictive temperatures (Fig. 2 and Fig. S4).

Cellular and nucleoid morphologies of the double mutant strains and their parents were examined. As expected, C600 and C600 Δ *topB::kan* cells showed no obvious defects (Figs. S5A and S5B) and had normal cell lengths (Fig. S5G) after 4 h of growth at 25 °C. Similarly, C600*parC1215* cells also appeared mostly normal, with a small fraction of filamentous cells (Figs. 5A and 5I). Surprisingly, deletion of *topB* in C600*parC1215* resulted in pronounced chromosome segregation defects, even at the favorable temperature of 25 °C. These cells exhibited large masses of unsegregated DNA and were filamentous (Figs. 5B and 5I). Likewise, after 2 h of growth at 30 °C, W3110, W3110*topB::kan*, and W3110*parE10* cells had normal cell sizes (Figs. S5H and S5J) and W3110 and W3110 Δ *topB::kan* cells had normal nucleoid morphology (Figs. S5C, S5D). The majority of the nucleoids in W3110*parE10* cells at 30 °C were normal (Fig. 5C). About one-quarter of the cells displayed a phenotype that we refer to as “nascent par,” where an unsegregated mass of DNA was centrally located in a normal-sized cell (Figs. 5C and 5J). This population

of nascent *par* cells doubled in the W3110*parE10ΔtopB::kan* double mutant strain. In addition, a significant population of anucleate cells, a phenotype often associated with chromosome segregation defects, was also observed in the double mutant (Figs. 5D and 5J).

The effect of the *ΔtopB* mutation could also be observed at the nonpermissive temperature of 42 °C in the W3110*parE10* strain. W3110 and W3110*ΔtopB::kan* cells grown for 2 h at 30 °C and then shifted to 42 °C for 1 h appeared wild-type (Fig. S5E and S5F) with average cell lengths (Fig. S5I and S5J). Although the *par* phenotype was obviously present in W3110*parE10* cells grown at 42 °C, the W3110*parE10ΔtopB::kan* cells were consistently more filamentous, with the fraction of cells >15 μm in length increasing by more than two-fold (Fig. 5K) and an increased appearance of anucleate cells (Fig. 5E and 5F). These results indicate that the growth defect of the W3110*parE10ΔtopB::kan* and C600*parC1215ΔtopB::kan* mutant strains is linked to a chromosome segregation defect.

To address whether the *ΔtopB*-mediated synthetic effect on chromosome segregation was specific to the Topo IV *ts* alleles, we tested for a synthetic relationship between the DNA gyrase *ts* allele *gyrBR136C P171S* [from strain N4177, *Cou^r ts* (Oram *et al.*, 1992)] and the *ΔtopB* mutation. Both the W3110*gyrBR136CP171S* (BP199) and W3110*gyrBR136CP171S ΔtopB::kan* (BP275) strains grew equally well on plates incubated at 30 °C and 37 °C, and, as anticipated, were less viable at 42 °C (Fig. S6D), although the W3110*gyrB^{ts}* strain was slightly more viable than the W3110*gyrB^{ts} ΔtopB::kan* strain at this temperature (Fig. S6D). Both W3110*gyrB^{ts}* and W3110*gyrB^{ts} ΔtopB::kan* cells cultured for 2 h at 30 °C showed no obvious cell or nucleoid morphology defects (Fig. S6A-C), whereas, after 1 h at 42 °C both strains amassed large unsegregated nucleoids and failed to divide (Fig. 5G, 5H, and 5L). Although W3110*gyrB^{ts} ΔtopB::kan* cells were slightly longer than W3110*gyrB^{ts}* cells at 42 °C (7.9 μm compared to 7.2 μm, Fig. 5L), it was clear that the *ΔtopB::kan* mutation had, at most, a minor effect on growth and chromosome segregation in the W3110*gyrB^{ts}* mutant strain. Thus, we conclude that the synthetic lethality of Topo IV *ts* mutations and the *ΔtopB::kan* mutation is specific to defects in Topo IV and Topo III.

The chromosome segregation defect and synthetic lethality of the *parE10ΔtopB::kan* double mutations does not arise from the failure to resolve toxic recombination intermediates

The Topo III and RecQ combination is clearly involved in the resolution of toxic recombination intermediates in eukaryotes by the dissolution of double Holliday junctions (Wu & Hickson, 2003). This has prompted others to investigate whether a similar paradigm holds in prokaryotes. For example, Lopez *et al.* (Lopez *et al.*, 2005) reported that removal of RecQ suppressed the synthetic lethality of the *parE10ΔtopB::kan* double mutation. Because we found that neither RecQ nor RecA was required for suppression of W3110*parE10* by overexpression of *dnaX(γ)* (Fig. 1), we tested whether the observed DNA segregation block and growth deficiency of the W3110*parE10ΔtopB::kan* strain could be alleviated by the introduction of either *ΔrecQ* or *recA938::cat*, a *recA* null mutation.

Comparison by microscopy showed that the lack of RecQ did not alleviate the chromosome segregation defect of W3110*parE10ΔtopB::kan* cells grown at 30 °C (Figs. 6B, and 6E). In addition, introduction of a *ΔrecQ* mutation to the W3110*parE10* strain did not phenocopy the effect of introducing *ΔtopB* to the W3110*parE10* strain (Figs. 6A and 6E). This observation indicates that linkage of Topo III and RecQ activity, as demonstrated to be required to resolve some forms of recombination intermediates (Oakley & Hickson, 2002), is not responsible for the defects in the W3110*parE10ΔtopB* strain. Similar results were observed when these strains were incubated at 30 °C for 2 h and then shifted to 42 °C for 1 h (Fig. S7), although the W3110*parE10ΔtopB::kanΔrecQ::cat* strain (CL015) showed a decrease in the proportion of very long filaments and an increase in the proportion of

anucleate cells compared to the W3110 *parE10ΔtopB::kan* strain (Figs. S7B, S7D, and S7E). Furthermore, the *ΔrecQ* mutation did not affect the viability of these strains at either 30 °C or 37 °C and did not rescue the synthetic lethality of the *parE10ΔtopB::kan* double mutations at the semipermissive temperature of 37 °C (Fig. 6F). Control experiments (Fig. S8) showed that the *ΔrecQ* mutation did not affect either cell or nucleoid morphology at either 30 °C or 42 °C or cell viability at either 30 °C or 37 °C, either in the presence or absence of Topo III in the parental W3110 strain (Figs. S8A, S8B, S8E, S8F, and S8I).

A general decrease in viability for all *recA938::cat* mutant strains, which is characteristic of mutants of *recA* (Fig. 6G and Fig. S8J), was observed. However, the presence of the *recA938::cat* mutation did not suppress the growth defect (Fig. 6G) and exacerbated the chromosome segregation defect of W3110 *parE10ΔtopB::kan*, with a dramatic increase in the population of anucleate cells and the appearance of a population of cells in which the nucleoid had become fragmented (Figs. 6C, 6D, 6E, 6G, and S7C, S7F, and S7G). The lack of a requirement for either RecQ or RecA for the *parE10ΔtopB* synthetic lethality argues strongly that the phenotype arises from a chromosome segregation defect and not the failure to resolve toxic recombination intermediates. Given that we rederived the W3110 *parE* strain, it is possible that the relief of synthetic lethality between the *parE10* and *ΔtopB* mutations afforded by mutations in *recQ* and *recA* observed by Lopez et al. (Lopez et al., 2005) is a result of unknown mutations in their strains. Indeed, our original lab stock of W3110 *parE10* filaments more extensively than the rederived strain at 42 °C.

Topoisomerase III mutants are sensitive to novobiocin

Our data supports the notion that Topo III provides either a supportive or backup role to Topo IV in the resolution of catenated daughter chromosomes. If so, then mutants of *topB* should be hypersensitive to drugs that block chromosome segregation and/or Topo IV. We therefore tested if *topB* mutants were sensitive to novobiocin, which targets the type II topoisomerases in *E. coli* (Gellert et al., 1976b, Mizuuchi et al., 1978, Levine et al., 1998). This proved to be the case.

The W3110 *ΔtopB::kan* mutant strain was approximately four times more sensitive to novobiocin than *topB⁺* strains. The wild-type W3110 strain had a MIC₅₀ of approximately 200 μg/ml of novobiocin, whereas the W3110 *ΔtopB::kan* strain had a MIC₅₀ of nearly 50 μg/ml. In fact, the W3110 *ΔtopB::kan* mutant strain failed to grow in the presence of 100 μg/ml of novobiocin (Fig. 7C). In control experiments using a superhelical DNA relaxation assay (Fig. S9), we confirmed that Topo III is unaffected by novobiocin (Srivenugopal et al., 1984).

To determine which type II topoisomerase was mediating the observed increased sensitivity to novobiocin, we examined drug response in the presence of a novobiocin-resistant allele of the GyrB subunit of DNA gyrase (the W3110 *gyrB^{ts}* strain (BP199) used in Fig. 5 is also novobiocin resistant). As expected, the W3110 *gyrB^{ts}* strain was overall more resistant to novobiocin (MIC₅₀ of ~ 750 μg/ml), but the W3110 *gyrB^{ts}ΔtopB::kan* mutant was still nearly two-fold more sensitive to the drug (MIC₅₀ of ~ 400 μg/ml) (Fig. 7D). Furthermore, overexpression of *topB* in the W3110 *gyrB^{ts}* strain conferred a growth advantage at high concentrations of novobiocin (Fig. S10). These data indicate that an increased demand for Topo IV was the cause of the increased sensitivity to novobiocin conferred by the *ΔtopB* allele and that overexpression of *topB* in the W3110 *gyrB^{ts}* strain relieved demand for Topo IV activity. If this is the case, we would expect to observe chromosome segregation defects in the W3110 *gyrB^{ts}ΔtopB::kan* strain in the presence of moderate levels of the drug.

After a 1.5-h incubation with 300 μg/ml of novobiocin, W3110 *gyrB^{ts}ΔtopB::kan* cells were noticeably more filamentous (average cell length 6.5 μm) compared to W3110 *gyrB^{ts}* cells

(average cell length 5.0 μm) (Figs. 7A, 7B and 7E). Additionally, W3110 *gyrB^{ts} Δ topB::kan* cells had accumulated larger unsegregated nucleoid masses (average area 3.1 μm^2) compared to W3110 *gyrB^{ts}* cells (average area of 1.3 μm^2) (Figs. 7A, 7B, and 7F), indicating that novobiocin causes a significant block in chromosome segregation in *topB* mutant strains.

If the observed novobiocin sensitivity of the *topB* mutant strains was indeed a result of increased challenge for Topo IV, then overexpression of Topo IV should result in bypass of this sensitivity. This proved to be the case. As expected, W3110 *Δ topB::kan* and W3110 *gyrB^{ts} Δ topB::kan* strains failed to grow as well as either the wildtype or *gyrB^{ts}* strains on novobiocin plates (Fig. 7G). However, overexpression of either Topo III or Topo IV allowed all *Δ topB::kan* mutant strains to grow (Fig. 7H). Hence, expression of Topo III *in trans* can complement the *Δ topB::kan* mutation for novobiocin hypersensitivity, as expected, while overproduction of Topo IV can bypass *topB*-mediated novobiocin sensitivity, suggesting that Topo III is important for chromosome segregation, and its requirement can be bypassed by producing more Topo IV.

Discussion

Topo III suppresses a Topo IV deficiency by decatenating chromosomes

To date, there has been no observed *topB* mutant phenotype in prokaryotes that has led to the elucidation of the function of Topo III in the cell. Our previously identified high-copy suppressors that rescue the temperature sensitivity of *parE* and *parC* mutant strains (Levine & Mariani, 1998, Espeli et al., 2003b, Nurse et al., 2003) have allowed us to dissect the role of Topo III in the cell. The *topB* gene was essential for the other previously identified high-copy suppressors including, *dnaX(γ)* (Fig. 1A), *dnaX(τ)*, and *setB* to rescue the W3110 *parE10* strain at restrictive temperatures. Based on these findings, we thought that the mechanism of suppression by the high-copy suppressors could be through a direct stimulation of the enzymatic activities of either Topo IV or Topo III. Although we have conducted numerous biochemical and protein-protein interaction studies to test this theory, we have been unable to support this idea. Alternatively, it was possible that the high-copy suppressors (with the exception of *topB*) worked in an independent fashion, as suggested by others (Grainge et al., 2007), and that Topo III acted in a supportive capacity to Topo IV by actively decatenating chromosomes.

To further support the notion that the decatenation activity of Topo III was essential for chromosome segregation and for the rescue of the W3110 *parE10* strain at restrictive temperatures, we analyzed the requirement for the catalytic activity of Topo III for suppression of the W3110 *parE10* growth defect at nonpermissive temperatures. We report that Topo III is the only topoisomerase in the cell that can efficiently rescue either W3110 *parE10* or C600 *parC1215* at the restrictive temperature. Overexpression of DNA gyrase has been shown to only partially rescue temperature-sensitive mutations of *parE* and *parC* (Kato et al., 1992), and we show here that overproduction of Topo I completely failed to rescue W3110 *parE10* at 42 °C. In addition, a catalytically dead mutant of *topB*, when overproduced, is unable to support suppression of either the *parE10* or *parC1215* mutations at 42 °C, suggesting that Topo III's decatenation function is indeed critical for rescue of the W3110 *parE10* and C600 *parC1215* strains at nonpermissive temperatures.

Even at permissive temperatures, in the absence of *topB*, W3110 *parE10* and C600 *parC1215* were not only compromised for viability, compared to their *topB⁺* counterparts, but had pronounced chromosome and cell division defects consistent with a severe *par* phenotype. The severity of the defect is emphasized by the fact that even when W3110 *parE10 Δ topB::kan* cells are grown at 30 °C, a significant proportion (average of

24%) of the cells in the population die, whereas mortality frequencies under the same growth conditions for W3110*parE10* were <1%. Moreover, the *par* phenotype displayed by the C600*parC1215ΔtopB::kan* mutant strain was consistently more severe than that of W3110*parE10ΔtopB::kan*, suggesting that the *par* phenotype in *topB* mutant strains worsen with increasing loss of Topo IV activity. These observations indicate that the well-studied W3110*parE10* and C600*parC1215* strains depend on Topo III for survival. Even when these *parE* and *parC* ts mutant strains are grown 30 °C, the Topo IV enzyme produced in either W3110*parE10* and C600*parC1215* mutant strains is inadequate to carry out efficient decatenation reactions for the timely DNA segregation that is critical for productive cell division and growth. Although the chromosomal levels of Topo III are significantly lower than that of the other DNA topoisomerases (DiGate & Mariani, 1989), Topo III's strong decatenase activity (DiGate & Mariani, 1988) is likely the key to its ability to partially substitute for a Topo IV defect in the cell.

The topB mutant phenotype is not the result of failure to resolve recombination intermediates

Numerous studies have shown biochemical and genetic associations between Topo III, RecQ, and cellular processes such as the resolution of converging replication fork complexes (Suski & Mariani, 2008) and the dissolution of double Holliday junctions (Mankouri & Hickson, 2007). Using our suppressor-based genetic assay, we demonstrated that *recQ* was not required for either Topo III or γ (when in excess) to rescue the W3110*parE10* strain. Rescue of W3110*parE10ΔrecQ* (CL004) by γ was Topo III-dependent, demonstrating that RecQ was not required for endogenous levels of Topo III to suppress the Topo IV deficiency and for successful DNA segregation. Consistent with these findings, there were no compounded effects on viability or chromosome segregation in W3110*parE10* strains with *ΔrecQ* mutations at either permissive or nonpermissive temperatures. Therefore, Topo III does not require its cognate helicase to unlink daughter chromosomes and to rescue W3110*parE10*, and the genetic evidence suggests that its substrate in vivo could be precatenanes.

Previous observations (Lopez et al., 2005) have suggested that the synthetic effect on viability of the *parE10ΔtopB* and *parC1215ΔtopB* double mutations was because of the accumulation of recombination intermediates, and that either introducing a *recQ* mutation or abolishing recombination (with a *recA* mutation) could suppress the described synthetic lethality. The authors used their observations to support a role for Topo III in recombinational repair of DNA lesions. This model is inconsistent with our findings.

Based on our biochemical assays, suppression of W3110*parE10* and C600*parC1215* temperature sensitivity by overexpression of *dnaX*(γ) and *setB* is unlikely to occur by direct stimulation of Topo III, we would then expect overexpression of *dnaX*(γ) or the other high-copy suppressors to suppress the formation of such toxic recombination intermediates in W3110*parE10ΔtopB*. Thus, failure of overexpression of *dnaX*(γ) to rescue W3110*parE10ΔtopB* is consistent with a role in chromosome segregation for Topo III. Of course, it is also possible that the high-copy suppressors potentiate Topo III-catalyzed decatenation of the daughter chromosomes in a manner that is undetected by our biochemical assays. However, we did not observe any differences in either growth or state of the nucleoids between W3110*parE10ΔtopB* or W3110*parE10ΔtopBΔrecQ*: both strains failed to grow at semipermissive temperatures and to segregate chromosomes properly. Overexpression of *dnaX*(γ) also failed to suppress the *parE10ΔtopBΔrecQ* triple mutation combination at either 37 °C or 42 °C. In addition, the pronounced chromosome segregation defects in the W3110*parE10ΔtopB* and C600*parC1215ΔtopB* mutant strains can be reversed by overexpression of Topo IV.

Finally, abolishing *recA*-dependent recombination had no effect on the described *parE10ΔtopB* phenotype of growth, rescue by high-copy suppressors, or chromosome segregation. RecA was not required for the *parE10ΔtopB* phenotype, and introduction of a *recA* null mutation neither permitted growth of W3110*parE10ΔtopB* nor alleviated the chromosome segregation block. Therefore, we conclude that the *parE10ΔtopB* synthetic lethality arises from a chromosome segregation problem, rather than because of the accumulation of toxic recombination intermediates.

Topo III participates in chromosome segregation

The findings in this study provide strong evidence that Topo III participates in chromosome segregation. To address whether Topo III was essential for chromosome segregation, we asked if *topB* mutants were more susceptible to agents that target Topo IV and block chromosome segregation. Mutants of *topB* treated with novobiocin were unable to grow in the presence of novobiocin at concentrations tolerated by *topB*⁺ strains, and these novobiocin-treated cells displayed aberrant chromosome segregation. This finding suggests that *topB* mutants are novobiocin sensitive because they cannot properly segregate their chromosomes in the presence of drug. The novobiocin sensitivity of the W3110*ΔtopB* mutant strain is the first identified phenotype of *topB* mutations that supports a clear role and requirement for Topo III in chromosome segregation when Topo IV activity is reduced, as herein by treatment with novobiocin. Furthermore, these observations are consistent with the phenotype displayed by the *parE10ΔtopB* double mutation. We consider it unlikely that enervation of Topo IV creates a new activity for Topo III and therefore suggest that the activities of both Topo IV and Topo III are necessary for segregation of the chromosomes in *E. coli*. The novobiocin sensitivity of the *topB* mutation is specific to a defect in chromosome segregation because overexpression of either Topo III or Topo IV can alleviate the novobiocin sensitivity of *topB* mutant strains, suggesting that an excess of Topo IV can suppress the requirement for Topo III. Moreover, the introduction of novobiocin-resistant alleles of *gyrB* did not suppress the novobiocin sensitivity of *topB* mutant strains.

Because Topo III rescues W3110*parE10* without the assistance of RecQ and in the absence of recombination, and requires single-stranded DNA gaps to decatenate, it is probable that its mechanism of action occurs during DNA replication by resolving precatenanes (Fig. 8). Therefore, we envision the action of three topoisomerases as being necessary during DNA replication to ensure complete topological disengagement of the parental strands: DNA gyrase acts ahead of the replication fork, removing positive supercoils. Any supercoils that diffuse behind the fork, leading to the formation of precatenanes, are resolved by Topo III prior to ligation of the Okazaki fragments. And catenanes that do form are resolved by Topo IV.

Because Topo IV and Topo III are the main decatenating enzymes, it is clear why the synthetic effect of the *topB* mutation on viability and chromosome segregation is specific to the *parE* and *parC* alleles. Combining the *gyrB* allele with a *topB* deletion mutation has only minor consequences for the cell because Topo IV is available to resolve the topological challenges. For decatenation, it is clear that Topo IV is the workhorse of the cell, because its function cannot be completely replaced by either DNA gyrase or Topo III. This could either be because of a persistent burden of catenanes that is too great of a challenge for either Topo III or DNA gyrase, or an additional essential role for Topo IV in other unrelated functions.

In consideration of our previous findings that Topo IV activity is temporally regulated and functions specifically at the later end stages of DNA replication (Espeli *et al.*, 2003a), one could predict that W3110*ΔtopB* mutant strains should have a delay in chromosome segregation. However, the interplay between Topo IV and Topo III and factors governing their regulation has not been explored. It is possible that the cell regulates the activities of

Topo III and Topo IV with respect to the demand for decatenation as it does the activities of DNA gyrase and Topo I to ensure the proper superhelical state of the DNA (Menzel and Gellert, 1983). Interestingly, reduced gene expression of *parE* in *Staphylococcus aureus* leads to an increase in *topB* transcript levels (Ince & Hooper, 2003).

Experimental procedures

Bacterial strains and growth conditions

E. coli strains are described in Table S1 and were cultured in either LB, Minimal A (Min A), or M9 medium supplemented with thiamine (40 µg/ml), threonine (40 µg/ml), and leucine (40 µg/ml). The following concentrations of antibiotics were used: ampicillin (Ap), 100 µg/ml; carbenicillin (Cb), 50 µg/ml; kanamycin (Km), 50 µg/ml; chloramphenicol (Cm), 34 µg/ml; novobiocin (Nov), 600 µg/ml. Unless indicated otherwise, wild-type strains were incubated at 37 °C, whereas strains with *ts* mutations were incubated at 30 °C. W3110 is the parental strain of W3110*parE10* (Kato et al., 1990) and C600 is the parental strain of C600*parC1215* (EJ812) (Kato et al., 1988).

DNA procedures

Plasmids (Table S2) were constructed by standard molecular genetic procedures. All plasmid preparations were done using Miniprep or Midiprep Spin Kits (Qiagen). DNA manipulations with restriction endonucleases and bacteriophage T4 DNA ligase were done according to the manufacturer's (NEB) recommendations. PCR amplifications were done with high fidelity ExTaq polymerase (TaKaRa), KOD HiFi polymerase (Novagen), or Phusion DNA polymerase (NEB) for 30 cycles, and all primers are listed in Table S3. To clone specific genes, either C600 or W3110 genomic DNA was used as template for PCR. The QiaexII kit (Qiagen) was used for agarose gel extraction. PCR products and constructed plasmids were confirmed by nucleotide sequencing at the MSKCC DNA Sequencing Facility.

Rederivation of the *parE10* and *parC1215* mutations in W3110

The technique of Link et al. was used (Link *et al.*, 1997).

pK03-*parE10*: Nucleotides 3171470-3173470 were amplified from genomic DNA prepared from our original lab stock of W3110*parE10* using primers that introduced *Bgl*III and *Sal*I restriction enzyme sites at the 5' and 3' ends of the amplified DNA fragment, respectively (*parE10*top and *parE10*bot, Table S3). This DNA fragment had 1 kbp of chromosomal DNA flanking either side of the *parE10* mutation. This DNA fragment was digested with these restriction enzymes and ligated to similarly digested pK03 DNA (Link et al., 1997) to give pK03-*parE10*.

pRC7-*parE*: The *parE* ORF was amplified from genomic DNA prepared from W3110 using primers that introduced *Eco*RI and *Hind*III restriction enzyme sites to the 5' and 3' ends of the amplified DNA fragment, respectively (*parE*top and *parE*bot, Table S3). This DNA fragment was digested with these restriction enzymes and ligated to similarly digested pRC7 DNA (de Boer *et al.*, 1989) to give pRC7-*parE*. This plasmid expresses an IPTG-inducible LacZ-ParE fusion and is lost very quickly from cells in the absence of ampicillin.

W3110 cells were transformed with pK03-*parE* by electroporation and selected on LB-Cm plates at 30 °C. A single transformant was grown in LB-Cm medium at 30 °C until mid-log, serially diluted in LB, plated on pre-warmed LB-Cm plates, and grown at 43 °C. A single co-integrate was picked from this plate, grown to mid-log and maintained in LB-Cm at 43 °C. These cells were transformed with pRC7-*parE* and plated on LB-Cm-Ap plates at 43 °C.

Colonies confirmed to have both plasmids established were then grown on LB plates supplemented with Cm, Ap, and 100 μ M IPTG to induce expression of the LacZ-ParE fusion protein. Several of the resulting colonies were dispersed into 1 ml of LB medium and cells were streaked on LB plates supplemented with Ap, 100 μ M IPTG, and 5% sucrose and grown at 30 °C. Clones were first checked for loss of pKO3-*parE10* by replica plating onto LB-Cm-Ap plates at 30 °C. Ap-resistant, Cm-sensitive clones were then replica plated onto LB-IPTG plates at 30 °C to check for loss of pRC7-*parE*. The resulting clones were checked for temperature sensitivity, as well as ensuring the loss of pRC7-*parE*, by growth on LB plates at 43 °C.

For the *parC1215* mutation the methodology was the same except that pKO3-*parC1215* carried nucleotides 3160813-3162822 amplified from chromosomal DNA of C600*parC1215* (EJ812) and pRC7-*parC* carried the ParC ORF amplified from W3110 genomic DNA and the primers used to introduce the ParC ORF to pRC7 added *SaI* and *HindIII* restriction sites to the PCR product.

Construction of the *TB28ΔtopB::FRTkan mutant*

The *ΔtopB::FRTkan* mutation was constructed by linear replacement (Datsenko & Wanner, 2000). FRT sites flank the *ΔtopB::kan* cassette and successful recombination of the allele resulted in the replacement of the *topB* ORF. The primers DtopB-1 and DtopB-2 (Table S3) were used to PCR amplify the *ΔtopB::FRTkan* cassette. The PCR fragments were transformed into TB28 as described previously (Datsenko & Wanner, 2000) and recombinants were selected in LB supplemented with Km. The *ΔtopB::FRTkan* mutation was verified by PCR and DNA sequencing.

Bacteriophage P1 transduction of mutant alleles

The following strains: TB28 *ΔtopB::FRTkan*, CRL31 (*ΔrecQ::FRTcat*), SS391 (*recA938::cat*), and N4177 (*gyrBR136C P171S*; *Cou^r ts*) served as P1 donors for the transduction of the *ΔtopB::FRTkan*, *ΔrecQ::FRTcat*, *recA938::cat*, and *gyrB^{ts}* alleles into strains described in Table S1.

Microscopy

Overnight cultures grown in LB were diluted to an OD₆₀₀ of 0.005 in LB medium and grown at the indicated temperatures. At the indicated times, cells were harvested by centrifugation at 1,500 × g for 10 min. Pellets were resuspended in 1 × PBS. To stain the cell membranes and DNA, FM4-64FX (Molecular Probes) at 1 μ g/ml and DAPI (Molecular Probes) at 1 μ g/ml, respectively, were added to the cell suspension. The suspensions were spread onto slides coated with 0.1% poly-L-lysine (Sigma). Cells were fixed by overlaying the slides with 1 ml of 2.5% paraformaldehyde, 0.03% glutaraldehyde, 1 × PBS for 30 min. Slides were washed three times with 1 × PBS and allowed to dry at room temperature. The cells were visualized with a Nikon Eclipse Ti fluorescence microscope. Nikon Elements software was used to determine cell length and nucleoid area; a minimum of 300 cells and 500 nucleoids, respectively, were measured.

Suppressor analysis

Strains were grown overnight in LB supplemented with Ap at 30 °C. The saturated cultures were then serially diluted and spotted (10 μ l) on either LB or Min A plates containing Ap and 100 μ M IPTG. Plates were incubated at 30 °C and 37 °C and/or 42 °C.

Novobiocin sensitivity plating assays

Cultures were grown overnight in LB at 37 °C for W3110 strains and 30 °C for *gyrBts^{Nov}* (BP199) strains. The cultures were diluted to an OD₆₀₀ of 0.001 in fresh LB, grown for 1 h at the appropriate temperature, and plated on LB plates containing varying concentrations of novobiocin. W3110 strains were incubated at 37 °C overnight, whereas *gyrBts^{Nov}* strains were incubated at 30 °C for approximately 30 h. Survival was determined by the following: average CFUs on novobiocin plates/ average CFUs on LB only plates. MIC₅₀ is defined as the minimum inhibitory concentration: The concentration of drug that is required to inhibit 50% of growth.

Topo III and Topo IV bypass of novobiocin sensitivity

W3110, W3110 Δ *topB::kan* (BP093), W3110*gyrBts^{Nov}* (BP199), and W3110*gyrBts^{Nov}* Δ *topB::kan* (BP275) were transformed with pLEX5BA, pLEX5BA-*parEparC*, and pLEX5BA-*topB* (pBP42). Transformants were grown and diluted back as described above. Prior to plating, strains were grown in LB with 100 μ M IPTG at 37 °C (for W3110 strains) or 30 °C (*gyrBts^{Nov}* strains) for 2 h. Cultures were titrated and spotted on LB plates supplemented with 100 μ M IPTG and novobiocin (50 μ g/ml for W3110 strains and 300 μ g/ml for *gyrBts^{Nov}* strains).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Olivier Espeli, John Petrini, and Rod Rothstein for comments on the manuscript, and Catherine Suski-Grabowski for strains. These studies were supported by NIH grant GM34558.

References

- Adams DE, Shekhtman EM, Zechiedrich EL, Schmid MB, Cozzarelli NR. The role of topoisomerase IV in partitioning bacterial replicons and the structure of catenated intermediates in DNA replication. *Cell*. 1992; 71:277–288. [PubMed: 1330320]
- Bates, BD.; Maxwell, A. DNA Topology. Oxford University Press Inc.; New York: 2005.
- Champoux JJ. DNA topoisomerases: structure, function, and mechanism. *Annu Rev Biochem*. 2001; 70:369–413. [PubMed: 11395412]
- Champoux, JJ.; Been, M. Topoisomerases and the Swivel Problem. In: Alberts, B., editor. *Mechanistic Studies of DNA Replication and Genetic Recombination*. Academic Press; New York: 1980. p. 809-815.
- Datsenko KA, Wanner BL. One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products. *Proc Natl Acad Sci U S A*. 2000; 97:6640–6645. [PubMed: 10829079]
- de Boer PA, Crossley RE, Rothfield LI. A division inhibitor and a topological specificity factor coded for by the minicell locus determine proper placement of the division septum in *E. coli*. *Cell*. 1989; 56:641–649. [PubMed: 2645057]
- DiGate RJ, Mariani KJ. Identification of a potent decatenating enzyme from *Escherichia coli*. *J Biol Chem*. 1988; 263:13366–13373. [PubMed: 2843517]
- DiGate RJ, Mariani KJ. Molecular cloning and DNA sequence analysis of *Escherichia coli* topB, the gene encoding topoisomerase III. *J Biol Chem*. 1989; 264:17924–17930. [PubMed: 2553698]
- Espeli O, Levine C, Hassing H, Mariani KJ. Temporal regulation of topoisomerase IV activity in *E. coli*. *Mol Cell*. 2003a; 11:189–201. [PubMed: 12535532]
- Espeli O, Nurse P, Levine C, Lee C, Mariani KJ. SetB: an integral membrane protein that affects chromosome segregation in *Escherichia coli*. *Mol Microbiol*. 2003b; 50:495–509. [PubMed: 14617174]

- Gellert M, Mizuuchi K, O'Dea MH, Nash HA. DNA gyrase: an enzyme that introduces superhelical turns into DNA. *Proc Natl Acad Sci U S A*. 1976a; 73:3872–3876. [PubMed: 186775]
- Gellert M, O'Dea MH, Itoh T, Tomizawa J. Novobiocin and coumermycin inhibit DNA supercoiling catalyzed by DNA gyrase. *Proc Natl Acad Sci U S A*. 1976b; 73:4474–4478. [PubMed: 794878]
- Grainge I, Bregu M, Vazquez M, Sivanathan V, Ip SC, Sherratt DJ. Unlinking chromosome catenanes in vivo by site-specific recombination. *EMBO J*. 2007; 26:4228–4238. [PubMed: 17805344]
- Harmon FG, DiGate RJ, Kowalczykowski SC. RecQ helicase and topoisomerase III comprise a novel DNA strand passage function: a conserved mechanism for control of DNA recombination. *Mol Cell*. 1999; 3:611–620. [PubMed: 10360177]
- Hiasa H, Marians KJ. Topoisomerase III, but not topoisomerase I, can support nascent chain elongation during theta-type DNA replication. *J Biol Chem*. 1994; 269:32655–32659. [PubMed: 7798272]
- Ince D, Hooper DC. Quinolone resistance due to reduced target enzyme expression. *J Bacteriol*. 2003; 185:6883–6892. [PubMed: 14617652]
- Kato J, Nishimura Y, Imamura R, Niki H, Hiraga S, Suzuki H. New topoisomerase essential for chromosome segregation in *E. coli*. *Cell*. 1990; 63:393–404. [PubMed: 2170028]
- Kato J, Nishimura Y, Suzuki H. *Escherichia coli* parA is an allele of the *gyrB* gene. *Mol Gen Genet*. 1989; 217:178–181. [PubMed: 2549366]
- Kato J, Nishimura Y, Yamada M, Suzuki H, Hirota Y. Gene organization in the region containing a new gene involved in chromosome partition in *Escherichia coli*. *J Bacteriol*. 1988; 170:3967–3977. [PubMed: 2842295]
- Kato J, Suzuki H, Ikeda H. Purification and characterization of DNA topoisomerase IV in *Escherichia coli*. *J Biol Chem*. 1992; 267:25676–25684. [PubMed: 1334483]
- Levine, C. Ph.D Thesis: The Identification and Analysis of High-copy Suppressors of Topo IV Mutations. Cornell University; 2000.
- Levine C, Hiasa H, Marians KJ. DNA gyrase and topoisomerase IV: biochemical activities, physiological roles during chromosome replication, and drug sensitivities. *Biochim Biophys Acta*. 1998; 1400:29–43. [PubMed: 9748489]
- Levine C, Marians KJ. Identification of *dnaX* as a high-copy suppressor of the conditional lethal and partition phenotypes of the *parE10* allele. *J Bacteriol*. 1998; 180:1232–1240. [PubMed: 9495763]
- Link AJ, Phillips D, Church GM. Methods for generating precise deletions and insertions in the genome of wild-type *Escherichia coli*: application to open reading frame characterization. *J Bacteriol*. 1997; 179:6228–6237. [PubMed: 9335267]
- Lopez CR, Yang S, Deibler RW, Ray SA, Pennington JM, Digate RJ, Hastings PJ, Rosenberg SM, Zechiedrich EL. A role for topoisomerase III in a recombination pathway alternative to RuvABC. *Mol Microbiol*. 2005; 58:80–101. [PubMed: 16164551]
- Mankouri HW, Hickson ID. The RecQ helicase-topoisomerase III-Rmi1 complex: a DNA structure-specific 'dissolvosome'? *Trends Biochem Sci*. 2007; 32:538–546. [PubMed: 17980605]
- Menzel R, Gellert M. Regulation of the genes for *E. coli* DNA gyrase: homeostatic control of DNA supercoiling. *Cell*. 1983; 34:105–113. [PubMed: 6309403]
- Mizuuchi K, O'Dea MH, Gellert M. DNA gyrase: subunit structure and ATPase activity of the purified enzyme. *Proc Natl Acad Sci U S A*. 1978; 75:5960–5963. [PubMed: 153529]
- Nurse P, Levine C, Hassing H, Marians KJ. Topoisomerase III can serve as the cellular decatenase in *Escherichia coli*. *J Biol Chem*. 2003; 278:8653–8660. [PubMed: 12509418]
- Oakley TJ, Hickson ID. Defending genome integrity during S-phase: putative roles for RecQ helicases and topoisomerase III. *DNA Repair (Amst)*. 2002; 1:175–207. [PubMed: 12509252]
- Oram M, Kuroda R, Fisher LM. *Escherichia coli* DNA gyrase: genetic analysis of *gyrA* and *gyrB* mutations responsible for thermosensitive enzyme activity. *FEBS Lett*. 1992; 312:61–65. [PubMed: 1330681]
- Peng H, Marians KJ. *Escherichia coli* topoisomerase IV. Purification, characterization, subunit structure, and subunit interactions. *J Biol Chem*. 1993; 268:24481–24490. [PubMed: 8227000]
- Peter BJ, Ullsperger C, Hiasa H, Marians KJ, Cozzarelli NR. The structure of supercoiled intermediates in DNA replication. *Cell*. 1998; 94:819–827. [PubMed: 9753328]

- Schofield MA, Agbunag R, Michaels ML, Miller JH. Cloning and sequencing of *Escherichia coli* mutR shows its identity to topB, encoding topoisomerase III. *J Bacteriol.* 1992; 174:5168–5170. [PubMed: 1321123]
- Srivenugopal KS, Lockshon D, Morris DR. *Escherichia coli* DNA topoisomerase III: purification and characterization of a new type I enzyme. *Biochemistry.* 1984; 23:1899–1906. [PubMed: 6326814]
- Sundin O, Varshavsky A. Terminal stages of SV40 DNA replication proceed via multiply intertwined catenated dimers. *Cell.* 1980; 21:103–114. [PubMed: 6250706]
- Sundin O, Varshavsky A. Arrest of segregation leads to accumulation of highly intertwined catenated dimers: dissection of the final stages of SV40 DNA replication. *Cell.* 1981; 25:659–669. [PubMed: 6269752]
- Suski C, Marians KJ. Resolution of converging replication forks by RecQ and topoisomerase III. *Mol Cell.* 2008; 30:779–789. [PubMed: 18570879]
- Uematsu N, Eda S, Yamamoto K. An *Escherichia coli* topB mutant increases deletion and frameshift mutations in the supF target gene. *Mutat Res.* 1997; 383:223–230. [PubMed: 9164483]
- Wang JC. DNA topoisomerases: why so many? *J Biol Chem.* 1991; 266:6659–6662. [PubMed: 1849888]
- Wang JC. Cellular roles of DNA topoisomerases: a molecular perspective. *Nat Rev Mol Cell Biol.* 2002; 3:430–440. [PubMed: 12042765]
- Wigley DB. Structure and mechanism of DNA topoisomerases. *Annu Rev Biophys Biomol Struct.* 1995; 24:185–208. [PubMed: 7663114]
- Wu L, Hickson ID. The Bloom's syndrome helicase suppresses crossing over during homologous recombination. *Nature.* 2003; 426:870–874. [PubMed: 14685245]
- Yi TM, Stearns D, Demple B. Illegitimate recombination in an *Escherichia coli* plasmid: modulation by DNA damage and a new bacterial gene. *J Bacteriol.* 1988; 170:2898–2903. [PubMed: 2838451]
- Zhang HL, Malpure S, DiGate RJ. *Escherichia coli* DNA topoisomerase III is a site-specific DNA binding protein that binds asymmetrically to its cleavage site. *J Biol Chem.* 1995; 270:23700–23705. [PubMed: 7559540]
- Zhu Q, Pongpech P, DiGate RJ. Type I topoisomerase activity is required for proper chromosomal segregation in *Escherichia coli*. *Proc Natl Acad Sci U S A.* 2001; 98:9766–9771. [PubMed: 11493711]

Fig. 1.

Inactivation of neither RecQ nor RecA relieves the requirement of *topB* for suppression of W3110*parE10* temperature sensitivity by overexpression of *dnaX*(γ). Shown are 10-fold serial dilutions of the strains.

A. Strains W3110, W3110 Δ *topB*::*kan*, W3110 Δ *recQ*::*FRTcat*, W3110 Δ *topB*::*kan* Δ *recQ*::*FRTcat*, W3110*parE10*, W3110*parE10* Δ *topB*::*kan*, W3110*parE10* Δ *recQ*::*FRTcat*, and W3110*parE10* Δ *topB*::*kan* Δ *recQ*::*FRTcat* carrying pLEX5BA-*dnaX*(γ) (strains BP116, BP098, BP271, BP272, CL012, CL013, CL014, and CL015, respectively) were grown overnight at 30 °C, diluted into fresh medium, spotted on LB plates containing Ap and 100 μ M IPTG, and incubated at the temperature indicated as described in Experimental Procedures. Additional vector controls can be found in Fig. S1.

B. Strains W3110*recA*^{398::cat}, W3110 Δ *topB*::*kanrecA*^{398::cat}, W3110*parE10**recA*^{398::cat}, and W3110*parE10* Δ *topB*::*kanrecA*^{398::cat} carrying pLEX5BA-*dnaX*(γ) (strains BP358, BP359, CL026, and CL027, respectively) were grown and analyzed as above

Fig. 2.

Overexpression of Topo III can rescue growth of the W3110*parE10ΔtopB* double mutant strain at 42 °C. Shown are 10-fold serial dilutions of the strains. Strains W3110, W3110*parE10*, W3110*ΔtopB*, and W3110*parE10ΔtopB* carrying either pLEX5BA (vector) or pLEX5BA-*topB* (pBP42) (*topB*) (strains BP115, BP120, CL016, CL021, BP097, BP355, CL017, and CL022, respectively) were grown overnight at 30 °C, diluted with fresh medium, and spotted on LB plates containing Ap and 100 μM IPTG. Plates were incubated at either 30 °C or 42 °C as described in Experimental Procedures.

Fig. 3.

The catalytic activity of Topo III is required for suppression of the Topo IV defect and Topo I can not replace Topo III.

Shown are 10-fold serial dilutions of the strains. The gene product overexpressed is indicated under each panel.

W3110, W3110*parE10*, C600, and C600*parC1215* carrying pLEX5BA (strains BP115, CL016, BP033, and BP069, respectively); pLEX5BA-*topA* (pBP59) (strains BP314, CL029, BP316, and BP317, respectively); pLEX5BA-*topB* (pBP42) (strains BP120, CL021, BP121, and BP126, respectively); and pLEX5BA-*topBY328F* (pBP47) (strains BP195, CL028, BP196, and BP198, respectively) were grown in LB medium overnight at 30 °C, diluted, spotted on LB plates containing Ap and 100 μM IPTG, and incubated at 42 °C as described in Experimental Procedures. Plating controls at 30 °C can be found in Fig. S2.

Fig. 4.

Neither RecQ nor RecA is required for suppression of W3110*parE10* temperature sensitivity by overexpression of *topB*. Shown are 10-fold dilutions of the strains.

A. W3110, W3110Δ*recQ*, W3110*parE10*, and W3110*parE10*Δ*recQ* carrying either pLEX5BA (vector) (strains BP115, BP174, CL016, and CL018, respectively) or pLEX5BA-*topB* (*topB*, pBP42) (strains BP120, BP175, CL021, and CL023, respectively) were grown overnight at 30 °C, diluted into fresh medium, spotted onto LB plates containing Ap and 100 μM IPTG, and the plates were incubated at the indicated temperatures as described in Experimental Procedures. B. W3110*recA*⁻ and W3110*parE10**recA*⁻ carrying either pLEX5BA (vector) (strains BP303 and CL020, respectively) or pLEX5BA-*topB* (*topB*, pBP42) (strains BP304 and CL024, respectively) were grown and analyzed as above.

Fig. 5.

Deletion of *topB* causes chromosome segregation defects at the permissive temperature in the C600*parC1215* and W3110*parE10* strains.

A - H. Shown are representative fluorescent images of strains treated with DAPI (DNA stain, blue) and FM4-64FX (membrane stain, red). Size bars are 5 μm.

(A) and (B), C600*parC1215* (EJ812) (A) and C600*parC1215ΔtopB* (BP079) (B) grown at 25 °C for 4 h.

(C) and (D), W3110*parE10* (CL001) (C) and W3110*parE10ΔtopB* (CL002) (D) grown at 30 °C for 2 h.

(E) - (H), W3110*parE10*(CL001) (E), W3110*parE10ΔtopB* (CL002) (F), W3110*gyrB^{ts}* (BP199) (G), and W3110*gyrB^{ts}ΔtopB* (BP275) (H) grown at 30 °C for 2 h and then shifted to 42 °C for 1 h.

I. Distribution of cell lengths for C600*parC1215* (EJ812) and C600*parC1215ΔtopB* (BP079) cells shown in panels A and B.

J. Distribution of cell morphologies for W3110*parE10*(CL001) and W3110*parE10ΔtopB* (CL002) cells shown in panels C and D.

K. Distribution of cell lengths for W3110*parE10*(CL001) and W3110*parE10ΔtopB* (CL002) cells shown in panels E and F.

L. Distribution of cell lengths for W3110*gyrB^{ts}* (BP199) and W3110*gyrB^{ts}ΔtopB* (BP275) cells shown in panels G and H.

M and N. Growth curves of C600*parC1215ΔtopB* (BP079) (M) and W3110*parE10ΔtopB* (CL002) (N). For the C600*parC1215ΔtopB* (BP079) experiment (M), C600 (pink), C600*ΔtopB* (BP105, green), C600*parC1215* (EJ812, red), and C600*parC1215ΔtopB* (BP079, blue) were grown overnight in LB at 25 °C, diluted to OD₆₀₀ of 0.0005, and growth was monitored at 30 °C. For the W3110*parE10ΔtopB* (CL002) experiment (N), W3110 (pink), W3110*ΔtopB* (BP093, green), W3110*parE10*(CL001, red), and W3110*parE10ΔtopB* (CL002, blue) were grown overnight in LB at 30 °C, diluted to OD₆₀₀ of 0.0005, and growth was monitored at 30 °C.

Fig. 6.

The chromosome segregation block in the W3110*parE10ΔtopB* mutant strain is not a result of toxic recombination intermediates.

A - D. Shown are representative fluorescent images of strains stained with DAPI (DNA stain, blue) and FM4-64FX (membrane stain, red). Size bars are 5 μm. (A) W3110*parE10ΔrecQ* (CL004); (B) W3110*parE10ΔtopBΔrecQ* (CL006); (C) W3110*parE10recA⁻* (CL008); and (D) W3110*parE10ΔtopBrecA⁻* (CL010) were incubated at 30 °C for 2 h.

E. Distribution of cell morphologies for the cells shown in panels (A) – (D).

F and G. W3110*parE10ΔtopBΔrecQ* (CL008) and *parE10ΔtopBrecA* (CL010) fail to grow at the semi-permissive temperature of 37 °C. (F) W3110*parE10* (CL001), W3110*parE10ΔtopB* (CL002), W3110*parE10ΔrecQ* (CL004) and W3110*parE10ΔtopBΔrecQ* (CL006) and (G) W3110*parE10* (CL001), W3110*parE10ΔtopB* (CL002), W3110*parE10recA*⁻ (CL008) and W3110*parE10ΔtopBrecA*⁻ (CL010) strains were grown at 30 °C, serially diluted 10-fold, spotted on LB plates containing AP and 100 μM IPTG, and incubated at 30 °C (left panel) and 37 °C (right panel).

Fig. 7.

W3110ΔtopB mutant strains are hypersensitive to novobiocin.

A and B. Shown are representative fluorescent images of strains stained with DAPI (DNA stain, blue) and FM4-64FX (membrane stain, red). Size bars are 5 μm. (A) *W3110gyrB^{tsNov}* (BP199) and (B) *W3110gyrB^{tsNov}ΔtopB* (BP275) treated with novobiocin (300 μg/ml) for 105 min at 30 °C. C and D. Novobiocin sensitivity of *topB* mutants. Strains were grown as described in Experimental Procedures and plated on LB plates supplemented with novobiocin. (C) *W3110* (red) and *W3110ΔtopB* (BP093; blue); (D) *W3110gyrB^{tsNov}* (BP199, red) and *W3110gyrB^{tsNov}ΔtopB* (BP275, blue).

E and F. Cell length distribution (E) and nucleoid area distribution (F) of cells shown in panels A and B. For panel F, bin size is 1 μm² and bin centers are shown.

G and H. Overexpression of Topo IV and Topo III can bypass novobiocin sensitivity. Shown are 10-fold serial dilutions of the strains. (G) *W3110* carrying pLEX5BA (vector, BP115) and *W3110ΔtopB* carrying pLEX5BA (vector, BP097), pLEX5BA-*parEparC* (parEC, BP093), and pLEX5BA-*topB* (pBP42) (topB, BP335) were plated on LB plates containing 50 μg/ml novobiocin and 100 μM IPTG. (H) *W3110gyrB^{tsNov}* carrying pLEX5BA (vector, BP204) and *W3110gyrB^{tsNov}ΔtopB* carrying pLEX5BA (vector, BP363), pLEX5BA-*parEparC* (parEC, BP364) and pLEX5BA-*topB* (pBP42) (topB, BP365) were plated on LB plates containing 300 μg/ml novobiocin and 100 μM IPTG.

Fig. 8.

Three topoisomerases contribute to the complete unlinking of the parental strands during DNA replication. At an active replication fork precatenanes form behind the replisome while positive supercoils accumulate ahead of it. Topo III (green oval) removes the precatenanes by acting near or at the point of nascent DNA synthesis where there is single-stranded template DNA available for it to bind, whereas gyrase acts ahead of the replication fork converting the positive supercoils into negative ones. Linkages between the parental strands that remain after replication is completed form catenated daughter chromosomes that are unlinked by the action of Topo IV.