Biology, Host Range, Pathogenesis and Diagnosis of *White spot syndrome virus*

Balakrishnan Pradeep · Praveen Rai ·
Seethappa A. Mohan · Mudagandur S. Shekhar ·
Indrani Karunasagar

Received: 27 January 2012 / Accepted: 26 June 2012 / Published online: 14 August 2012
© Indian Virological Society 2012

Abstract *White spot syndrome virus* (WSSV) is the most serious viral pathogen of cultured shrimp. It is a highly virulent virus that can spread quickly and can cause up to 100 % mortality in 3–10 days. WSSV is a large enveloped double stranded DNA virus belonging to genus *Whispovirus* of the virus family *Nimaviridae*. It has a wide host range among crustaceans and mainly affects commercially cultivated marine shrimp species. The virus infects all age groups causing large scale mortalities and the foci of infection are tissues of ectodermal and mesodermal origin, such as gills, lymphoid organ and cuticular epithelium. The whole genome sequencing of WSSV from China, Thailand and Taiwan have revealed minor genetic differences among different strains. There are varying reports regarding the factors responsible for WSSV virulence which include the differences in variable number of tandem repeats, the genome size and presence or absence of different proteins. Aim of this review is to give current information on the status, host range, pathogenesis and diagnosis of WSSV infection.

Keywords WSSV · Shrimp · Host range · Pathogenesis · Diagnosis · PCR

B. Pradeep (✉)
Krishi Vigyan Kendra, Indian Institute of Spices Research,
Peruvannamuzhi, Kozhikode 673528, Kerala, India
e-mail: nairpradeep2@rediffmail.com;
nairpradeep2@gmail.com

P. Rai · S. A. Mohan · I. Karunasagar
Department of Fishery Microbiology, College of Fisheries,
Karnataka Veterinary, Animal and Fisheries Sciences
University, Mangalore 575 002, India

M. S. Shekhar
Genetics and Biotechnology Unit, Central Institute of
Brackishwater Aquaculture, Chennai, India

Introduction

Shrimp culture covers a major portion of the aquaculture industry and its export forms the main source of revenue of many countries. Global aquaculture of penaeid shrimp has grown rapidly during the last two decades and its culture is being undertaken even in saline aquifers, marine, fresh and brackish water areas. The farmed shrimps contributed 42.2 % of total global shrimp production of 6.67 million tones in 2009. This alarming increase was possible due to the intensification of farming practices, expansion of culture and the scientific inputs provided to farmers. Due to the intensification and specialization of shrimp farming, the contribution from farm-raised shrimp to total world supply of shrimp grew by approximately 400 % over one decade during 1985–1995. The shrimp farming sector has grown at an annual average growth rate of over 18.8 % since 1970 [23]. Approximately 75 % of global shrimp production is from aquaculture which is now almost entirely dominated by two species—the black tiger shrimp (*Penaeus monodon*) and the white Pacific shrimp (*Litopenaeus vannamei*) [134].

In India, commercial shrimp farming started gaining roots only during the mid-eighties. The boom period of commercial-scale shrimp culture in India started in 1990 and the worst came in 1995–1996, with the outbreak of viral disease. The fact that most of the coastal States in India were new to commercial-scale shrimp farming, the general ignorance of good farming practices, and the lack of suitable extension services, led to a series of problems [150]. In India, a variety of farming techniques are practiced like traditional, improved traditional, extensive, semi-intensive and intensive. The scientific shrimp farming has shown phenomenal growth since early nineties, and at present about 150,000 ha is under shrimp farming which is about 15 % of the total potential brackishwater area available in

the country. Development of shrimp farming in the country has been restricted mainly to east coast, due to availability of shrimp broodstock, seed and location of hatcheries. Shrimp farming in India is mainly restricted to monoculture of black-tiger shrimp, *P. monodon*, which accounts for more than 95 % of the production [39]. Farmed shrimp yield in India has decreased to 75,997 tones in 2009–2010. The decline was around 28.4 % in production and 10.9 % in area compared to previous year as per MPEDA owing to recent disease outbreaks. Despite the decline in production, an increase in shrimp export was recorded, which may be due to high global demand for shrimp and a consequence of lessening of economic crisis. Intensive shrimp cultivation, inadequate sanitation, poor pond management and irresponsible movement of broodstock and larvae have enhanced the disease incidence and its spread [55].

Since 1981, several new viral pathogens have emerged successively, causing mass mortalities and threatening the economic sustainability of the industries in Asia and the Americas [132]. With expansion of the shrimp industry, the number of viral pathogens of penaeid shrimp has also increased, reaching a new high of 22 [94]. Among the known viruses of shrimp, white spot disease caused by *White spot syndrome virus* (WSSV) has had the greatest impact on shrimp culture and continues to be an obstacle to sustainable shrimp farming [135]. The Government of India has recently given approval for culturing *L. vannamei* in India. This shrimp can be cultured only in those farms which have got approval from Coastal Aquaculture Authority (CAA). Many farms in East coast have taken up its culture. The projected production of *L. vannamei* in India is about 50,000 tonnes. Although the farmers got good crops initially there are recent reports of WSSV outbreak in these farms. It should be borne in mind that *L. vannamei* culture crashed in Latin American countries such as Ecuador due to WSSV. This review mainly focuses on current information on the status, host range, pathogenesis and diagnosis of WSSV infection in crustaceans.

Geographical Distribution

Since the discovery of WSSV in China (Fujian) in 1992, the virus spread rapidly to all the major shrimp farming

countries of the world due to intensification in shrimp farming and inadequate sanitation [134]. By 1996 the disease had spread to major shrimp farming regions of East Asia, South East Asia, Indonesia and India. In 1992, it was first detected in Taiwan and then spread to all Asian countries [100, 132]. In Japan, white spot disease was observed in *Penaeus japonicus* in 1992 [28, 40]. In 1995 it was first diagnosed in the United States of America (Texas and South-Carolina) [93]. During 1998, shrimp exhibiting gross signs and histopathology of white spot disease (WSD) was reported from India and Korea [45, 46, 49]. In early 1999, WSSV had spread to Latin America and was reported in Central and South-America [94]. In 2002, the virus was detected in Europe (France) and the Middle East (Iran) [95]. In 2004, it was observed in *P. indicus* from Choebdeh area in Iran [116]. Australia is the only shrimp farming country free of WSSV, where shrimp farming is being practiced, which is a result of the strict quarantine measures adopted by the Australian authorities. The original source of virus still remains a mystery [133].

Clinical Signs and Transmission

In cultured shrimp, WSSV infection can result in a cumulative mortality of up to 100 % within 3–10 days [55]. Infected animals show lethargic behavior, reduction in food consumption, reduced preening activities, anorexia, loose cuticle and reddish to pink body discoloration [12]. A characteristic of the syndrome due to the virus include white spots on the exoskeleton especially on the carapace and last abdominal segment (Fig. 1). These spots are the result of calcified deposits that range in size from a few mm to 1 cm or more in diameter [12]. However, in the case of acute (experimental) infections, the only signs of WSSV infection observed are lethargy and lack of appetite. In *P. indicus*, body turns to reddish colour and it is necessary to remove the carapace to confirm WSSV infection [90].

The modes of transmission of WSSV in natural environment are mainly by vertical and/or horizontal route. Horizontal transmissions is by ingestion of dead infected shrimp, by contact with water containing infected animals or free virus particles. Infection by the latter is thought to occur primarily through the gills, but may occur via other

Fig. 1 *White spot syndrome virus* infected *Penaeus monodon* showing white spot symptoms on carapace (a) and last abdominal segment (b)

body surfaces as well [6, 12, 13]. Lotz and Soto demonstrated that, transmission by cohabitation is of a lower magnitude than ingestion transmission [62]. Vertical transmission occurs from the mother to offspring through viral particles being shed at the time of spawning and then ingested by larvae at first feeding, although it is not clear whether the WSSV virions are present inside the shrimp eggs [35, 58, 84, 117]. By using in situ hybridization and TEM, WSSV viral particles were located in the connective tissue layer surrounding the seminiferous tubules in testes and in the case of ovary, viral particles were detected in follicle cells, oogonia, oocytes and connective tissue cells. In addition, no infection was found in mature eggs, which may imply that infected eggs were killed by the virus before maturation [58]. No penaeid shrimp species are known to be resistant to WSSV infection [55, 61]. Even specific pathogen free (SPF) shrimps were observed to be asymptomatic carriers of WSSV which was attributed to viral latency. The latent infection of WSSV is reported to be associated with three WSSV genes (ORFs 366, 151 and 427) [47].

Taxonomic Affiliation

White spot syndrome virus (WSSV) was earlier known by different names until 2005 when the virus got its internationally approved name [131]. WSSV was originally called a baculovirus based on its cylindrical morphology and on the histological lesions that resembled those of “non-occluded” baculoviruses [146]. The other names by which it was known earlier, includes hypodermal and hematopoietic necrosis baculovirus (HHNBV) [36], rod-shaped nuclear virus of *P. japonicus* (RV-PJ) [113], Chinese baculovirus (CBV), systemic ectodermal and mesodermal baculovirus (SEMBV) [92], penaeid rod-shaped DNA virus (PRDV) [41], white spot baculovirus (WSBV) [57] and now as white spot syndrome virus (WSSV) [55]. The disease caused by WSSV is often indicated as white spot disease (WSD). The taxonomic position of WSSV was assigned by the International Committee of Taxonomy of Viruses (ICTV) to new genus, *Whispovrius*, under the family *Nimaviridae* based on its unique morphology and genomic composition [67, 68, 130]. Since WSSV has a thread-like polar extension at one end of the virus particle, WSSV became the sole species of a new family *Nimaviridae* [53, 100].

Host Range

WSSV has a wide host range which includes all cultured, wild marine shrimps, crabs, lobsters, crayfishes, squilla,

copepods and freshwater cultures species of *Macrobrachium rosenbergii* [5, 7, 10, 16, 22, 24, 32, 43, 57, 63, 71, 83, 91, 98, 106, 111, 112, 138, 160]. WSSV is potentially lethal to most of the commercially important species of penaeid shrimps including *P. monodon*, *P. vannamei*, *P. indicus*, *P. japonicus*, *P. chinensis*, *P. penicillatus*, *P. azteus*, *P. merguensis*, *F. duorarum*, *P. stylirostris* [79]. Table 1 shows the list of host species for WSSV. The susceptibility to WSSV differs significantly between hosts. In some species, WSSV results in a non-lethal or latent infection, making these species potential virus reservoirs and important sources of infection in shrimp ponds. In addition, the occurrence of WSSV in pond soil was detected by nested-PCR [75]. Hence, soil may be an important aspect of WSSV ecology and may have an implication for viral transmissibility. The study conducted in 2006 shows that, rotifer (*Brachionus urceus*) serve as a vector of WSSV to infect the shrimps [159]. By PCR-dot blot hybridization, WSSV was detected in rotifer resting eggs from shrimp culture-pond sediments. Detection of WSSV from disinfected eggs suggests the presence of WSSV within eggs and indicates vertical transmission of WSSV [151]. Studies need to be taken up to know, further on the host range of WSSV as it will be helpful in preventing the spread of disease and also to know the reasons behind the variation in host susceptibility to infection.

Histopathology and Cytopathology

WSSV infects most tissues originating from both ectoderm and mesoderm. Histological signs of WSSV infection include enlarged nuclei in tissues. These include subcuticular epithelium, gills, lymphoid organs, antennal gland, hematopoietic tissues, connective tissue, ovary and the ventral nerve cord [146]. The most convenient tissue for diagnosis is the subcuticular epithelium in tissue sections. The subcuticular epithelium of the stomach usually provides excellent views revealing pathognomonic enlarged nuclei containing basophilic inclusions which are surrounded by vacant cytoplasm. Nuclei at the early stage of infection show marginated chromatin separated from a reddish inclusion by a ring of unstained nucleoplasm. These are called Cowdry A-type inclusions [25]. During early infection stages, the stomach, gills, cuticular epidermis and the connective tissue of the hepatopancreas gets infected. At later stages of infection, the lymphoid organ, antennal gland, muscle tissue, hematopoietic tissue, heart, hindgut and parts of the midgut also become infected. The nervous system and the compound eyes are infected very late. The stomach, gills, cuticular epidermis, lymphoid organ, hematopoietic tissue and antennal gland are all

Table 1 The known host species reported to be naturally or experimentally infected with *White spot syndrome virus*

Scientific Name	Common Name	Type	References
A. Shrimps			
<i>Alpheus brevicristatus</i>	Snapping shrimp	N	[112]
<i>Alpheus lobidens</i>	Apping shrimp	N	[112]
<i>Aristeus</i> sp.	Red shrimp	N	[5]
<i>Callinassa</i> sp.	Mud shrimps	N	[57]
<i>Exopalaemon orientalis</i>	Oriental prawn	N, E	[7, 24, 138]
<i>Farfantepenaeus aztecus</i>	Northern brown shrimp	E	[55]
<i>Farfantepenaeus duorarum</i>	Pink shrimp	E	[55]
<i>Fenneropenaeus penicillatus</i>	Red tail shrimp	N	[12, 57, 138]
<i>Fenneropenaeus chinensis</i>	Fleshy shrimp	N	[24]
<i>Heterocarpus</i> sp.		N	[5]
<i>Litopenaeus vannamei</i>	Whiteleg shrimp	N, E	[55]
<i>Litopenaeus setiferus</i>	Northern white shrimp	E	[55]
<i>Macrobrachium rosenbergii</i>	Giant freshwater shrimp	N, E	[5, 7, 24, 32, 57, 91, 138]
<i>Macrobrachium idella</i>	Sunset shrimp	E	[91]
<i>Macrobrachium lamarrei</i>	Kuncho river prawn	E	[100]
<i>Marsupenaeus japonicus</i>	Kuruma shrimp	N, E	[57, 113, 138]
<i>Metapenaeus ensis</i>	Greasyback shrimp	N, E	[7, 57, 138]
<i>Metapenaeus dobsoni</i>	Kadal shrimp	N, E	[5, 32, 91]
<i>Metapenaeus lysianassa</i>	Bird shrimp	N	[100]
<i>Exopalaemon orientis</i>	Oriental prawn	E	[7]
<i>Metapenaeus monoceros</i>	Speckled shrimp	E	[91]
<i>Metapenaeus elegans</i>	Fine shrimp	N	[5]
<i>Metapenaeus brevicornis</i>	Yellow prawn		[70]
<i>Metapenaeus affinis</i>	Jinga shrimp		[70]
<i>Palaemon adspersus</i>	Baltic prawn	E	[16]
<i>Palaemon styliferus</i>	Grass shrimp	N	[24, 57]
<i>Parapenaeopsis stylifera</i>	Kiddi shrimp	N	[5, 32]
<i>Penaeus monodon</i>	Giant tiger shrimp	N, E	[5, 12, 57, 147]
<i>Penaeus indicus</i>	Indian white prawn	N, E	[24, 91]

Table 1 continued

Scientific Name	Common Name	Type	References
<i>Penaeus merguensis</i>	Banana prawn	N	[24]
<i>Penaeus semiculcatus</i>	Green tiger prawn	N, E	[57, 91, 138]
<i>Penaeus canaliculatus</i>	Striped prawn	N	[70]
<i>Penaeus aztecus</i>	Brown shrimp		[55]
<i>Penaeus schmitti</i>	Southern white shrimp	E	[100]
<i>Penaeus chinensis</i>	Chinese white shrimp	N/E	[2]
<i>Penaeus duorarum</i>	Northern pink shrimp	N/E	[55]
<i>Penaeus setiferus</i>	Northern white shrimp	N/E	[55]
<i>Penaeus stylirostris</i>	Blue shrimp	E	[55]
<i>Solenocera crassicornis</i>	Coastal mud shrimp	N	[32]
<i>Solenocera indica</i>		N	[32]
<i>Seranopsis longipes</i>		N	[70]
<i>Solenocera choprai</i>	Ridgeback shrimp	N	[70]
<i>Squilla</i> sp.	Mantis shrimp	N, E	[5, 32]
<i>Trachypenaeus curvirostris</i>	Southern rough shrimp	E	[7, 138]
B. Crabs			
<i>Atergatis integerrimus</i>	Bashful crab	E	[96]
<i>Cancer pagurus</i>	Edible or rock crab	E	[16]
<i>Calappa lophos</i>	Box crab	N, E	[5, 24]
<i>Calappa philargius</i>	Box crab	E	[98]
<i>Callinectes arcuatus</i>	Swimming crab	N	[26]
<i>Callinectes sapidus</i>	Blue crab	N	[8]
<i>Carcinus maenas</i>	Littoral crab	E	[16]
<i>Charybdis annulata</i>	Swimming crab	N, E	[32, 98]
<i>Charybdis cruciata</i>	Red sea crab	N	[5, 32]
<i>Charybdis feriata</i>	Coral crab	N	[57]
<i>Charybdis granulata</i>	Swimming crab	E	[7]
<i>Charybdis hoplites</i>	Swimming crab	N	[5]
<i>Charybdis lucifera</i>	Swimming crab	N, E	[5, 98]
<i>Charybdis natator</i>	Hairyback crab	N	[24]
<i>Charybdis riversandersoni</i>		N	[70]
<i>Demania splendida</i>		E	[96]

Table 1 continued

Scientific Name	Common Name	Type	References
<i>Doclea hybrida</i>		E	[98]
<i>Gelasimus marionis nitidus</i>		N	[32]
<i>Grapsus albolineatus</i>	Rock crab	E	[98]
<i>Halimede ochtodes</i>	Hairy crab	E	[98]
<i>Helice tridens</i>	Shore crab	N	[57]
<i>Liagore rubromaculata</i>		E	[98]
<i>Liocarcinus depurator</i>	Harbour crab	E	[16]
<i>Liocarcinus puber</i>	Velvet swimming crab	E	[16]
<i>Lithodes maja</i>	Deepsea king crab	E	[98]
<i>Macrophthalmus sulcatus</i>	Ghost/fiddler crab	N	[32]
<i>Mantura</i> sp.		N	[24]
<i>Matuta miersi</i>	Moon crab	E	[98]
<i>Matuta planipes</i>	Moon crab	N	[81]
<i>Metopograpsus messor</i>	Purple climber crab	N, E	[32, 91]
<i>Menippe rumphii</i>	Stone crab	E	[96]
<i>Paradorippe granulata</i>		E	[98]
<i>Parthenope prensor</i>	Elbow crab	E	[98]
<i>Parathelphusa hydrodomous</i>		E	[97]
<i>Parathelphusa pulvinata</i>		E	[97]
<i>Philyra syndactyla</i>	Purse crab	E	[98]
<i>Podophthalmus vigil</i>	Long-eyed swimming crab	E	[98]
<i>Portunus pelagicus</i>	Sand crab	N, E	[5, 57, 111]
<i>Portunus sanguinolentus</i>	Blood spot crab	N, E	[5, 57, 98]
<i>Pseudograpsus intermedius</i>	Mosaic crab	N	[5]
<i>Scylla serrata</i>	Mud crab	N, E	[5, 10, 57, 91, 98, 111]
<i>Scylla tranquebarica</i>	Mangrove crab	E,N	[27, 91]
<i>Scylla olivacea</i>	Orange mud crab	E	[108]
<i>Scylla paramamosain</i>	Green mud crab	E	[108]
<i>Sesarma</i> sp.	Marsh crabs	N, E	[24, 91]
<i>Somannia-thelpusa</i> sp.	Black rice crab	E	[24]
<i>Thalamita danae</i>	Swimming crab	N, E	[24, 98]
C. Lobsters			
<i>Acetes</i> sp.	Krill	E	[111]

Table 1 continued

Scientific Name	Common Name	Type	References
<i>Panulirus homarus</i>	Scalloped spiny lobster	E	[91]
<i>Panulirus longipes</i>	Longlegged spiny lobster	E	[24]
<i>Panulirus ornatus</i>	Ornata spiny lobster	E	[24, 91]
<i>Panulirus penicillatus</i>	Pronghorn spiny lobster	E	[7]
<i>Panulirus polyphagus</i>	Mud spiny lobster	E	[91]
<i>Panulirus versicolor</i>	Painted spiny lobster	E	[7]
<i>Scyllarus arctus</i>	Small European locust lobster	E	[16]
D. Crayfish			
<i>Astacus leptodactylus</i>	Turkish crayfish	E	[16]
<i>Astacus astacus</i>	Broad-fingered crayfish	E	[44]
<i>Cherax destructor albidus</i>	Yabby	E	[22]
<i>Cherax quadricarinatus</i>	Australian redclaw	E	[105]
<i>Orconectes limosus</i>	Spinycheek crayfish	E	[16]
<i>Orconectes punctimanus</i>	Spothanded Crayfish	N	[141]
<i>Pacifastacus leniusculus</i>	Signal crayfish	E	[43]
<i>Procambarus clarkia</i>	Red swamp crayfish	E	[7, 138]
E. Insects			
<i>Ephydriidae</i> sp.	Shore fly	N	[57]
<i>Gerries</i>		N	[70]
<i>Notonecta</i>	Backswimmer	N	[70]
<i>Nepa</i>		N	[70]
<i>Ranatra</i>	Water scorpion	N	[70]
<i>Dragon nymphfly</i>		N	[70]
F. Copepod, Polychaeta, Rotifera and Strmatopoda			
<i>Schmackeria dubia</i>	Copepoda	N	[57]
<i>Squilla mantis</i>	Mantis shrimp	N	[97]
<i>Marphysa gravelyi</i>	Polychaeta	N	[129]
<i>Brachionus urceus</i>	Rotifera	NE	[151, 159]
G. Brine shrimps			
<i>Artemia</i> sp.		E	[98]
<i>A. franciscana</i>		E	[54]

heavily infected with WSSV at late stages of infection and become necrotic [6, 58]. Cells of the hepatopancreatocytes and epithelial cells of the midgut have never been shown to be infected with WSSV. A significant reduction in the total

hemocyte count is observed after shrimp are infected with WSSV [30, 43, 121]. This is probably caused by infection of the hemocytes themselves as well as by apoptosis in the WSSV infected hematopoietic tissue, from which the hemocytes derive [145]. WSSV infected hemocytes either undergo apoptosis or are removed from the circulation by attaching to host tissues. As a consequence abnormal low amount of hemocytes weaken the shrimp defenses.

WSSV replication initially occurs in the nucleus and early signs of infection are characterized by the appearance of homogeneous hypertrophied nuclei and marginated chromatin [55, 140, 145]. Virus morphogenesis is initiated by the formation of viral envelopes in the nucleoplasm. The formation of the nucleocapsids begins with extended, empty, long tubules, which break up into fragments of 12–14 rings to form empty nucleocapsid shells. Subsequently, the empty capsids are surrounded by the envelope leaving at one end an open extremity. The nucleoproteins, possibly together with the viral DNA, enter the empty capsid through its open end. Mature virions are obtained after narrowing of the open end and formation of a tail-like extension of the envelope [21, 140]. There are two postulates regarding the transmission of WSSV virions from one cell to another. These are either by budding or by rupture of the nuclear envelope or cell membrane of infected cell.

Morphology and Ultrastructure

Electron microscopy studies of ultra thin sections and WSSV viral suspensions obtained from infected shrimp revealed that, the virion are enveloped, rod shaped, nucleocapsids with a bacilliforms to ovoidal particles of about 275 nm in length and 120 nm in width. These virions have a tail-like appendage at one end, of which neither the function nor the composition is known [21, 130]. Electron microscopic examination of white spot-diseased *P. monodon* and *P. indicus* revealed the occurrence of a non-occluded, rod-shaped, double-enveloped WSSV some of which possessed a tube-like or branched extension and empty capsids. The virions measured 154 ± 2 by 373 ± 5 nm and nucleocapsids 111 ± 1.5 by 293 ± 7 nm [90]. The isolated nucleocapsid have cross hatched appearance and a size about 300×70 nm formed by stacks of rings (about 14 in total), which are in turn formed by regular spaced globular subunits of about 8 nm in diameter, arranged in two parallel rows [21, 72]. The virion contains at least 5 major polypeptides ranging from 15 to 28 kDa [130]. The nucleocapsid contains the viral genome and mainly consists of the WSSV encoded proteins VP664, VP26, VP24 and VP15 [52, 123, 124, 127]. VP664, a remarkable large protein of around 664 kDa, is thought to be the major core protein, responsible for the striated

appearance of the nucleocapsids [52]. VP15, a highly basic protein with no hydrophobic regions, is a histon-like, double-stranded DNA-binding protein [144]. The function of VP26 and VP24 in the nucleocapsid are unknown. Furthermore, about 40 WSSV encoded minor virion proteins were identified by protein sequencing of individual bands after applying purified WSSV virions on a SDS-PAGE gel [37, 38, 52, 120, 123, 124, 127, 158]. The various geographic isolates of WSSV, which have been characterized, are very similar if not identical in morphology and proteome [72, 142].

WSSV Genome

The WSSV genome constitutes a circular, supercoiled, double-stranded (ds) DNA of approximately 300 kilobase pairs (kbp) [21, 72, 146]. Sequence analysis of WSSV genome and comparison of sequence data have shown, it to be unique in not showing any homology with any known viruses [58]. Hence, it has been assigned to a new family (*Nimaviridae*) and genus (*Whispovirus*) [122, 130]. It is one of the largest animal virus that has been completely sequenced so far [122, 152]. The various geographical isolates of WSSV identified are very similar in morphology and protein profile [19]. The restriction fragment length polymorphism (RFLP) patterns of isolates show only limited differences, suggesting a high degree of genomic stability [19, 60, 65, 72, 140, 141]. Two complete genome sequences of WSSV isolates were published in 2001, from Thailand (WSSV-TH, AF369029) [126] and China (WSSV-CN, AF332093) [152]. A third complete genome sequence was published in 2002 from Taiwan (WSSV-TW, AF440570) [118]. The complete genome sequences of these three isolates show a considerable difference in genome size of 292,967, 305,107 and 307,287 bp, for the Thailand, China, and Taiwan isolates respectively, but with an overall identity of 99.32 %. The WSSV-CN genome has a total GC content of 41 % [152]. A total of 684, 531 and 507 putative ORFs have been identified by sequence analysis on the WSSV-TH, WSSV-CN, and WSSV-TW isolates, respectively. Among these putative ORFs, only 184 (WSSV-TH) and 181 (WSSV-CN) are likely to encode functional proteins [126, 152]. Several studies have been carried out on WSSV isolates from different parts of Asia on WSSV genome. The variations associated with the three minisatellite ORF 94, ORF 75 and ORF125 were used for genotyping WSSV and for epidemiological purpose [87, 88, 102, 103, 136, 147]. Further studies on genotyping can be carried out based on the microsatellites associated with WSSV genome. Characterization of variable genomic regions of WSSV from different parts of the world need to be done which will be helpful to elucidate the genetic

evolution of WSSV and to know the variations that are taking place in WSSV genome over a period of time and its relevance to virulence.

Virulence

Virulence is the degree of pathogenicity within a group or species. Preliminary studies have shown, slight difference in virulence between different strains of WSSV, but no direct comparisons were made [51, 139]. The first attempt to compare virulence of six geographical origin (China, India, Thailand, Texas, South Carolina, as well as from crayfish maintained at the USA National Zoo) WSSV isolates in *L. vannamei* (PL) and *Farfantepenaeus duorarum* (juveniles) was made by Wang et al. [139]. The study revealed that, all six isolates of WSSV were infectious and highly virulent to *L. vannamei* (PL), while *F. duorarum* (juveniles) showed moderate resistant to WSSV infection. Slight differences of virulence among six geographic isolates of WSSV were detected; with the crayfish WSSV isolate as the least virulent and Texas WSSV isolate as the most virulent forms [139]. Further studies compared the virulence between WSSV-TH-96-II (largest WSSV genome reported to date and considered as the common ancestral to the WSSV isolates described till date) and WSSV-TH (isolate containing the smallest genome) which suggested that the virus with the smallest genome was more virulent and fit than the one with the largest [66]. Furthermore, a comparative study was conducted between six isolates of WSSV from farms experiencing outbreaks (virulent WSSV; vWSSV) and three isolates of WSSV from farms that had infected shrimps but no outbreaks (non-virulent WSSV; nvWSSV). Both vWSSV and nvWSSV isolates resulted in 100 % mortality within 5 days on challenge in shrimps [110].

In 2009, a study was conducted to compare the virulence and fitness between three different Indian genotypes of WSSV by infecting shrimps and crabs. The three different strains of WSSV were isolated from disease outbreak ponds from West and East coast of India. The results showed that, there was no difference in virulence between the three genotypes of WSSV used for infection, as all three strains caused mortality in shrimps and crabs within the same time period and also there were no significant differences between the viral loads ($P < 0.05$) as indicated by real-time PCR. When three strains were mixed and used, only the strains with a smaller genome size (ORF 94, 3 RU and ORF 94, 6 RU) caused disease. This was further supported by the observation that in the subsequent passage only the strain with ORF 94, 3 RU caused disease. In the case of crabs, when a mixture of viruses was used for infection, only the strain with the smallest genome size

(ORF 94, 3 RU) caused disease. This result suggests that, although all the strains of WSSV used were equally virulent, a difference in fitness was observed [89]. A recent study by Hoa et al. [31] showed that WSSV mixed-genotype infections correlate with lower outbreak incidence and that disease outbreaks correlate with single-genotype infections. Further studies need to be taken to understand the functions of various WSSV genes, which in turn will explain the reasons for differential virulence of various WSSV strains and viral latency in asymptomatic carriers.

Techniques for WSSV Detection

It is extremely important to understand that diagnosis for WSSV infection cannot be based only on the gross signs of white inclusions in the cuticle. Studies have shown that bacterial infections of the cuticle can also be associated with the formation of white inclusions, in the absence of WSSV infection [143]. The white spots produced closely resemble those caused by WSSV. They are associated with the presence of rod shaped bacteria revealed by scanning electron microscopy. Thus, white spots in the cuticle are unreliable for diagnosis of WSSV. Histopathology has been one of the pioneer techniques for the detection of WSSV. Other techniques which are developed include conventional PCR, real-time PCR, in situ hybridization, isothermal amplification and monoclonal antibody based detection methods are currently in use. Histology has been the gold coin for detection of WSSV in cases when clinical signs and histological changes are observed but other sensitive diagnostic methods like PCR are helpful even when histopathological changes are not seen. A large number of commercial kits, based on in situ hybridization, PCR and immunodetection, are also available (EBTL, DiagXotics, IQ2000, BIOTEC) for detection of WSSV.

DNA Based Diagnostic Methods

DNA Probe Based Methods

Non-radioactively labeled digoxigenin (DIG) DNA probe has been used in dot blot [22], in situ hybridization [9, 20, 77] and Southern blot hybridization for detection and analysis of WSSV [141]. Shekhar et al. [104] on comparing the sensitivity of DNA dot blot and PCR suggested that although PCR was found more sensitive to detect WSSV, DNA dot blot had an advantage over PCR in allowing use of those tissues which prevented PCR amplification due to inhibiting factors. DNA dot blot was found successful in detecting WSSV from the crude DNA extracted from

infected tissue samples such as eye stalk, eye stalk with eye, gills, cuticle, pleopod, periopods, uropods and telson.

Polymerase Chain Reaction (PCR)

Several PCR methods have been described for WSSV either in single [33, 57, 114, 115, 128] or together with other viruses in a multiplex PCR [48, 119]. For detection of WSSV in carrier animals such as wild caught marine shrimp, two-step nested PCR has been developed [32]. In India, PCR has been used for detection WSSV in cultured and captured crustaceans [128]. In Taiwan, Tsai et al. [119] designed a multiplex RT PCR for detection of WSSV and TSV in penaeids shrimp. A multiplex reverse transcription-polymerase chain reaction (mRT-PCR) was developed for simultaneously detection of six major shrimp viruses including WSSV, YHV, TSV, HPV, IHNV and MBV [48]. Recently multiplex PCR was developed for simultaneous detection of four DNA viruses including WSSV, IHNV, MBV and HPV [69]. The protocol described by Lo et al. [57] is the standard method used by the International Organization for Animal Health, although a recent publication suggests that under some conditions, this test may give false positive results with the Australian crayfish *Cherax quadricarinatus* [15]. A rapid and modified PCR assay has been developed based on the nested PCR procedure described by Lo et al. [57] for detection of WSSV [78].

A rapid and highly sensitive real-time PCR detection and quantification method for dual detection of WSSV and IHNV infecting penaeid shrimp has been developed using SYBR Green chemistry [18]. A recent comparison using real-time PCR as the gold standard clearly showed that, nested PCR tests were generally superior to one-step PCR tests in that they could reliably detect approximately 10–50 virions per reaction test vial [109]. Recently in South Korea a TaqMan real-time polymerase chain reaction was used to quantify and compare infection of WSSV in cultured *P. (Fenneropenaeus) chinensis* [42]. A real-time multiplex polymerase chain reaction (rtm-PCR) assay was developed for simultaneous detection of WSSV, IHNV and TSV by using TaqMan probes [149]. A multiplex real-time PCR and high-resolution melting (HRM) analysis has been developed to detect simultaneously three of the major viruses of penaeid shrimp including WSSV, YHV and *Penaeus monodon* densovirus (*PmDNV*) [82]. Recently real-time PCR assay was developed for detection of WSSV infection in *P. monodon* with sensitivity of 10 copies/ μ l [69].

Though PCR has been one of the popular detection methods, there have been reports of inter laboratory variations as evident from the ring testing conducted in Thailand [109], India and Indonesia [29]. The main reasons for

false positive results could be cross contamination and for false negative could be the primers designed targeting the deletion region. Marks et al. [65] showed that, the probe designed by Nunan and Lightner [77] for in situ hybridization gave false negative results as the sequence of probe was in the deletion region, which is absent in few strains of WSSV. This suggests that, the primers and probe should be designed only targeting conserved region of virus.

Serological Based Detection Methods

As an alternative to PCR, the progress in the use of immuno-based detection methods for shrimp viruses has been rather slow. This is primarily because of the difficulties associated with the availability of a purified antigen for immunization and availability of purified antibodies. Immunologically-based diagnosis methods using both polyclonal antibodies (PABs) and monoclonal antibodies (MAbs) have been developed for the detection of WSSV [1, 3, 4, 17, 56, 64, 74, 80, 85, 86, 105, 124, 153, 154, 156, 157] with a diagnostic sensitivity comparable to single step PCR. VP28 is one of the most abundant envelope proteins of WSSV and located on the surface of the viral particles and is crucial for virus entry [125]. This protein has been extensively used for development of MAbs [1, 34, 56]. The monoclonal antibodies specific to VP28 of WSSV could detect 500 pg protein by immunoperoxidase dot blotting [1] and 625 pg by the immunocomb detection method [64] whereas, about 76 % of the nested PCR-positive samples could be detected by immunodot blot using polyclonal antibodies specific to recombinant VP28 protein of WSSV [105]. Major advantage of immuno based detection methods is the ability to detect and grade the severity of viral infections as grading of infections is one of the important criteria required for the proper management of WSSV in culture systems [59].

A simple and rapid dot-blot nitrocellulose enzyme immunoassay (DB-NC-EIA) was developed by using HRP-conjugated virus-specific antibodies for the detection of WSV and YHV in infected shrimp [73]. Another sensitive, specific and simple immunodot assay was developed for the rapid detection of WSSV using polyclonal antiserum specific to the 27.5 kDa envelope protein of WSSV. The immunodot assay detected WSSV in a 1:1,000 dilution of a 10 % gill homogenate and approximately the equivalent of 5 ng total protein [153]. In 2004, a reversed passive latex agglutination (RPLA) assay was developed for detection of WSSV from stomach tissue homogenate of the kuruma shrimp, using high-density latex particles and specific polyclonal antibody in Japan [80]. Enzyme-linked immunosorbent assay (ELISA) and immunodot blot assay was developed for detection of WSSV in shrimp samples using

MAbs against WSSV as primary antibody, and goat anti-mouse Ig serum conjugated with alkaline phosphatase (AP) as secondary antibody [155]. A cost effective, a latex agglutination assay was developed using anti-18 kDa antiserum to detect WSSV in shrimps and crabs and detected the virus as early as 24 h post infection in shrimps [101]. A rapid, single step, sensitive and specific immunochromatographic test (ICT) was described for detecting WSSV using anti-WSSV MAbs, labeled with colloidal gold as a detection reagent and another anti-WSSV MAb as a capture antibody immobilized on nitrocellulose membrane (NCM) [137]. An immunoblot assay using an immunocomb was developed to detect WSSV infection from field samples and had an analytical sensitivity of 625 pg of purified antigen [64]. A sensitive and rapid lateral-flow immunoassay (LFIA) was developed for detection of WSSV using colloidal gold as an indicator with polyclonal antibody raised against recombinant VP (19+8) fusion protein [11]. A study was conducted to compare two chromogenic substrates (3,3',5,5'-tetramethylbenzidine (TMB) and 3–3' diaminobenzidine tetrahydrochloride (DAB)) by immunodot blot assay for detection of WSSV. Result suggested that TMB is more sensitive than the DAB method [105]. A immunochromatographic strip test for the simultaneous detection of WSSV and YHV was described using MAbs specific to the WSSV major envelope protein VP28 and the YHV nucleocapsid protein p20 [107]. Recently antibody-based microarray was developed for detection of WSSV in multiple samples [148].

Other Diagnostic Methods

Loop-mediated isothermal amplification (LAMP) method [76] is a novel, sensitive and rapid technique which can be applied for diagnosis of WSSV in aquaculture. It is a highly specific and sensitive diagnostic system using set of four primers, two outer and two inner primers. The detection limit using the LAMP method was found up to 1 fg, when compared to 10 fg by nested PCR [50]. Recently by combining LAMP and amplicon-specific fluorescence energy transfer-based probes, a real-time target amplicon-specific assay was successfully developed for accurate diagnosis of WSSV [14]. A highly sensitive method, capacitive biosensor was developed for the quantitative detection of WSSV in shrimp pond water samples with a linearity that ranged from 1 to 1×10^5 copies/ μ l [99]. Further studies need to be taken up to know the proportion or ratio of concentration of WSSV DNA and host DNA for getting up a positive reaction for each of the DNA based diagnostics methods. Studies also need to be carried out to know the effect of various preservatives used for storing the tissue samples and period for which the viral DNA is stable in them.

Concluding Remarks

WSSV is now enzootic and widespread in the culture environment, i.e., it is present in many crabs, wild shrimp and plankton crustaceans. Despite the application of numerous strategies, the problems associated with WSSV infection still remains unabated. Fitness and virulence studies on different genotypes of WSSV suggest that, all the strains of WSSV are equally virulent, but the one with smaller genomic size is the fittest. Studies also have shown that, mixed genotype infection of WSSV correlates with lower disease outbreaks. Till date, as there are no treatments available for WSSV infection, good management practices including stocking PCR tested WSSV negative PL may help to prevent the occurrence of WSSV infection in shrimp culture systems. Further research need to be carried out to deploy effective strategies to overcome the spread of WSD by developing different vaccines based on various WSSV proteins and also to develop a cost effective and practical mode for administering it. The changes occurring in WSSV genome due to various environmental stresses which acts as a trigger for WSSV outbreak need studied.

Acknowledgments The authors are gratefully acknowledging the funding support from the Department of Biotechnology, Government of India.

References

1. Anil TM, Shankar KM, Mohan CV. Monoclonal antibodies developed for sensitive detection and comparison of white spot syndrome virus isolates in India. *Dis Aquat Organ*. 2002;51: 67–75.
2. Cai S, Huang J, Wang C, Song X, Sun X, Yu J, Yan Z, Coughai Y. Epidemiological studies on explosive epidemic disease of prawn 1993–1994. *J Fish China*. 1995;19:112–7.
3. Chaivisuthangkura P, Tangkhabuanbutra J, Longyant S, Sithigornkul W, Rukpratanporn S, Menasveta P, Sithigornkul P. Monoclonal antibodies against a truncated viral envelope protein (VP28) can detect white spot syndrome virus (WSSV) infections in shrimp. *Sci Asia*. 2004;30:359–63.
4. Chaivisuthangkura P, Longyant S, Rukpratanporn S, Srisuk C, Sridulyakul P, Sithigornkul P. Enhanced white spot syndrome virus (WSSV) detection sensitivity using monoclonal antibody specific to heterologously expressed VP19 envelope protein. *Aquaculture*. 2010;299:15–20.
5. Chakraborty A, Otta SK, Joseph B, Kumar S, Hossain MS, Karunasagar I, Venugopal MN, Karunasagar I. Prevalence of white spot syndrome virus in wild crustaceans along the coast of India. *Curr Sci*. 2002;82:1392–7.
6. Chang PS, Lo CF, Wang YC, Kou GH. Identification of white spot syndrome associated baculovirus (WSBV) target organs in the shrimp *Penaeus monodon* by in situ hybridization. *Dis Aquat Organ*. 1996;27:131–9.
7. Chang PS, Chen HC, Wang YC. Detection of white spot syndrome associated baculovirus in experimentally infected wild shrimp, crab and lobsters by in situ hybridization. *Aquaculture*. 1998;164:231–3.

8. Chang YS, Peng SE, Wang HC, Hsu HC, Ho CH, Wang CH, Wang SY, Lo CF, Kou GH. Sequencing and amplified restriction fragment length polymorphism analysis of ribonucleotide reductase large subunit gene of the white spot syndrome virus in blue crab (*Callinectes sapidus*) from American coastal waters. *Mar Biotechnol*. 2001;3:163–71.
9. Chapman RW, Craig LB, Suzanne S, Sarah P, Wenner E. Sampling and evaluation of white spot syndrome virus in commercially important Atlantic penaeid shrimp stocks. *Dis Aquat Organ*. 2004;59:179–85.
10. Chen LL, Lo CF, Chiu YL, Chang CF, Kou GH. Natural and experimental infection of white spot syndrome virus (WSSV) in benthic larvae of mud crab *Scylla serrata*. *Dis Aquat Organ*. 2000;40:157–61.
11. Cheng QY, Meng XL, Xu JP, Lu W, Wang J. Development of lateral-flow immunoassay for WSSV with polyclonal antibodies raised against recombinant VP (19+28) fusion protein. *Virol Sin*. 2009;22:61–7.
12. Chou HY, Huang CY, Wang CH, Chiang HC, Lo CF. Pathogenicity of a baculovirus infection causing White spot syndrome in cultured penaeid shrimp in Taiwan. *Dis Aquat Organ*. 1995;23:165–73.
13. Chou HY, Huang CY, Lo CF, Kou GH. Studies on transmission of White spot syndrome associated baculovirus (WSBV) in *Penaeus monodon* and *P. japonicus* via waterborne contact and oral ingestion. *Aquaculture*. 1998;164:263–76.
14. Chou PH, Lin YC, Teng PH, Chen CL, Lee PY. Real-time target-specific detection of loop-mediated isothermal amplification for white spot syndrome virus using fluorescence energy transfer-based probes. *J Virol Methods*. 2011;173:67–74.
15. Claydon K, Cullen B, Owens L. OIE white spot syndrome virus PCR gives false-positive results in *Cherax quadricarinatus*. *Dis Aquat Organ*. 2004;62:265–8.
16. Corbel V, Zuprizal Shi Z, Huang C, Sumartono Arcier JM, Bonami JR. Experimental infection of European crustaceans with white spot syndrome virus (WSSV). *J Fish Dis*. 2001;24:377–82.
17. Dai H, Gao H, Zhao X, Dai L, Zhang X, Xiao N, Zhao R, Hemmingsen SM. Construction and characterization of a novel recombinant single-chain variable fragment antibody against white spot syndrome virus from shrimp. *J Immunol Methods*. 2003;279:267–75.
18. Dhar AK, Roux MM, Klimpel KR. Detection and quantification of infectious hypodermal and hematopoietic necrosis virus and white spot virus in shrimp using real-time quantitative PCR and SYBR green chemistry. *J Clin Microbiol*. 2001;39:2835–45.
19. Dieu BTM, Marks H, Siebenga JJ, Goldbach RW, Zuidema D, Duong TP, Vlcek JM. Molecular epidemiology of white spot syndrome virus within Vietnam. *J Gen Virol*. 2004;85:3607–18.
20. Durand S, Lightner DV, Nunan LM, Redman RM, Mari J, Bonami JR. Application of gene probes as a diagnostic tool for the white spot baculovirus (WSBV) of penaeid shrimp. *Dis Aquat Organ*. 1996;27:59–66.
21. Durand S, Lightner DV, Redman RM, Bonami JR. Ultrastructure and morphogenesis of white spot syndrome baculovirus (WSSV). *Dis Aquat Organ*. 1997;29:205–11.
22. Edgerton BF. Susceptibility of the Australian freshwater crayfish *Cherax destructor albidus* to white spot syndrome virus (WSSV). *Dis Aquat Organ*. 2004;59:187–93.
23. FAO. FAO Fisheries Department, Fishery information, data and statistics unit. FISHSTAT Plus: Universal software for fishery statistical time series. Version 2.30. 25; 2001.
24. Flegel TW. Major viral diseases of the black tiger prawn (*Penaeus monodon*) in Thailand. *World J Microbiol Biotechnol*. 1997;13:433–42.
25. Flegel TW. Detection of major penaeid shrimp viruses in Asia, a historical perspective with emphasis on Thailand. *Aquaculture*. 2006;258:1–33.
26. Galaviz-Silva L, Molina-Garza ZJ, Alcocer-Gonzalez JM, Rosales-Encinas JL, Ibarra-Gamez C. White spot syndrome virus genetic variants detected in Mexico by a new multiplex PCR method. *Aquaculture*. 2004;242:53–68.
27. Gopalakrishnan A, Rajkumar M, Sun J, Wang M, Kumar SK. Mud crab, *Scylla tranquebarica* (Decapoda: Portunidae), a new host for the white spot syndrome virus. *Aquac Res*. 2011;42:308–12.
28. Griffith DRW. White spot and Taura syndrome virus: disease as drivers in the shrimp farming industry. In: Alday-Sanz V, editor. *The shrimp book*. Nottingham: Nottingham University Press; 2010. p. 429–41.
29. Gudkovs N, Pradeep B, Stalin Raj V, Slater J, Sittidilokratna N, Mohan CV, Walker PJ. Application of PCR for improved shrimp health management in the Asian region. In: *FRDC aquatic animal health subprogram scientific conference*. Cairns; 2007.
30. Hennig O, Itami T, Maeda M, Kondo M, Natsukari Y, Takahashi Y. Analyses of hemolymph immunoparameters in kuruma shrimp infected with penaeid rod-shaped DNA virus. *Fish Pathol*. 1998;33:389–93.
31. Hoa TT, Zwart MP, Phuong NT, Oanh DT, Mart CM, Vlcek JM. Mixed-genotype white spot syndrome virus infections of shrimp are inversely correlated with disease outbreaks in ponds. *J Gen Virol*. 2011;92:675–80.
32. Hossain MS, Chakraborty A, Joseph B, Otta SK, Karunasagar I, Karunasagar I. Detection of new hosts for white spot syndrome virus of shrimp using nested polymerase chain reaction. *Aquaculture*. 2001;198:1–11.
33. Hossain MS, Khadijah S, Kwang J. Characterization of ORF89—a latency-related gene of white spot syndrome virus. *Virology*. 2004;325:106–15.
34. Hou C, Cao Y, Xie R, Wang Y, Du H. Characterization and diagnostic use of a monoclonal antibody for VP28 envelope protein of white spot syndrome virus. *Virol Sin*. 2011;26:260–6.
35. Hsu HC, Lo CF, Lin SC, Liu KF, Peng SE, Chang YS, Chen LL, Liu WJ, Kou GH. Studies on effective PCR screening strategies for white spot syndrome virus (WSSV) detection in *Penaeus monodon* brooders. *Dis Aquat Organ*. 1999;39:13–9.
36. Huang J, Cai S, Song X, Wang C, Yu J, Yang C. Study on artificial infection for *Penaeus chinensis* by the pathogen of explosive endemic disease of shrimp. *Mar Fish Res*. 1995;16:51–8.
37. Huang C, Zhang X, Lin Q, Xu X, Hew CL. Characterization of a novel envelope protein (VP281) of shrimp white spot syndrome virus by mass spectrometry. *J Gen Virol*. 2002;83:2385–92.
38. Huang C, Zhang X, Lin Q, Xu X, Hew CL. Proteomic analysis of shrimp white spot syndrome viral proteins and characterization of a novel envelope protein VP466. *Mol Cell Proteomics*. 2002;1:223–31.
39. ICAR. Shrimp farming. In: *Handbook of fisheries and aquaculture*. New Delhi: Indian Council of Agricultural Research; 2006. p. 392–403.
40. Inouye K, Miwa S, Oseko N, Nakano H, Kimura T, Momoyama K, Hiraoka M. Mass mortality of cultured kuruma shrimp *Penaeus japonicus* in Japan in 1993: electron microscopic evidence of the causative virus. *Fish Pathol*. 1994;29:149–58.
41. Inouye K, Yamano K, Ikeda N, Kimura T, Nakano H, Momoyama K, Kobayashi J, Miyajima S. The penaeid rod-shaped DNA virus (PRDV), which causes penaeid acute viremia (PAV). *Fish Pathol*. 1996;31:39–45.
42. Jang IK, Kim JS, Kim BR, Meng XH. Comparison of white spot syndrome virus quantification of fleshy shrimp *Fenneropenaeus*

- chinensis* in outdoor ponds between different growing seasons by TaqMan real-time polymerase chain reaction. *Aquac Res.* 2011;42:1869–77.
43. Jiravanichpaisal P, Bangyeekhun E, Soderhall K, Soderhall I. Experimental infection of white spot syndrome virus in freshwater crayfish *Pacifastacus leniusculus*. *Dis Aquat Organ.* 2001;47:151–7.
 44. Jiravanichpaisal P, Soderha K, Soderha I. Characterization of white spot syndrome virus replication in in vitro-cultured haematopoietic stem cells of freshwater crayfish, *Pacifastacus leniusculus*. *J Gen Virol.* 2006;87:847–54.
 45. Karunasagar I, Otta SK, Karunasagar I. Histopathological and bacteriological study of white spot syndrome in *Penaeus monodon* along the west coast of India. *Aquaculture.* 1997;153:9–13.
 46. Karunasagar I, Otta SK, Karunasagar I. Disease problems affecting cultured penaeid shrimp in India. *Fish Pathol.* 1998;33:413–9.
 47. Khadijah S, Ying Neo S, Hossain MS, Miller LD, Mathavan S, Kwang J. Identification of white spot syndrome virus latency-related genes in specific-pathogen-free shrimps by use of a microarray. *J Virol.* 2003;77:10162–7.
 48. Khawsak P, Deesukon W, Chaivisuthangkura P, Sukhumsiri-chart W. Multiplex RT-PCR assay for simultaneous detection of six viruses of penaeid shrimp. *Mol Cell Probes.* 2008;22:177–83.
 49. Kim CK, Kim PK, Sohn SG, Sim DS, Park MA, Heo MS, Lee TH, Lee JD, Jun HK, Jang KL. Development of a polymerase chain reaction (PCR) procedure for the detection of baculovirus associated with white spot syndrome (WSBV) in penaeid shrimp. *J Fish Dis.* 1998;21:11–7.
 50. Kono T, Savan R, Sakai M, Itami T. Detection of white spot syndrome virus in shrimp by loop-mediated isothermal amplification. *J Virol Methods.* 2004;115:59–65.
 51. Lan Y, Lu W, Xu X. Genomic instability of prawn white spot bacilliform virus (WSBV) and its association to virus virulence. *Virus Res.* 2002;90:269–74.
 52. Leu JH, Tsai JM, Wang HC, Wang AH, Wang CH, Kou GH, Lo CF. The unique stacked rings in the nucleocapsid of the white spot syndrome virus virion are formed by the major structural protein VP664, the largest viral structural protein ever found. *J Virol.* 2005;79:140–9.
 53. Leu JH, Yang F, Zhang X, Xu X, Kou GH, Lo CF. Whispovirus. In: Van Etten JL, editor. *Lesser known large dsDNA viruses. Current topics in microbiology and immunology.* Berlin: Springer; 2009. p. 328.
 54. Li Q, Zhang J, Chen Y, Yang F. White spot syndrome virus (WSSV) infectivity for *Artemia* at different developmental stages. *Dis Aquat Organ.* 2003;57:261–4.
 55. Lightner DV. A handbook of pathology and diagnostic procedures for diseases of penaeid shrimp. Baton Rouge: Special publication of the World Aquaculture Society; 1996.
 56. Liu W, Wang YT, Tian DS, Yin ZC, Kwang J. Detection of white spot syndrome virus (WSSV) of shrimp by means of monoclonal antibodies (MAbs) specific to an envelope protein (28 kDa). *Dis Aquat Organ.* 2002;49:11–8.
 57. Lo CF, Leu JH, Ho CH, Chen CH, Peng SE, Chen YT, Chou CM, Yeh PY, Huang CJ, Chou HY, Wang CH, Kou GH. Detection of baculovirus associated with white spot syndrome (WSBV) in penaeid shrimps using polymerase chain reaction. *Dis Aquat Organ.* 1996;25:133–41.
 58. Lo CF, Ho CH, Chen CH, Liu KF, Chiu YL, Yeh PY, Peng SE, Hsu HC, Liu HC, Chang CF, Su MS, Wang CH, Kou GH. Detection and tissue tropism of White spot syndrome baculovirus (WSBV) in captured brooders of *Penaeus monodon* with a special emphasis on reproductive organs. *Dis Aquat Organ.* 1997;30:53–72.
 59. Lo CF, Chang YS, Cheng CT, Kou H. Monitoring cultured shrimp for white spot syndrome virus (WSSV) infection during their growth period by polymerase chain reaction. In: Flegel TW, editor. *Advances in shrimp biotechnology.* Bangkok: National Center for Genetic Engineering and Biotechnology; 1998. p. 281–6.
 60. Lo CF, Hsu HC, Tsai MF, Ho CH, Peng SE, Kou GH, Lightner DV. Specific genomic DNA fragment analysis of different geographical clinical samples of shrimp white spot syndrome virus. *Dis Aquat Organ.* 1999;35:175–85.
 61. Lotz JM. Viruses, biosecurity and specific pathogen free stocks in shrimp aquaculture. *World J Microbiol Biotechnol.* 1997;13:405–13.
 62. Lotz JM, Soto MA. Model of white spot syndrome virus (WSSV) epidemics in *Litopenaeus vannamei*. *Dis Aquat Organ.* 2002;50:199209.
 63. Maeda M, Itami T, Furumoto A, Hennig O, Imamura T, Kondo M, Hirano I, Takashi A, Takahashi Y. Detection of penaeid rod-shaped DNA virus (PRDV) in wild-caught shrimp and other crustaceans. *Fish Pathol.* 1998;33:373–80.
 64. Makesh M, Koteeswaran A, Daniel Joy Chandran N, Murali Manohar B, Ramasamy V. Development of monoclonal antibodies against VP28 of WSSV and its application to detect WSSV using immunocomb. *Aquaculture.* 2006;261:64–71.
 65. Marks H, Goldbach RW, Vlak JM, van Hulten MCW. Genetic variation among isolates of white spot syndrome virus. *Arch Virol.* 2004;149:673–97.
 66. Marks H, van Duijse JJ, Zuidema D, van Hulten MC, Vlak JM. Fitness and virulence of an ancestral white spot syndrome virus isolate from shrimp. *Virus Res.* 2005;110:9–20.
 67. Mayo MA. Virus taxonomy—Houston 2002. *Arch Virol.* 2002;147:1071–6.
 68. Mayo MA. A summary of taxonomic changes recently approved by ICTV. *Arch Virol.* 2002;14:1655–6.
 69. Mohan SA. Development of molecular based diagnostics for simultaneous detection of viruses of crustaceans. PhD Thesis, Department of Fishery Microbiology, Karnataka Veterinary, Animal and Fisheries Sciences University, College of Fisheries, Mangalore; 2012.
 70. Mohan SA. Study on natural reservoirs of shrimp pathogenic viruses. MSc Thesis, Department of Fishery Microbiology, Karnataka Veterinary, Animal and Fisheries Sciences University, College of Fisheries, Mangalore; 2008.
 71. Musthaq SS, Sudhakaran R, Balasubramanian G, Sahul Hameed AS. Experimental transmission and tissue tropism of white spot syndrome virus (WSSV) in two species of lobsters *Panulirus homarus* and *Panulirus ornatus*. *J Invertebr Pathol.* 2006;93:75–80.
 72. Nadala ECB, Loh PC. A comparative study of three different isolates of white spot virus. *Dis Aquat Organ.* 1998;33:231–4.
 73. Nadala ECB, Loh PC. Dot-blot nitrocellulose enzyme immunoassays for the detection of white-spot virus and yellow-head virus of penaeid shrimp. *J Virol Methods.* 2000;84:175–9.
 74. Nadala ECB, Tapay LM, Cao S, Loh PC. Detection of yellow head virus and Chinese baculovirus in penaeid shrimp by western blot technique. *J Virol Methods.* 1997;69:39–44.
 75. Natividad KDT, Nomura N, Matsumura M. Detection of white spot syndrome virus DNA in pond soil using a 2-step nested PCR. *J Virol Methods.* 2008;149:28–34.
 76. Notomi T, Okayama H, Masubuchi H, Yonekawa T, Watanabe K, Amino N, Hase T. Loop-mediated isothermal amplification of DNA. *Nucleic Acids Res.* 2000;28:e63.
 77. Nunan LM, Lightner DV. Development of a non-radioactive gene probe by PCR for detection of white spot syndrome virus (WSSV). *J Virol Methods.* 1997;63:193–201.

78. Nunan LM, Lightner DV. Optimized PCR assay for detection of white spot syndrome virus (WSSV). *J Virol Methods*. 2011; 171:318–21.
79. OIE. Diagnostic manual for aquatic animal diseases. 4th ed. Paris: Office International des Epizooties (OIE); 2003. p. 358.
80. Okumura T, Nagai F, Yamamoto S, Yamano K, Oseko N, Inouye K, Oomura H, Sawada H. Detection of white spot syndrome virus from stomach tissue homogenate of the kuruma shrimp (*Penaeus japonicus*) by reverse passive latex agglutination. *J Virol Methods*. 2004;119:11–6.
81. Otta SK, Shubha G, Joseph B, Chakraborty A, Karunasagar I, Karunasagar I. Polymerase chain reaction (PCR) detection of white spot syndrome virus (WSSV) in cultured and wild crustaceans in India. *Dis Aquat Organ*. 1999;38:67–70.
82. Panichareon B, Khawsak P, Deesukon W, Sukhumsirichart W. Multiplex real-time PCR and high-resolution melting analysis for detection of white spot syndrome virus, yellow-head virus, and *Penaeus monodon* densovirus in penaeid shrimp. *J Virol Methods*. 2011;178:16–21.
83. Peng SE, Lo CF, Ho CH, Chang CF, Kou GH. Detection of white spot baculovirus (WSBV) in giant freshwater prawn, *Macrobrachium rosenbergii* using polymerase chain reaction. *Aquaculture*. 1998;164:253–62.
84. Peng SE, Lo CF, Lin SC, Chen LL, Chang YS, Liu KF, Su MS, Kou GH. Performance of WSSV-infected and WSSV-negative *Penaeus monodon* postlarvae in culture ponds. *Dis Aquat Organ*. 2001;46:165–72.
85. Poulos BT, Pantoja CR, Bradley-Dunlop D, Aguilar J, Lightner DV. Development and application of monoclonal antibodies for the detection of white spot syndrome virus of penaeid shrimp. *Dis Aquat Organ*. 2001;47:13–23.
86. Powell JWB, Burge EJ, Browdy CL, Shepard EF. Efficiency and sensitivity determination of shrimples, an immunochromatographic assay for white spot syndrome virus (WSSV), using quantitative real-time PCR. *Aquaculture*. 2006;257:167–72.
87. Pradeep B, Shekar M, Gudkovs N, Karunasagar I, Karunasagar I. Genotyping of white spot syndrome virus prevalent in shrimp farms of India. *Dis Aquat Organ*. 2008;78:189–98.
88. Pradeep B, Shekar M, Karunasagar I, Karunasagar I. Characterization of variable genomic regions of Indian white spot syndrome virus. *Virology*. 2008;376:24–30.
89. Pradeep B, Karunasagar I, Karunasagar I. Fitness and virulence of different strains of white spot syndrome virus. *J Fish Dis*. 2009;32:801–5.
90. Rajan PR, Ramasamy P, Purushothaman V, Brennan GP. White spot baculovirus syndrome virus in the Indian shrimp *Penaeus monodon* and *Penaeus indicus*. *Aquaculture*. 2000; 184:31–44.
91. Rajendran KV, Vijayan KK, Santiago TC, Krol RM. Experimental host range and histopathology of white spot syndrome virus (WSSV) infection in shrimp, prawns, crabs and lobsters from India. *J Fish Dis*. 1999;22:183–91.
92. Ramasamy P. Diseases of shrimp in aquaculture systems: diagnosis and therapeutic measures. Chennai: Vanitha Publications; 1995. p. 1–99.
93. Rosenberry B. World shrimp farming 1996. San Diego: Shrimp News International; 1996.
94. Rosenberry B. World shrimp farming 2000. San Diego: Shrimp News International; 2000.
95. Rosenberry B. World shrimp farming 2002. San Diego: Shrimp News International; 2002.
96. Sahul-Hameed AS, Charles MX, Anikumar M. Tolerance of *Macrobrachium rosenbergii* to white spot syndrome virus. *Aquaculture*. 2000;183:207–13.
97. Sahul-Hameed AS, Yoganandhan K, Sathish S, Rasheed M, Murugan V, Jayarama K. White spot syndrome virus (WSSV) in two species of freshwater crabs (*Paratelphusa hydrodomus* and *P. pulvinata*). *Aquaculture*. 2001;201:179–86.
98. Sahul-Hameed AS, Balasubramanian G, Syed Musthaq S, Yoganandhan K. Experimental infection of twenty species of Indian marine crabs with white spot syndrome virus (WSSV). *Dis Aquat Organ*. 2003;57:157–61.
99. Samanman S, Kanatharana P, Chotigeat W, Deachamag P, Thavarungkul P. Highly sensitive capacitive biosensor for detecting white spot syndrome virus in shrimp pond water. *J Virol Methods*. 2011;173:75–84.
100. Sanchez-Paz A. White spot syndrome virus: an overview on an emergent concern. *Vet Res*. 2010;41:3.
101. Sathish S, Selvakumar C, Sahul Hameed AS, Narayanan RB. 18-kDa protein as a marker to detect WSSV infection in shrimps. *Aquaculture*. 2004;238:39–50.
102. Shekar M, Karunasagar I, Karunasagar I. A computer-based identification of variable number tandem repeats in white spot syndrome virus genomes. *Curr Sci*. 2005;89:882–7.
103. Shekhar MS, Ravichandran P. Comparison of white spot syndrome virus structural gene sequences from India with those at GenBank. *Aquac Res*. 2007;38:321–4.
104. Shekhar MS, Azad IS, Ravichandran P. Comparison of dot blot and PCR diagnostic techniques for detection of white spot syndrome virus in different tissues of *Penaeus monodon*. *Aquaculture*. 2006;261:1122–7.
105. Shekhar MS, DilliKumar, Gopikrishna G. Development of immunodot blot assay for the detection of white spot syndrome virus infection in shrimps (*Penaeus monodon*). *Aquac Res*. 2010;41:1683–90.
106. Shi Z, Huang C, Zhang J, Chen D, Bonami JR. White spot syndrome virus (WSSV) experimental infection of the freshwater crayfish, *Cherax quadricarinatus*. *J Fish Dis*. 2000; 23:285–8.
107. Sithigorngula P, Rukpratanporn S, Chaivisuthangkura P, Sridulyakul P, Longyant S. Simultaneous and rapid detection of white spot syndrome virus and yellow head virus infection in shrimp with a dual immunochromatographic strip test. *J Virol Methods*. 2011;173:85–91.
108. Somboonna N, Mangkalan S, Udompetcharaporn A, Krittanai C, Sritunyalucksana K, Flegel TW. Mud crab susceptibility to disease from white spot syndrome virus is species-dependent. *BMC Res Notes*. 2010;3:315.
109. Sritunyalucksana K, Srisala J, Mccoll K, Nielsen L, Flegel TW. Comparison of PCR testing methods for white spot syndrome virus (WSSV) infections in penaeid shrimp. *Aquaculture*. 2006;255:95–104.
110. Stalin Raj V, Vijayan KK, Sanjuktha M, Balasubramanian CP, Alavandi SV, Santiago TC. Virulence status, viral accommodation and structural protein profiles of white spot syndrome virus isolates in farmed *Penaeus monodon* from the southeast coast of India. *Aquac Res*. 2009;40:129–38.
111. Supamattaya K, Hoffmann RW, Boonyaratpalin S, Kanchanaphum P. Experimental transmission of white spot syndrome virus (WSSV) from black tiger shrimp *Penaeus monodon* to the sand crab *Portunus pelagicus*, mud crab *Scylla serrata* and krill *Acetes* sp. *Dis Aquat Organ*. 1998;32:79–85.
112. Takahashi Y, Fukuda K, Kondo M, Chongthaleong A, Nishi K, Nishimura M, Ogata K, Shinya I, Takise K, Fujishima K, Matsumura M. Detection and prevention of WSSV infection in cultured shrimp. *Asian Aquaculture Magazine*; 2003. p. 25–7.
113. Takahashi Y, Itami T, Kondom M, Maeda M, Fujii R, Tomonaga S, Supamattaya K, Boonyaratpalin S. Electron microscopic evidence of bacilliform virus infection in Kuruma shrimp (*Penaeus japonicus*). *Fish Pathol*. 1994;29:121–5.
114. Takahashi Y, Itami T, Maeda M, Suzuki N, Kasornchandra J, Supamattaya K, Khongpradit R, Boonyaratpalin S, Kondo M,

- Kawai K, Kusuda R, Hirono I, Aoki T. Polymerase chain reaction (PCR) amplification of bacilliform virus (RV-PS) DNA in *Penaeus japonicus* Bate and systemic ectodermal and mesodermal baculovirus (SEMBV) DNA in *Penaeus monodon* Fabricius. *J Fish Dis*. 1996;19:399–403.
115. Tang KFJ, Lightner DV. Quantification of white spot syndrome virus DNA through a competitive polymerase chain reaction. *Aquaculture*. 2000;189:11–21.
 116. Tokhmafshan M, Akbari S, Tamjidi M, Laloi F, Soltani M. Occurrence of white spot syndrome disease in farmed *P. indicus* in Iran. *Appl Fish Aquac*. 2004;4:42–7.
 117. Tsai MF, Kou GH, Liu HC, Liu KF, Chang CF, Peng SE, Hsu HC, Wang CH, Lo CF. Long-term presence of white spot syndrome virus (WSSV) in a cultivated shrimp population without disease outbreaks. *Dis Aquat Organ*. 1999;38:107–14.
 118. Tsai MF, Yu HT, Tzeng HF, Leu JH, Chou CM, Huang CJ, Wang CH, Lin JY, Kou GH, Vlo CF. Identification and characterization of a shrimp white spot syndrome virus (WSSV) gene that encodes a novel chimeric polypeptide of cellular-type thymidine kinase and thymidylate kinase. *Virology*. 2000;277:100–10.
 119. Tsai JM, Shiau LJ, Lee HH, Chan PW, Lin CY. Simultaneous detection of white spot syndrome virus (WSSV) and Taura syndrome virus (TSV) by multiplex reverse transcription-polymerase chain reaction (RT-PCR) in pacific white shrimp *Penaeus vannamei*. *Dis Aquat Organ*. 2002;50:9–12.
 120. Tsai JM, Wang HC, Leu JH, Hsiao HH, Wang AH, Kou GH, Lo CF. Genomic and proteomic analysis of thirty-nine structural proteins of shrimp White spot syndrome virus. *J Virol*. 2004;78:11360–70.
 121. van De Braak CB, Botterblom MH, Huisman EA, Rombout JH, Van Der Knaap WP. Preliminary study on haemocyte response to White spot syndrome virus infection in black tiger shrimp *Penaeus monodon*. *Dis Aquat Organ*. 2002;51:149–55.
 122. van Hulten MCW, Vlak JM. Identification and phylogeny of a protein kinase gene of white spot syndrome virus. *Virus Genes*. 2001;22:201–20.
 123. van Hulten MCW, Goldbach RW, Vlak JM. Three functionally diverged major structural proteins of white spot syndrome virus evolved by gene duplication. *J Gen Virol*. 2000;81:2525–9.
 124. van Hulten MCW, Westenberg M, Goodall SD, Vlak JM. Identification of two major virion protein genes of white spot syndrome virus of shrimp. *Virology*. 2000;266:227–36.
 125. van Hulten MCW, Witteveldt J, Snippe M, Vlak JM. White spot syndrome virus envelope protein VP28 is involved in the systemic infection of shrimp. *Virology*. 2001;285:228–33.
 126. van Hulten MCW, Witteveldt J, Peters S, Kloosterboer N, Tarchini R, Fiers M, Sandbrink H, Lankhorst RK, Vlak JM. The white spot syndrome virus DNA genome sequence. *Virology*. 2001;286:7–22.
 127. van Hulten MCW, Reijns M, Vermeesch AM, Zandbergen F, Vlak JM. Identification of VP19 and VP15 of white spot syndrome virus (WSSV) and glycosylation status of the WSSV major structural proteins. *J Gen Virol*. 2002;83:257–65.
 128. Vaseeharan B, Jayakumar R, Ramasamy P. PCR-based detection of white spot syndrome virus in cultured and captured crustaceans in India. *Lett Appl Microbiol*. 2003;37:443–7.
 129. Vijayan KK, Raj VS, Balasubramanian CP, Alavandi SV, Sekhar VT, Santiago TC. Polychaete worms—a vector for white spot syndrome virus (WSSV). *Dis Aquat Organ*. 2005;63:107–11.
 130. Vlak JM, Bonami JR, Flegel TW, Kou GH, Lightner DV, Lo CF, Loh PC, Walker PJ. *Nimaviridae* A new virus family infecting aquatic invertebrates. In: XIIIth international congress of virology. 2002.
 131. Vlak JM, Bonami JR, Flegel TW, Kou GH, Lightner DV, Lo CF, Loh C, Walker PW. *Nimaviridae*. In: Fauquet CM, Mayo MA, Maniloff J, Desselberger U, Ball LA, editors. VIIIth report of the international committee on taxonomy of viruses. Elsevier. 2005. p. 187–92.
 132. Walker PJ, Mohan CV. Viral disease emergence in shrimp aquaculture: origins, impacts and the effectiveness of health management strategies. *Rev Aquac*. 2009;1:125–54.
 133. Walker PJ. Application of PCR in management of shrimp viral diseases. PCR training workshop manual, Chennai; 2005. p. 1–5.
 134. Walker PJ, Winton JR. Emerging viral diseases of fish and shrimp. *Vet Res*. 2010;41:51.
 135. Walker PJ, Gudkovs N, Mohan CV, Stalin Raj V, Pradeep B, Sergeant E, Chandra Mohan AB, Ravibabu G, Karunasagar I, Santiago TC. Longitudinal disease studies in small-holder black tiger shrimp (*Penaeus monodon*) farms in Andhra Pradesh, India. II. Multiple WSSV genotypes associated with disease outbreaks in ponds seeded with uninfected postlarvae. *Aquaculture*. 2011;319:18–24.
 136. Walker PJ, Gudkovs N, Pradeep B, Stalin Raj V, Evan Sergeant AB, Chandra Mohan AB, Ravibabu G, Umesh NR, Karunasagar I, Santiago TC, Mohan CV. Longitudinal disease studies in small-holder black tiger shrimp (*Penaeus monodon*) farms in Andhra Pradesh, India. III. A complex dynamic of WSSV infection and WSSV genotype distribution in farmed shrimp and wild crustaceans. *Aquaculture*. 2011;319:319–27.
 137. Wang X, Zhan W. Development of an immunochromatographic test to detect white spot syndrome virus of shrimp. *Aquaculture*. 2006;255:196–200.
 138. Wang YC, Lo CF, Chang PS, Kou GH. Experimental infection of white spot baculovirus in some cultured and wild decapods in Taiwan. *Aquaculture*. 1998;164:221–31.
 139. Wang Q, White BL, Redman RM, Lightner DV. Per os challenge of *Litopenaeus vannamei* postlarvae and *Farfantepenaeus duorarum* juveniles with six geographic isolates of White spot syndrome virus. *Aquaculture*. 1999;170:179–94.
 140. Wang CH, Yang HN, Tang CY, Lu CH, Kou GH, Lo CF. Ultrastructure of White spot syndrome virus development in primary lymphoid organ cell cultures. *Dis Aquat Organ*. 2000;41:91–104.
 141. Wang Q, Nunan LM, Lightner DV. Identification of genomic variations among geographic isolates of white spot syndrome virus using restriction analysis and southern blot hybridization. *Dis Aquat Organ*. 2000;43:175–81.
 142. Wang Q, Poulos BT, Lightner DV. Protein analysis of geographic isolates of shrimp White spot syndrome virus. *Arch Virol*. 2000;145:263–74.
 143. Wang YG, Lee KL, Najiah M, Shariff M, Hassan MD. A new bacterial white spot syndrome (BWSS) in cultured tiger shrimp *Penaeus monodon* and its comparison with white spot syndrome (WSS) caused by virus. *Dis Aquat Organ*. 2000;41:9–18.
 144. Witteveldt J, Vermeesch AM, Langenhof M, de Lang A, Vlak JM, van Hulten MCW. Nucleocapsid protein VP15 is the basic DNA binding protein of white spot syndrome virus of shrimp. *Arch Virol*. 2005;150:1121–33.
 145. Wongprasert K, Khanobdee K, Glunukarn SS, Meeratana P, Withyachumnarnkul B. Time course and levels of apoptosis in various tissues of black tiger shrimp *Penaeus monodon* infected with white spot syndrome virus. *Dis Aquat Organ*. 2003;55:3–10.
 146. Wongteerasupaya C, Vickers JE, Sriurairatana S, Nash GL, Akarajamorn A, Boonsaeng V, Panyim S, Tassanakajon A, Withyachumnarnkul B, Flegel TW. A non-occluded, systemic baculovirus that occurs in cells of ectodermal and mesodermal origin and causes high mortality in the black tiger prawn *Penaeus monodon*. *Dis Aquat Organ*. 1995;21:69–77.

147. Wongteerasupaya C, Pungchai P, Withyachumnarnkul B, Boonsaeng V, Panyim S, Flegel TW, Walker PJ. High variation in repetitive DNA fragment length for white spot syndrome virus (WSSV) isolates in Thailand. *Dis Aquat Organ*. 2003;54:253–7.
148. Xiaoli X, Xiuzhen S, Wenbin Z. Development and application of antibody microarray for white spot syndrome virus detection in shrimp. *Chin J Oceanol Limnol*. 2011;29:930–41.
149. Xie Z, Xie L, Pang Y, Lu Z, Xie Z, Sun J, Deng X, Liu J, Tang X, Khan M. Development of a real-time multiplex PCR assay for detection of viral pathogens of penaeid shrimp. *Arch Virol*. 2008;153:2245–51.
150. Yadava YS. Shrimp farming in India—lessons and challenges in sustainable development. *Aquac Auth News*. 2002;1:1–4.
151. Yan DC, Dong SL, Huang J, Yu XM, Feng MY. White spot syndrome virus (WSSV) detected by PCR in rotifers and rotifer resting eggs from shrimp pond sediments. *Dis Aquat Organ*. 2004;59:69–73.
152. Yang F, He J, Lin XH, Li Q, Pan D, Zhang X, Xu X. Complete genome sequence of the shrimp white spot bacilliform virus. *J Virol*. 2001;75:11811–20.
153. You Z, Nadala ECB, Yang J, van Hulten MCW, Loh PC. Production of polyclonal antiserum specific to the 27.5 kDa envelope protein of white spot syndrome virus. *Dis Aquat Organ*. 2002;51:77–80.
154. Zhan WB, Chen J, Zhang ZD, Zhou L, Fukuda H. Elimination of shrimp endogenous peroxidase background in immunodot blot assays to detect white spot syndrome virus (WSSV). *Dis Aquat Organ*. 2003;53:263–5.
155. Zhan W, Wang X, Chen J, Xing J, Fukuda H. Elimination of shrimp endogenous alkaline phosphatase background and development of enzyme immunoassays for the detection of white spot syndrome virus (WSSV). *Aquaculture*. 2004;239:15–21.
156. Zhang X, Huang C, Xu X, Hew CL. Identification and localization of a prawn white spot syndrome virus gene that encodes an envelope protein. *J Gen Virol*. 2002;83:1069–74.
157. Zhang X, Huang C, Xu X, Hew CL. Transcription and identification of an envelope protein gene (p22) from shrimp white spot syndrome virus. *J Gen Virol*. 2002;83:471–7.
158. Zhang X, Huang C, Tang X, Zhuang Y, Hew CL. Identification of structural proteins from shrimp white spot syndrome virus (WSSV) by 2DE-MS. *Proteins*. 2004;55:229–35.
159. Zhang JS, Dong SL, Tian XL, Dong YW, Liu XY, Yan DC. Studies on the rotifer (*Brachionus urceus* Linnaeus, 1758) as a vector in white spot syndrome virus (WSSV) transmission. *Aquaculture*. 2006;261:1181–5.
160. Zhang JS, Dong SL, Dong YW, Tian XL, Hou CQ. Bioassay evidence for the transmission of WSSV by the harpacticoid copepod *Nitocra* sp. *J Invertebr Pathol*. 2007;97:33–9.