available at www.sciencedirect.com

journal homepage: www.elsevier.com/locate/jus

Role of color Doppler sonography in post-transplant surveillance of vascular complications involving pancreatic allografts[☆]

L. Morelli^{*}, G. Di Candio, A. Campatelli, F. Vistoli, M. Del Chiaro, E. Balzano, C. Croce, C. Moretto, S. Signori, U. Boggi, F. Mosca

General and Transplantation Surgery, University of Pisa, Cisanello Hospital, Pisa, Italy

KEYWORDS

Pancreas;
Transplantation;
Color Doppler
ultrasonography;
Vascular;
Complications.

Abstract *Purpose:* To evaluate the role of color Doppler ultrasonography in the postoperative surveillance of the vascular complications involving pancreas allografts.

Methods: A retrospective analysis of a consecutive series of 223 pancreas transplantations was performed. All recipients received antithrombotic prophylaxis, which was tailored to the individual's estimated risk of thrombosis. All patients were monitored with daily color Doppler ultrasonography during the first post-transplant week and thereafter whenever clinically indicated. Vascular complications were defined as all thrombotic events requiring: increased anticoagulant therapy, angiography with fibrinolytic therapy, or repeat surgery.

Results: The overall patient survival rates at one, three, and five years after transplantation were 94.7%, 93.3%, and 91%, respectively. The overall graft survival rates at the same time points were 87.4%, 79.6%, and 75.6%, respectively. In 28 of the 223 cases (12.5%) graft thromboses were diagnosed with Doppler ultrasound within the first 10 days after transplantation. In 3 cases, graft pancreatectomies were performed because of a complete loss of blood flow in the parenchyma. An attempt to rescue the graft was made in 18 patients. Fourteen of these grafts were saved and are still functioning (77.7%); and 4 rescue attempts failed and the grafts were subsequently explanted (32.3%).

Conclusion: Color Doppler ultrasound is a suitable tool for postoperative surveillance of pancreas transplant recipients. Its use can lead to early diagnosis and timely treatment of vascular complications.

Sommario *Scopo:* Valutare il ruolo del monitoraggio postoperatorio precoce con eco-color Doppler nel paziente sottoposto a trapianto di pancreas.

Materiali e metodi: Uno studio retrospettivo è stato effettuato su una serie consecutiva di 223 trapianti di pancreas effettuati presso il Centro Trapianti di Pisa. L'eco-color Doppler è stato sempre eseguito giornalmente i primi 7 giorni dopo il trapianto e ogni qualvolta le variazioni

[☆] EUROSON–SIUMB 2006 — SIUMB Award for the best oral communication presented at the 18th European Congress of Ultrasound held in conjunction with the XVII National Congress of the SIUMB.

^{*} Corresponding author. General and Transplantation Surgery, AO Universitaria Pisana, Ospedale Cisanello, via Paradisa 2, 56124 Pisa, Italy.

E-mail address: lu.morelli@ao-pisa.toscana.it (L. Morelli).

del quadro clinico e biochimico abbiano determinato il sospetto di una complicanza. Nel sospetto ecografico di una trombosi l'esame di secondo livello effettuata è stata la angio-TC, seguita, in caso di conferma da tre possibili opzioni terapeutiche: incremento della terapia anticoagulante, angiografia con fibrinolisi e intervento chirurgico.

Risultati: La sopravvivenza globale dei pazienti a 1, 3 e 5 anni è stata del 94,7%, 93,3% e 91% rispettivamente. La sopravvivenza a 1, 3 e 5 anni dell'organo è stata del 87,4%, 79,6% e 75,6%. La diagnosi eco-color doppler di trombosi del graft è stata fatta entro i primi 10 giorni dal trapianto in 28 casi (12,5%). Un tentativo di salvataggio dell'organo è stato effettuato in 18 pazienti. Tre pancreas sono stati espianati direttamente perché privi completamente di vascolarizzazione. Complessivamente sono stati salvati e sono tuttora funzionanti 14 pancreas (77,7%), mentre 4 (32,3%) sono stati successivamente espianati.

Conclusione: L'eco-color doppler è un'indagine idonea alla sorveglianza dei pancreas trapiantati permettendo la diagnosi precoce nel periodo più a rischio di trombosi vascolari e quindi il trattamento tempestivo delle complicanze più frequenti.

© 2007 Elsevier Masson Srl. All rights reserved.

Introduction

Pancreas transplantation is widely recognized as the most effective therapy for type 1 diabetes. It is the only treatment with the potential to render patients insulin-independent [1] and to reverse or stabilize complications associated with the disease [2,3]. In recent years, there has been a steady increase in the number of pancreas transplantations performed. Significant advances involving surgical technique and postoperative management have increased graft survival rates and the overall number of successful transplantations [4]. A new technique for pancreas transplantation has recently been developed in our institution, which involves retroperitoneal placement of the graft with portal-enteric drainage [5].

Despite these advances, clinical experience indicates that some postoperative complications are still common. Vascular thrombosis is one of these: in particularly severe cases, it can necessitate transplant pancreatectomy. Close postoperative monitoring is essential to detect vascular dysfunctions at an early stage. Sonography plays a key role in this process. It is the first-line method for postoperative evaluation of the pancreatic graft and in the detection of postoperative complications [6]. Color Doppler flow imaging has greatly enhanced the ability to evaluate the intra- and extra-parenchymal vasculature of pancreatic grafts. In fact, with this approach the Y-graft vascular anastomosis can be easily visualized, and in many cases the splenic artery and vein can be identified.

Materials and methods

Between May 1996 and March 2006, a total of 223 pancreas transplantations were performed in our institution. These included 115 simultaneous pancreas and kidney transplants (SPK); 27 simultaneous cadaver pancreas–living kidney transplants (SPLK); 66 involving pancreas transplant alone (PTA); and 15 pancreas after kidney transplants (PAK). The pancreas transplantations were performed using different techniques, including systemic-bladder drainage, systemic-enteric drainage, and retroperitoneal portal-enteric drainage. All recipients received antithrombotic prophylaxis, which was tailored to the individual's estimated risk of thrombosis. SPK recipients were generally managed with subcutaneous heparin calcium

for one week followed by oral aspirin therapy, which was started on post-transplant day three and continued for six months. SPT, SPLK, PAK and PTA recipients received heparin infusions for five days, followed by oral warfarin sodium for six months.

All patients were monitored on a daily basis with color Doppler ultrasonography during the first post-transplant week, and thereafter whenever clinically indicated. Whenever the results were

Fig. 1 (a) B-mode ultrasonographic scan reveals the presence of echogenic material within the splenic vein; and (b) color Doppler scan confirmed the presence of splenic thrombosis.

suggestive of venous thrombosis, a computed tomography (CT) was performed. If the presence of thrombosis was confirmed, 3 therapeutic strategies were used: (1) increased anticoagulant therapy; (2) angiography with fibrinolytic therapy; and (3) repeat surgery.

Results

The overall patient survival rates one, three, and five years after transplantation were 94.7%, 93.3%, and 91%, respectively. Corresponding rates for graft survival were 87.4%, 79.6%, and 75.6%, respectively. Twenty-eight graft thromboses (12.5%) were diagnosed with color Doppler ultrasound within the first 10 days after transplantation; and in 21 of these cases the presence of thrombosis was confirmed by CT-angiography (75%). In 14/21 cases, there was partial venous (Fig. 1) or arterial thrombosis (Fig. 2); and in the remaining 7 cases, the thrombosis produced full vessel occlusion.

Fig. 2 (a) A case of arterial thrombosis of a pancreatic graft. (b) Computed tomographic angiography confirmed the color Doppler diagnosis of arterial thrombosis.

An attempt to rescue the graft was made in 18 patients (14 with partial thrombosis and 4 with complete occlusive thrombosis). In 3 cases, graft pancreatectomy was performed because of complete loss of blood flow within the parenchyma. Three patients were treated with infusions of sodium heparin; 13 patients were managed with angiography, fibrinolysis, and heparin infusions; and 2 patients were treated with fibrinolysis, heparin infusion, and surgery (Figs. 3 and 4). Fourteen of the 18 rescue attempts (77.7%) were successful, and the grafts are still functioning. In the other 4 cases (32.3%), rescue failed, and the grafts were subsequently explanted.

Discussion

Pancreas allografts are particularly prone to surgical complications [7–9]. The pancreas graft is uniquely susceptible to vascular thrombosis due to its intrinsically low blood flow, which can be further diminished by ischemia–reperfusion injury [10].

Overall, more than 60% of the technical failures associated with pancreas transplantation are due to vascular thrombosis [11]. Although several risk factors for this type of complication have been identified, no effective prophylactic regimen or surgical strategy has been developed to eliminate post-transplant thrombosis. Regardless of their severity, the surgical complications associated with pancreas transplantation may be silent in the early stages [12]. We therefore decided to implement a standard protocol of post-transplant surveillance based on daily Doppler ultrasonographic assessments of the graft for the first week, which is the period in which most surgical complications, especially graft thrombosis, occur [13,14]. Close

Fig. 3 Angiography confirmed the complete thrombosis of the arterial axis.

Fig. 4 Fibrinolysis of a venous thrombus.

monitoring of the graft was also deemed necessary to improve rescue rates in cases of thrombosis. Indeed the rate of successful allograft rescue proved to be significantly higher when the treatment was performed in the presence of minimal parenchymal damage. Moreover, Doppler ultrasonography plays a dual role: it can identify the need for immediate treatment or provide information regarding the necessity for further investigation [15,16]. Color Doppler ultrasonography may be considered the most suitable tool for postoperative surveillance of patients undergoing pancreas transplantation. It is easy to perform and noninvasive, features that make it particularly useful for the early diagnosis of vascular thrombosis during the period of highest risk after transplantation. It therefore represents a key diagnostic tool as well as the first step towards timely therapeutic intervention, which in most cases leads to successful graft rescue.

References

- [1] Stratta RJ, Taylor RJ, Bynon JS, et al. Surgical treatment of diabetes mellitus with pancreas transplantation. *Ann Surg* 1994;220:809.

- [2] Brayman KL, Weber M, Sutherland DER. Pancreatic and islet cell transplantation. In: Trede M, Carter DC, editors. *Surgery of the pancreas*. New York: Churchill Livingstone; 1997. p. 637.
- [3] Giannarelli R, Coppelli A, Sartini MS, et al. Early improvement of unstable diabetes retinopathy after solitary pancreas transplantation. *Diabetes Care* 2002;25:2358–9.
- [4] Gruessner AC, Sutherland DER. Report for the international pancreas transplant registry-2000. *Transplant Proc* 2001;33:1643–6.
- [5] Boggi U, Vistoli F, Signori S, et al. A technique for retroperitoneal pancreas transplantation with portal-enteric drainage. *Transplantation* 2005;79:1137–42.
- [6] Nikolaidis P, Amin RS, Hwang CM, et al. Role of sonography in pancreatic transplantation. *Radiographics* 2003;23:939–49.
- [7] Humar A, Kandaswamy R, Granger D, Gruessner RW, Gruessner AC, Sutherland DE. Decreased surgical risks of pancreas transplantation in the modern era. *Ann Surg* 2000;231:269–75.
- [8] Reddy KS, Stratta RJ, Shokouh-Amiri MH, Alloway R, Egidi MF, Gaber AO. Surgical complications after pancreas transplantation with portal-enteric drainage. *J Am Coll Surg* 1999;189:305–13.
- [9] Troppmann C, Gruessner AC, Dunn DL, Sutherland DE, Gruessner RW. Surgical complications requiring early relaparotomy after pancreas transplantation. A multivariate risk factor and economic analysis of the cyclosporine era. *Ann Surg* 1998;227:255–68.
- [10] Ciancio G, Cespedes M, Olson L, Miller J, Burke GW. Partial venous thrombosis of the pancreas allografts after simultaneous pancreas–kidney transplantation. *Clin Transplant* 2000;14:464–71.
- [11] International Pancreas Transplant Registry Newsletter. August 31, 2004; Volume 16, Number 1.
- [12] Ciancio G, Julian JF, Fernandez L, Miller J, Burke GW. Successful surgical salvage of pancreas allograft after complete venous thrombosis. *Transplantation* 2000;70:126–31.
- [13] Boudreaux JP, Corry RJ, Dickerman R, Sutherland DE. Combined experience with immediate pancreas transplantation. *Transplant Proc* 1991;23:1628–9.
- [14] Kuo PC, Wong J, Schweitzer EJ, Johnson LB, Lim JW, Bartlett ST. Outcome after splenic vein thrombosis in the pancreas allograft. *Transplantation* 1997;64:933–5.
- [15] Gilibert R, Fernandez-Cruz L, Real MI, Ricart MJ, Astudillo E, Montana X. Treatment and outcome of pancreatic venous graft thrombosis after kidney–pancreas transplantation. *Br J Surg* 2002;89:355–60.
- [16] Eschwege P, Rocher L, Rondeau Y, et al. Complications of pancreas transplantation: radiological evaluation. *Transplant Proc* 1998;30:2795.