Published in final edited form as:

J Acquir Immune Defic Syndr. 2010 December ; 55(Suppl 2): S141–S143. doi:10.1097/QAI.0b013e3181fbc76.

Lessons Learned from Africa

Miriam Rabkin, MD, MPH^{*,†,‡}, Kevin M. De Cock, MD, FRCP (UK), DTM&H[§], and Wafaa M. El-Sadr, MD, MPH^{*,†,‡}

^{*}International Center for AIDS Care and Treatment Programs, Columbia University Mailman School of Public Health

[†]Department of Medicine, Columbia University College of Physicians and Surgeons

[‡]Department of Epidemiology, Columbia University Mailman School of Public Health

[§]U.S. Centers for Disease Control and Prevention

Abstract

Delayed access to HIV care and treatment in sub-Saharan Africa meant that the early years of HIV scale-up were characterized by a largely North-to-South transfer of knowledge and resources. Clinicians from wealthy countries were amongst the first to gain experience with antiretroviral treatment and care of people living with HIV, and shared key lessons with their colleagues in sub-Saharan Africa when widespread access to HIV services became available. Ten years later, South-to-North lessons learned from the remarkable achievements of HIV programs in Africa now have the potential to inform the response to the US domestic epidemic.

Keywords

HIV scale-up; program implementation; lessons; Africa

INTRODUCTION

The global response to the HIV epidemic has often been perceived as a North-to-South transfer of expertise, as well as resources. Because HIV care and treatment, including antiretroviral therapy (ART), were first widely available in wealthy countries, practitioners from the United States, western Europe and similarly well-resourced settings were among the first to gain experience treating people living with HIV (PLWH), the first to develop clinical guidelines, and the first to design and implement HIV service delivery programs. The delayed initiation of HIV care and treatment programs in sub-Saharan Africa created a “lost decade,” in which millions of PLWH were deprived of life-saving interventions and only a few African clinicians gained experience with HIV care and treatment. When resources finally became available, key insights from years of experience in the North informed the scale-up in the South. Examples include the importance of combination ART and the perils of sequential monotherapy, the critical role of retention in care and adherence to treatment, the need for multidisciplinary teams of providers, and the key role for PLWH

Correspondence and reprint requests to: Wafaa El-Sadr, International Center for AIDS Care and Treatment Programs (ICAP), 722 West 168th street, room 715, New York, New York 10032 (mr84@columbia.edu), 212/342-0532, 212/342-1824 (fax)..

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

in shaping the response to the HIV epidemic. These principles are now accepted as fundamental to effective HIV programs.

Times have changed, however, and the flow of information and experience is no longer a one way street. Clinicians and public health practitioners in many lower-income countries have rapidly acquired proficiency in the prevention, care, and treatment of HIV, and now lead the development and implementation of their national guidelines and trainings. The HIV epidemic has also redefined the concept of sustainability in development and highlighted the fact that global health requires global financing. Fiscal sustainability is unlikely to be achieved in the near future, but technical sustainability is much more promising. While funds and technical assistance from the global North are still welcome and necessary in many settings, local solutions and expertise often provide the most effective interventions. This new equilibrium highlights the fact that innovations and best practices are generated worldwide – and that South-South and South-North transfers and technical assistance have important roles to play. Indeed, despite marked differences in the magnitude and characteristics of the HIV epidemics in the United States and in sub-Saharan Africa, the response to the US HIV epidemic may be enhanced by some of the lessons learned from the enormous success of HIV program scale-up in Africa.

CONTRASTING REALITIES

Sub-Saharan Africa has a generalized HIV epidemic with high seroprevalence rates in many countries, although recent data indicate that specific subpopulations are more severely impacted. For example, young women in southern Africa are at particular risk for HIV infection when contrasted with young men of the same age, and there are high rates of HIV infection among drug users and men who have sex with men in some countries.¹ In contrast, overall HIV prevalence in the US is much lower, and the US epidemic is a localized one, with foci of high prevalence rivaling that in some parts of sub-Saharan Africa, particularly among men who have sex with men and African Americans.²

Another contrast is the state of health systems in the US compared to those in sub-Saharan Africa. In 2007, per capita total expenditure on health was \$7,285 in the US, compared with \$819 in South Africa, \$79 in Zambia, and \$39 in Mozambique.³ Out of pocket expenditures represent 12% of total health expenditures in the US, compared to 33% in the African region as a whole.⁴ There is a glaring difference in the availability of health workers, with approximately 27 physicians and 560 health workers per 10,000 inhabitants in the US compared to 2 and 23/10,000 respectively for the African region.^{3,4} The US has 31 hospital beds per 10,000 people; compared to 1.8 in Ethiopia, 4 in Côte d'Ivoire, and 13 in Lesotho.⁵ In 2007, healthy life expectancy at birth was 70 years in the US and 45 years in the African region.³ And while precise data are lacking, PLWH in the US have far greater access to HIV care and treatment than in Africa. However, the US response to the HIV epidemic has not been without its challenges including continued HIV transmission and disparities in access to HIV services.⁶

LESSONS FROM AFRICA

The response to HIV in Africa is as diverse and heterogeneous as the epidemic itself, with important within-country and between-country differences. Although generalizations about the 47 different countries in sub-Saharan Africa are not always helpful, several recurrent lessons have emerged. We note that these innovations are not limited to the African region, and that additional important lessons are available from around the world.

- *National plans:* Since 2004, UNAIDS has emphasized the importance of a single national AIDS Coordinating Authority and a single national HIV/AIDS Action

Framework as part of its “Three Ones” approach.⁷ Countries such as Kenya and Ethiopia have had national plans for years, with targets and budgets supporting concrete objectives and activities. In contrast, the US released its first-ever National HIV/AIDS Strategic Plan in 2010.⁶

- *Targets and accountability:* In marked contrast to the US, many African countries have developed specific numeric targets for HIV testing, care, and treatment based on the best available information about local prevalence and need. Enrollment and coverage targets have been developed for the site, district, province, and national levels and progress against targets is reported and shared. In Ethiopia, for example, the Federal Ministry of Health (FMOH) and the Federal HIV/AIDS Prevention and Control Office (FHAPCO) publish monthly data on enrollment in HIV care and ART for each region – and for every health care facility providing HIV services in the country. ART coverage is an indicator in WHO's *World Health Statistics* report, which publishes annual estimates of national ART coverage among people with advanced HIV infection, and PMTCT coverage among HIV-infected pregnant women in more than 100 countries; these data are difficult to ascertain for the US.
- *Algorithmic guidelines:* The development of simple algorithmic HIV treatment guidelines in response to physician shortages has enabled not only rapid scale-up of HIV services, but the standardization of first- and second-line treatment, streamlined drug procurement, and task-shifting from physicians to nurses and other health workers. In many settings, these guidelines have also averted the “pharmacologic chaos” seen in some resource-rich settings, in which personalized prescribing and disregard of guidelines can devolve into unnecessarily complex care, over-use of laboratory investigations, and even widespread misuse of ART.⁸
- *“Know your epidemic”:* Although national disease surveillance systems (especially those based on case reporting) are typically much more robust in resource-rich settings than in sub-Saharan Africa, the availability of population-based data regarding HIV incidence and prevalence in the US is surprisingly scant. With the exception of data from the National Health and Nutrition Examination Survey (NHANES), the US lacks the information found in Demographic Health Surveys (DHS), Multiple Indicator Cluster Surveys (MICS), antenatal sentinel surveillance, and initiatives such as the 2007 Kenya AIDS Indicator Survey, which provides a detailed and profoundly useful snapshot of the epidemic in that country.⁹
- *Task-shifting:* Driven by a critical shortage of physicians and nurses^{10,11} the HIV response in sub-Saharan Africa has been characterized by health workforce innovations, task-shifting, and task-sharing.^{12, 13} For example, in some countries, nurses and clinical officers can now prescribe and/or refill antiretroviral medications,^{14, 15} while in the US these activities are limited to physicians and nurse practitioners.
- *Diagnostics and case-finding:* While countries in sub-Saharan Africa have a long way to go in achieving broad access to HIV testing for their populations, these same countries were quick to adopt rapid HIV testing assays, utilizing algorithms that do not include Western blot assays. They also pioneered the provision of such testing by lay workers trained to perform rapid HIV tests in mobile, community, and even home-based settings. At health facilities, provider initiated point-of-service screening for HIV enables rapid testing at multiple venues, from antenatal care, to TB clinics, to outpatient departments.
- *Integration of services:* While access to integrated service delivery varies from site to site and country to country, some types of integrated services are often more developed and more accessible in African programs than in the US. The integration

of HIV and TB services – including TB screening for patients with HIV and HIV testing for patients with TB – is one example; the use of antenatal care programs for the provision of prevention of mother-to-child transmission (PMTCT) and the identification and enrollment of women and families with HIV is another.

- *Research:* Studies conducted in sub-Saharan Africa have contributed substantially to the world's knowledge base on HIV/AIDS, directly influencing practice elsewhere. After productive research in descriptive and analytic epidemiology focusing on natural history, risk factors and disease associations¹⁶, Africa is now a vibrant setting for clinical trials of relevance worldwide, in areas such as prevention and treatment of HIV-associated tuberculosis, HIV prevention, microbicides, and other interventions.^{17,18,19}

CONCLUSIONS

Despite the horrific impact of the HIV epidemic, it has also triggered an outpouring of solidarity within and between nations. Again and again, we see the impact of generous support from individuals, communities, and countries. While common wisdom suggests that financial and technical inputs from resource-rich settings can and should continue to inform the design and implementation of HIV programs in the global South when appropriate, the potential benefits of South-South and South-North contributions should also be acknowledged. Remarkable innovations have germinated in Africa in the process of scaling up HIV prevention, care, and treatment programs; many of these can inform the response to the HIV epidemic. As the US continues its quest to control the epidemic in its midst, we should pause and consider what we can learn from Africa.

Acknowledgments

The authors wish to acknowledge their valued colleagues in the US and in Africa who have contributed so much to the response to the HIV epidemic, as well as the patients and families who have trusted us with their care. We also acknowledge contributions by Shannon Hader and Chris Collins.

REFERENCES

1. The Joint United Nations Programme on HIV/AIDS (UNAIDS). World Health Organization (WHO). AIDS Epidemic Update: November 2009. UNAIDS; Geneva: 2009.
2. El-Sadr WM, Mayer KH, Hodder SL. AIDS in America – forgotten but not gone. *N Engl J Med*. 2010; 362:967–970. [PubMed: 20147707]
3. World Health Organization (WHO). World Health Statistics 2010. WHO; Geneva: 2010. Available at: http://www.who.int/whosis/whostat/EN_WHS10_Full.pdf
4. World Health Organization (WHO). The World Health Report 2006: Working Together for Health. WHO; Geneva: 2006. Available at <http://www.who.int/whr/2006/en/index.html>
5. WHO Global Health Observatory. Geneva: 2010. [database online] <http://apps.who.int/ghodata/>
6. The White House Office of National AIDS Policy. National HIV/AIDS Strategy for the United States. Washington, DC: Jul. 2010 Available at: <http://www.whitehouse.gov/sites/default/files/uploads/NHAS.pdf>
7. The Joint United Nations Programme on HIV/AIDS (UNAIDS). The “Three Ones”: Principles for the coordination of national AIDS responses. Available at: <http://www.unaids.org/en/CountryResponses/MakingTheMoneyWork/ThreeOnes/default.asp>
8. Cocohoba J, Wang QJ, Cox C, et al. Consistency of initial antiretroviral therapy with HIV treatment guidelines in a US cohort of HIV-infected women. *JAIDS*. 2008; 47:377–383. [PubMed: 18176324]
9. National AIDS and STI Control Programme (NASCOP). Kenya AIDS Indicator Survey 2007: Final Report. Nairobi: Sep. 2009 Available at: http://www.aidskenya.org/public_site/webroot/cache/article/file/Official_KAIS_Report_20091.pdf

10. Chen L, Evan T, Anand S, et al. Human resources for health: overcoming the crisis. *Lancet*. 2004; 364:1984–1990. [PubMed: 15567015]
11. Joint Learning Initiative. Human resources for health: overcoming the crisis. Harvard University Press; Cambridge (MA): 2004.
12. Sherr K, Pfeiffer J, Mussa A, et al. The role of nonphysician clinicians in the rapid expansion of HIV care in Mozambique. *JAIDS*. 2009; 52:S20–S23. [PubMed: 19858931]
13. Samb B, Celletti F, Holloway J, et al. Rapid expansion of the health workforce in response to the HIV epidemic [Sounding Board]. *N Engl J Med*. 2007; 357:2510–2514. [PubMed: 18077816]
14. World Health Organization (WHO). Task Shifting: Rational Redistribution of Tasks Among Health Workforce Teams. Global Recommendations and Guidelines. WHO; Geneva, Switzerland: 2008. Available at: <http://www.who.int/healthsystems/TTR-TaskShifting.pdf>
15. Dohrn J, Nzuama B, Murrman M. The impact of HIV scale-up on the role of nurses in South Africa: Time for a new approach. *JAIDS*. 2009; 52:S27–S29. [PubMed: 19858933]
16. De Cock KM, Ekpini E, Gnaore E, et al. The public health implications of AIDS research in Africa. *JAMA*. 1994; 272:481–486. [PubMed: 8040984]
17. Abdool Karim SS, Naidoo K, Grobler A, et al. Timing of initiation of antiretroviral drugs during tuberculosis therapy. *N Engl J Med*. Feb 25; 2010 362(8):697–706. [PubMed: 20181971]
18. Lingappa JR, Baeten JM, Wald A, et al. Daily acyclovir for HIV-1 disease progression in people dually infected with HIV-1 and herpes simplex virus type 2: a randomised placebo-controlled trial. *Lancet*. Mar 6; 2010 375(9717):824–33. [PubMed: 20153888]
19. Abdool Karim Q, Abdool Karim SS, Frohlich J, et al. Effectiveness and safety of tenofovir gel, an antiretroviral microbicide, for the prevention of HIV infection in women. *Science*. Jul 19.2010 Epub ahead of print.