CORRESPONDENCE

Recognizing and Treating Peripartum Depression

by Dr. rer. medic. Bettina Hübner-Liebermann, Dr. jur. Helmut Hausner,
Dr. med. Markus Wittmann in volume 24/2012

Awareness Should Be Raised Among Doctors

In times of compulsive control, demand, and perfectionism in the realm of limitless possibilities, almost every pregnancy turns into a crisis with long latency periods until exhaustion and ambivalence are integrated or resolved. This pathway includes symptoms such as hyperemesis, premature labor, hypertension, dystocia of labor, breastfeeding and bonding problems, and sometimes depression. The latter does not have to be used as the umbrella term even if ambivalence always entails an element of depression.

Independently of these factors I share the following considerations regarding possible strategies: awareness needs to be improved in gynecologists and midwives in order for them to address ambivalences, identify these, and initiate the correct treatment. Networks to support parents with psychosocial burdens (for example, www.mutterkindgesundheits.de) are also suitable instruments. The high prevalence is acknowledged by the fact that gynecologists have to acquire a qualification in basic psychosomatic care. This should actually be included in the further training curriculum for every medical specialty. The German Society of Psychosomatic Obstetrics and Gynecology (DGPF, Deutsche Gesellschaft für Psychosomatische Geburtshilfe und Gynäkologie) does not only provide curricula for psychosomatic basic care on a nationwide basis, but it also arranges specialty-specific psychotherapy and psychosomatic further professional education for midwives and psychologists.

DOI: 10.3238/arztebl.2013.0011a

REFERENCES

1. Hübner-Liebermann B, Hausner H, Wittmann M: Recognizing and treating peripartum depression. *Dtsch Arztebl Int* 2012; 109(24): 419–24.

Dr. med. Wolf Lütje

Frauenklinik ev. Amalie Sieveking Krankenhaus Hamburg
Vizepräsident DGPF (Deutsche Gesellschaft für Psychosomatische
Geburtshilfe und Geburtshilfe)
wluetje@googlemail.com

Conflict of interest statement

The author declares that no conflict of interest exists.

Somatic Causes Were Omitted

In their description of the causes the authors go into great detail about psychopathological and psychosocial causes but completely omit any mention of somatic

causes of peripartum depression. In addition to rare endocrine disorders, such as primary hyperparathyroidism or endogenous hypercortisolism, postpartum thyroiditis is a clinically relevant disorder that may be accompanied by symptoms of depression. In 5–7% of all births worldwide, women develop autoimmune postpartum thyroiditis, with higher incidence rates in women with pre-existing and confirmed raised thyroid antibody concentrations. The disorder usually starts in the third or fourth month post partum. 25–30% of women develop functional disorders of the thyroid gland, mostly hypothyroidism, and more rarely, initial hyperthyroidism that resolves spontaneously. Not only patients with functional disorders have clinical symptoms; patients with euthyroidism also feel worse than healthy controls. Endocrine causes of postpartum depression should always be considered since excellent therapeutic options are available.

DOI: 10.3238/arztebl.2013.0011b

REFERENCES

1. Lazarus JH: Clinical manifestations of postpartum thyroid disease. *Thyroid* 1999; 9: 685–9.
2. Stagnaro-Green A: Approach to the patient with postpartum thyroiditis. *J Clin Endocrinol Metab* 2012; 97: 334–42.
3. Pilhatsch M, Marxen M, Winter C, Smolka MN, Bauer M: Hypothyroidism and mood disorders: integrating novel insights from brain imaging techniques *Thyroid Res.* 2011; 4(Suppl 1): 3.
4. Hübner-Liebermann B, Hausner H, Wittmann M: Recognizing and treating peripartum depression. *Dtsch Arztebl Int* 2012; 109(24): 419–24.

Priv.-Doz. Dr. med. Joachim Feldkamp

Klinik für Allgemeine Innere Medizin, Endokrinologie und Diabetologie,
Pneumologie und Infektiologie, Klinikum Bielefeld
Joachim.Feldkamp@klinikumbielefeld.de

Prof. Dr. med. Matthias Schott

Klinik für Endokrinologie und Diabetologie, Rheumatologie
Heinrich-Heine-Universität Düsseldorf

Conflict of interest statement

The authors declare that no conflict of interest exists.

Evidence Needs to Be Presented

It is important in this context to present evidence from controlled studies, which is by its very nature limited. With this background, it is worth mentioning a randomized controlled study by Wisner et al (1), in which 22 pregnant women without current depression, who had a medical history of at least one postpartum major depression, were treated with sertraline (initially 25–50 mg, target dose 75 mg) for 20 weeks immediately after giving birth. 1 in 14 patients taking sertraline had a recurrence compared with 4 out of 8 patients taking placebo; the difference reached significance.

In spite of methodological problems (small number of cases, no statistical adjustment for the number of

depressive episodes) pharmacological prophylaxis should be considered in pregnant women whose medical history includes depressive episodes. The risk of recurrence in a new pregnancy after previous postpartum depression is high, at 25% or more (2). According to the Embryotox database, sertraline passes into the neonatal plasma only in traces in breastfed infants (3). In terms of balancing risks and benefits, prophylactic treatment with sertraline is therefore a justifiable option.

DOI: 10.3238/arztebl.2013.0011c

REFERENCES

1. Wisner KL, Perel JM, Peindl KS, Hanusa BH, Piontek CM, Findling RL: Prevention of postpartum depression: a pilot randomized clinical trial. *Am J Psychiatry* 2004; 161: 1290–2.
2. Wisner KL, Perel JM, Peindl KS, Hanusa BH, Findling RL, Rappaport DJ: Prevention of recurrent postpartum depression: a randomized clinical trial. *J Clin Psychiatry* 2001; 62: 82–6.
3. www.embryotox.de/sertralin.html; last accessed on 20.06.2012
4. Hübner-Liebermann B, Hausner H, Wittmann M: Recognizing and treating peripartum depression. *Dtsch Arztebl Int* 2012; 109(24): 419–24.

Dr. med. David Zilles

Universitätsmedizin Göttingen, Georg-August-Universität
Abteilung Psychiatrie und Psychotherapie
david.zilles@med.uni-goettingen.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Oxytocin Concentrations Are Crucial

The causes of postnatal depression are a mesh of physical, hormonal, biochemical, psychological, social, and societal factors. This means that successful treatment will have to take into consideration all these factors and include them in the treatment concept.

Additionally, oxytocin is of great importance during the entire period of pregnancy, birth, and puerperium. A low oxytocin concentration in the last four weeks of the pregnancy seems to be associated with an increased risk of developing symptoms of postpartum depression within the first two weeks after the birth. Good bonding between mother and baby after the birth, such as is standard in certified “baby-friendly hospitals” (www.babyfreundlich.org) increases the concentration of oxytocin. Concentrations of oxytocin as well as prolactin are higher in breastfeeding women; breast feeding therefore reduces the psychological stress reaction (1, 3).

Very few publications consider the difference between non-breastfeeding women and breastfeeding women. Extensive discussions about the influence of hormone imbalances ignore lactation and nature’s potential protective mechanisms owing to high concentrations of prolactin and oxytocin during the breastfeeding period (1, 3).

Breast feeding problems may be an indication of postpartum depression. Hypothyroidism is also associated with severe “baby blues” or fatigue, in addition to

reduced lactation, and should therefore also be excluded (1).

With regard to pharmacotherapy, it is worth drawing attention to the 8th edition of *Arzneimittel in Schwangerschaft und Stillzeit* (2012) by Schaefer et al. It is also worth pointing out that medication-led weaning using prolactin inhibitors can increase depression and should therefore be avoided at all cost. If weaning is necessary physiological weaning without drugs is the preferred option (2).

DOI: 10.3238/arztebl.2013.0012a

REFERENCES

1. Lawrence R: Breastfeeding. A Guide for the Medical Profession. 7th edition Maryland heights, Missouri: Elsevier, Mosby, Inc., 2011.
2. Schaefer C, Spielmann H, Vetter K, Weber-Schöndorfer C: Arzneimittel in Schwangerschaft und Stillzeit. 8th edition. München: Urban und Fischer 2012.
3. Uvnäs-Moberg K: The Oxytocin factor: Tapping the hormone of calm, love, and healing. Cambridge: Da Capo Press 2003.
4. Hübner-Liebermann B, Hausner H, Wittmann M: Recognizing and treating peripartum depression. *Dtsch Arztebl Int* 2012; 109(24): 419–24.

Gudrun von der Ohe

Ärztin, Hamburg
Postfach@stillberatung.info

Conflict of interest statement

The author declares that no conflict of interest exists.

Metabolic Derailment Should Be Considered

A maternal deficiency in ornithine transcarbamylase (OTC) is a rare but sometimes very impressive cause of peripartum depression in mothers. This is an X-linked inherited metabolic disorder of the urea cycle. Male patients are usually severely affected, while symptoms in hemizygous females are often mild. Female patients complain of headaches and difficulties in concentration; stress—such as pregnancy or giving birth—may trigger metabolic derailment, which is facilitated by a catabolic state. This may manifest as depression or psychosis.

This metabolic disorder can be diagnosed by measuring blood concentrations of ammonia, which are notably raised. Orotic acid concentrations in blood/urine are also typically elevated.

Specific treatment is available for this particular type of postpartum depression. The treatment consists of ‘high energy’ nutrition (via glucose/fat) in order to trigger an anabolic state, substitution of citrulline, and administration of “scavengers” that bind nitrogen and eliminate this via urine independently of the urea cycle. In most cases, peripartum depression can be controlled using this approach.

DOI: 10.3238/arztebl.2013.0012b

REFERENCES

1. Hübner-Liebermann B, Hausner H, Wittmann M: Recognizing and treating peripartum depression. *Dtsch Arztebl Int* 2012; 109(24): 419–24.

Prof. Dr. med. A.M. Das

Klinik für Pädiatrische Nieren-, Leber- und Stoffwechselerkrankungen
Zentrum Kinderheilkunde und Jugendmedizin
Medizinische Hochschule Hannover
das.anibh@mh-hannover.de

Conflict of interest statement

Professor Das has received honoraria for acting as an adviser and expert referee, honoraria for conducting commissioned clinical studies, and funding for a research project he initiated, all from Cytonet. He has received moneys from Swedish Orphan Biovitrum for preparing scientific advanced training events.

Successful Bonding Is Important

The comprehensive article on risk factors for postpartum depression has omitted important factors that can trigger or increase depression on the one hand, and counteract it on the other hand.

Risk factors for postpartum depression include weaning and separation of mother and baby. Neither was mentioned in the article. There is sufficient evidence that successful bonding and a good start to breast feeding counteract peripartum depression (2). Baby-friendly hospitals support bonding between parents and babies and promote breast feeding. Uninterrupted initial encounters, skin contact immediately after the birth, and gradual build-up of familiarity are vital for establishing a bond. Women who do not breast feed are shown in baby-friendly hospitals how to give the baby a bottle while remaining physically close. Successful bonding increases oxytocin concentrations and therefore counteracts postpartum depression. Fewer breastfeeding mothers seem to be affected by postpartum depression than non-breastfeeding mothers (1).

Baby-friendly hospitals therefore make an important contribution to the prevention of postpartum depression.

DOI: 10.3238/arztebl.2013.0013a

REFERENCES

1. Ip S, Chung M, Raman G, et al.: Breastfeeding and maternal and infant health outcomes in developed countries. Evid Rep Technol Assess (Full Rep) 2007; 1–186.
2. Ip S et al.: Breastfeeding and maternal and infant health outcomes in developed countries. Evid Rep Technol Assess 2007; 153: 1–186.
3. Hübner-Liebermann B, Hausner H, Wittmann M: Recognizing and treating peripartum depression. Dtsch Arztebl Int 2012; 109(24): 419–24.

Gisela Meese M.A., Geschäftsführerin

WHO/UNICEF-Initiative „Babyfreundlich“
Bergisch Gladbach
info@babyfreundlich.org

Conflict of interest statement

The author sits on the executive board of the WHO/UNICEF Baby Friendly Initiative.

In Reply:

We thank our correspondents for their critical comments and ideas. The comments mostly relate to risk factors and differential diagnoses that we did not mention in the article; rare causes were mentioned that should certainly be included in the differential diagnostic evaluations, but common somatic causes, such as postpartum thyroiditis, were also included. For reasons of space we were not able to include all aspects of postpartum depression in our article, and the comments therefore provided particularly valuable and important information.

Psychological problems in pregnancy and during the breastfeeding period should obviously not be unnecessarily and wholesale psychopathologized; we agree with Lütje in this respect. The prevalence rates in our article for severe prenatal and postnatal depression requiring treatment, the suicide rate associated with this disorder, and the low uptake of professional help are, however, enough of a reason to warn against trivializing these symptoms.

DOI: 10.3238/arztebl.2013.0013b

REFERENCES

1. Hübner-Liebermann B, Hausner H, Wittmann M: Recognizing and treating peripartum depression. Dtsch Arztebl Int 2012; 109(24): 419–24.

Dr. med. Markus Wittmann

Psychiatrische Institutsambulanz
Bezirksklinikum Mainkofen/Deggendorf
m.wittmann@mainkofen.de

Conflict of interest statement

Dr Wittmann has received consultancy fees from Bristol Myers Squibb and Servier. He has received honoraria for authorship in the context of a publication from Servier, which was relevant to the topic of the article. He has been reimbursed the costs for congress participation, travel, and accommodation, and fees for the preparation of scientific training courses from AstraZeneca, Servier, Wyeth, Lilly, Janssen-Cilag, GlaxoSmithKline, Pfizer, Eisai, and Lundbeck. He has accepted funds into a third-party account for performing commissioned clinical studies from Servier.