Published in final edited form as:

Ann Behav Med. 2013 February ; 45(Suppl 1): 162–172. doi:10.1007/s12160-012-9417-y.

Locations of Joint Physical Activity in Parent-Child Pairs Based on Accelerometer and GPS Monitoring

Genevieve Fridlund Dunton, Ph.D, MPH¹, Yue Liao, MPH¹, Estela Almanza, MPH², Micheal Jerrett, Ph.D², Donna Spruijt-Metz, Ph.D¹, and Mary Ann Pentz, Ph.D¹

¹Department of Preventive Medicine, University of Southern California

²Environmental Health Sciences, University of California, Berkeley

Abstract

Background—Parental factors may play an important role in influencing children's physical activity levels.

Purpose—This cross-sectional study sought to describe the locations of joint physical activity among parents and children.

Methods—Parent-child pairs (N = 291) wore an Actigraph GT2M accelerometer and GlobalSat BT-335 Global Positioning Systems (GPS) device over the same 7-day period. Children were ages 8–14 years. Joint behavior was defined by a linear separation distance of less than 50m between parent and child. Land use classifications were assigned to GPS data points.

Results—Joint physical activity was spread across residential locations (35%), and commercial venues (24%), and open spaces/parks (20%). Obese children and parents performed less joint physical activity in open spaces/parks than under/normal weight children and parents (p 's < .01).

Conclusions—Understanding where joint parent-child physical activity naturally occurs may inform location-based interventions to promote these behaviors.

Keywords

moderate-to-vigorous physical activity; sedentary behavior; parents; children; global positioning systems; environments

Introduction

Evidence suggests that parental factors may play an important role in influencing children's physical activity levels (1,2). Parents may help promote children's physical activity by providing support, purchasing equipment, paying fees, providing transportation, watching and supervising activities, offering encouragement, and discussing benefits (3,4). Evidence is less consistent about whether parents' physical activity levels influence the amount of physical activity that their children perform (5–8). Research in this area typically does not distinguish between parents' physical activity, performed *separately from* versus *together with* their children. Recent work suggests that only a small proportion of parents' overall

Correspondence: Genevieve Fridlund Dunton, Ph.D, MPH, Assistant Professor, Department of Preventive Medicine, University of Southern California, Soto Street Building (SSB), 200 Soto N. Street, 3rd floor, MC 9239, Los Angeles, CA 90033-9045, Phone: 323-442-8224, Fax: 323-442-8201, dunton@usc.edu.

Conflict of Interest Statement

The authors have no conflict of interest to disclose.

physical activity is done with their children (9,10). Increasing joint parent-child physical activity could provide opportunities for positive parental role modeling of active behaviors while health benefits for children and parents alike. Information about the locations where parents and children engage in joint physical activity can inform location-based programs and policies to promote joint physical activity in parents and children (11,12).

A growing body of research uses new methods such as Global Positioning Systems (GPS) and Ecological Momentary Assessment (EMA) to investigate the locations of children's physical activity. Studies have found that most of children's moderate-to-vigorous intensity physical activity occurs at school and at home (13–15). Research has shown that only small amounts (approximately 2 – 10%) of children's physical activity takes place in parks and green spaces (14–18). Fewer studies have examined where adults' physical activity takes place. Troped and colleagues (19) found that most of adults' moderate-to-vigorous physical activity took place more than 1km from home. Time use data suggest that more of adults' sports and exercise occur outdoors (25%) and at home (25%) than at a gym/health club (8%) or at work (3%) (20). However, no known studies to date have sought to identify the locations of physical activity that parents and children perform together.

This study examined the locations of joint physical activity and sedentary behavior in parent-child pairs who wore an accelerometer and GPS device over the same 7-day period. The first objective was to classify the locations of (a) joint parent-child moderate-to-vigorous physical activity (MVPA), (b) joint parent-child sedentary behavior, (c) child MVPA accompanied by parent sedentary behavior, and (d) parent MVPA accompanied by child sedentary behavior according to primary land use type (e.g., residential, commercial, open space). Evidence suggests that overweight and obese children and adults may have different patterns, types, and locations of physical activity than their normal weight counterparts (18, 21, 22). Thus, our second objective was to determine whether weight status of the parent and child are associated with the location of joint activity for each joint behavior type. Based on prior work (18), we hypothesized that overweight and obese children and parents would perform more joint physical activity in residential locations than open space and parks.

Methods

Sample

This study used data from a subgroup of children and parents enrolled in a 4-year project examining the effects of smart growth community design on family obesity risk. Participants included a convenience sample of fourth through eighth grade children (ages 8–14 years) and their parents. Families lived in Chino, CA or surrounding communities within 30 minutes driving time from Chino. The city of Chino is located within the greater Los Angeles Basin, about 35 miles East of downtown Los Angeles. According to the 2010 U.S. Census, Chino has a population of 77,983 persons, with a population density of 2,630 people per square mile. About 50% of residents are of Latino or Hispanic origin. The median annual household income is \$71,000 (23). Participants were recruited through informational flyer, posters, and letters. All recruitment materials included the study recruitment hotline phone number and email address. A telephone recruiter called all interested families and screened for eligibility. Inclusion criteria consisted of the following: (a) child currently enrolled in the 4–8th grade, (b) living in Chino, CA or a surrounding community, and (c) annual household income less than \$210,000. Extremely high income households were excluded because the goal of the study was to focus on children from families who have elevated obesity risk. Parents and children who met the eligibility criteria were scheduled for a data collection appointment at a local community site or their home. Written informed consent and minor assent was obtained from participants. This research was reviewed and

approved by the Institutional Review Board at the University of Southern California and the Committee on the Protection of Human Subjects, University of California, Berkeley.

Study Design

Baseline data from the 4-year study were used. Data were collected from March 2009–December 2010. No data collection took place from late July–August and during January due to typically adverse temperatures and weather conditions that limit outside activity in that part of Southern California. Within the parent-child pairs, both wore an accelerometer and GPS device over the same 7-day period.

Measures

Physical activity—Physical activity was assessed using an Actigraph, Inc. GT2M model activity monitor (firmware v06.02.00), although only data from the vertical axis were utilized. Participants wore the device on the right hip attached to an adjustable belt for seven continuous days. The devices were not worn when sleeping, bathing, or swimming. Activity counts were recorded in 30-second epochs in both parents and children to facilitate matching among the pairs. Cut-points for moderate-to-vigorous physical activity (MVPA) were applied from studies of national surveillance data (24,25). For adults, the MVPA threshold was 2020 counts per minute (equivalent to 3 METs). MVPA for children was defined using age-specific thresholds generated from the Freedson prediction equation (26,27). A threshold for moderate activity of 4 METs was used for children (as opposed to a 3 METs moderate activity cut-off for adults) to account for higher resting energy expenditure in children and youth (28,29). For children and adults, sedentary activity was defined as less than 100 counts per minute (30,31).

Location Monitoring—Geographic locations were recorded through portable GPS devices worn by both children and parents. Data were gathered for a 7-day period with the BT-335 Bluetooth GPS data logger device by GlobalSat Technology Corp (Taipei) attached to a belt worn around the waist along with the accelerometer. The BT-335 (16M bit, 1575.42 MHz) consists of a GPS receiver and data logger with Bluetooth PC interface. This device records time, date, speed, altitude, and GPS location at preset intervals. It is Wide Area Augmentation System (WAAS) enabled and uses a SiRF star III chipset for accurate position tracking (up to 3 m accuracy outdoors) and improved indoor signal acquisition. The recording interval was set to a 30-second epoch to match the accelerometer specifications. A battery charger was provided, and participants were instructed to recharge the battery each night. After the GPS devices were returned, all recorded information was downloaded to a computer where the recorded longitudinal and latitudinal data and speed were downloaded to a CSV file format. Linear distance between the parent and child for each 30-sec. epoch was calculated using geographic coordinates from the GPS. The GPS devices and accelerometers were marked with separate color-coded stickers for the parent and child to avoid any mistaken cross usage.

Land use classifications—GPS data points for joint parent-child behavior (at the level of the 30-sec. epoch) were given land use classification in a geographic information system (GIS) using a Southern California Association of Governments (SCAG) database (32). Data points were assigned the land use classification of the area nearest to the point that had land use data. Most of the GPS data points (93.7%) fell in an area that had been classified in the SCAG database. However, some GPS data points had small distances to their nearest land use classified area (6% had distances greater than 0 but less than 15 m). GPS data points that were greater than 50 meters away from the nearest area that had a classification were excluded from the analyses (0.08%). For analyses, land uses were grouped into six major categories: residential (e.g., houses, apartments, condos), commercial (e.g., retail,

restaurants, office use, manufacturing), open space (e.g., vacant lots, parks, golf courses, gardens, beaches), educational (e.g., schools and school grounds), public facilities (e.g., community centers, churches, libraries), and other (e.g., military, mixed uses, airports, freeways, roads, utilities).

Height and weight—Parents' and children's height and weight were measured in duplicate using an electronically calibrated digital scale (Tanita WB-110A) and professional stadiometer (PE-AIM-101) to the nearest 0.1 kg and 0.1 cm, respectively. Body Mass index (BMI) was calculated (kg/m^2). Parents' weight status was classified using the following BMI categories: under or normal weight ($\text{BMI} < 25$), overweight ($\text{BMI} = 25\text{--}29.9$), and obese ($\text{BMI} \geq 30$). Children's weight status was classified according to CDC age- and sex-specific BMI percentile cut-offs (under or normal weight $< 85^{\text{th}}$ percentile, overweight = 85^{th} to less than the 95^{th} percentile, obese $\geq 95^{\text{th}}$ percentile).

Demographic variables—Age, sex, and ethnicity were assessed through child and parent self-report surveys. Ethnicity was coded as Hispanic versus non-Hispanic. Parents reported annual household income, which was divided into quartiles (Less than or equal to \$30,000; \$30,001–\$60,000; \$60,001–\$100,000 and greater than \$100,000).

Data Merging and Processing

Accelerometer and GPS files were imported into the R version 2.9.2 programming language interface. Date and time stamps to the nearest 30-second epoch were used to match all accelerometer and GPS records within each parent-child pair. In the numerous cases where concurrent accelerometer and GPS were unavailable for either the parent or the child, we used a missing data code (NA) for designating the accelerometer or GPS values for these epochs. Overnight (11pm–5am) and school (8am–3pm on weekdays during the school season) hours were removed from the analyses. Strings of consecutive readings of 0 activity counts lasting 60 minutes or more were considered accelerometer non-wear and excluded from analyses. Activity outliers were identified as records with greater than 16,383 counts per 30-second epoch (33). Motorized activity was also excluded from the analyses, which were identified by records with speeds greater than 32 kph since typical bicycling speeds range from 15 to 30 kph (9.32 to 18.64 mph). Once these records were removed, parent-child pairs were determined to have sufficient data for inclusion in the analysis if they had a minimum of 2 valid days (any combination of weekdays or weekend days) of matched data—where a valid weekday was defined as having a minimum of two hours of matched accelerometer and GPS data points for the pair (between 3pm–11pm), and a valid weekend day was defined as having a minimum of four hours of matched accelerometer and GPS data points for the pair (between 5am–11pm). “Joint” behaviors were defined as behaviors that occurred at the same time and in the same location ($< 50\text{m}$. apart). A maximum separation of less than 50m between the parent and child was selected because this distance is approximately equivalent to the length of a ball court (e.g., basketball, volleyball, racquetball) or large residential yard. Whether the joint behavior occurred inside of the pair's residential neighborhood was defined as the GPS coordinates falling within a 500m buffer around the home address. Four types of joint parent-child behavior were examined: (a) joint MVPA, (b) joint sedentary behavior, (c) child MVPA accompanied by parent sedentary behavior, and (d) parent MVPA accompanied by child sedentary behavior.

Data Analyses

Two multilevel multinomial logistic regression models were fit for each of the four types of joint parent-child behavior to predict the likelihood of occurring in each land use type as compared with residential locations (reference group). The first model tested the effect of child weight status (i.e., BMI category). The second model tested the effect of parent weight

status (i.e., BMI category). Each of the four types of joint behavior (joint MVPA, child MVPA/parent sed., parent MVPA/child sed., joint sed.) was tested in a separate model. The level of analysis was the 30-sec. epoch. Only pairs with at least one minute (two 30-sec. epochs) of the joint behavior were included in each model. Models testing the effect of child weight status controlled for child age, sex, ethnicity (Hispanic versus non-Hispanic) and annual household income. Models testing the effects of parent weight status controlled for parent age, sex, ethnicity (Hispanic versus non-Hispanic) and annual household income. Predicted margins (i.e., standardized predicted probabilities adjusting for all of the other model covariates) were calculated using the multinomial logistic regressions (34). All models used the generalized estimation equation (GEE) model-fitting method to adjust standard errors for the clustering of observations within each parent-child pair. We only interpreted weight status differences for land use categories where at least 5% of the joint behavior occurs for both weight status groups, as categories with < 5% of the behavior would have little clinical or policy relevance. All analyses were conducted using SUDAAN 10.0 (RTI International, Research Triangle Park, NC).

Results

Descriptive Statistics

A total of 1,023 parent-child pairs responded to the recruitment materials of which 667 pairs were reached for eligibility screening. Among those who were screened, 447 pairs met eligibility criteria of which 407 pairs were further consented to participate in the study. From this group, 387 pairs completed the baseline data collection of which 363 pairs provided some accelerometer and GPS data. Among these pairs, 291 had a sufficient amount of matched accelerometer and GPS data to be included in the data analyses. Demographic characteristics for parent-child pairs with sufficient data are shown in Table 1 (N = 291). Children were 52.2% female and 43.0% Hispanic. The distribution of children's weight status was as follows: 63.9% under or normal weight, 15.5% overweight, 20.6% obese. The majority of parents participating in this study were female (87.6%) with a mean age around 40.0 years (SD = 6.0 years, range = 26 – 62 years). Parents' weight status was as follows: 27.2% under or normal weight, 41.2% overweight, 31.6% obese. About one quarter of the households in the sample earned \$30,000 annually. Parent-child pairs, who were excluded due to insufficient GPS and accelerometer data, differed from the analytic sample on a few factors. Parents in the excluded sample were younger and more likely to be male than parents in the analytic sample (p 's < .05). Children from pairs excluded due to insufficient data had a higher mean BMI percentile than the analytic sample (p < .05).

On average, parent-child pairs had 4.5 (SD = 1.6) valid days of matched accelerometer-GPS data over the 7-day period (M = 310.8 minutes, SD = 75.2 minutes per day)—after removing records representing overnight hours (11pm–5am), school time (8am–3pm on weekdays), time spent in motorized transit (> 32 kph), missing accelerometer and GPS data and outliers, and accelerometer non-wear for either the parent or the child. Information about missing, outlying and non-wear data for the GPS and accelerometer are reported elsewhere (9). Missing GPS data (due to power loss or signal loss) and accelerometer non-wear accounted for the majority of the unavailable data.

Children performed an average of 19.5 minutes (SD = 15.5) of total MVPA and 170.7 minutes (SD = 53.53) of total sedentary behavior each day during non-school waking hours. Parents engaged in 11.7 minutes (SD = 11.7 minutes) of total MVPA and 191.0 minutes (SD = 55.5 minutes) of total sedentary behavior.

Parent-child pairs spent an average of 233.6 minutes (SD = 80.0) per day in the same location (less than 50m apart) during non-school waking hours together, not accounting for

time spent together in the car. Of that time, parents and children spent 2.4 minutes (SD = 4.1, Median = 0.9) per day performing MVPA together and 92.9 minutes (SD = 40.1, Median = 89.3) per day engaging in sedentary behavior together during non-school, waking hours. Parents performed 7.4 minutes (SD = 7.2, Median = 5.3) per day of sedentary behavior while their child was engaging in MVPA nearby (< 50m). Conversely, children engaged in 1.9 minutes (SD = 2.3, Median = 1.1) per day of sedentary behavior while their parent was performing MVPA nearby (< 50m).

Of the 291 parent-child pairs in the analytic sample, 229 pairs (79%) had at least one minute of joint MVPA (6,566 joint MVPA epochs across the entire analytic sample), 273 pairs (94%) had at least one minute of child MVPA accompanied by parent sedentary behavior (14,433 total epochs across the entire sample), 279 pairs (96%) had at least one minute of parent MVPA accompanied by child sedentary behavior (4,681 total epochs across the entire sample), and all of them had at least one minute of joint sedentary behavior (2,465,559 total epochs across the entire sample). These values represent the sizes of the subsamples analyzed in the current paper. Other descriptive information about joint parent-child behaviors observed in this study is reported elsewhere (9).

Locations of Joint Parent-Child Behaviors

Figure 1 shows the proportion of joint MVPA, child MVPA accompanied by parent sedentary behavior, parent MVPA accompanied by child sedentary behavior, and joint sedentary behavior occurring in each land use type during non-school waking hours. The largest proportion of joint parent-child MVPA occurred in residential locations (35%), followed by commercial land uses (e.g., retail stores, restaurants, personal services, private health club/gym, motels) (24%), open space (e.g., parks, gardens, wildlife preserves), (20%), educational institutions (e.g., schools) (14%), public facilities (e.g., government, health care, religious, libraries, community centers) (7%), and mixed/other land uses (e.g., roads, water) (1%). Figure 2 shows an example of a parent and child performing MVPA together at a location classified as open space. When joint parent-child MVPA occurred in an open space, 35% of these instances took place within the pair's neighborhood and 65% occurred outside of their neighborhood. The proportion of joint parent-child MVPA occurring within the neighborhood was even less for commercial (17%), educational (13%) and public facility (10%) settings. During non-school waking hours, the majority of child MVPA accompanied by parent sedentary behavior, parent MVPA accompanied by child sedentary behavior, and joint sedentary behavior occurred in residential locations (~75%), with smaller proportions taking place in the other land use categories. Open spaces served as the location for 13% of child MVPA accompanied by parent sedentary behavior, 10% of parent MVPA accompanied by child sedentary behavior, and 8% of joint sedentary behavior during non-school waking hours.

Locations of Joint Behaviors by Child Weight Status

Table 2 shows that after adjusting child age, sex, Hispanic origin, and annual household income, the locations of joint parent-child MVPA differed by child weight status (Adj. Wald $F = 3.45$, $df = 10$, $p < .01$). As compared with children who were under/normal weight and overweight, obese children were less likely to perform joint MVPA with their parent on land classified as open space than in residential locations (see Figure 3). Differences in the land use type of parent MVPA accompanied by child sedentary behavior were by child weight status were also observed (Adj. Wald $F = 2.79$, $df = 10$, $p < .01$). However, this difference occurred in educational settings versus residential locations, and education settings accounted for less than 5% of parent MVPA/child sedentary behavior for normal weight, overweight, and obese children. The locations of child MVPA accompanied by parent

sedentary behavior and joint sedentary behavior during non-school waking hours did not differ by child weight status.

Locations of Joint Behaviors by Parent Weight Status

Table 3 show that the locations of joint parent-child MVPA differed by parent weight status (Adj. Wald $F = 3.27$, $df = 10$, $p < .01$) after adjusting parent age, sex, Hispanic origin, and annual household income. As compared with parents who were under and normal weight, overweight and obese parents were less likely to perform joint MVPA with their child on land classified as open space as compared with residential settings (see Figure 4). Differences in the land use type of child MVPA accompanied by parent sedentary behavior were also observed by child weight status (Adj. Wald $F = 2.20$, $df = 10$, $p < .05$). However, these differences occurred in commercial and other settings versus residential locations, and commercial and other settings each represented less than 5% of child MVPA/parent sedentary behavior for normal weight, overweight, and obese parents. The locations of parent MVPA accompanied by child sedentary behavior and joint sedentary behavior during non-school waking hours did not differ by parent weight status.

Discussion

The study represents one of the first attempts to use objective measures (GPS and accelerometer) to understand where children and their parents engage in physical activity and sedentary behavior together during non-school waking hours. Results indicated that joint parent-child physical activity occurred fairly evenly across residential locations, open spaces/parks, and commercial venues. The majority of the open spaces/parks and commercial venues used for joint parent-child MVPA were outside of the pair's immediate neighborhood. In contrast, the vast majority of joint parent-child sedentary behavior occurred in residential settings. Instances where child MVPA was accompanied by parent sedentary behavior and parent MVPA was accompanied by child sedentary behavior was also more likely in residential locations, with the former more likely than the latter to take place in open spaces/parks. The locations of joint parent-child behavior showed some differences by weight status.

Results indicating that open spaces and parks offered a common setting for joint physical activity performed by parents and children add to a growing body of literature highlighting the importance of these types of settings for providing physical activity opportunities (35–37). Previous studies examining children's total physical activity have found smaller percentages to take place in parks and green spaces (only 2–10%) (14–18) than the percentage of child's physical activity with a parent occurring in open spaces in the current study (almost 20%). A possible explanation is that parks and open spaces provide more opportunities for physical activities involving both children and their parents (e.g., playing catch, sports, bicycling), whereas joint parent-child physical activities may be restricted in other settings (e.g., health clubs, dance or karate studios), which may offer primarily single-age group classes and activities. Other research suggests that children with greater access to parks during childhood have lower BMI when they reach age 18 (38). Interestingly, results from the current study indicate that a notable amount of sedentary behavior occurs in parks and open spaces. On average during non-school waking hours, parents and children engaged in 8 minutes of joint sedentary behavior per day in open spaces and parks. There was also about 1–2 minutes per day, on average, when children were physically active at a park or open space with a sedentary parent nearby. If both of these instances of parental sedentary behavior at a park or open space were converted to physical activity, parents could obtain an additional hour or so more of physical activity per week, which would account for over a third of the total recommended amount (39).

The likelihood of joint parent-child physical activity taking place at parks or other types of open space, however, varied by child and parent weight status. As hypothesized, normal and underweight children and parents were more likely to engage in joint physical activity at a park or open space than obese children and adults. In contrast, obese children and parents, engaged in a greater percentage of their joint physical activities on school grounds, public facilities (including community centers and churches), and commercial locations (including retail stores and restaurants). In a previously published paper reporting results for this same study sample, the average number of joint parent-child physical MVPA minutes per day did not differ by parent or child weight status (9). Also, post-hoc analyses of interactions by joint parent-child physical activity level found that among parent-child pairs who engaged in more than 2.0 minutes per day of joint MVPA, those pairs comprised of normal weight performed a greater percentage of joint MVPA at parks and open space (32%) than pairs comprised of obese parents (7%) ($p = .051$). These results suggest that the manner but not the amount of joint parent-child physical activity differs between weight groups. Normal and overweight parents and children may accumulate their joint physical activity through recreational sports and exercise performed at parks, beaches, or other open spaces. Obese parents and children, on the other hand, may perform more of their joint physical activity through walking behaviors, possibly due to the fact that other types of exercise are perceived as too strenuous. In fact, a recent study found that obese individuals performed almost half of their moderate intensity activity through lower intensity, longer duration walking (22). Walking activities performed together by obese children and parents (whether for recreational or transportation purposes) may be more likely to occur in residential, commercial, public, and educational land use types than parks and open space. Regardless, there may be reasons to encourage obese parents and children to engage in more joint parent-child physical activity at parks and open spaces. Primarily, evidence suggests that physical activity occurring in green spaces is more enjoyable (40,41) and has a greater likelihood of reaching moderate-to-vigorous intensity (33) than activity occurring elsewhere. Therefore, encouraging park-based joint physical activity for obese children may offer health benefits in the long term by supporting the maintenance of higher energy expending activities.

Despite the use of objective physical activity and location measures, this study has a few limitations. It should be noted that there was an imbalance in the proportion of mothers (88%) versus fathers (12%) comprising the parent-child pair. Thus, father-child physical activity and sedentary behavior is underrepresented. The accelerometer and GPS technology have inherent weaknesses. The GPS units have differential measurement error depending on available satellites, meteorology, and physical obstructions, with errors often less than 5m, but also frequently much larger (i.e. > 15m), which could have caused over- or underestimation of joint activity when the linear distance of parent-child proximity was close to the specific 50m. GPS measurement error could also have resulted in misclassification of land use types when the individual was near the border separating two different land use parcels. Also, a greater amount of measurement error and missing data is expected with indoor compared to outdoor wear, although the SiRF star III chipset is designed to improve the performance of the GPS device in indoor environments. Rates of data loss were similar to rates observed in other studies using accelerometer (42,43) and GPS devices (44–46). However, the number of missing analysis units was compounded by the deletion of matched parent-child minutes when either accelerometer or GPS observations were missing in either the parent or the child. While there is some evidence to suggest that 2 days is a reliable indicator of MVPA (25,47), whether 2 days offers a reliable indicator of joint physical activity and sedentary behavior is unknown. Statistical analyses adjusted for the clustering of observations within parent-child pairs, which is important as these observations are likely to be correlated. The analyses, however, did not adjust for spatial autocorrelation because we expect observations closer together in space to have an increased

likelihood of the same land use type. Furthermore, the methodology for land use classification may not accurately represent active transportation or recreational trips. The land use of joint behaviors taking place on sidewalks or roads could be misclassified as the use category of the adjacent land parcel (e.g., commercial, educational) instead of the other category, which includes roads. Also, we were unable to control for potential differences in park access because many of the participants lived in a new community for which complete park GIS data are not yet available. Furthermore, data were not available on the nature of the social interactions between parents and children during joint physical activity and sedentary behavior (e.g., interacting behavior versus parallel but non-interacting behavior). Lastly, joint behaviors performed by pairs with younger, male parents and children with a higher BMI may not be fully represented because they were more likely to be excluded due to insufficient data. Also, results may not be generalizable to urban populations, since the data were primarily captured in suburban and exurban communities.

Parks and other types of open spaces may offer opportunities for unstructured joint parent-child physical activity, which may be restricted in other settings such as organized classes and lessons. The lower likelihood of parks and open spaces to serve as locations for joint parent-child physical activity among obese parents and children highlights an area for future investigation and intervention. Whether land use patterns for joint parent-child activity represent preferences for types of activities or amenities should be the focus of further research.

Acknowledgments

Support for this project was provided by National Cancer Institute #R01-CA-123243 (Pentz, PI) and the American Cancer Society Mentored Research Scholar Grant 118283-MRSGT-10-012-01-CPPB (Dunton, PI). We thank the Active Living Research Accelerometer Loan Program. Acknowledgments also go to Robert Gomez, B.A. and Keito Kawabata, B.A. of the University of Southern California for their assistance in managing this project. The clinical trials registration number for this study is NCT00986011.

References

1. Eriksson M, Nordqvist T, Rasmussen F. Associations between parents' and 12-year-old children's sport and vigorous activity: the role of self-esteem and athletic competence. *J Phys Act Health*. 2008; 5:359–373. [PubMed: 18579915]
2. Wagner A, Klein-Platat C, Arveiler D, Haan MC, Schlienger JL, Simon C. Parent-child physical activity relationships in 12-year old French students do not depend on family socioeconomic status. *Diabetes Metab*. 2004; 30:359–366. [PubMed: 15525880]
3. Beets MW, Cardinal BJ, Alderman BL. Parental social support and the physical activity-related behaviors of youth: a review. *Health Educ Behav*. 2010; 37:621–644. [PubMed: 20729347]
4. Davison KK, Cutting TM, Birch LL. Parents' activity-related parenting practices predict girls' physical activity. *Med Sci Sports Exerc*. 2003; 35:1589–1595. [PubMed: 12972881]
5. Trost S, Loprinzi P. Parental Influences on Physical Activity Behavior in Children and Adolescents: A Brief Review. *AM J Lifestyle Med*. 2011; 5:171–181.
6. Welk GJ, Wood K, Morss G. Parental influences on physical activity in children: An exploration of potential mechanisms. *Ped Exerc Sci*. 2003; 15:19–33.
7. Kimiecik JC, Horn TS. Parental beliefs and children's moderate-to-vigorous physical activity. *Res Q Exerc Sport*. 1998; 69:163–175. [PubMed: 9635330]
8. Eriksson M, Nordqvist T, Rasmussen F. Associations between parents' and 12-year-old children's sport and vigorous activity: the role of self-esteem and athletic competence. *J Phys Act Health*. 2008; 5:359–373. [PubMed: 18579915]
9. Dunton GF, Almanzo E, Liao Y, et al. Joint physical activity and sedentary behavior in parent-child pairs. *Med Sci Sports Exerc*. 2012; 44:1473–1480. [PubMed: 22367744]

10. Thompson JL, Jago R, Brockman R, Cartwright K, Page AS, Fox KR. Physically active families - de-bunking the myth? A qualitative study of family participation in physical activity. *Child Care Health Dev.* 2010; 36:265–274. [PubMed: 20047594]
11. Ransdell LB, Dratt J, Kennedy C, O'Neill S, DeVoe D. Daughters and mothers exercising together (DAMET): a 12-week pilot project designed to improve physical self-perception and increase recreational physical activity. *Women & Health.* 2001; 33:101–116.
12. Cutting TM, Fisher JO, Grimm-Thomas K, Birch LL. Like mother, like daughter: familial patterns of overweight are mediated by mothers' dietary disinhibition. *Am J Clin Nut.* 1999; 69:608–613.
13. Maddison R, Jiang Y, Vander Hoorn S, Exeter D, Mhurchu CN, Dorey E. Describing patterns of physical activity in adolescents using global positioning systems and accelerometry. *Pediatr Exerc Sci.* 2010; 22:392–407. [PubMed: 20814035]
14. Oreskovic NM, Blossom J, Field AE, Chiang SR, Winickoff JP, Kleinman RE. Combining global positioning system and accelerometer data to determine the locations of physical activity in children. *Geospat Health.* 2012; 6:263–272. [PubMed: 22639128]
15. Rainham DG, Bates CJ, Blanchard CM, Dummer TJ, Kirk SF, Shearer CL. Spatial classification of youth physical activity patterns. *Am J Prev Med.* 2012; 42:e87–e96. [PubMed: 22516507]
16. Wheeler BW, Cooper AR, Page AS, Jago R. Greenspace and children's physical activity: a GPS/GIS analysis of the PEACH project. *Prev Med.* 2010; 51:148–152. [PubMed: 20542493]
17. Quigg R, Gray A, Reeder AI, Holt A, Waters DL. Using accelerometers and GPS units to identify the proportion of daily physical activity located in parks with playgrounds in New Zealand children. *Prev Med.* 2010; 50:235–240. [PubMed: 20153361]
18. Dunton GF, Kawabata K, Intille S, Wolch J, Pentz MA. Assessing the social and physical contexts of children's leisure-time physical activity: an ecological momentary assessment study. *Am J Health Promot.* 2012; 26:135–142. [PubMed: 22208410]
19. Troped PJ, Wilson JS, Matthews CE, Cromley EK, Melly SJ. The built environment and location-based physical activity. *Am J Prev Med.* 2010; 38:429–438. [PubMed: 20307812]
20. Dunton GF, Berrigan D, Ballard-Barbash R, Graubard BI, Atienza AA. Social and physical environments of sports and exercise reported among adults in the American Time Use Survey. *Prev Med.* 2008; 47:519–524. [PubMed: 18675293]
21. Dorsey KB, Herrin J, Krumholz HM. Patterns of moderate and vigorous physical activity in obese and overweight compared with non-overweight children. *Int J Pediatr Obes.* 2011; 6:e547–e555. [PubMed: 20883127]
22. Spees CK, Scott JM, Taylor CA. Differences in amounts and types of physical activity by obesity status in US adults. *Am J Health Behav.* 2012; 36:56–65. [PubMed: 22251783]
23. U.S. Census Bureau. State and County QuickFacts. 2012. Available at <http://quickfacts.census.gov/qfd/states/06/0613210.html>.
24. Troiano RP, Berrigan D, Dodd KW, Masse LC, Tilert T, McDowell M. Physical activity in the United States measured by accelerometer. *Med Sci Sports Exerc.* 2008; 40:181–188. [PubMed: 18091006]
25. Byrd-Williams C, Belcher B, Spruijt-Metz D, Davis J, Ventura E, Kelly L, et al. Increased Physical Activity and Reduced Adiposity in Overweight Hispanic Adolescents. *Med Sci Sports Exerc.* 2010; 42:478–484. [PubMed: 19952807]
26. Freedson P, Melanson E, Sirard J. Calibration of the computer science and applications, Inc. accelerometer. *Med Sci Sports Exerc.* 1997; 30:777–781. [PubMed: 9588623]
27. Freedson P, Pober D, Janz K. Calibration of accelerometer output for children. *Med Sci Sports Exerc.* 2005; 37(11 Suppl):S523–S530. [PubMed: 16294115]
28. Harrell JS, McMurray RG, Baggett CD, Pennell ML, Pearce PF, Bangdiwala SI. Energy costs of physical activities in children and adolescents. *Med Sci Sports Exerc.* 2005; 37:329–336. [PubMed: 15692331]
29. Roemmich JN, Clark PA, Walter K, Patrie J, Weltman A, Rogol AD. Pubertal alterations in growth and body composition. V. Energy expenditure, adiposity, and fat distribution. *Am J Physiol Endocrinol Metab.* 2000; 279:E1426–E1436. [PubMed: 11093932]

30. Treuth MS, Schmitz K, Catellier DJ, McMurray RG, Murray DM, Almeida MJ, Going S, Norman JE, Pate R. Defining accelerometer thresholds for activity intensities in adolescent girls. *Med Sci Sports Exerc.* 2004; 36:1259–1266. [PubMed: 15235335]
31. Healy GN, Dunstan DW, Salmon J, Cerin E, Shaw JE, Zimmet PZ, Owen N. Breaks in sedentary time: beneficial associations with metabolic risk. *Diabetes Care.* 2008; 31:661. [PubMed: 18252901]
32. SCAG, Southern California Association of Governments. Existing Land Use Database. 2008
33. Almanza E, Jerrett M, Dunton G, Seto E, Pentz MA. Health Place. A study of community design, greenness, and physical activity in children using satellite, GPS and accelerometer data. 2012; 18:46–54.
34. Korn, EL.; Graubard, BI. Analysis of Health Surveys. New York: John Wiley and Sons; 1999.
35. Kaczynski AT, Henderson KA. Parks and recreation settings and active living: a review of associations with physical activity function and intensity. *J Phys Act Health.* 2008; 5:619–632. [PubMed: 18648125]
36. Cummins S, Fagg J. Does greener mean thinner? Associations between neighbourhood greenspace and weight status among adults in England. *Int J Obes (Lond).* 2011
37. Cohen DA, Ashwood JS, Scott MM, et al. Public parks and physical activity among adolescent girls. *Pediatrics.* 2006; 118:1381–1389.
38. Wolch J, Jerrett M, Reynolds K, et al. Childhood obesity and proximity to urban parks and recreational resources: A longitudinal cohort study. *Health Place.* 2010 Oct 15.
39. Physical Activity Guidelines Advisory Committee Report, 2008. Washington, DC: US Department of Health and Human Services; 2008. Physical Activity Guidelines Advisory Committee.
40. Thompson Coon J, Boddy K, Stein K, Whear R, Barton J, Depledge MH. Does participating in physical activity in outdoor natural environments have a greater effect on physical and mental wellbeing than physical activity indoors? A systematic review. *Environ Sci Technol.* 2011; 45:1761–1772. [PubMed: 21291246]
41. Dunton GF, Liao Y, Intille S, Wolch J, Pentz MA. Physical and social contextual influences on children's leisure-time physical activity: an ecological momentary assessment study. *J Phys Act Health.* 2011; 8:S103–S108. [PubMed: 21350250]
42. Colley R, Gorber SC, Tremblay MS. Quality control and data reduction procedures for accelerometry-derived measures of physical activity. *Health Rep.* 2010; 21:63–69. [PubMed: 20426228]
43. Van Coevering P, Harnack L, Schmitz K, Fulton JE, Galuska DA, Gao S. Feasibility of using accelerometers to measure physical activity in young adolescents. *Med Sci Sports Exerc.* 2005; 37:867–871. [PubMed: 15870643]
44. Cooper AR, Page AS, Wheeler BW, et al. Mapping the walk to school using accelerometry combined with a global positioning system. *Am J Prev Med.* 2010; 38:178–183. [PubMed: 20117574]
45. Oliver M, Badland H, Mavoa S, Duncan MJ, Duncan S. Combining GPS, GIS, and accelerometry: methodological issues in the assessment of location and intensity of travel behaviors. *J Phys Act Health.* 2010; 7:102–108. [PubMed: 20231761]
46. Rodriguez DA, Brown AL, Troped PJ. Portable global positioning units to complement accelerometry-based physical activity monitors. *Med Sci Sports Exerc.* 2005; 37:S572–S581. [PubMed: 16294120]
47. Hart TL, Swartz AM, Cashin SE, Strath SJ. How many days of monitoring predict physical activity and sedentary behaviour in older adults? *Int J Behav Nutr Phys Act.* 2011; 16(8):62. [PubMed: 21679426]

Figure 1.

Joint parent-child behaviors during non-school waking hours by land use categories. The x-axis indicates the types of joint parent-child behavior. The y-axis indicates the percent of time the joint behavior occurred in each land use category. Land use categories were as follows: residential (e.g., houses, apartments, condos), commercial (e.g., retail, restaurants, office use, manufacturing), open space (e.g., vacant lots, parks, golf courses, gardens, beaches), educational (e.g., schools and school grounds), facilities (e.g., community centers, churches, libraries), and other (e.g., military, mixed uses, airports, freeways, roads, utilities). Joint behavior was defined by a linear separation distance of less than 50m between the parent and child. MVPA = Moderate-to-Vigorous Physical Activity. Sed. = Sedentary behavior.

Figure 2. Geovisualization of Global Positioning System (GPS) monitoring points for a parent-child pair show joint moderate-to-vigorous physical activity (MVPA) occurring within a land use categorized as open space. Joint behavior was defined by a linear separation distance of less than 50m between the parent and child.

Figure 3.

Predicted probability of joint parent-child moderate-to-vigorous physical activity (MVPA) during non-school waking hours in different land use categories by children's Body Mass Index (BMI) category. Land use categories were as follows: residential (e.g., houses, apartments, condos), commercial (e.g., retail, restaurants, office use, manufacturing), open space (e.g., vacant lots, parks, golf courses, gardens, beaches), educational (e.g., schools and school grounds), facilities (e.g., community centers, churches, libraries), and other (e.g., military, mixed uses, airports, freeways, roads, utilities). Error bars on the graph represent standard errors. Joint behavior was defined by a linear separation distance of less than 50m between the parent and child.

Figure 4.

Predicted probability of joint parent-child moderate-to-vigorous physical activity (MVPA) during non-school waking hours in different land use categories by parents' Body Mass Index (BMI) category. Land use categories were as follows: residential (e.g., houses, apartments, condos), commercial (e.g., retail, restaurants, office use, manufacturing), open space (e.g., vacant lots, parks, golf courses, gardens, beaches), educational (e.g., schools and school grounds), facilities (e.g., community centers, churches, libraries), and other (e.g., military, mixed uses, airports, freeways, roads, utilities). Error bars on the graph represent standard errors. Joint behavior was defined by a linear separation distance of less than 50m between the parent and child.

Table 1

Demographic Characteristics for Parent-Child Pairs with Sufficient Data (N = 291)

Child	
Sex (n, %)	
Female	152, 52.2%
Male	139, 47.8%
Age (mean, range)	11.2, 8 – 14
BMI Category (n, %)	
Under/Normal Weight	186, 63.9%
Overweight	45, 15.5%
Obese	60, 20.6%
Race (n, %)	
Caucasian/White	76, 26.1%
African Am./Black	11, 3.8%
Hispanic	125, 43.0%
Asian	27, 9.3%
Other	52, 17.9%
Parent	
Sex (n, %)	
Female	255, 87.6%
Male	36, 12.4%
Age (mean, range)	39.6, 26 – 62
BMI Category (n, %)	
Under/Normal Weight	79, 27.2%
Overweight	120, 41.2%
Obese	92, 31.6%
Annual Household Income (n, %)	
\$30,000	73, 26.5%
\$30,001 – \$60,000	61, 22.1%
\$60,001 – \$100,000	82, 29.7%
> \$100,000	60, 21.7%

Note. Sample sizes vary due to missing data.

Table 2

Results of Multinomial Logistic Regression Predicting Land Use Type for Joint Parent-Child Behaviors During Non-School Waking Hours by Child Body Mass Index (BMI) Categories (Odds Ratio, 95% CI).

Covariate	Commercial	Open Space	Educational	Public Facilities	Other	Adj. Wald <i>F</i>
Joint MVPA						
Child BMI Category						3.45**
Under/Normal Weight	1.00	1.00	1.00	1.00	1.00	
Overweight	1.49 (.75 – 2.98)	.88 (.36 – 2.15)	.84 (.18 – 4.00)	1.25 (.51 – 3.03)	1.11 (.25 – 4.87)	
Obese	2.57 (1.16 – 5.72)	.43 (.18 – 1.02)	3.37 (1.49 – 7.62)	4.05 (1.39 – 11.79)	.49 (.05 – 4.66)	
Child MVPA/Parent Sed.						
Child BMI Category						1.76
Under/Normal Weight	1.00	1.00	1.00	1.00	1.00	
Overweight	.78 (.40 – 1.53)	1.17 (.50 – 2.78)	.90 (.30 – 2.67)	.72 (.25 – 2.07)	.10 (.01 – 1.01)	
Obese	.48 (.20 – 1.16)	.91 (.42 – 2.00)	.49 (.16 – 1.53)	1.99 (.89 – 4.46)	2.46 (.50 – 12.15)	
Parent MVPA/Child Sed.						
Child BMI Category						2.79**
Under/Normal Weight	1.00	1.00	1.00	1.00	1.00	
Overweight	.49 (.20 – 1.22)	2.48 (.79 – 7.85)	.32 (.08 – 1.26)	1.30 (.36 – 4.71)	1.88 (.28 – 12.71)	
Obese	1.18 (.41 – 3.41)	1.17 (.32 – 4.33)	.12 (.03 – .43)	1.65 (.59 – 4.55)	6.47 (.57 – 73.66)	
Joint Sed.						
Child BMI Category						1.01
Under/Normal Weight	1.00	1.00	1.00	1.00	1.00	
Overweight	1.21 (.80 – 1.83)	1.55 (.61 – 3.95)	.61 (.21 – 1.79)	.89 (.49 – 1.61)	.52 (.18 – 1.55)	
Obese	.97 (.59 – 1.59)	.59 (.20 – 1.77)	.39 (.15 – 1.01)	.99 (.53 – 1.86)	2.31 (.45 – 11.77)	

MVPA = Moderate-to-Vigorous Physical Activity. Sed = Sedentary behavior. The unit of analysis is the 30-sec. epoch. All models control for child age, sex, ethnicity (Hispanic vs. non-Hispanic), and annual household income. Reference group for land use categories = residential (e.g., houses, apartments, condos). Land use categories were as follows: commercial (e.g., retail, restaurants, office use, manufacturing), open space (e.g., vacant lots, parks, golf courses, gardens, beaches), educational (e.g., schools and school grounds), facilities (e.g., community centers, churches, libraries), and other (e.g., military, mixed uses, airports, freeways, roads, utilities). Joint behavior was defined by a linear separation distance of less than 50m between the parent and child.

**
 $p < .01$.

Table 3

Results of Multinomial Logistic Regression Predicting Land Use Type for Joint Parent-Child Behaviors During Non-School, Waking Hours by Parent Body Mass Index (BMI) Categories (Odds Ratio, 95% CI).

Covariate	Commercial	Open Space	Educational	Public Facilities	Other	Adj. Wald F
Joint MVPA						
Parent BMI Category						3.27**
Under/Normal Weight	1.00	1.00	1.00	1.00	1.00	
Overweight	.99 (.46 – 2.14)	.55 (.24 – 1.24)	1.27 (.44 – 3.64)	1.91 (.62 – 5.88)	.33 (.06 – 1.84)	
Obese	1.64 (.69 – 3.88)	.28 (.12 – .64)	3.79 (1.41 – 10.23)	2.34 (.51 – 10.78)	.77 (.07 – 8.11)	
Child MVPA/Parent Sed.						
Parent BMI Category						2.20*
Under/Normal Weight	1.00	1.00	1.00	1.00	1.00	
Overweight	.62 (.29 – 1.32)	.69 (.33 – 1.42)	.61 (.24 – 1.58)	1.07 (.41 – 2.80)	.08 (.01 – .53)	
Obese	.37 (.15 – .95)	1.05 (.42 – 2.63)	.38 (.13 – 1.12)	1.81 (.65 – 5.07)	1.95 (.33 – 11.46)	
Joint Sed.						
Parent BMI Category						1.02
Under/Normal Weight	1.00	1.00	1.00	1.00	1.00	
Overweight	1.17 (.77 – 1.79)	1.11 (.44 – 2.81)	1.11 (.46 – 2.67)	.95 (.53 – 1.72)	.28 (.10 – .78)	
Obese	.92 (.60 – 1.40)	.85 (.31 – 2.29)	.89 (.30 – 2.67)	.74 (.35 – 1.55)	1.38 (.39 – 4.87)	

MVPA = Moderate-to-Vigorous Physical Activity. Sed = Sedentary behavior. The unit of analysis is the 30-sec. epoch. All models control for parent age, sex, ethnicity (Hispanic vs. non-Hispanic), and annual household income. Reference group for land use categories = residential (e.g., houses, apartments, condos). Land use categories were as follows: residential (e.g., houses, apartments, condos), commercial (e.g., retail, restaurants, office use, manufacturing), open space (e.g., vacant lots, parks, golf courses, gardens, beaches), educational (e.g., schools and school grounds), facilities (e.g., community centers, churches, libraries), and other (e.g., military, mixed uses, airports, freeways, roads, utilities). Joint behavior was defined by a linear separation distance of less than 50m between the parent and child. Results for the model predicting the land use type for Parent MVPA/Child Sed. are not presented because the *p*-value was approaching 0 due to singularities in data.

* *p*<.05.

** *p*<.01.