Hypertrophic Cardiomyopathy Associated with Mid-cavity Obstruction and High Left Intraventricular Pressure

Ramush A. Bejiqi¹, Ragip J. Retkoceri¹, Hana Sh. Bejiqi²

Department of Cardiology, Pediatrics Clinic, University Clinical Centre of Kosova, Prishtina, Kosova¹

Clinic for Obstetric and Gynecology, University Clinical Centre of Kosova, Prishtina, Kosova²

Care report

SUMMARY

We report a case of a child, with a rare form of the idiopathic hypertrophic cardiomyopathy, associated with mid-cavity obstruction

and high intraventricular peak pressure. Cardiomyopathy, diagnosed antenatally, was followed postnatally and, despite of a lot echocardiographic findings - the growing, development and clinical signs are minimal.

Key words: Hypertrophic cardiomyopathy, intraventricular pressure, left ventricle outlet tract obstruction, sudden death, echocardiography.

1. INTRODUCTION

The most characteristic morphological abnormality in children with hypertrophic cardiomyopathy (HCM) is the excessive hypertrophy of and no dilated left ventricle with absence of other cardiac or systemic diseases that could produce left ventricular hypertrophy (1, 2, 3). Often, it is asymmetric in nature, with a preference for ventricular septum and occurs either in sporadic or familial forms. From the genetics point of view the disease is highly variable with respect to the specific gene mutation and degree of penetration. Basically, there is no correlation between the severity of the disease and symptomatology. Often, children lack symptoms for a long period of time and frequently, the disease is detected in case of presence murmur or arrhythmia. Classically, symptoms include pulmonary congestion, fatigue, palpitations, chest pain, syncope and congestive heart failure. A number of pathophysiological components and process are identified: systolic dysfunction and left ventricular outflow tract obstruction, diastolic dysfunction, coronary artery abnormalities, leading to myocardial ischemia, mitral regurgitation, arrhythmias and sudden cardiac death (2, 3). M-mode

and 2-D echocardiography are the primary screening and evaluation of HCM tools, whereas Doppler imaging can fully delineate the entire spectrum of hemodynamic abnormalities.

2. CASE REPORT

During the routine echosonographic screening by an obstetrician hypertrophy of the left ventricle was registered, otherwise, until then, a normal pregnancy of 36 weeks. Left ventricular hypertrophy was present at the second level of fetal echocardiographic examination. Especially hypertrophic was presented interventricular septum, with separate cavity of left ventricle in two parts. Apical part has been seen as separated from the outlet and inlet part, acquiring a diverticular shape. This part was clearly presented that actively take part in left ventricle contractility. By continuous Doppler waves between the two parts high velocity was measured consistent with the maximum hemodynamic gradient of 37 mmHg at a mid left ventricle level. The Color flow Doppler imaging showed a narrow area of the left ventricular cavity in systole and a trivial mitral regurgitation was noted. This raised doubts that it is more likely that this is a case of

suspected hypertrophic cardiomyopathy then a case of diverticulum of the left ventricle.

Pregnancy went to full term, with a vaginal delivery of a male infant weighing 3280 gr, Appgar score 8/9. Complete clinical and cardiological examination was obtained. An arterial blood gas showed normal range. Clinical examination presented a quite precordium, normal first heart sound, very short midsystolic murmur on the apex and single second heart sound. Electrocardiogram showed non-specific changes, presenting normal sinus rhythm, normal frequency and left heart deviation with signs of biventricular hypertrophy. A chest radiogram revealed an enlargement cardiac silhouette with a narrow mediastinum. The pulmonary vasculature was normal in appearance, and there were no infiltrates seen. Echocardiography demonstrated a normal systemic and pulmonary vein connection, with small interatrial communication. There was normal atrio-ventricular and ventriculo-arterial connection. Cross-sectional echocardiography from apical four-chamber view, during the diastole presented hypertrophy of the left ventricle walls, especially interventricular septum. During the sys-

tola left ventricular walls form obstruction at the middle level of the left ventricle, causing two “separate chambers” where apical part actively participates during the contraction. The Color flow Doppler imaging shows a very narrow and constricted area at the middle ventricle level in systole whereas continuous Doppler imaging presented a high systolic velocity with the peak gradient of 53 mmHg. Other echocardiography findings, including aortic valve, aortic arch and coronary artery, were normal. Blood pressure was within normal limits. The child’s growth and development was completely normal, there was no sweating or fatigue during physical activities. Last examination was done at the age of 11 months when the child weighted 9450 g (50 %), and clinically manifested only sideropenic anemia. Final echocardiographic examination registered increasing mid-cavity obstruction at the papillary muscles level and increasing intracavitary pressure, with peak gradient of 105.1 mmHg. In good basic condition child is continually under Propranolol therapy, 1 mg/kg divided in three doses.

3. DISCUSSION

An atypical form of hypertrophic cardiomyopathy is presented with mid-cavitary obstruction of left ventricle and fast increasing intracavitary pressure. This form involves selective hypertrophy and obstruction at the mid-left ventricular level. In adults, these findings may represent effects of a long-standing hypertension with relatively small left ventricular cavities in some individuals (2, 8). In children this type of hypertrophic cardiomyopathy is very rare and it is quite likely that there is a distinct anatomic subtype of hypertrophic cardiomyopathy, with still clearly unidentified etiology. Over the recent years, knowledge about HCM has evolved enormously, mainly through advances in molecular genetics and in understanding of pathophysiological mechanisms, as well as our awareness of the great variability in its expression. HCM is now recognized as a genetic cardiac disease with an autosomal dom-

Figure 1. Turbulent flow presented by color Doppler image from apical view of a intracavitary mid-level obstruction of left ventricle

inant pattern of inheritance, but with variable penetrance and expression albeit that sporadic case occurs (7). There is genetic heterogeneity, with more than one gene being associated with the clinical manifestation and there is a great phenotypic variability, not only among unrelated families, but also within the same family (1).

All forms of HCM have in common an inappropriate left ventricular hypertrophy, often with abnormal myofibril orientation and represent a diverse spectrum of disease with varying degrees of hypertrophic expression even among a given family. There is, however, wide variation of expression within affected

families. It is believed that the penetrance of the disease is incomplete during childhood and adolescence, increasing with age to nearly complete penetrance in adulthood. In general, there is no correlation between the severity of the disease and the symptomatology (3, 6). Children often lack symptoms, and as a rule, HCM does not interfere with the physical development of the patient except in its severe forms. Affected children may remain asymptomatic for a long period of time. The clinical manifestation of the classic form of HCM result from systolic dysfunction and dynamic left ventricular outflow tract obstruction, diastolic dysfunction, mitral regurgita-

Figure 2. High peak pressure measured by continuous Doppler from apical view

tion, a high prevalence of arrhythmias and sudden cardiac death (4).

Our case is specific in many aspects including early antenatal diagnosis with normal growth without other manifestations during pregnancy, normal at term delivery and asymptomatic neonatal and postnatal period. Our case is especially interesting because of the increasing intracavitary pressure for a short time – during delivery it was 37 mmHg and at the 11th month of life the peak pressure was 105.1 mmHg.

When HCM is suspected, the antenatal detection is relatively easy after elimination of other cardiac diseases caused by the left ventricle hypertrophy (coronary anomalies, congenital heart obstruction like aortic valvular stenosis, aortic coarctation etc.) (5). According to the literature, it is still a small number of children antenatally diagnosed with mid-cavity HCM and possibility of miss diagnosis is high. Also, our case is a rare one because of the existing such high peak intracavitary

pressure without any clinical manifestations.

Conflict of interest: non declared.

REFERENCES

1. Ciro E, Marion BJ, Bonow RO, Cannon RO, Epstein SE. Relation between marked changes in left ventricular outflow tract gradient and disease progression in hypertrophic cardiomyopathy. *Am J Cardiol.* 1984; 53: 1103-1109.
2. Feigenbaum H, Armstrong WF, Ayan T. Feigenbaum's Echocardiography, Lippincott Williams and Wilkins, Philadelphia, 2005; 552-553.
3. Lakkis NM, Nagueh SF, Kleiman NS. Et al. Echocardiography-guided ethanol septal reduction for hypertrophic obstructive cardiomyopathy. *Circulation.* 1998; 98: 1750-1755.
4. Ross J, Jr, Braunwald E, Gault JH et al. The mechanism of the intraventricular pressure gradient in idiopathic hypertrophic subarotic stenosis. *Circulation.* 1966; 34(4) 558-578.
5. Shapiro LM, McKenna WJ. Distribution of left ventricular hypertrophy in hypertrophic cardiomyopathy: a two dimensional echocardiographic study. *J Am Coll Cardiol.* 1983; 2: 437-444.
6. Spirito P, Maron BJ et al. Relation between extent of left ventricular hypertrophy and occurrence of sudden cardiac death in hypertrophic cardiomyopathy *J Am Coll Cardiol.* 1990; 15: 1521-1526.
7. Watkins H, Ashrafian H, McKenna WJ, The genetics of hypertrophic cardiomyopathy: *Heart.* 2008; 94; 10: 1264-1269.
8. Webb JG, Sasson Z, Rakowski H. et al. Apical hypertrophic cardiomyopathy: clinical follow-up and diagnostic correlates. *J Am Coll Cardiol.* 1990; 15: 83-90.

Corresponding author: Ramush Bejqi, MD. Department of Abdominal Surgery, University Clinical Centre of Kosovo, Prishtina, Republic of Kosovo. E-mail: rbejqi@gmail.com

Quality of Life through Quality of Information

XXIV Conference of the European Federation for Medical Informatics

MIE

2012

PISA - ITALY
26 - 29 AUGUST
www.mie2012.it

- Home
- Stay informed: Register as Friend
- Call for Submissions
- Instructions for Authors
- Online Submissions
- Conference Registration
- Committees
- Venue
- Hotels
- Travel to Pisa
- General Information
- Tourist Information

Welcome!

On behalf of the Italian Medical Informatics Association (AIIM), the e-health Italian Community and the European Federation for Medical Informatics (EFMI), we invite you to the 24th European Medical Informatics Conference - MIE2012 - in Pisa, August 26th - 29th, 2012.

The theme of the conference is Quality of Life through Quality of Information.

We intend to focus on the role of e-health in pursuing benefits in terms of quality of life for patients, health-care personnel, citizens and the society. The topic can be analyzed from many points of view, and MIE2012 will provide a unique platform for fruitful exchange of ideas and experiences among the actors and stakeholders of ICT supported healthcare. Both methodologies and examples of applications are the base for the discussion, and all of us will learn not only from success stories in the field, but also from failures. The opinion and experiences of both the designers/planners and the addressees/users of e-health systems are of great importance when the target is the global improvement of the quality of life, at personal and public level. For this reason we do encourage health professionals, policy makers, and also patient/citizen associations to take an active part in the conference and share their knowledge.

Keynotes, scientific sessions, thematic workshops and panels, short communications and tutorials will constitute the traditional backbone of the conference. In parallel, special "partnership for innovation" tracks and round tables will be organized to get the best from the synergy between the scientific / industrial / political world. Various kinds of contributions are solicited and welcome in order to provide MIE2012 attendees a stage for active participation.

In addition MIE2012 will offer national, regional and European projects and initiatives the opportunity to demonstrate their results in the "Village of the Future" that will be hosted during the conference.