CORRESPONDENCE

Prevalence of Wine Intolerance: Results of a Survey From Mainz, Germany

by Dr. rer. nat. Petra Wigand, Prof. Dr. rer. nat. Maria Blettner,
Prof. Dr. med. Joachim Saloga, Prof. Dr. rer. nat. Heinz Decker in volume
25/2012

Substances Used in Wine Clarification as Potential Allergens

Potential allergens also include proteins and substances used in the clarification of wine: fish gelatin or isinglass, protein from chicken eggs (ovalbumin), milk (casein) products, rubber arabicum, lysozyme, pectinase, cellulose, glucosidase, urease, and aroma enzymes (1). Furthermore, moulds—especially *Botrytis cinerea*, which is responsible for noble rot in grapes—as well as insect proteins (for example, wasp venom or bee venom), which may have contaminated the mash during the pressing process, play an allergenic part that should not be underestimated (1, 2).

Triggers for allergic intolerance reactions such as headaches or migraine after ingestion of red wine may also be the phenolic flavonoids that are present in the skin of the grapes (anthocyanins and catechins), because they inhibit catechol-O-methyltransferase in the intestinal wall as well as the phenol sulfotransferase. Other non-organic ingredients, such as the ethanol metabolite acetaldehyde and acetic acid, also play a part in wine intolerance—often in the shape of urticaria (1).

The intolerance to alcohol in general that was observed by the authors is supported by a representative survey in adults in Copenhagen (4142/6000=70.7% response rate), which showed a prevalence of hypersensitivity reactions after alcohol consumption (especially red wine) of 8% (3, 4).

In Bavaria, for example, all superior quality wines have to be examined by the Bavarian Health and Food Safety Agency, in order to ensure that the legally defined upper limits of sulfur dioxide are adhered to. In order to protect consumers from intolerance reactions to sulfites, this type of state controlled wine examination in addition to the official testing number on the label would also be desirable for other German states.

DOI: 10.3238/arztebl.2013.0039a

REFERENCES

1. Wüthrich B: Weinallergien und -intoleranzen. *Allergologie* 2011; 34: 427–36.
2. Armentia A, Pineda F, Fernández S: Wine-induced anaphylaxis and sensitization to hymenoptera venom. *N Engl J Med* 2007; 357: 719–20.
3. Linneberg A, Berg ND, Gonzalez-Quintela A, Vidal C, Elberling J: Prevalence of self-reported hypersensitivity symptoms following intake of alcoholic drinks. *Clin Exp Allergy* 2008; 38: 145–51.

4. Wigand P, Blettner M, Saloga J, Decker H: Prevalence of wine intolerance: results of a survey from Mainz, Germany. *Dtsch Arztebl Int* 2012; 109(25): 437–44.

Dr. oec. troph. Martin Hofmeister

Verbraucherzentrale Bayern e.V., Referat Lebensmittel und Ernährung,
München
hofmeister@vzbayern.de

Conflict of interest statement

The author is employed by the Consumer Centre of the German Federal State of Bavaria.

Types of Grapes May Be Triggers

Intolerance reactions are not only associated generally with white, red, and rosé wines but also with individual grape varieties.

A case in point is the effect of the Merlot grape, which has become fashionable among wine drinkers in recent years. In patients with latent or manifest hypertension, a notable increase in blood pressure was observed after drinking Merlot, and this was accompanied by tachyarrhythmias. The reactions usually occur at night or in the early morning and are accompanied by symptoms resembling those of myocardial infarction. What is particularly striking is that even minute doses can trigger such reactions, and this also applies to blended wines consisting of a combination of several grapes, which often contain only 10–20% of Merlot.

It is therefore recommended that these associations should be explained to patients with latent or manifest hypertension.

DOI: 10.3238/arztebl.2013.0039b

REFERENCES

1. Wigand P, Blettner M, Saloga J, Decker H: Prevalence of wine intolerance: results of a survey from Mainz, Germany. *Dtsch Arztebl Int* 2012; 109(25): 437–44.

Dr. med. Nikolaus W. Klehr

Facharzt für Dermatologie und Venerologie
Oberanger 30
80331 München

Conflict of interest statement

The author declares that no conflict of interest exists.

Declaring Allergens Has Become Mandatory

In cases of allergic or intolerance reactions after drinking wine, some thought should be given to possible associations with substances used during enological treatments. Some substances with a high allergenic potential, such as sulfites or sulfur dioxide, were actually mentioned in the article. These substances, if used in quantities in excess of 10 mg per kg or liter, have to be declared in foods as well as in wine.

However, the use of other enological treatments, such as lysozyme or ovalbumin or casein from cow's

milk, is less commonly known. The presence of such substances in wine can accordingly trigger undesirable reactions in patients with existing allergies to cow's milk or egg protein. According to Implementing Regulation (EU) No. 579/2012 of the European Commission of 29 June 2012, these substances will now have to be declared in wine, in the same way as in other foods, if they are present in the final product.

This is helpful for affected patients, but it means that in future, the declaration of allergens on the wine label will have to be read even more carefully.

DOI: 10.3238/arztebl.2013.0039c

REFERENCES

1. Wigand P, Blettner M, Saloga J, Decker H: Prevalence of wine intolerance: results of a survey from Mainz, Germany. *Dtsch Arztebl Int* 2012; 109(25): 437–44.

Dr. med. Klaus Richter

Bundesinstitut für Risikobewertung
Fachgruppe Ernährungsrisiken, Allergien und Neuartige Lebensmittel
Abteilung Lebensmittelsicherheit, Berlin
Klaus.Richter@bfr.bund.de

Conflict of interest statement

The author declares that no conflict of interest exists.

In Reply:

We thank our correspondents for their great interest in our study, and special thanks go to Dr Richter from the Federal Institute for Risk Assessment (Bundesinstitut für Risikobewertung) and Dr Hofmeister from the Consumer Centre of the German Federal State of Bavaria for their comments, which we are in complete agreement with. With regard to Dr Klehr's letter, we have to admit that we do not have any data—neither from our own investigations nor from those of others—that confirm a special case for Merlot wines.

Real allergic reactions and/or other intolerance reactions can in principle be triggered by any ingredient within wine, but the likelihood varies. Such ingredients include substances used in the clarification and stabilization of wine and their known allergenic potential, such as mentioned in the letters from Dr Hofmeister and Dr Richter. In modern vinification processes these are hardly of any relevance, at least not in Rheinhessen, but they have had to be declared on the label since 2012

if they can be confirmed within the wine. Another class of factors that may trigger intolerance reactions include the phenols and phenolic flavonoids. Intolerance reactions are often reported for red wines, and their causes are not easily identifiable, especially if the wine is a blend of different grapes. We don't think that any generalization is possible about whether Merlot is particularly important in this setting, as Dr Klehr remarks. As a rule, any reactions are person-specific and can be avoided only by not drinking wine and alcohol or changing to a wine that is more easily tolerated.

Of course acetaldehyde, a product of alcohol metabolism, also needs to be borne in mind, as this can trigger intolerance reactions.

Even though sulfite is used in winemaking since antiquity, in order to protect the wine against oxidation and micro-organisms, the well-known intolerance reaction and allergic reaction owing to sulfite needs to be mentioned in this context. Sulfite now has to be declared on the wine label. We are currently studying the molecular foundations for the interactions with different wine ingredients, in order to be able to better understand their effects on people's health.

Dr Hofmeister rightly mentioned the Copenhagen study (1), which reported a similar prevalence of symptoms of intolerance after ingestion of alcohol, especially red wine, as our own study.

DOI: 10.3238/arztebl.2013.0040

REFERENCES

1. Linneberg A, Berg ND, Gonzalez-Quintela A, Vidal C, Elberling J: Prevalence of self-reported hypersensitivity symptoms following intake of alcoholic drinks. *Clin Exp Allergy* 2008; 38: 145–51.
2. Wigand P, Blettner M, Saloga J, Decker H: Prevalence of wine intolerance: results of a survey from Mainz, Germany. *Dtsch Arztebl Int* 2012; 109(25): 437–44.

Dr. rer. nat. Petra Wigand

Prof. Dr. rer. nat. Heinz Decker

Institut für Molekulare Biophysik, Johannes Gutenberg-Universität Mainz
hdecker@uni-mainz.de

Prof. Dr. rer. nat. Maria Blettner

Institut für Medizinische Biostatistik, Epidemiologie und Informatik,
Universitätsmedizin der Johannes Gutenberg-Universität Mainz

Prof. Dr. med. Joachim Saloga

Hautklinik und Poliklinik,
Universitätsmedizin der Johannes Gutenberg-Universität Mainz

Conflict of interest statement

The authors declare that no conflict of interest exists.