Figure 2 Assembled equipment for seroma drainage

ment, seroma fluid can be briskly evacuated into the 400ml vacuum drain. This can be disconnected and replaced via the Luer lock connector should further drainage be required, without re-siting the needle or compromising sterility of the closed system. If a sterile fluid sample is required, this can be taken from the drain following aspiration.

DISCUSSION

The causes of seromas are not fully understood although increased age, weight, lymphatic disruption and use of extensive electrocautery have been implicated.^{1,2} Numerous treatments for seromas have been trialled including fibrin-based glues,³ pressure garments, and local application of corticosteroids⁴ and sclerosants such as erythromycin.⁵ None eliminate occurrence conclusively.

Our method for drainage avoids the need for awkward syringe aspiration, which can be time consuming and uncomfortable for the patient. The technique is versatile and can also be used for other procedures such as conservative aspiration of a large scrotal hydrocoele.

References

1. Agrawal A, Ayantunde AA, Cheung KL. Concepts of seroma formation and prevention in breast cancer surgery. *ANZ J Surg* 2006; **76**: 1,088–1,095.
2. Porter KA, O'Connor S, Rimm E, Lopez M. Electrocautery as a factor in seroma formation following mastectomy. *Am J Surg* 1998; **176**: 8–11.
3. Carless PA, Henry DA. Systematic review and meta-analysis of the use of fibrin sealant to prevent seroma formation after breast cancer surgery. *Br J Surg* 2006; **93**: 810–819.
4. Taghizadeh R, Shoaib T, Hart AM, Weiler-Mithoff EM. Triamcinolone reduces seroma re-accumulation in the extended latissimus dorsi donor site. *J Plast Reconstr Aesthet Surg* 2008; **61**: 636–642.
5. Ali-Khan AS, Orlando A, Kenealy J. Erythromycin sclerotherapy in the management of seroma. *J Plast Reconstr Aesthet Surg* 2009; **62**: e55–58.

External rotation of the draining vein improves flow in arteriovenous fistulas

P Yates, A Mahmood

University Hospital, Coventry, UK

CORRESPONDENCE TO

Phillip Yates, E: pjy4@hotmail.co.uk

BACKGROUND

Arteriovenous (AV) fistulas were first described in 1966 by Brescia *et al.*¹ High success rates have resulted in the technique of anastomosis remaining largely unchanged since this time. Here we describe a refinement on this technique that optimises venous length and helps prevent kinking of the draining AV fistula vein.

TECHNIQUE

Often the vein and artery are closely related, allowing side-to-side anastomosis or end-to-side anastomosis with a chamfered vein. In some instances there is a significant distance between artery and vein, especially in the antecubital fossa, when an almost 90° AV anastomosis may be required. In this situation maintenance of venous length is pivotal to avoid anastomotic tension and twisting that may preclude fistula maturation.

Figure 1 (A) Vein and artery running parallel; anterior aspect of vessels denoted by dashed/solid black line. (B) Vein mobilised and anastomosed to artery 'as it lies' without rotation (solid and dashed lines continuous); length disparity between heel and toe of anastomosis leads to kinking at point X. (C) Vein rotated by 90° from 'as it lies' position; reduces disparity in length between heel and toe of anastomosis and reduces kinking (solid line at 90° to dashed line)

In closely related vessels, chamfering of the vein minimises kinking with a minimal loss of venous length. However, with widely spaced vessels chamfering alone does not prevent kinking and may serve to significantly shorten the draining vein. Figure 1B shows that when the vein is anastomosed directly 'as it lies', without rotation, the length disparity between the heel and toe of the anastomosis leads to kinking of the vein at point X, the apex of the fistula.

DISCUSSION

Figure 1C shows that by rotating the vein through 90°, from the 'as it lies' position, prior to anastomosis, heel to toe length disparities are minimised, thereby avoiding kinking of the draining vein and maintaining flow and patency.

Reference

1. Brescia MJ, Cimino JE, Appel K, Hurwich BJ. Chronic hemodialysis using venipuncture and a surgically created arteriovenous fistula. *N Engl J Med* 1966; **275**: 1,089–1,092.