Published in final edited form as:

Inorg Chem. 2012 December 3; 51(23): 12959–12967. doi:10.1021/ic302084g.

Reactivity of Diphenylpropynone Derivatives Toward Metal-Associated Amyloid- β Species

Amit S. Pithadia^{1,‡}, Akiko Kochi^{1,‡}, Molly T. Soper¹, Michael W. Beck¹, Yuzhong Liu¹, Sanghyun Lee², Alaina S. DeToma¹, Brandon T. Ruotolo^{1,*}, and Mi Hee Lim^{1,2,*}

¹Department of Chemistry, University of Michigan, Ann Arbor, Michigan 48109

²Life Sciences Institute, University of Michigan, Ann Arbor, Michigan 48109

Abstract

In Alzheimer's disease (AD), metal-associated amyloid- β (metal- $A\beta$) species have been suggested to be involved in neurotoxicity; however, their role in disease development is still unclear. To elucidate this aspect, chemical reagents have been developed as valuable tools for targeting metal- $A\beta$ species, modulating the interaction between the metal and $A\beta$, and subsequently altering metal- $A\beta$ reactivity. Herein, we report the design, preparation, characterization, and reactivity of two diphenylpropynone derivatives (**DPP1** and **DPP2**) composed of structural moieties for metal chelation and $A\beta$ interaction (bifunctionality). The interactions of these compounds with metal ions and $A\beta$ species were confirmed by UV-Vis, mass spectrometry, and docking studies. The effects of these bifunctional molecules on the control of in vitro metal-free and metal-induced $A\beta$ aggregation were investigated and monitored by gel electrophoresis and transmission electron microscopy (TEM). Both **DPP1** and **DPP2** showed reactivity toward metal- $A\beta$ species over metal-free $A\beta$ species to different extents. In particular, **DPP2**, which contains a dimethylamino group, exhibited greater reactivity with metal- $A\beta$ species than **DPP1**, suggesting a structure-reactivity relationship. Overall, our studies present a new bifunctional scaffold that could be utilized to develop chemical reagents for investigating metal- $A\beta$ species in AD.

Introduction

Alzheimer's disease (AD) is a fatal neurodegenerative disease that affects over 5 million people in the United States.^{1, 2} A pathological hallmark of the diseased brain is an accumulation of misfolded amyloid- β ($A\beta$) aggregates.^{2–8} Monomeric $A\beta$ peptides, generated from the proteolytic cleavage of the transmembrane amyloid precursor protein, can further aggregate to produce oligomers, protofibrils, and eventually fibrils. It is still not completely understood which conformation of $A\beta$ is associated with AD neuropathogenesis; however, recent evidence has proposed that soluble oligomers might be the toxic species due to their ability to interrupt neurotransmission.^{4, 5, 9} In addition to $A\beta$ species, elevated concentrations of transition metals, such as Fe, Cu and Zn, have been observed within the

*To whom correspondence should be addressed: bruotolo@umich.edu (B. T. R.) and mhlum@umich.edu (M. H. L.).

‡These authors contributed equally.

ASSOCIATED CONTENT

Supporting Information

Table S1 and Figures S1 – S6 are available in the supporting information. This material is available free of charge via the Internet at <http://pubs.acs.org>.

Notes

The authors declare no competing financial interest.

deposits of A β aggregates.^{2, 7, 8, 10–15} The possible relationship between metal-associated A β species (metal–A β species) and neurotoxicity has been suggested based on observations that upon binding to A β , metal ions facilitate peptide aggregation as well as enhance oxidative stress caused by overproduction of reactive oxygen species;^{2, 4, 5, 7, 8, 12–21} however, this connection has not been clearly revealed.

To gain a better understanding of the involvement of metal–A β species in AD pathogenesis, recent advancements in the development of chemical reagents to specifically target metal–A β species and modulate their interaction and reactivity have been made.^{2, 7, 8, 21–33} Among them, rationally-designed small molecules with both metal chelation and A β interaction properties (defined as bifunctionality) have been devised. Some of the compounds have been fashioned based on the incorporation approach (Figure 1), where a metal chelation site is directly inserted into a known A β imaging agent with minimal structural modifications, along with consideration of criteria for possible brain uptake.^{8, 21–25, 27–35} These molecules have been shown to be able to control the interactions and reactivity of metal–A β species in vitro and/or in living cells, suggesting that the incorporation approach (Figure 1) could be considered as a promising design strategy to fashion suitable chemical reagents for uncovering the potential role of metal–A β species in AD development.

Herein, we report the design and preparation of a new class of bifunctional small molecules (**DPP1** and **DPP2**, Figure 1), composed of a diphenylpropynone framework, as chemical reagents for targeting and regulating metal–A β species, the characterization of their interactions with metal ions and A β species, as well as the investigation of their in vitro reactivity with metal–A β species. From our overall results and observations, the diphenylpropynone scaffold could be a structural feature for constructing chemical reagents for investigating metal–A β species. Moreover, the dimethylamino functionality in **DPP2** was observed to be an important moiety for potentially enhancing A β interaction of the compound, as reported previously,^{29, 31, 36} offering noticeable reactivity with metal–A β species.

Experimental Section

Materials and Procedures

All reagents were purchased from commercial suppliers and used as received unless otherwise stated. The compound, 3-phenyl-1-(pyridin-2-yl)prop-2-yn-1-one (**DPP1**) was prepared following previously reported methods.^{37–39} A β _{1–40} (DAEFRHDSGYEVHHQKLVFFAEDVGSNKGAIIGLMVGGVV) was purchased from AnaSpec (Fremont, CA). An Agilent 8453 UV-Visible (UV-Vis) spectrophotometer was used to measure the optical spectra. Transmission electron microscopy (TEM) images were recorded with a Philips CM-100 transmission electron microscope. A SpectraMax M5 microplate reader (Molecular Devices, Sunnyvale, CA) was employed for the measurement of absorbance for 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) and parallel artificial membrane permeability (PAMPA) assays. Nuclear magnetic resonance (NMR) spectra of small molecules and for Zn²⁺ binding studies were obtained by a Varian 400 MHz NMR spectrometer. Mass spectrometric measurements for compounds were conducted by a Micromass LCT Electrospray Time-of-Flight mass spectrometer. Mass spectrometric studies for investigating the interaction of **DPP1** and **DPP2** with the peptide were carried out on a Waters Synapt G2 ion-mobility mass spectrometer (Milford, MA).

Preparation of 3-(4-(dimethylamino)phenyl)-1-(pyridin-2-yl)-2-propyn-1-one (**DPP2**)—

DPP2 was synthesized by slight modifications to a previously reported procedure.³⁹ To a solution of 4'-dimethylaminophenyl acetylene (0.42 g, 2.9 mmol) in dry tetrahydrofuran (THF, 5 mL) was added *n*-butyllithium (0.72 mL, 2.9 mmol, 2.5 M solution

in hexanes) dropwise by a syringe over 5 min at -40°C . The reaction mixture was allowed to stir at -40°C for 10 min and then warmed to -15°C . After 30 min, the mixture was cooled to -78°C (dry ice/acetone) and a solution of Weinreb's amide (0.40 g, 2.4 mmol in 9 mL of dry THF) was introduced through a syringe. The reaction mixture was allowed to react at -78°C for 10 min and then warmed to room temperature and followed by 1 h stirring. The reaction was quenched by adding saturated aqueous NaHCO_3 (2 mL), diluted with EtOAc (10 mL), and washed with brine (2×25 mL). The aqueous layer was extracted with EtOAc (2×15 mL), and the combined organic solutions were dried over MgSO_4 and filtered, followed by removal of the solvent in vacuo. The crude product was purified by silica gel chromatography (CH_2Cl_2 :EtOAc = 9:1) to yield an orange product (202 mg, 0.81 mmol, 28%). ^1H NMR (400 MHz, CDCl_3)/ δ (ppm): 8.77 (d, J = 4.8 Hz, 1 H), 8.14 (d, J = 7.6 Hz, 1 H), 7.82 (td, J = 7.6, 1.6 Hz, 1 H), 7.57 (d, J = 8.4 Hz, 2 H), 7.44 (m, 1 H), 6.59 (d, J = 8.4 Hz, 2 H), 2.98 (s, 6 H). ^{13}C NMR (100 MHz, CDCl_3)/ δ (ppm): 177.6, 153.7, 151.8, 149.7, 136.8, 135.6, 126.9, 123.3, 111.4, 105.5, 100.2, 89.3, 39.9. HRMS: Calcd for $[\text{M} + \text{H}]^+$, 251.1179; Found, 251.1176.

Parallel Artificial Membrane Permeability Assay Adapted for Blood-Brain Barrier (PAMPA-BBB)—PAMPA-BBB experiments were carried out using the PAMPA Explorer kit (μION Inc., Billerica, MA) with modification to previously reported protocols.^{29, 40–42} Each stock solution was diluted with pH 7.4 Prisma HT buffer (μION) to a final concentration of 25 μM (1% v/v final DMSO concentration) and 200 μL were added to the wells of the donor plate (number of replicates = 12). BBB-1 lipid formulation (5 μL μION) was used to coat the polyvinylidene fluoride (PVDF, 0.45 μM) filter membrane on the acceptor plate. The acceptor plate was placed on top of the donor plate forming a - sandwich- and the brain sink buffer (BSB, 200 μL , μION) was added to each well of the acceptor plate. The sandwich was incubated for 4 h at ambient temperature without stirring. UV-Vis spectra of the solutions in the reference, acceptor, and donor plates were measured using a microplate reader. The PAMPA Explorer software v. 3.5 (μION) was used to calculate the $-\log P_c$ values for the compounds. CNS+/- designations were assigned by comparison to compounds that were identified in previous reports.^{29, 40–42}

Determination of Solution Speciation for DPP1, DPP2, and the Cu^{2+} -DPP2 Complex—The $\text{p}K_a$ values for **DPP1** and **DPP2** were determined by UV-Vis variable-pH titrations as previously reported.^{27, 29–33, 43} To establish the $\text{p}K_a$ values, a solution (100 mM NaCl, 10 mM NaOH, pH 12) of **DPP1** (40 μM) or **DPP2** (20 μM) was titrated with small aliquots of HCl. At least 30 spectra were recorded in the range of pH 2–10. Similarly, a solution containing **DPP2** (20 μM) and CuCl_2 in a ratio of 2:1 was titrated with small additions of HCl and at least 30 spectra were recorded over the range of pH 2–7. The acidity and stability constants were calculated by using the HypSpec program (Protonic Software, UK).⁴⁴ Speciation diagrams for **DPP1**, **DPP2**, and Cu^{2+} -**DPP2** complexes were modeled by the HySS2009 program (Protonic Software).⁴⁵

Metal Binding Studies—The interaction of **DPP1** and **DPP2** with Cu^{2+} and Zn^{2+} was determined by UV-Vis and ^1H NMR spectroscopy, respectively, based on previously reported procedures.^{28–33, 46} A solution of ligand (20 μM in EtOH) was prepared, treated with 1 to 20 equiv of CuCl_2 , and incubated at room temperature for 2.5 h (for **DPP1**) or 5 min (for **DPP2**). The optical spectra of the resulting solutions were measured by UV-Vis. The interaction of **DPP1** or **DPP2** with ZnCl_2 was observed by ^1H NMR. ZnCl_2 (1 equiv) was added to a solution of **DPP1** or **DPP2** (4 mM) in CD_3CN . The metal selectivity of both compounds was investigated by measuring the optical changes upon addition of 1 equiv of CuCl_2 to a solution of ligand (**DPP1** = 40 μM ; **DPP2** = 20 μM in EtOH) containing 1 or 25 equiv of a divalent metal chloride salt (MgCl_2 , CaCl_2 , MnCl_2 , FeCl_2 , CoCl_2 , NiCl_2 , or

ZnCl₂). The Fe²⁺ samples were prepared anaerobically (all solutions were purged with N₂). Quantification of the metal selectivity was calculated by comparing and normalizing the absorption values of metal-ligand complexes at 360 nm (for **DPP1**) or 580 nm (for **DPP2**) to the absorption at these wavelengths before and after the addition of CuCl₂ (A_M/A_{Cu}). Cu²⁺ binding of **DPP2** in the presence of A β was examined by UV-Vis. A β (25 μ M) was treated with CuCl₂ (25 μ M in 20 mM HEPES (2-[4-(2-hydroxyethyl)piperazin-1-yl]ethanesulfonic acid), pH 6.6, 150 mM NaCl for 2 min at room temperature. **DPP2** (50 μ M) was added to the resulting solution, followed by 0.5–4 h incubation. For comparison, the optical spectra of **DPP2** (50 μ M) were measured in the absence and presence of CuCl₂ (25 μ M; 0.5–4 h incubation) without A β at pH 6.6.

A β Interaction of DPP1 and DPP2 by Mass Spectrometry—The interaction of **DPP1** or **DPP2** with A β _{1–40} was investigated by nano-electrospray ionization-mass spectrometry (nESI-MS) that was carried out on a Synapt G2 quadrupole-ion mobility-mass spectrometry system. Samples were prepared by mixing stock solutions of **DPP1** or **DPP2** (prepared in DMSO) and A β _{1–40} (dissolved in 100 mM ammonium acetate, pH 6.8) to generate desired final concentrations of the peptide and the compound. Mixtures were incubated on ice or at room temperature for 2 or 4 h, respectively and then analyzed. To produce protein complex ions, an aliquot of the sample (*ca.* 5 μ L) was sprayed from the nESI emitter using a capillary voltage of 1.4 kV, with the source operating in positive ion mode and the sample cone operated at 50 V. In order to normalize nESI-MS data for non-specific and electrospray artifact interactions which could occur at high concentrations, data were acquired for A β samples containing thioflavin-T (ThT), a compound known to have no affinity for soluble forms of the A β peptide,⁴⁷ under identical concentration conditions as our **DPP1** and **DPP2** experiments. Any ThT binding observed was assumed to be due to either non-specific binding or the electrospray process, and subtracted from the intensities of the **DPP1** and **DPP2** interactions observed.⁴⁸ This procedure was performed over a broad range of concentrations. The mass spectra were acquired with the following settings and tuned to avoid ion activation and to preserve non-covalent protein-ligand complexes:⁴⁹ backing pressure, 7.3 mbar; IMS pressure reading, 3.09 mbar; ToF analyzer pressure, 1.14×10^{-6} mbar.

Docking Studies—Flexible ligand docking studies using AutoDock Vina⁵⁰ for **DPP1** and **DPP2** were conducted against the A β _{1–40} monomer from the previously determined aqueous solution NMR structure (PDB 2LFM).⁵¹ Ten conformations were selected from 20 conformations within the PDB file (1, 3, 5, 8, 10, 12, 13, 16, 17, and 20). The MMFF94 energy minimization in ChemBio3D Ultra 11.0 was used to optimize the structures of the ligands for the docking studies. The structural files of **DPP1**, **DPP2**, and the peptide were generated by AutoDock Tools and imported into PyRx,⁵² which were used to run AutoDock Vina. The search space dimensions were set to contain the entire peptide. The exhaustiveness for the docking runs was set at 1024. Docked poses of the ligands were visualized with A β using Pymol.

Amyloid- β (A β) Peptide Experiments—A β _{1–40} was used in all A β experiments. A β _{1–40} peptide (1 mg) was dissolved with ammonium hydroxide (NH₄OH, 1% v/v, aq), aliquoted, lyophilized, and stored at –80 °C. A stock solution (*ca.* 200 μ M) was prepared by redissolving A β with NH₄OH (1% v/v, aq, 10 μ L) followed by dilution with ddH₂O. All A β solutions were prepared following previously reported procedures.^{28–32} The buffered solutions (20 μ M HEPES, pH 6.6 (for Cu²⁺ samples) or pH 7.4 (for metal-free and Zn²⁺ samples), 150 μ M NaCl) were used for both inhibition and disaggregation studies. For the inhibition experiment, A β (25 μ M) was first treated with a metal chloride salt (CuCl₂ or ZnCl₂, 25 μ M) for 2 min followed by addition of a compound (**DPP1** or **DPP2**, 50 μ M in

DMSO, 1% v/v final DMSO concentration). The resulting samples were incubated at 37 °C for 4, 8, or 24 h with constant agitation. For the disaggregation experiment, A β and a metal chloride salt (CuCl₂ or ZnCl₂) were initially incubated at 37 °C for 24 h with steady agitation. The compound was added afterwards followed by additional 4, 8, or 24 h incubation at 37 °C with constant agitation.

Native Gel Electrophoresis/Sodium Dodecyl Sulfate-Polyacrylamide Gel Electrophoresis (SDS-PAGE) with Western Blotting—The A β peptide experiments (as described above) were analyzed by both native gel electrophoresis and SDS-PAGE with Western blotting using an anti-A β antibody (6E10).^{28–32, 53} Each sample containing 25 μ M A β (10 μ L) was separated using either a 10–20% gradient Tris-tricine gel (Invitrogen, Grand Island, NY) or SDS gel (4% stacking gel; 10% resolving gel; non-reducing conditions). The gel was transferred to a nitrocellulose membrane and blocked overnight with bovine serum albumin (BSA, 3% w/v, Sigma, St. Louis, MO) dissolved in Tris-buffered saline (TBS, Fisher, Pittsburgh, PA) containing 0.1% Tween-20 (TBS-T, Sigma). The membrane was treated with 6E10 (1:2,000; 2% BSA in TBS-T, Covance, Princeton, NJ) for 4 h at room temperature. The membrane was probed with a horseradish peroxidase-conjugated goat anti-mouse secondary antibody (1:5,000; Cayman Chemical, Ann Arbor, MI) in 2% BSA in TBS-T solution for 1 h at room temperature. The protein bands were visualized using the Thermo Scientific Supersignal West Pico Chemiluminescent Substrate (Rockford, IL).

Transmission Electron Microscopy (TEM)—Samples for TEM were prepared following a previously reported method.^{28–31, 33, 53} Glow-discharged grids (Formar/Carbon 300-mesh, Electron Microscopy Sciences, Hatfield, PA) were treated with samples from either inhibition or disaggregation experiments (5 μ L) for 2 min at room temperature. Excess sample was removed with filter paper and washed with ddH₂O five times. Each grid was stained with uranyl acetate (1% w/v, ddH₂O, 5 μ L) for 1 min. Uranyl acetate was blotted off and grids were dried for 15 min at room temperature. Images of samples were taken by a Philips CM-100 transmission electron microscope (80 kV, 25,000x magnification).

Cytotoxicity (MTT Assay)—The murine neuro-2a (N2a) neuroblastoma cell line was purchased from the American Type Culture Collection (ATCC, Manassas, VA). The cell line was maintained in media containing 45% Dulbecco's modified Eagle's medium (DMEM) and 50% OPTI-MEM (GIBCO, Grand Island, NY), supplemented with 5% fetal bovine serum (FBS, Sigma), 2 mM glutamine, 100 U/ml penicillin, and 100 mg/ml streptomycin (GIBCO). The cells were grown and maintained at 37 °C in a humidified atmosphere with 5% CO₂. Cell viability upon treatment of compounds was determined using a MTT assay (Sigma). N2a cells were seeded in a 96 well plate (15,000 cells in 100 μ L per well) and treated with various concentrations of **DPP1** and **DPP2** (2.5–50 μ M, final 1% v/v DMSO). After 24 h incubation at 37 °C, 25 μ L MTT (5 mg/mL in phosphate buffered saline, PBS, pH 7.4, GIBCO) was added to each well and the plates were incubated for 4 h at 37 °C. Formazan produced by the cells was dissolved overnight at room temperature by addition of a solution (100 μ L) containing *N,N*-dimethylformamide (DMF, 50% v/v, aq, pH 4.5) and sodium dodecyl sulfate (SDS, 20% w/v). A microplate reader was used to measure the absorbance (A₆₀₀).

Results and Discussion

Design Consideration, Preparation, and Characterization of Diphenylpropynone Derivatives for Targeting and Modulating Metal–A β Species

The diphenylpropynone scaffold was shown to have high binding affinity (*ca.* 6 nM) for A β aggregates; thus, it has been utilized for the design of an A β plaque imaging probe (Figure

1).⁵⁴ Based on the incorporation approach, a nitrogen donor atom was installed into this framework to generate a metal chelation site with an oxygen donor atom of the carbonyl group, which afforded two bifunctional molecules (**DPP1** and **DPP2**, Figure 1). Furthermore, a minor structural difference between **DPP1** and **DPP2** (*i.e.*, a dimethylamino functionality, suggested to be critical for A β interaction)^{29, 31, 36} was made to understand a structure-reactivity relationship. **DPP1** was synthesized as previously established.^{37–39} The new compound, **DPP2**, was prepared by slight modifications to a previously reported method (Scheme 1).³⁹

In order to predict potential drug-like and BBB penetration properties of the structural scaffolds of **DPP1** and **DPP2**, values of Lipinski's rules and logBB were calculated.^{7, 34, 35, 40, 41} As shown in Table 1, the theoretical values indicate that both compounds are drug-like and possibly BBB permeable. To verify this prediction for BBB penetration of compounds, in vitro PAMPA-BBB was performed following a previously reported procedure.^{29, 40–42} Permeability values ($-\log P_e$) were measured to be 4.2 (± 0.1) for both **DPP1** and **DPP2** (Table 1). Based on empirical classification of BBB-permeable molecules (*i.e.*, verapamil),^{29, 40–42} **DPP1** and **DPP2** would also be likely to cross the BBB.

Along with BBB permeability, the solution speciation of **DPP1** and **DPP2** was determined through UV-Vis variable-pH titration experiments ($I = 0.10$ M NaCl; room temperature; pH 2 – 10).^{27, 29–33, 43} As summarized in Figure 2, titration results indicated a single acidity constant (pK_a) for **DPP1** ($pK_a = 2.035(5)$) and two pK_a values for **DPP2** ($pK_{a1} = 7.106(1)$ and $pK_{a2} = 2.959(4)$). These pK_a values suggest that monoprotonated, diprotonated, and neutral species exist in solution of **DPP2** depending on pH (from 2 to 10), while the monoprotonated and neutral forms of **DPP1** are present in this pH range (Figure 2). In addition, the generated solution speciation diagrams of **DPP1** and **DPP2** exhibit that their neutral forms are relatively predominant at physiologically relevant pH (*i.e.*, 7.4) (**DPP1**, 100%; **DPP2**, *ca.* 65%).

Metal Binding Properties of DPP1 and DPP2

Metal binding properties of **DPP1** and **DPP2** (specifically, Cu^{2+} and Zn^{2+}) were studied by UV-Vis and NMR spectroscopy. Upon the addition of CuCl_2 (1–20 equiv) to a solution of **DPP1** and **DPP2** in EtOH, new optical bands were observed, indicative of Cu^{2+} binding to the ligand (Figure 3a). In particular, in the presence of Cu^{2+} , **DPP2**, which has a dimethylamino group, showed a distinguishable optical shift from 407 to 525 nm. NMR was employed to investigate the interaction of **DPP1** or **DPP2** with Zn^{2+} . When 1 equiv of ZnCl_2 was introduced in a solution of **DPP1** or **DPP2** (in CD_3CN), noticeable downfield chemical shifts of the pyridyl protons were recorded (Figure 3b), demonstrating that Zn^{2+} binding to the ligand occurred through the pyridyl N-donor atom.⁴⁶ Overall, both UV-Vis and NMR studies confirmed Cu^{2+} or Zn^{2+} binding to **DPP1** and **DPP2**.

To further identify binding stoichiometry and affinity, the solution speciation investigation of the Cu^{2+} -**DPP2** complexes was carried out through UV-Vis variable-pH titration experiments (1:2 [Cu^{2+}]/[**DPP2**]; $I = 0.10$ M NaCl, room temperature). Based on the pK_a values of **DPP2** and these titration results, stability constants for the Cu^{2+} -**DPP2** complexes were obtained (Figure 4, $\text{M} + \text{LH} \rightleftharpoons \text{M}(\text{LH})$ ($\log \beta_1 = 12.99(9)$); $\text{M} + \text{L} \rightleftharpoons \text{ML}$ ($\log \beta_2 = 5.85(3)$); $\text{M} = \text{Cu}^{2+}$, $\text{L} = \text{DPP2}$). A solution speciation diagram was generated from these stability constants, suggesting that the major species at pH 7 are a mixture of $\text{Cu}(\text{LH})$ and CuL complexes in a ratio of 3:2. Free Cu^{2+} was shown up to pH 7, indicating $p\text{Cu} = 6.6$ at pH 6.6 ($p\text{Cu} = -\log[\text{Cu}_{\text{unchelated}}]$) (Figure 4).^{27, 29, 31–33, 43} The $p\text{Cu}$ value suggests the approximate dissociation constant (K_d) of Cu^{2+} -**DPP2** to be *ca.* high nanomolar. When compared to the reported K_d values of Cu^{2+} -A β (picomolar to nanomolar),^{2, 5, 7, 8, 12, 14, 21}

DPP2 may interact with Cu^{2+} from soluble Cu^{2+} -A β species. In order to test if Cu^{2+} binding of ligand occurs in the presence of A β , a solution containing **DPP2** with Cu^{2+} -treated A β was monitored by UV-Vis. The new spectral features coincided with those of the Cu^{2+} -**DPP2** complex without A β were observed, suggesting an interaction of **DPP2** with Cu^{2+} in the presence of A β (Figure S1). Cu^{2+} binding of **DPP2** with A β occurred more slowly than that without A β , proposing that A β might interfere with metal binding to the ligand. Taken together, our spectroscopic studies present the capability of **DPP1** and **DPP2** to chelate Cu^{2+} and Zn^{2+} , as well as the potential interaction of **DPP2** with Cu^{2+} in the presence of A β species, which may be associated with its noticeable reactivity toward metal-A β species (vide infra).

The metal selectivity of **DPP1** and **DPP2** was also determined by competitive reactions with Cu^{2+} over biologically relevant divalent metal ions (Mg^{2+} , Ca^{2+} , Mn^{2+} , Fe^{2+} , Co^{2+} , Ni^{2+} , and Zn^{2+}), which was monitored by UV-Vis. As depicted in Figure S2, **DPP1** and **DPP2** displayed selectivity for Cu^{2+} over Mg^{2+} , Ca^{2+} , Mn^{2+} , and Zn^{2+} . Binding of both ligands to Co^{2+} and Ni^{2+} was also observed. Considering the lower abundance of Co^{2+} and Ni^{2+} than Cu^{2+} in biological systems,^{55, 56} the overall metal selectivity of **DPP1** and **DPP2** may be sufficient to be used for targeting and interacting with Cu^{2+} -A β species in heterogeneous biological environments like the brain.

A β Interaction with **DPP1** and **DPP2** Studied by MS and Docking Studies

The interaction of **DPP1** and **DPP2** with A β_{1-40} in the absence of metal ions was probed by electrospray ionization-mass spectrometry (ESI-MS), tuned to preserve non-covalent protein-ligand interactions.⁴⁹ At a low A β concentration (10 μM), a small signal corresponding to the interaction between **DPP2** (30 μM) and the A β monomer in the 3+ charge state could be detected, while no interaction between **DPP1** (60 μM) and the peptide was observed under these conditions (Figure S3). At high concentrations of the peptide (100 μM) and compounds (600 μM) both **DPP1** and **DPP2** interacted with A β species to different extents (Figure 5a). Data for **DPP1** indicated that the molecule interacted broadly with A β monomers and oligomers in 1:1, 2:1, and 3:1 A β to ligand ratios. In the case of **DPP2**, a stronger preference toward larger A β oligomers was shown, but with similar stoichiometries as **DPP1**. The total bound intensities recorded from MS data, and those from individual oligomeric species were shown in Figure 5b and Table S1. The intensities shown were normalized for both non-specific interactions and artifactual complexes formed during the electrospray process using A β :ThT binding data as a control, and ion mobility separation was used to separate oligomers that overlapped in m/z .^{6, 49, 57} From these data, it was clear that, at high concentrations, a higher proportion of **DPP1** was bound to A β species than **DPP2**, but that both could be classified as having weak A β affinity in solution (low mM K_d). Therefore, a weak A β /compound interaction was captured by MS. Normalized intensity MS data suggest that **DPP2** binding was almost exclusively driven through A β multimer interactions (Figure 5b). Overall, our MS results suggest that while both compounds could interact with A β species at high concentrations, **DPP2** was able to bind A β species at both low and high concentrations.

In order to visualize the potential interaction between A β and **DPP1/DPP2**, docking studies by AutoDock Vina⁵⁰ were performed using the previously determined NMR structure (PDB 2LFM)⁵¹ of A β_{1-40} monomer. Typically, both compounds were positioned between the α -helix and the unstructured *N*-terminal side of A β (Figures 5c and S4). Most docked structures showed a non-specific orientation of the ligand with respect to the surface features of A β . Our preliminary docking studies support the potential interaction of the compounds with A β monomer.

Effects of DPP1 and DPP2 on Metal-free and Metal-induced A β Aggregation In Vitro

Confirming metal binding and A β interaction properties (bifunctionality) of **DPP1** and **DPP2**, their influence on in vitro metal-free and metal-induced A β aggregation were studied.^{26–33} Two different experiments (inhibition and disaggregation) were performed to investigate whether **DPP1/DPP2** can control the formation of metal-free and metal-induced A β aggregates (inhibition, Figure 6) or transform preformed metal-free and metal-induced A β aggregates (disaggregation, Figure S5). Various-sized A β species from both studies were monitored by native gel electrophoresis and SDS-PAGE followed by Western blotting with an anti-A β antibody (6E10), while morphological changes were identified by transmission electron microscopy (TEM).^{26, 28–33, 53}

The inhibition studies, as shown in Figure 6, demonstrated a different reactivity of **DPP1** or **DPP2** toward metal-induced A β species over metal-free A β species. In the case of metal-involved A β aggregation, A β species with a wide range of MW were indicated with **DPP2** for both Cu²⁺- and Zn²⁺-treated samples upon longer incubation (Figure 6a, lanes 6 and 9). In the samples containing **DPP1** (lanes 5 and 8), less intense gel bands were detected across the longer incubation time, suggesting that further A β aggregation may have occurred. The reaction of **DPP1** or **DPP2** with metal-free A β also exhibited a different distribution of various-sized A β species (Figure 6a, lanes 2 and 3). A β species formed with compounds in both metal-mediated and metal-free conditions were not completely denatured by SDS implying that these molecules may generate different A β assemblies (Figure 6b). TEM images of metal-induced A β species incubated with **DPP2** for 24 h revealed smaller amorphous aggregates compared to **DPP1**; some of the metal-free A β species treated with **DPP1** and **DPP2** presented similar morphology to those untreated with compounds (Figure 6c). Overall, **DPP1** and **DPP2** displayed their ability to recognizably modulate metal-induced A β aggregation to different extents.

Furthermore, for the disaggregation experiment (Figure S5), **DPP2**-treated metal-triggered A β aggregates presented different-sized A β species than **DPP1**-treated samples, indicating that **DPP2** could alter the properties of preformed A β aggregates to a greater extent than **DPP1** (Figure S5a). In the metal-free conditions, more various-sized A β species were indicated in the presence of **DPP2** than **DPP1** (in particular, at 4 h, Figure S5a, lanes 2 and 3). The A β species generated with compounds in both metal-triggered and metal-free conditions were relatively stable in the presence of SDS (Figure S5b). As shown in Figure S5c, **DPP2** was able to reorganize preformed structured metal–A β aggregates to amorphous species more noticeably than **DPP1**. Taken together, the results from both the inhibition and disaggregation experiments presented that **DPP1** and **DPP2** could regulate metal-involved A β aggregation over metal-free aggregation in vitro differently. Moreover, the structural variation (*i.e.*, dimethylamino functionality) may enhance the contact with A β species via hydrogen bonding and/or hydrophobic interactions,^{29, 31, 36} which may afford greater reactivity toward metal–A β species.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This study was supported by the Alzheimer's Art Quilt Initiative, the Alzheimer's Association (NIRG-10-172326), and the Ruth K. Broad Biomedical Foundation (to M.H.L.), as well as start-up funding from the University of Michigan (to B.T.R.). A.K., M.T.S., and A.S.D. are grateful for the Research Excellence Fellowship from the Department of Chemistry at the University of Michigan, the NIH Training grant (T32 CA140044), and the NSF Graduate Research fellowship, respectively. We thank Dr. Xiaoming He for synthetic assistance and Nathan Merrill for the NMR measurements for Zn²⁺-binding studies.

References

1. Alzheimer's Association. *Alzheimers Dement.* 2012; 8:131–168. [PubMed: 22404854]
2. Kepp KP. *Chem. Rev.* 2012; 112:5193–5239. [PubMed: 22793492]
3. Hardy JA, Higgins GA. *Science.* 1992; 256:184–185. [PubMed: 1566067]
4. Jakob-Roetne R, Jacobsen H. *Angew. Chem. Int. Ed.* 2009; 48:3030–3059.
5. Rauk A. *Chem. Soc. Rev.* 2009; 38:2698–2715. [PubMed: 19690748]
6. Teplow DB, Lazo ND, Bitan G, Bernstein S, Wytttenbach T, Bowers MT, Baumketner A, Shea JE, Urbanc B, Cruz L, Borreguero J, Stanley HE. *Acc. Chem. Res.* 2006; 39:635–645. [PubMed: 16981680]
7. Scott LE, Orvig C. *Chem. Rev.* 2009; 109:4885–4910. [PubMed: 19637926]
8. DeToma AS, Salamekh S, Ramamoorthy A, Lim MH. *Chem. Soc. Rev.* 2012; 41:608–621. [PubMed: 21818468]
9. Haass C, Selkoe DJ. *Nat. Rev. Mol. Cell. Biol.* 2007; 8:101–112. [PubMed: 17245412]
10. Lovell MA, Robertson JD, Teesdale WJ, Campbell JL, Markesbery WR. *J. Neurol. Sci.* 1998; 158:47–52. [PubMed: 9667777]
11. Frederickson CJ, Koh J-Y, Bush AI. *Nat. Rev. Neurosci.* 2005; 6:449–462. [PubMed: 15891778]
12. Gaggelli E, Kozlowski H, Valensin D, Valensin G. *Chem. Rev.* 2006; 106:1995–2044. [PubMed: 16771441]
13. Faller P, Hureau C. *Dalton Trans.* 2009:1080–1094. [PubMed: 19322475]
14. Faller P. *ChemBioChem.* 2009; 10:2837–2845. [PubMed: 19877000]
15. Bourassa MW, Miller LM. *Metallomics.* 2012; 4:721–738. [PubMed: 22797194]
16. Zhu X, Su B, Wang X, Smith MA, Perry G. *Cell. Mol. Life Sci.* 2007; 64:2202–2210. [PubMed: 17605000]
17. Zatta P, Drago D, Bolognin S, Sensi SL. *Trends Pharmacol. Sci.* 2009; 30:346–355. [PubMed: 19540003]
18. Hureau C, Faller P. *Biochimie.* 2009; 91:1212–1217. [PubMed: 19332103]
19. Bonda DJ, Lee H-g, Blair JA, Zhu X, Perry G, Smith MA. *Metallomics.* 2011; 3:267–270. [PubMed: 21298161]
20. Jomova K, Valko M. *Toxicology.* 2011; 283:65–87. [PubMed: 21414382]
21. Pithadia AS, Lim MH. *Curr. Opin. Chem. Biol.* 2012; 16:67–73. [PubMed: 22366383]
22. Hureau C, Sasaki I, Gras E, Faller P. *ChemBioChem.* 2010; 11:950–953. [PubMed: 20401891]
23. Braymer JJ, DeToma AS, Choi J-S, Ko KS, Lim MH. *Int. J. Alzheimers Dis.* 2011; 2011 Article ID 623051.
24. Perez LR, Franz KJ. *Dalton Trans.* 2010; 39:2177–2187. [PubMed: 20162187]
25. Rodríguez-Rodríguez C, Telpoukhovskaia M, Orvig C. *Coord. Chem. Rev.* 2012; 256:2308–2332.
26. Sharma AK, Pavlova ST, Kim J, Finkelstein D, Hawco NJ, Rath NP, Kim J, Mirica LM. *J. Am. Chem. Soc.* 2012; 134:6625–6636. [PubMed: 22452395]
27. Rodríguez-Rodríguez C, Sánchez de Groot N, Rimola A, Álvarez-Larena Á, Lloveras V, Vidal-Gancedo J, Ventura S, Vendrell J, Sodupe M, González-Duarte P. *J. Am. Chem. Soc.* 2009; 131:1436–1451. [PubMed: 19133767]
28. Hindo SS, Mancino AM, Braymer JJ, Liu Y, Vivekanandan S, Ramamoorthy A, Lim MH. *J. Am. Chem. Soc.* 2009; 131:16663–16665. [PubMed: 19877631]
29. Choi J-S, Braymer JJ, Nanga RPR, Ramamoorthy A, Lim MH. *Proc. Natl. Acad. Sci. U. S. A.* 2010; 107:21990–21995. [PubMed: 21131570]
30. Choi J-S, Braymer JJ, Park SK, Mustafa S, Chae J, Lim MH. *Metallomics.* 2011; 3:284–291. [PubMed: 21210061]
31. Braymer JJ, Choi J-S, DeToma AS, Wang C, Nam K, Kampf JW, Ramamoorthy A, Lim MH. *Inorg. Chem.* 2011; 50:10724–10734. [PubMed: 21954910]
32. He X, Park HM, Hyung S-J, DeToma AS, Kim C, Ruotolo BT, Lim MH. *Dalton Trans.* 2012; 41:6558–6566. [PubMed: 22437427]

33. Jones MR, Service EL, Thompson JR, Wang MC, Kimsey IJ, DeToma AS, Ramamoorthy A, Lim MH, Storr T. *Metallomics*. 2012; 4:910–920. [PubMed: 22825244]
34. Lipinski CA, Lombardo F, Dominy BW, Feeney PJ. *Adv. Drug Deliv. Rev.* 2001; 46:3–26. [PubMed: 11259830]
35. Clark DE, Pickett SD. *Drug Discov. Today*. 2000; 5:49–58. [PubMed: 10652455]
36. Leuma Yona R, Mazères S, Faller P, Gras E. *ChemMedChem*. 2008; 3:63–66. [PubMed: 17926318]
37. Seregin IV, Schammel AW, Gevorgyan V. *Org. Lett.* 2007; 9:3433–3436. [PubMed: 17637023]
38. Harkat H, Blanc A, Weibel JM, Pale P. *J. Org. Chem.* 2008; 73:1620–1623. [PubMed: 18193886]
39. Friel DK, Snapper ML, Hoveyda AH. *J. Am. Chem. Soc.* 2008; 130:9942–9951. [PubMed: 18588297]
40. Di L, Kerns EH, Fan K, McConnell OJ, Carter GT. *Eur. J. Med. Chem.* 2003; 38:223–232. [PubMed: 12667689]
41. Avdeef A, Bendels S, Di L, Faller B, Kansy M, Sugano K, Yamauchi Y. *J. Pharm. Sci.* 2007; 96:2893–2909. [PubMed: 17803196]
42. pION Inc. BBB protocol and test compounds. Woburn (MA): 2009.
43. Storr T, Merkel M, Song-Zhao GX, Scott LE, Green DE, Bowen ML, Thompson KH, Patrick BO, Schugar HJ, Orvig C. *J. Am. Chem. Soc.* 2007; 129:7453–7463. [PubMed: 17511455]
44. Gans P, Sabatini A, Vacca A. *Ann. Chim.* 1999; 89:45–49.
45. Alderighi L, Gans P, Ienco A, Peters D, Sabatini A, Vacca A. *Coord. Chem. Rev.* 1999; 184:311–318.
46. Braymer JJ, Merrill NM, Lim MH. *Inorg. Chim. Acta*. 2012; 380:261–268.
47. Reinke AA, Gestwicki JE. *Chem. Biol. Drug Des.* 2011; 77:399–411. [PubMed: 21457473]
48. Sun J, Kitova EN, Wang W, Klassen JS. *Anal. Chem.* 2006; 78:3010–3018. [PubMed: 16642987]
49. Hernández H, Robinson CV. *Nat. Protoc.* 2007; 2:715–726. [PubMed: 17406634]
50. Trott O, Olson AJ. *J. Comput. Chem.* 2010; 31:455–461. [PubMed: 19499576]
51. Vivekanandan S, Brender JR, Lee SY, Ramamoorthy A. *Biochem. Biophys. Res. Commun.* 2011; 411:312–316. [PubMed: 21726530]
52. Wolf LK. *Chem. Eng. News*. 2009; 87:31.
53. Reinke AA, Seh HY, Gestwicki JE. *Bioorg. Med. Chem. Lett.* 2009; 19:4952–4957. [PubMed: 19640715]
54. Ono M, Watanabe H, Watanabe R, Haratake M, Nakayama M, Saji H. *Bioorg. Med. Chem. Lett.* 2011; 21:117–120. [PubMed: 21145738]
55. Lippard, SJ.; Berg, JM. California: University Science Books; 1994.
56. Gray, HB.; Stiefel, EI.; Valentine, JS.; Bertini, I. California: University Science Books; 2007.
57. Bernstein SL, Dupuis NF, Lazo ND, Wyttenbach T, Condrón MM, Bitan G, Teplow DB, Shea JE, Ruotolo BT, Robinson CV, Bowers MT. *Nat. Chem.* 2009; 1:326–331. [PubMed: 20703363]

Summary and Perspective

Following the incorporation approach, we have developed two bifunctional small molecules (**DPP1** and **DPP2**) composed of a metal chelation site and a diphenylpropynone framework (for A β interaction), which could possibly serve as chemical reagents to target and modulate metal–A β species in vitro. Their bifunctionality (metal chelation and A β interaction) was confirmed by physical methods and preliminary docking studies. Biochemical and TEM studies revealed that **DPP1** and **DPP2** could modulate metal-induced A β aggregation in vitro. Notably, **DPP2**, which has a dimethylamino group, exhibited more apparent reactivity toward metal–A β species, compared to **DPP1**. This suggests that the interaction and reactivity of molecules with metal–A β species can be tuned by such structural variations, proposing a structure–interaction–reactivity relationship. **DPP1** and **DPP2**, however, would be limited in their biological applications since they displayed cytotoxicity in living cells at low micromolar concentrations (Figure S6). Overall, the promising in vitro reactivity of these potentially BBB-permeable molecules toward metal–A β species warrants pursuit of structural modifications that would improve the viability of diphenylpropynone derivatives in biological settings, followed by more detailed characterization by MS and molecular modeling. This study has demonstrated the capability of two diphenylpropynone derivatives to target metal–A β species and modulate their interaction and reactivity in vitro, which can be further optimized toward the development of future chemical reagents for investigating metal–A β species in biological systems.

Figure 1.

Incorporation approach (top) and structures of small molecules (bottom). Left to right: 3-(4-(dimethylamino)phenyl)-1-(4-iodophenyl)-2-propyn-1-one; **DPP1** = 3-phenyl-1-(pyridin-2-yl)prop-2-yn-1-one; **DPP2** = 3-(4-(dimethylamino)phenyl)-1-(pyridin-2-yl)-2-propyn-1-one.

		pK_a^a	
		DPP1	DPP2
LH_2	$\rightleftharpoons LH + H$	–	7.106(1)
LH	$\rightleftharpoons L + H$	2.035(5)	2.959(4)

Figure 2.

Solution speciation studies of **DPP1** and **DPP2**. Top: UV-Vis spectra of **DPP1** (40 μ M, left) and **DPP2** (20 μ M, right) in the range of pH 2–10. Middle: Solution speciation diagrams for **DPP1** (left) and **DPP2** (right) (F_L = fraction of compound with given protonation). Bottom: Acidity constants (pK_a) of L (L = **DPP1** or **DPP2**). Charges are omitted for clarity. ^a Error in the parentheses is shown in the last digit. Conditions: I = 0.10 M NaCl; room temperature.

Figure 3.

Cu^{2+} or Zn^{2+} binding of **DPP1** and **DPP2**. (a) UV-Vis spectra of **DPP1** (left) and **DPP2** (right) with CuCl_2 (1–20 equiv) in EtOH at room temperature (incubation for 2.5 h (for **DPP1**) and 5 min (for **DPP2**)). (b) ^1H NMR spectra of **DPP1** (left, black) or **DPP2** (right, black) with ZnCl_2 (red) in CD_3CN at room temperature ([compound] = 4 mM; [ZnCl_2] = 4 mM).

Figure 4.

Solution speciation investigation of the Cu²⁺-DPP2 complexes. Top left: UV-Vis spectra (pH 2–7) for the Cu²⁺-DPP2 complexes ([Cu²⁺]/[L] = 1:2; [Cu²⁺]_{total} = 10 μM; 7 h incubation with ligand (L) prior to pH titration, L = DPP2; room temperature). Top right: Solution speciation diagram of the Cu²⁺-DPP2 complexes (F_{Cu} = fraction of free Cu and Cu complexes). Bottom: Stability constants (log β) of the Cu²⁺-DPP2 complexes. Charges are omitted for clarity. ^aError in the parentheses is shown in the last digit. ^bThe species containing CuL₂ was introduced into the calculation model yielding a good fit to the data.

Figure 5.

Aβ interactions of **DPP1** and **DPP2**. (a) MS data for the complexes of Aβ₁₋₄₀ and **DPP1** or **DPP2** ([Aβ] = 100 μM; [compound] = 600 μM; M = monomer, D = dimer, and T = trimer). Many binding stoichiometries were detected, including 1:1 (star), 2:1 (square), and 3:1 (triangle), (b) A histogram showing the total bound MS signal intensity, normalized for non-specific interactions and ESI-MS artifacts, for each binding stoichiometry observed in (a), (c) Docking studies of **DPP1** (orange) and **DPP2** (green) with Aβ₁₋₄₀ (PDB 2LFM) by AutoDock Vina. Poses for both compounds were overlapped in this conformation (other conformations, Figure S4). The helical region of Aβ (H13-D23) is highlighted in color (tan) in both the cartoon (left) and surface (right) representations.

Figure 6.

Inhibition experiment (scheme, top). Analysis of various-sized $A\beta$ species by (a) native gel electrophoresis and (b) SDS-PAGE (non-reducing conditions) with Western blot using an anti- $A\beta$ antibody (6E10). (c) TEM images of the 24 h incubated samples. Conditions: $[A\beta] = 25\ \mu$ M; $[CuCl_2 \text{ or } ZnCl_2] = 25\ \mu$ M; $[compound] = 50\ \mu$ M; pH 6.6 (for Cu^{2+} samples) or 7.4 (for metal-free and Zn^{2+} samples); 4, 8, or 24 h incubation; $37^\circ C$; constant agitation.

Scheme 1.
Synthesis of **DPP2**.

Table 1Values (MW, $\text{clog}P$, HBA, HBD, PSA, $\log BB$, and $-\log P_e$) of **DPP1** and **DPP2**.

Calculation ^a	DPP1	DPP2	Lipinski's rules & others
MW	207	250	450
$\text{clog}P$	2.47	2.63	5.0
HBA	2	3	10
HBD	0	0	5
PSA	30.0	33.2	90 Å ²
$\log BB$	0.0620	0.0384	> 0.3 (readily cross the BBB) < -1.0 (poorly distributed in the brain)
$-\log P_e$ ^b	4.2 ± 0.1	4.2 ± 0.1	
CNS+/- prediction ^c	CNS+	CNS+	$-\log P_e < 5.4$ (CNS+) $-\log P_e > 5.7$ (CNS-)

^aMW: Molecular weight; $\text{clog}P$: Calculated logarithm of the octanol-water partition coefficient; HBA: Hydrogen-bond acceptor atoms; HBD: Hydrogen-bond donor atoms; PSA: Polar surface area; $\log BB = -0.0148 \times \text{PSA} + 0.152 \times \text{clog}P + 0.130$.

^bThe values of $-\log P_e$ were measured by the Parallel Artificial Membrane Permeability assay (PAMPA).

^cCompounds categorized as CNS+ have the ability to permeate through the BBB and target the CNS. In the case of compounds assigned as CNS-, they have poor permeability through the BBB and therefore, their bioavailability into the CNS is considered to be minimal.