Published in final edited form as:

Org Lett. 2013 February 1; 15(3): 444–447. doi:10.1021/ol303040r.

Synthesis of Isoquinuclidines from Highly Substituted Dihydropyridines via the Diels-Alder Reaction

Rhia M. Martin[†], Robert G. Bergman^{‡,*}, and Jonathan A. Ellman^{*,†}

[†]Department of Chemistry, Yale University, New Haven, Connecticut, 06520 United States

[‡]Division of Chemical Sciences, Lawrence Berkeley National Laboratory, and Department of Chemistry, University of California, Berkeley, Berkeley, California 94720, United States

Abstract

A stereo- and regioselective Diels-Alder reaction for the synthesis of highly substituted isoquinuclidines from dihydropyridines and electron deficient alkenes has been developed. While reactions with activated dienophiles proceed readily under thermal conditions, the use of Lewis acid additives is necessary to facilitate cycloadditions for less reactive alkenes. This procedure affords the target compounds in high yields and diastereoselectivities.

The synthesis of nitrogen heterocycles is an important area of research due to their prevalence in natural products and drugs.^{1,2} Recently, we reported Rh (I)-catalyzed β -C–H bond alkenylation of α , β -unsaturated imines followed by in situ electrocyclization to give *N*-alkyl and aryl 1,2-dihydropyridines (Scheme 1A).³ These highly substituted dihydropyridines would be difficult to prepare by alternative methods and have proven to be versatile intermediates to other classes of heterocycles.⁴ For example, aromatization to the corresponding pyridines can be accomplished in an overall one pot procedure (Scheme 1A).^{3a,5} Alternatively, 1,2,3,6-tetrahydropyridines can be obtained with high diastereoselectivities via protonation of the enamine followed by in situ reduction of the resulting iminium.^{3b}

In our efforts to further harness the 1,2-dihydropyridine intermediates generated through Rh-catalyzed C–H activation, we became interested in the potential of generating isoquinuclidines utilizing a Diels-Alder reaction (Scheme 1B). Isoquinuclidines have been used as intermediates in the preparation of tetrahydroisoquinoline alkaloids,⁶ piperidines,⁷ Iboga alkaloids⁸ and the related Cantharanthus alkaloids,⁹ which have been developed as cancer therapeutics. Isoquinuclidines are most often prepared by Diels-Alder reaction of 1,2-dihydropyridines generated by *N*-acylation followed by nucleophilic addition to pyridines.^{10–13} However, the incorporation of an acyl protecting group attenuates the nucleophilicity of the dihydropyridine substrates.¹¹ Moreover, reliance on pyridine inputs generally results in dihydropyridine products lacking a high degree of substitution.⁴ Herein,

*jonathan.ellman@yale.edu

*rbergman@berkeley.edu

Supporting Information Available. Full experimental procedures, characterization data and NMR spectra for all new compounds. This material is available free of charge via the internet at <http://pubs.acs.org>.

we report that isoquinuclidines with unprecedented substitution levels are obtained in good yield and with high regio- and stereoselectivities by Diels-Alder reaction between highly substituted *N*-alkyl and aryl-1,2-dihydropyridines and electron deficient alkenes.

Diels-Alder reactions for a range of dienophiles with differential relative reactivity were first investigated (products **3a–d**, Scheme 2). The most reactive dienophile, *N*-phenyl maleimide, underwent efficient cycloaddition to give isoquinuclidine **3a** at room temperature within 16 h with only the endo isomer detected by ¹H and ¹³C NMR analysis of the unpurified reaction mixture. Methyl acrylate and acrylonitrile required more forcing conditions, but provided the desired products **3b** and **3c**, respectively, in high yield when run neat with heating to 105 °C. Isoquinuclidines **3b** and **3c** were each produced as a single regioisomer, and for **3b** only the endo product was observed while for **3c** a 93:7 endo/exo ratio was obtained.

Identifying effective conditions for coupling the less reactive crotonaldehyde required significant optimization (Table 1). Performing the reaction at 0.1 or 0.5 M in CH₂Cl₂ with heating to 50 °C provided little if any conversion (entries 1 and 2). Even when the reaction was performed neat with excess crotonaldehyde with heating at 105 °C only trace amounts of product was detected (entry 3). Lewis acid additives have proven to be effective for increasing the rate of Diels-Alder reactions for dienes incorporated within nitrogen heterocycles.^{6a,e,12,14} We therefore examined their affect upon our substrate combination (entries 4–6). The aluminum-based Lewis acids AlEtCl₂ and AlCl₃ led to modest yields of the isoquinuclidine **3d** (entries 4 and 5). Employing ZnCl₂ as a Lewis acid additive at 0 °C provided even better results, with **3d** obtained as a single diastereomer in 59% yield (entry 6). BF₃·OEt₂ was also investigated, but did not yield the desired isoquinuclidine **3d**.

As illustrated in Scheme 2, a range of differently substituted 1,2-dihydropyridines underwent Diels-Alder reactions in high yields. A variety of nitrogen substituents were well-tolerated, including *N*-benzyl (**3a–d**, **h–p**), branched *N*-alkyl (**3e**) and *N*-phenyl (**3f–g**) groups. Different substitution patterns at other sites within the dihydropyridine ring were also acceptable. The tetracyclic isoquinuclidine **3h** was prepared as a single diastereomer in high yield. Dihydropyridines substituted with the branched isopropyl group were also effective coupling partners (**3i**, **j** and **o**). Even a dihydropyridine bearing the electronically deactivating methyl ester reacted with *N*-phenyl maleimide to provide **3i** in good yield and as a single isomer. A 4-phenyl substituted dihydropyridine also underwent Diels Alder reaction to provide isoquinuclidine **3m** as a single isomer. Additionally, as should be expected, isoquinuclidines can be prepared in high yields and excellent selectivities from 1,2-dihydropyridines with modestly lower substitution levels (**3n–p**).

The relative configuration of isoquinuclidine **3g** was established by X-ray crystallography of the corresponding ammonium salt (Figure 1). By analogy, we assigned the endo configuration to the other dihydropyridine products.

The 1,2-dihydropyridines that result from the rhodium-catalyzed C-H alkenylation of α,β-unsaturated imines and subsequent electrocyclization are versatile intermediates for the preparation of highly functionalized nitrogen heterocycles. The Diels-Alder reaction with a variety of dienophiles provides access to isoquinuclidines with unprecedented substitution levels in high yields and with excellent regio- and stereoselectivities.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by the NIH grant GM069559 (to J.A.E.). R.G.B. acknowledges funding from The Director, Office of Energy Research, Office of Basic Energy Sciences, Chemical Sciences Division, DOE, under Contract DE-AC02-05CH11231. R.M.M. is grateful for a NSF Predoctoral Fellowship and a UC, Berkeley Chancellor's Fellowship. The authors are grateful to the Keck Center of Yale University for collecting the high resolution mass spectrometry (HRMS) data and to Dr. Michael Takase at the Chemical and Biophysical Instrumentation Center at Yale University for solving the structure of **3g**.

References

- Carey JS, Laffan D, Thomson C, Williams MT. *Org Biomol Chem*. 2006; 4:2337. [PubMed: 16763676]
- For recent reviews on heterocycle synthesis via C–H activation, see: Yamaguchi J, Yamaguchi AD, Itami K. *Angew Chem Int Ed*. 2012; 51:2. Engle KM, Mei TS, Wasa M, Yu JQ. *Acc Chem Res*. 2012; 45:788. [PubMed: 22166158] Yeung CS, Dong VM. *Chem Rev*. 2011; 111:1215. [PubMed: 21391561] Wencel-Delord J, Dröge T, Liu F, Glorius F. *Chem Soc Rev*. 2011; 40:4740. [PubMed: 21666903] McMurray L, O'Hara F, Gaunt MJ. *Chem Soc Rev*. 2011; 40:1885. [PubMed: 21390391] Lyons TW, Sanford MS. *Chem Rev*. 2010; 110:1147. [PubMed: 20078038] Colby DA, Bergman RG, Ellman JA. *Chem Rev*. 2010; 110:624. [PubMed: 19438203] Xu LM, Li BJ, Yang Z, Shi ZJ. *Chem Soc Rev*. 2010; 39:712. [PubMed: 20111789] Satoh T, Miura M. *Chem –Eur J*. 2010; 16:11212. [PubMed: 20740508] Li CJ. *Acc Chem Res*. 2009; 42:335. [PubMed: 19220064] Seregin IV, Gevorgyan V. *Chem Soc Rev*. 2007; 36:1173. [PubMed: 17576484]
- (a) Colby DA, Bergman RG, Ellman JA. *J Am Chem Soc*. 2008; 130:3645. [PubMed: 18302381] (b) Duttwyler S, Lu C, Rheingold AL, Bergman RG, Ellman JA. *J Am Chem Soc*. 2012; 134:4064. [PubMed: 22356093]
- For a recent review on the synthesis of dihydropyridines from pyridines, see: Bull JA, Mousseau JJ, Pelletier G, Charette AB. *Chem Rev*. 2012; 112:2642. [PubMed: 22352938] For leading references on alternative methods for accessing substituted 1,2-dihydropyridines, see: Harrison DP, Sabat M, Myers WH, Harman WD. *J Am Chem Soc*. 2010; 132:17282. Harschneck T, Kirsch SF. *J Org Chem*. 2011; 76:2145. [PubMed: 21381702] Tejedor D, M_endez-Abt G, García-Tellado F. *Chem –Eur J*. 2010; 16:428. [PubMed: 19921718] Padwa A, Dean DC, Osterhout MH, Precado L, Semones MA. *J Org Chem*. 1994; 59:5347.
- For alternative Rh-catalyzed C–H bond functionalization methods for the synthesis of pyridines from 3,3-unsaturated oximes and alkynes, see: Martin RM, Bergman RG, Ellman JA. *J Org Chem*. 2012; 77:2501. [PubMed: 22332639] Parthasarathy K, Jeganmohan M, Cheng CH. *Org Lett*. 2008; 10:325. [PubMed: 18092796] Too PC, Chua SH, Wong SH, Chiba S. *J Org Chem*. 2011; 76:6159. [PubMed: 21688786] Hyster TK, Rovis T. *Chem Commun*. 2011; 47:11846.
- (a) MaGee DI, Lee ML. *Tetrahedron Lett*. 2001; 42:7177. (b) Comins DL, Brooks CA, Al-awar RS, Goehring RR. *Org Lett*. 1999; 1:229. [PubMed: 10822560] (c) MaGee DI, Lee ML. *Synlett*. 1997; 7:786. (d) Comins DL, Al-war RS. *J Org Chem*. 1992; 57:4098. (e) Baxter EW, Labaree D, Chao S, Mariano PS. *J Org Chem*. 1989; 54:2893. (f) Wender PA, Schaus JM, White AW. *J Am Chem Soc*. 1980; 102:6157. (g) Wender P, Schaus J, Torney D. *Tetrahedron Lett*. 1979; 20:2485. (h) Mariano PS, Dunaway-Mariano D, Huesmann PL. *J Org Chem*. 1979; 44:124. (i) Mariano PS, Dunaway-Mariano D, Huesmann PL, Beamer RL. *Tetrahedron Lett*. 1977; 18:4299.
- (a) Hardin AN, Sarpong R. *Org Biomol Chem*. 2012; 10:70. [PubMed: 22072189] (b) Lemonnier G, Charette AB. *J Org Chem*. 2010; 75:7465. [PubMed: 20945853] (c) Barbe G, St-Onge M, Charette AB. *Org Lett*. 2008; 10:5497. [PubMed: 19006316] (d) Maison W. *Eur J Org Chem*. 2007:2276. (e) Sales M, Charette AB. *Org Lett*. 2005; 7:5773. [PubMed: 16354063] (f) Arakawa Y, Murakami T, Ozawa F, Arakawa Y, Yoshifuji S. *Tetrahedron*. 2003; 59:7555. (g) Reding MT, Kaburagi Y, Tokuyama H, Fukuyama T. *Heterocycles*. 2002; 56:313. (h) Polniaszek RP, Dillard LW. *J Org Chem*. 1992; 57:4103.
- (a) Jana GK, Sinha S. *Tetrahedron*. 2012; 68:7155. (b) Jana GK, Sinha S. *Tetrahedron Lett*. 2012; 53:1671. (c) Paul S, Pattanayak S, Sinha A. *Tetrahedron Lett*. 2011; 52:6166. (d) Jana GK, Sinha S. *Tetrahedron Lett*. 2010; 51:1994. (e) Hodgson DM, Galano JM. *Org Lett*. 2005; 7:2221. [PubMed: 15901174] (f) Sundberg RJ, Cherney RJ. *J Org Chem*. 1990; 55:6028. (g) Krow GR, Shaw DA, Lynch B, Lester W, Szczepanski SW, Raghavachari R, Derome AE. *J Org Chem*. 1988; 53:2258. (h)

- Sundberg RJ, Amat M, Fernando AM. *J Org Chem.* 1987; 52:3151.(i) Raucher S, Bray BL. *J Org Chem.* 1985; 50:3236.(j) Raucher S, Lawrence RF. *Tetrahedron Lett.* 1983; 24:2927.(k) Sundberg RJ, Bloom JD. *J Org Chem.* 1981; 46:4836.(l) Marazano C, Fourrey JL, Das BC. *J Chem Soc Chem Commun.* 1981:37.
9. Sundberg RJ, Hong J, Smith SQ, Sabat M. *Tetrahedron.* 1998; 54:6259.
10. For recent syntheses of isoquinuclidines via thermal Diels-Alder reactions, see: Comins DL, Bharathi P, Sahn JJ. *Tetrahedron Lett.* 2012; 53:1347.Oshima K, Ohmura T, Suginome M. *J Am Chem Soc.* 2012; 134:3699. [PubMed: 22320932] Chou SS, Wang HC, Chen PW, Yang C-H. *Tetrahedron.* 2008; 64:5291.Songis O, Géant PY, Sautrey G, Martinez J, Calmès M. *Eur J Org Chem.* 2007:308.Hoshino M, Matsuzaki H, Fujita R. *Heterocycles.* 2008; 76:267.Krow GR, Huang Q, Szczepanski SW, Hausheer FH, Caroll. *J Org Chem.* 2007; 72:3458. [PubMed: 17394357]
11. For select examples of the preparation of isoquinuclidines from N-alkyl 1,2-dihydropyridines and alkenes, see: Taylor SJ, Taylor AM, Schreiber SL. *Angew Chem Int Ed.* 2004; 43:1681.Reine I, Muceniece VL, Kemme A. *Heterocycl Commun.* 2002; 8:173.dos Santos DC, de Freitas Gil RP, Gil L, Marazano C. *Tetrahedron Lett.* 2001; 42:6109.Mehmandoust M, Marazano C, Singh R, Gillet B, Césario M, Fourrey J-L, Das BC. *Tetrahedron Lett.* 1988; 29:4423.Cardellini M, Cingolani GM, Claudi F, Gulini U, Cantalamessa F, Venturi F. *Eur J Med Chem.* 1987; 22:1.
12. For recent asymmetric syntheses of isoquinuclidines via Diels-Alder reactions, see: Suttibut C, Kohari Y, Igarashi K, Nakano H, Hiram M, Seki C, Matsuyama H, Uwai K, Takano N, Okuyama Y, Osone K, Takeshita M, Kwon E. *Tetrahedron Lett.* 2011; 52:4745.Nakano H, Osone K, Takeshita M, Kwon E, Seki C, Matsuyama N, Kohari Y. *Chem Commun.* 2010; 46:4827.Hiram M, Kato Y, Matsuyama H, Oshikiri N, Iyoda M. *Chem Lett.* 2008; 37:924.Nakano H, Tsugawa N, Takahashi K, Okuyama Y, Fujita R. *Tetrahedron.* 2006; 62:10879.Takenaka N, Huang, Rawal VH. *Tetrahedron.* 2002; 58:8299.
13. For recent syntheses of isoquinuclidines via Lewis Acid mediated Diels-Alder reaction, see: Seki C, Hiram M, Hutabarat R, Takada J, Suttibut C, Takahashi H, Takaguchi T, Kohari Y, Nakano H, Uwai K, Takano N, Yasui M, Okuyama Y, Takeshita M, Matsuyama H. *Tetrahedron Lett.* 2012; 68:1774.Harrison DP, Iovan DA, Myers WH, Sabat M, Wang S, Zottig VE, Harman WD. *J Am Chem Soc.* 2011; 133:18378. [PubMed: 21961580] Barbe G, Charette AB. *J Am Chem Soc.* 2008; 130:13873. [PubMed: 18821760] Graham PM, Delafuente DA, Lui W, Myers WH, Sabat M, Harman WD. *J Am Chem Soc.* 2005; 127:10568. [PubMed: 16045344] Matsumura Y, Nakamura Y, Maki T, Onomura O. *Tetrahedron Lett.* 2000; 41:7685.
14. For recent Lewis Acid mediated Diels-Alder reactions of pyrroles, see: Kimura H, Fujiwara T, Katoh T, Nishide K, Node M. *Chem Pharm Bull.* 2006; 54:399. [PubMed: 16508202] Nishide K, Ichihashi S, Kimura H, Katoh T, Node M. *Tetrahedron Lett.* 2001; 42:9237.

Figure 1.
Relative Configuration and Ball-and-Stick Representation of **3g**.

Colby, D. A., *et. al. J. A. Chem. Soc.* **2008**, 130, 3645.

Duttwyler, S., *et. al. J. Am. Chem. Soc.* **2012**, 134, 4064.

B. This Work

Scheme 1.

Rh-Catalyzed C-H Functionalization Provides Versatile 1,2-Dihydropyridine Intermediates

Scheme 2. Substrate Scope of Diels-Alder Reactions^a

^a Yields correspond to the yields of isolated products. The diastereoselectivities were determined by ^1H NMR analysis of unpurified material. ^b ZnCl_2 (1.1 equiv), 0.5 M CH_2Cl_2 , 0 °C, 6 h. ^c *N*-phenyl maleimide (1.05 equiv), 0.1 M CH_2Cl_2 , rt. ^d Neat, 105 °C. ^e 24 h. ^f 72 h. ^g *N*-phenyl maleimide (1.05 equiv), 0.5 M CH_2Cl_2 , 50 °C. ^h 48 h.

Table 1Optimization of Diels-Alder Reaction with Crotonaldehyde.^a

entry	Lewis acid	concentration (M)	temperature (°C)	yield ^b (%)
1	-	0.1	50	trace
2	-	0.5	50	trace
3	-	-	105	trace
4	AlEtCl ₂	0.5	0	10 ^c
5	AlCl ₃	0.5	0	43
6	ZnCl ₂	0.5	0	59

^a All reactions were performed using 0.05 mmol of dihydropyridine **1a** and 0.5 mmol of crotonaldehyde.^b Yields were determined by NMR relative to 1,3,5-trimethoxybenzene.^c Reaction was performed in 1:1 CH₂Cl₂:hexanes.