CORRESPONDENCE

The Indications for Liver Biopsy

by Prof. Dr. med. Andrea Tannapfel, Prof. Dr. med. Hans-Peter Dienes,
Prof. Dr. med. Ansgar W. Lohse in volume 27–28/2012

Thrombocyte Aggregation Inhibitors Are Important

In the review article entitled “The indications for liver biopsy” (1), the recommendations for coagulation say: “For safety, the platelet count should be above 50/nL, and the Quick value over 50% (e4).” Unfortunately the authors do not make any mention of platelet aggregation inhibitors.

In context of an expert report, the case of a patient was worked up in which, after a biopsy of a hepatic nodule with continued intake of acetylsalicylic acid, a massive hemorrhage occurred at the puncture site. The insurers declared non-liability because no guidelines or binding regulations exist in this scenario. The German-language literature does, however, provide recommendations according to which platelet aggregation inhibitors should be stopped in good time before a liver biopsy is undertaken (2, 3).

It would therefore be of general interest to find out where the authors stand with regard to this question.

DOI: 10.3238/arztebl.2013.0059a

REFERENCES

1. Tannapfel A, Dienes HP, Lohse AW: The indications for liver biopsy. Dtsch Arztebl Int 2012; 109(27–28): 477–83.
2. Glaser J, Pausch J: Perkutane Leberbiopsie – Empfehlungen der Deutschen Gesellschaft für Verdauungs- und Stoffwechselkrankheiten (DGVS) für die Durchführung endoskopischer Untersuchungen. 3rd edition. Demeter-Verlag Stuttgart 2002; 296–9. www.dgvs.de/media/6.1.Leberbiopsie.pdf.
3. Semmo N, et al.: Die perkutane Leberbiopsie. Dtsch Med Wochenschr 2007; 132: 2643–5.

Prof. Dr. med. Georg Schultze
Villingen-Schwenningen
GSchultze.VS@t-online.de

Conflict of interest statement

The author has received honoraria for consultancy services from the medical service of the health insurance companies in Germany (Medizinischer Dienst der Krankenversicherung, MDK) Saarland.

Biopsy and Puncture Track Metastasis

In their review article, the authors provide an overview of the indications for liver biopsy. They discuss biopsy

of focal solitary and multiple hepatic lesions and mention the problem of biopsy-track metastases (1). The relevant data are heterogeneous, as the authors themselves pointed out.

This complication has to be considered and included early on in the clinical process of making a diagnosis, especially in patients in whom surgical-curative treatment is an option (2–4). It seems important to refer such patients early on—that is, before the planned biopsy—to a multidisciplinary team for further investigation, such as is common practice in so called tumor boards (2).

The authors mention that in a 2008 meta-analysis, biopsy-track metastases did not affect patients’ survival. The literature (for example, a 2009 review) indicates that, in addition to repeated treatment, the biopsy of malignant hepatic lesions can also have a negative effect on patients’ survival (2). It was also mentioned that biopsy does not necessarily confer a diagnostic advantage compared with the combination of non-invasive imaging procedures and measuring of tumor markers.

A retrospective study found that the possibility of biopsy-track metastases also existed in patients with well differentiated primary tumors (4). This may indicate that metastasis after percutaneous biopsy is independent of the degree of differentiation of the primary malignancy.

Laparoscopic biopsy is an alternative option to avoid this complication, and it offers the twin advantages of being able to assess intraperitoneal organs and structures and to immediately initiate an intervention if complications occur during the tissue removal.

DOI: 10.3238/arztebl.2013.0059b

REFERENCES

1. Tannapfel A, Dienes HP, Lohse AW: The indications for liver biopsy. Dtsch Arztebl Int 2012; 109(27–28): 477–83.
2. Cresswell AB, Welsh FK, Rees M: A diagnostic paradigm for resectable liver lesions: to biopsy or not to biopsy? HPB (Oxford). 2009; 11: 533–40.
3. Takamori R, Wong LL, Dang C, Wong L: Needle-tract implantation from hepatocellular cancer: is needle biopsy of the liver always necessary? Liver Transpl 2000; 6: 67–72.
4. Kosugi C, Furuse J, Ishii H, et al.: Needle tract implantation of hepatocellular carcinoma and pancreatic carcinoma after ultrasound-guided percutaneous puncture: clinical and pathologic characteristics and the treatment of needle tract implantation. World J Surg 2004; 28: 29–32. Epub 2003 Dec 5.

Dr. med. Sammy A. Baierlein
Bayreuth
sammybaierlein@web.de

Conflict of interest statement

The author declares that no conflict of interest exists.

In Reply:

The question to which extent liver biopsy is allowed in a setting of changed coagulation, reduced platelet count, and medication for platelet aggregation inhibition, cannot be answered in a scientifically based way because almost no studies exist of this clinically relevant situation. The consensus recommendation of a Quick value of over 50% and a thrombocyte count of over 50 is consistent with general guidance but has not been sufficiently validated for liver biopsy. The Quick value in particular is likely to have a much less important role than is assumed in current clinical practice. Especially in hepatic disorders the synthesis is reduced not only in the pro-coagulation factors but also in the anti-coagulation factors, so that plasmatic coagulation is barely changed and in some cases even activated (3).

The platelet count and function seems more important in context of the problem of the post-interventional risk of hemorrhage (1, 2). It has therefore become common practice to stop ASS and other platelet aggregation inhibitors five days before the procedure, but the risk of a break in medication has to be weighed up against the risk of post-interventional hemorrhage. Biopsy should not be conducted in dual platelet aggregation inhibition. If a patient cannot stop taking platelet aggregation inhibitors we recommend obtaining a liver specimen by the transjugular route or via mini-laparoscopy, while giving thrombocyte replacement and possibly fresh frozen plasma or coagulants. This is also our method of choice if plasma coagulation is poor and in pronounced thrombopenia. Serious hemorrhages occur almost exclusively in patients with several risk factors.

The comments regarding the subject of hepatic nodules underline the importance of interdisciplinary cooperation in the clinical management of hepatic nodules. Independent of the tumor histology, liver biopsy can of course be dispensed with altogether if surgical intervention is undertaken anyway. We wish to emphasize once again the importance of discussion within an interdisciplinary tumor board, which should include colleagues from all relevant specialties. This problem will increase in importance in the future because the updated guidelines for colorectal cancer stipulates tumor board discussions of hepatic metastases of colorectal cancers, for example, which involve a surgeon with experience of liver disorders (4). A “biopsy algorithm” will be an essential part of the S3 guideline on hepatocellular carcinoma, which is currently going through the consensus process. We agree with the authors that a decision needs to be made on a case-by-case

basis as to whether invasive diagnostic evaluation can be dispensed with, even though blood based biomarkers and tumor markers may be used. However, a paradigm shift has occurred in the diagnosis of hepatocellular cancer, and raised alpha-fetoprotein (AFP) is not included any longer in the diagnostic algorithm and biopsy algorithm (5). The degree of differentiation of a malignant tumor (the so-called grading), as mentioned in one of the letters, is unlikely to have any effect on the development of possible puncture track metastases—but systematic, valid studies in this setting are lacking.

In conclusion we wish to underline once more our correspondents’ recommendations—a whole range of complications after liver biopsy can be avoided by using direct visualizing methods, such as (mini-)laparoscopy.

DOI: 10.3238/arztebl.2013.0060

REFERENCES

1. Denzer U, Helmreich-Becker I, Galle PR, Lohse AW: Liver assessment and biopsy in patients with marked coagulopathy: value of mini-laparoscopy and control of bleeding. *Am J Gastroenterol* 2003; 98: 893–900.
2. Frenzel C, Koch J, Lorenzen V, Werner T, Lohse AW, Denzer UW: Complications and risk factors in 2731 diagnostic mini-laparoscopies in patients with liver disease. *Liver Int* 2012; 32: 970–6.
3. Tripodi A, Mannucci PM: The coagulopathy of chronic liver disease. *N Engl J Med* 2011; 365: 147–56.
4. Schmiegel W, Pox C, Arnold D, et al.: Colorectal carcinoma: the management of polyps, (neo)adjuvant therapy, and the treatment of metastases. *Dtsch Arztebl Int* 2009; 106(51–52): 843–8.
5. Bruix J, Sherman M: Management of hepatocellular carcinoma. *Hepatology*; 2010: 1–35.
6. Tannapfel A, Dienes HP, Lohse AW: The indications for liver biopsy. *Dtsch Arztebl Int* 2012; 109(27–28): 477–83.

Prof. Dr. med. Andrea Tannapfel
Institut für Pathologie
Ruhr-Universität Bochum
Andrea.Tannapfel@rub.de

Prof. Dr. med. (emer.) Dienes
Institut für Pathologie, Köln

Prof. Dr. med. Lohse
I. Medizinische Klinik und Poliklinik, UKE Hamburg

Conflict of interest statement

Professor Tannapfel has received honoraria for consulting from Merck, Sanofi, Falk, Pfizer, Amgen, AstraZeneca, Lilly, and Novartis. She has received travel expenses and honoraria for speaking from Sanofi, Falk, Pfizer, Amgen, and Apda.

Professor Lohse has received honoraria for speaking from Falk-Foundation, MSD, and Roche. He has received third-party funding from Boehringer Ingelheim and Roche, among others.

Professor Dienes declares that no conflict of interest exists.